


Sensitivity of N_e/N to generation time L

Model: Ne_clonal_Y2000 | Population: Fritillaria Miz




V is held constant across L values.
 $N_e/N = 2 / (V * L)$, so the relationship is hyperbolic.

help("Ne_sensitivity_L")

Sensitivity of N_e/N to clonal reproductive variance


Focal stage: stage_3 | Population: mixed herb



Other stages fixed at $V_c/c_{\text{bar}} = 1$. Note: $V_{\text{term}3} = 0$ when $V_c/c_{\text{bar}} = V_k/k_{\text{bar}} = 1$ (Poisson).
Red dotted line = N_e/N at Poisson default.

Sensitivity of N_e/N to sexual reproductive variance


Focal stage: stage_3 | Population: hypothetical herb



Other stages fixed at $V_k/k_{\text{bar}} = 1$. Red dotted line = N_e/N at Poisson default.

Sensitivity of N_e/N to clonal fraction d_i

Focal stage: stage_3 | Population: mixed herb (Poisson)




Other stage d values fixed. Dashed line = base d value (0.5).

Flat curve = Poisson defaults active (supply $V_c \text{ over } c \neq 1$ to see effect).

help("Ne_sensitivity_d")

Sensitivity of N_e/N to clonal fraction d_i

Focal stage: stage_3 | Population: mixed herb ($V_c/c_{\text{bar}} = 3$ in stage 3)



Other stage d values fixed. Dashed line = base d value (0.5).
Flat curve = Poisson defaults active (supply $V_c/c_{\text{bar}} \neq 1$ to see effect).

help("Ne_sensitivity_d")