

# Lecture # 33

## Matrix Factorization (LU Decomposition)

---

- Two basic classes of solving methods: direct and iterative
- **Direct methods** : assume that the exact solutions exists and find the precise solution in a final number of steps
  - Gauss-Jordan method (simple and accurate)
  - Gaussian elimination method (calculatively efficient)
  - Cholesky method (in case of nonsingular symmetrical matrix)
- **Iterative methods** : starting from some initial approximation value, construct a series of solution approximations such that it converges to the exact solution of a system
  - Jacobi method
  - Gauss-Seidel method
  - SOR (Successive Over-Relaxation) method

Suppose that  $A$  has been factored into the triangular form  $A = LU$ , where  $L$  is lower triangular and  $U$  is upper triangular. Then we can solve for  $\mathbf{x}$  more easily by using a two-step process.

- First we let  $\mathbf{y} = U\mathbf{x}$  and solve the lower triangular system  $L\mathbf{y} = \mathbf{b}$  for  $\mathbf{y}$ . Since  $L$  is triangular, determining  $\mathbf{y}$  from this equation requires only  $O(n^2)$  operations.
- Once  $\mathbf{y}$  is known, the upper triangular system  $U\mathbf{x} = \mathbf{y}$  requires only an additional  $O(n^2)$  operations to determine the solution  $\mathbf{x}$ .

Solving a linear system  $A\mathbf{x} = \mathbf{b}$  in factored form means that the number of operations needed to solve the system  $A\mathbf{x} = \mathbf{b}$  is reduced from  $O(n^3/3)$  to  $O(2n^2)$ .

**Example 1** Compare the approximate number of operations required to determine the solution to a linear system using a technique requiring  $O(n^3/3)$  operations and one requiring  $O(2n^2)$  when  $n = 20$ ,  $n = 100$ , and  $n = 1000$ .

**Solution** Table 6.3 gives the results of these calculations. ■

**Table 6.3**

$n$	$n^3/3$	$2n^2$	% Reduction
10	$3.\bar{3} \times 10^2$	$2 \times 10^2$	40
100	$3.\bar{3} \times 10^5$	$2 \times 10^4$	94
1000	$3.\bar{3} \times 10^8$	$2 \times 10^6$	99.4

As the example illustrates, the reduction factor increases dramatically with the size of the matrix. Not surprisingly, the reductions from the factorization come at a cost; determining the specific matrices  $L$  and  $U$  requires  $O(n^3/3)$  operations. But once the factorization is determined, systems involving the matrix  $A$  can be solved in this simplified manner for any number of vectors  $\mathbf{b}$ .

**Example 2** (a) Determine the  $LU$  factorization for matrix  $A$  in the linear system  $A\mathbf{x} = \mathbf{b}$ , where

$$A = \begin{bmatrix} 1 & 1 & 0 & 3 \\ 2 & 1 & -1 & 1 \\ 3 & -1 & -1 & 2 \\ -1 & 2 & 3 & -1 \end{bmatrix} \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} 1 \\ 1 \\ -3 \\ 4 \end{bmatrix}.$$

**Solution** (a) The original system was considered in Section 6.1, where we saw that the sequence of operations  $(E_2 - 2E_1) \rightarrow (E_2)$ ,  $(E_3 - 3E_1) \rightarrow (E_3)$ ,  $(E_4 - (-1)E_1) \rightarrow (E_4)$ ,  $(E_3 - 4E_2) \rightarrow (E_3)$ ,  $(E_4 - (-3)E_2) \rightarrow (E_4)$  converts the system to the triangular system

$$\begin{aligned} x_1 + x_2 + 3x_4 &= 4, \\ -x_2 - x_3 - 5x_4 &= -7, \\ 3x_3 + 13x_4 &= 13, \\ -13x_4 &= -13. \end{aligned}$$

The multipliers  $m_{ij}$  and the upper triangular matrix produce the factorization

$$A = \begin{bmatrix} 1 & 1 & 0 & 3 \\ 2 & 1 & -1 & 1 \\ 3 & -1 & -1 & 2 \\ -1 & 2 & 3 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ 3 & 4 & 1 & 0 \\ -1 & -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 & 3 \\ 0 & -1 & -1 & -5 \\ 0 & 0 & 3 & 13 \\ 0 & 0 & 0 & -13 \end{bmatrix} = LU.$$



(b) Then use the factorization to solve the system

$$\begin{aligned} x_1 + x_2 + 3x_4 &= 8, \\ 2x_1 + x_2 - x_3 + x_4 &= 7, \\ 3x_1 - x_2 - x_3 + 2x_4 &= 14, \\ -x_1 + 2x_2 + 3x_3 - x_4 &= -7. \end{aligned}$$

$$A\mathbf{x} = LU\mathbf{x} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ 3 & 4 & 1 & 0 \\ -1 & -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 & 3 \\ 0 & -1 & -1 & -5 \\ 0 & 0 & 3 & 13 \\ 0 & 0 & 0 & -13 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 8 \\ 7 \\ 14 \\ -7 \end{bmatrix},$$

we first introduce the substitution  $\mathbf{y} = U\mathbf{x}$ . Then  $\mathbf{b} = L(U\mathbf{x}) = L\mathbf{y}$ . That is,

$$L\mathbf{y} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 \\ 3 & 4 & 1 & 0 \\ -1 & -3 & 0 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \end{bmatrix} = \begin{bmatrix} 8 \\ 7 \\ 14 \\ -7 \end{bmatrix}.$$

This system is solved for  $\mathbf{y}$  by a simple forward-substitution process:

$$\begin{aligned} y_1 &= 8; \\ 2y_1 + y_2 &= 7, & \text{so } y_2 &= 7 - 2y_1 = -9; \\ 3y_1 + 4y_2 + y_3 &= 14, & \text{so } y_3 &= 14 - 3y_1 - 4y_2 = 26; \\ -y_1 - 3y_2 + y_4 &= -7, & \text{so } y_4 &= -7 + y_1 + 3y_2 = -26. \end{aligned}$$

We then solve  $U\mathbf{x} = \mathbf{y}$  for  $\mathbf{x}$ , the solution of the original system; that is,

$$\begin{bmatrix} 1 & 1 & 0 & 3 \\ 0 & -1 & -1 & -5 \\ 0 & 0 & 3 & 13 \\ 0 & 0 & 0 & -13 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 8 \\ -9 \\ 26 \\ -26 \end{bmatrix}.$$

Using backward substitution we obtain  $x_4 = 2, x_3 = 0, x_2 = -1, x_1 = 3$ . ■



**ALGORITHM  
6.4**

### LU Factorization

To factor the  $n \times n$  matrix  $A = [a_{ij}]$  into the product of the lower-triangular matrix  $L = [l_{ij}]$  and the upper-triangular matrix  $U = [u_{ij}]$ ; that is,  $A = LU$ , where the main diagonal of either  $L$  or  $U$  consists of all ones:

**INPUT** dimension  $n$ ; the entries  $a_{ij}$ ,  $1 \leq i, j \leq n$  of  $A$ ; the diagonal  $l_{11} = \dots = l_{nn} = 1$  of  $L$  or the diagonal  $u_{11} = \dots = u_{nn} = 1$  of  $U$ .

**OUTPUT** the entries  $l_{ij}$ ,  $1 \leq j \leq i$ ,  $1 \leq i \leq n$  of  $L$  and the entries,  $u_{ij}$ ,  $i \leq j \leq n$ ,  $1 \leq i \leq n$  of  $U$ .

**Step 1** Select  $l_{11}$  and  $u_{11}$  satisfying  $l_{11}u_{11} = a_{11}$ .  
If  $l_{11}u_{11} = 0$  then OUTPUT ('Factorization impossible');  
STOP.

**Step 2** For  $j = 2, \dots, n$  set  $u_{1j} = a_{1j}/l_{11}$ ; (First row of  $U$ )  
 $l_{j1} = a_{j1}/u_{11}$ . (First column of  $L$ .)

**Step 3** For  $i = 2, \dots, n-1$  do Steps 4 and 5.

**Step 4** Select  $l_{ii}$  and  $u_{ii}$  satisfying  $l_{ii}u_{ii} = a_{ii} - \sum_{k=1}^{i-1} l_{ik}u_{ki}$ .  
If  $l_{ii}u_{ii} = 0$  then OUTPUT ('Factorization impossible');  
STOP.

**Step 5** For  $j = i+1, \dots, n$

$$\text{set } u_{ij} = \frac{1}{l_{ii}} \left[ a_{ij} - \sum_{k=1}^{i-1} l_{ik}u_{kj} \right]; \quad (\text{ith row of } U.)$$

$$l_{ji} = \frac{1}{u_{ii}} \left[ a_{ji} - \sum_{k=1}^{i-1} l_{jk}u_{ki} \right]. \quad (\text{ith column of } L.)$$

**Step 6** Select  $l_{nn}$  and  $u_{nn}$  satisfying  $l_{nn}u_{nn} = a_{nn} - \sum_{k=1}^{n-1} l_{nk}u_{kn}$ .  
(Note: If  $l_{nn}u_{nn} = 0$ , then  $A = LU$  but  $A$  is singular.)

**Step 7** OUTPUT ( $l_{ij}$  for  $j = 1, \dots, i$  and  $i = 1, \dots, n$ );  
OUTPUT ( $u_{ij}$  for  $j = i, \dots, n$  and  $i = 1, \dots, n$ );  
STOP.

**Theorem 6.19** If Gaussian elimination can be performed on the linear system  $A\mathbf{x} = \mathbf{b}$  without row interchanges, then the matrix  $A$  can be factored into the product of a lower-triangular matrix  $L$  and an upper-triangular matrix  $U$ , that is,  $A = LU$ , where  $m_{ji} = a_{ji}^{(i)} / a_{ii}^{(i)}$ ,

$$U = \begin{bmatrix} a_{11}^{(1)} & a_{12}^{(1)} & \cdots & a_{1n}^{(1)} \\ 0 & a_{22}^{(2)} & \cdots & \vdots \\ \vdots & \vdots & \ddots & a_{n-1,n}^{(n-1)} \\ 0 & \cdots & 0 & a_{nn}^{(n)} \end{bmatrix}, \quad \text{and} \quad L = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ m_{21} & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ m_{n1} & \cdots & m_{n,n-1} & 1 \end{bmatrix}.$$

An  $n \times n$  **permutation matrix**  $P = [p_{ij}]$  is a matrix obtained by rearranging the rows of  $I_n$ , the identity matrix. This gives a matrix with precisely one nonzero entry in each row and in each column, and each nonzero entry is a 1.

## Recall Elementary Matrices!

**Illustration** The matrix

$$P = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

is a  $3 \times 3$  permutation matrix. For any  $3 \times 3$  matrix  $A$ , multiplying on the left by  $P$  has the effect of interchanging the second and third rows of  $A$ :

$$PA = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{31} & a_{32} & a_{33} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}.$$

- $P^{-1}$  exists and  $P^{-1} = P^t$ .

$$PA\mathbf{x} = P\mathbf{b}$$

can be solved without row interchanges. As a consequence, this matrix  $PA$  can be factored into

$$PA = LU,$$

where  $L$  is lower triangular and  $U$  is upper triangular. Because  $P^{-1} = P^t$ , this produces the factorization

$$A = P^{-1}LU = (P^tL)U.$$

The matrix  $U$  is still upper triangular, but  $P^tL$  is not lower triangular unless  $P = I$ .

**Example 3** Determine a factorization in the form  $A = (P^t L)U$  for the matrix

$$A = \begin{bmatrix} 0 & 0 & -1 & 1 \\ 1 & 1 & -1 & 2 \\ -1 & -1 & 2 & 0 \\ 1 & 2 & 0 & 2 \end{bmatrix}.$$

**Solution** The matrix  $A$  cannot have an  $LU$  factorization because  $a_{11} = 0$ . However, using the row interchange  $(E_1) \leftrightarrow (E_2)$ , followed by  $(E_3 + E_1) \rightarrow (E_3)$  and  $(E_4 - E_1) \rightarrow (E_4)$ , produces

$$\begin{bmatrix} 1 & 1 & -1 & 2 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 1 & 1 & 0 \end{bmatrix}.$$

Then the row interchange  $(E_2) \leftrightarrow (E_4)$ , followed by  $(E_4 + E_3) \rightarrow (E_4)$ , gives the matrix

$$U = \begin{bmatrix} 1 & 1 & -1 & 2 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 3 \end{bmatrix}.$$



The permutation matrix associated with the row interchanges  $(E_1) \leftrightarrow (E_2)$  and  $(E_2) \leftrightarrow (E_4)$  is

$$P = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix},$$

and

$$PA = \begin{bmatrix} 1 & 1 & -1 & 2 \\ 1 & 2 & 0 & 2 \\ -1 & -1 & 2 & 0 \\ 0 & 0 & -1 & 1 \end{bmatrix}.$$

Gaussian elimination is performed on  $PA$  using the same operations as on  $A$ , except without the row interchanges. That is,  $(E_2 - E_1) \rightarrow (E_2)$ ,  $(E_3 + E_1) \rightarrow (E_3)$ , followed by  $(E_4 + E_3) \rightarrow (E_4)$ . The nonzero multipliers for  $PA$  are consequently,

$$m_{21} = 1, \quad m_{31} = -1, \quad \text{and} \quad m_{43} = -1,$$

and the  $LU$  factorization of  $PA$  is

$$PA = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & -1 & 2 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 3 \end{bmatrix} = LU.$$

Multiplying by  $P^{-1} = P^t$  produces the factorization

$$A = P^{-1}(LU) = P^t(LU) = (P^tL)U = \begin{bmatrix} 0 & 0 & -1 & 1 \\ 1 & 0 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 & -1 & 2 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 3 \end{bmatrix}. \quad \blacksquare$$

Do Q 1,2,3 & 6 from Ex#6.5

Doolittle form  
Obtained by  
Gaussian elimination

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix} \begin{bmatrix} u_{11} & u_{12} & u_{13} \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{bmatrix}$$

# Method: $[A]$ Decomposes to $[L]$ and $[U]$

$$[A] = [L][U] = \begin{bmatrix} 1 & 0 & 0 \\ \ell_{21} & 1 & 0 \\ \ell_{31} & \ell_{32} & 1 \end{bmatrix} \begin{bmatrix} u_{11} & u_{12} & u_{13} \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{bmatrix}$$

$[U]$  is the same as the coefficient matrix at the end of the forward elimination step.

$[L]$  is obtained using the *multipliers* that were used in the forward elimination process

---

# Finding the $[U]$ matrix

Using the Forward Elimination Procedure of Gauss Elimination

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix}$$

$$\text{Step 1: } \frac{64}{25} = 2.56; \quad \text{Row2} - \text{Row1}(2.56) = \begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 144 & 12 & 1 \end{bmatrix}$$

$$\frac{144}{25} = 5.76; \quad \text{Row3} - \text{Row1}(5.76) = \begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & -16.8 & -4.76 \end{bmatrix}$$

---

# Finding the [U] Matrix

Matrix after Step 1: 
$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & -16.8 & -4.76 \end{bmatrix}$$

Step 2:  $\frac{-16.8}{-4.8} = 3.5$ ;  $Row3 - Row2(3.5) = \begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix}$

$$[U] = \begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix}$$

---



# Finding the [L] matrix

$$\begin{bmatrix} 1 & 0 & 0 \\ \ell_{21} & 1 & 0 \\ \ell_{31} & \ell_{32} & 1 \end{bmatrix}$$

Using the multipliers used during the Forward Elimination Procedure

From the first step  
of forward  
elimination

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \quad \ell_{21} = \frac{a_{21}}{a_{11}} = \frac{64}{25} = 2.56$$
$$\ell_{31} = \frac{a_{31}}{a_{11}} = \frac{144}{25} = 5.76$$

---

# Finding the [L] Matrix

From the second step of forward elimination

$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & -16.8 & -4.76 \end{bmatrix} \quad \ell_{32} = \frac{a_{32}}{a_{22}} = \frac{-16.8}{-4.8} = 3.5$$

$$[L] = \begin{bmatrix} 1 & 0 & 0 \\ 2.56 & 1 & 0 \\ 5.76 & 3.5 & 1 \end{bmatrix}$$

---

**Does  $[L][U] = [A]$ ?**

$$[L][U] = \begin{bmatrix} 1 & 0 & 0 \\ 2.56 & 1 & 0 \\ 5.76 & 3.5 & 1 \end{bmatrix} \begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix} = ?$$



# Using LU Decomposition to solve SLEs

Solve the following set of linear equations using LU Decomposition

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix}$$

Using the procedure for finding the  $[L]$  and  $[U]$  matrices

$$[A] = [L][U] = \begin{bmatrix} 1 & 0 & 0 \\ 2.56 & 1 & 0 \\ 5.76 & 3.5 & 1 \end{bmatrix} \begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix}$$

---

# Example

Set  $[L][y] = [b]$

$$\begin{bmatrix} 1 & 0 & 0 \\ 2.56 & 1 & 0 \\ 5.76 & 3.5 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix}$$

Solve for  $[y]$

$$y_1 = 10$$

$$2.56y_1 + y_2 = 177.2$$

$$5.76y_1 + 3.5y_2 + y_3 = 279.2$$

---

# Example


Complete the forward substitution to solve for  $[Z]$

$$y_1 = 106.8$$

$$\begin{aligned} y_2 &= 177.2 - 2.56y_1 \\ &= 177.2 - 2.56(106.8) \\ &= -96.2 \end{aligned}$$

$$\begin{aligned} y_3 &= 279.2 - 5.76y_1 - 3.5y_2 \\ &= 279.2 - 5.76(106.8) - 3.5(-96.21) \\ &= 0.735 \end{aligned}$$

$$[y] = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ -96.21 \\ 0.735 \end{bmatrix}$$





# Example

$$\text{Set } [U][X] = [y] \quad \begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ -96.21 \\ 0.735 \end{bmatrix}$$

Solve for  $[X]$       The 3 equations become

$$25x_1 + 5x_2 + x_3 = 106.8$$

$$-4.8x_2 - 1.56x_3 = -96.21$$

$$0.7x_3 = 0.735$$

# Example

From the 3<sup>rd</sup> equation

$$0.7x_3 = 0.735$$

$$x_3 = \frac{0.735}{0.7}$$

$$x_3 = 1.050$$

Substituting in  $a_3$  and using the second equation

$$-4.8x_2 - 1.56x_3 = -96.21$$

$$x_2 = \frac{-96.21 + 1.56x_3}{-4.8}$$

$$x_2 = \frac{-96.21 + 1.56(1.050)}{-4.8}$$

$$x_2 = 19.70$$

---

# Example

Substituting in  $a_3$  and  $a_2$  using the first equation

$$25x_1 + 5x_2 + x_3 = 106.8$$

$$\begin{aligned} x_1 &= \frac{106.8 - 5a_2 - a_3}{25} \\ &= \frac{106.8 - 5(19.70) - 1.050}{25} \\ &= 0.2900 \end{aligned}$$

Hence the Solution Vector is:

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0.2900 \\ 19.70 \\ 1.050 \end{bmatrix}$$

# Lecture # 38-40

## Matrix Factorization (LU Decomposition)

---

# Special Types of Matrices

## 1. Diagonally Dominant Matrices

Each main diagonal entry in a strictly diagonally dominant matrix has a magnitude that is strictly greater than the sum of the magnitudes of all the other entries in that row.

# 1. Diagonally Dominant Matrices

The  $n \times n$  matrix  $A$  is said to be **diagonally dominant** when

$$|a_{ii}| \geq \sum_{\substack{j=1, \\ j \neq i}}^n |a_{ij}| \quad \text{holds for each } i = 1, 2, \dots, n. \quad (6.10)$$

A diagonally dominant matrix is said to be **strictly diagonally dominant** when the inequality in (6.10) is strict for each  $n$ , that is, when

$$|a_{ii}| > \sum_{\substack{j=1, \\ j \neq i}}^n |a_{ij}| \quad \text{holds for each } i = 1, 2, \dots, n. \quad \blacksquare$$



Consider the matrices

$$A = \begin{bmatrix} 7 & 2 & 0 \\ 3 & 5 & -1 \\ 0 & 5 & -6 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 6 & 4 & -3 \\ 4 & -2 & 0 \\ -3 & 0 & 1 \end{bmatrix}.$$

The nonsymmetric matrix  $A$  is strictly diagonally dominant because

$$|7| > |2| + |0|, \quad |5| > |3| + |-1|, \quad \text{and} \quad |-6| > |0| + |5|.$$

The symmetric matrix  $B$  is not strictly diagonally dominant because, for example, in the first row the absolute value of the diagonal element is  $|6| < |4| + |-3| = 7$ . It is interesting to note that  $A'$  is not strictly diagonally dominant, because the middle row of  $A'$  is  $[2 \ 5 \ 5]$ , nor, of course, is  $B'$  because  $B' = B$ . □

**Theorem 6.21** A strictly diagonally dominant matrix  $A$  is nonsingular. Moreover, in this case, Gaussian elimination can be performed on any linear system of the form  $A\mathbf{x} = \mathbf{b}$  to obtain its unique solution without row or column interchanges, and the computations will be stable with respect to the growth of round-off errors. ■

## 2. Positive Definite Matrices

The name positive definite refers to the fact that the number  $\mathbf{x}'\mathbf{A}\mathbf{x}$  must be positive whenever  $\mathbf{x} \neq \mathbf{0}$ .

## 2. Positive Definite Matrices

**Definition 6.22** A matrix  $A$  is **positive definite** if it is symmetric and if  $\mathbf{x}'A\mathbf{x} > 0$  for every  $n$ -dimensional vector  $\mathbf{x} \neq \mathbf{0}$ . ■

**Example 1** Show that the matrix

$$A = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$$

is positive definite

**Solution** Suppose  $\mathbf{x}$  is any three-dimensional column vector. Then

$$\begin{aligned} \mathbf{x}^T A \mathbf{x} &= [x_1, x_2, x_3] \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \\ &= [x_1, x_2, x_3] \begin{bmatrix} 2x_1 & -x_2 & 0 \\ -x_1 & 2x_2 & -x_3 \\ 0 & -x_2 & 2x_3 \end{bmatrix} \\ &= 2x_1^2 - 2x_1x_2 + 2x_2^2 - 2x_2x_3 + 2x_3^2. \end{aligned}$$

$$= 2x_1^2 - 2x_1x_2 + 2x_2^2 - 2x_2x_3 + 2x_3^2.$$

Rearranging the terms gives

$$\begin{aligned}\mathbf{x}'\mathbf{A}\mathbf{x} &= x_1^2 + (x_1^2 - 2x_1x_2 + x_2^2) + (x_2^2 - 2x_2x_3 + x_3^2) + x_3^2 \\ &= x_1^2 + (x_1 - x_2)^2 + (x_2 - x_3)^2 + x_3^2,\end{aligned}$$

which implies that

$$x_1^2 + (x_1 - x_2)^2 + (x_2 - x_3)^2 + x_3^2 > 0$$

unless  $x_1 = x_2 = x_3 = 0$ .

**Theorem 6.23** If  $A$  is an  $n \times n$  positive definite matrix, then

- (i)  $A$  has an inverse;
- (ii)  $a_{ii} > 0$ , for each  $i = 1, 2, \dots, n$ ;
- (iii)  $\max_{1 \leq k, j \leq n} |a_{kj}| \leq \max_{1 \leq i \leq n} |a_{ii}|$ ;
- (iv)  $(a_{ij})^2 < a_{ii}a_{jj}$ , for each  $i \neq j$ . ■

**Definition 6.24** A **leading principal submatrix** of a matrix  $A$  is a matrix of the form

$$A_k = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1k} \\ a_{21} & a_{22} & \cdots & a_{2k} \\ \vdots & \vdots & & \vdots \\ a_{k1} & a_{k2} & \cdots & a_{kk} \end{bmatrix},$$

for some  $1 \leq k \leq n$ . ■

**Theorem 6.25** A symmetric matrix  $A$  is positive definite if and only if each of its leading principal submatrices has a positive determinant. ■

**Example 2** In Example 1 we used the definition to show that the symmetric matrix

$$A = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$$

is positive definite. Confirm this using Theorem 6.25.



**Solution** Note that

$$\det A_1 = \det[2] = 2 > 0,$$

$$\det A_2 = \det \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} = 4 - 1 = 3 > 0,$$

and

$$\begin{aligned} \det A_3 &= \det \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix} = 2 \det \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} - (-1) \det \begin{bmatrix} -1 & -1 \\ 0 & 2 \end{bmatrix} \\ &= 2(4 - 1) + (-2 + 0) = 4 > 0. \end{aligned}$$

in agreement with Theorem 6.25. ■

**Theorem 6.26** The symmetric matrix  $A$  is positive definite if and only if Gaussian elimination without row interchanges can be performed on the linear system  $A\mathbf{x} = \mathbf{b}$  with all pivot elements positive. Moreover, in this case, the computations are stable with respect to the growth of round-off errors. ■

**Corollary 6.27** The matrix  $A$  is positive definite if and only if  $A$  can be factored in the form  $LDL^t$ , where  $L$  is lower triangular with 1s on its diagonal and  $D$  is a diagonal matrix with positive diagonal entries. ■

**Corollary 6.28** The matrix  $A$  is positive definite if and only if  $A$  can be factored in the form  $LL^t$ , where  $L$  is lower triangular with nonzero diagonal entries. ■

## $LDL^t$ Factorization

**Corollary 6.29** Let  $A$  be a symmetric  $n \times n$  matrix for which Gaussian elimination can be applied without row interchanges. Then  $A$  can be factored into  $LDL^t$ , where  $L$  is lower triangular with 1s on its diagonal and  $D$  is the diagonal matrix with  $a_{11}^{(1)}, \dots, a_{nn}^{(n)}$  on its diagonal. ■

**Example 3** Determine the  $LDL^t$  factorization of the positive definite matrix

$$A = \begin{bmatrix} 4 & -1 & 1 \\ -1 & 4.25 & 2.75 \\ 1 & 2.75 & 3.5 \end{bmatrix}.$$

**Example 3** Determine the  $LDL'$  factorization of the positive definite matrix

$$A = \begin{bmatrix} 4 & -1 & 1 \\ -1 & 4.25 & 2.75 \\ 1 & 2.75 & 3.5 \end{bmatrix}.$$

**Solution** The  $LDL'$  factorization has 1s on the diagonal of the lower triangular matrix  $L$  so we need to have

$$A = \begin{bmatrix} a_{11} & a_{21} & a_{31} \\ a_{21} & a_{22} & a_{32} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix} \begin{bmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{bmatrix} \begin{bmatrix} 1 & l_{21} & l_{31} \\ 0 & 1 & l_{32} \\ 0 & 0 & 1 \end{bmatrix}$$

**Solution** The  $LDL^T$  factorization has 1s on the diagonal of the lower triangular matrix  $L$  so we need to have

$$A = \begin{bmatrix} a_{11} & a_{21} & a_{31} \\ a_{21} & a_{22} & a_{32} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix} \begin{bmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{bmatrix} \begin{bmatrix} 1 & l_{21} & l_{31} \\ 0 & 1 & l_{32} \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 4 & -1 & 1 \\ -1 & 4.25 & 2.75 \\ 1 & 2.75 & 3.5 \end{bmatrix} = \begin{bmatrix} d_1 & d_1 l_{21} & d_1 l_{31} \\ d_1 l_{21} & d_2 + d_1 l_{21}^2 & d_2 l_{32} + d_1 l_{21} l_{31} \\ d_1 l_{31} & d_1 l_{21} l_{31} + d_2 l_{32} & d_1 l_{31}^2 + d_2 l_{32}^2 + d_3 \end{bmatrix}$$

Thus

$$a_{11} : 4 = d_1 \implies d_1 = 4,$$

$$a_{31} : 1 = d_1 l_{31} \implies l_{31} = 0.25,$$

$$a_{32} : 2.75 = d_1 l_{21} l_{31} + d_2 l_{32} \implies l_{32} = 0.75,$$

$$a_{21} : -1 = d_1 l_{21} \implies l_{21} = -0.25$$

$$a_{22} : 4.25 = d_2 + d_1 l_{21}^2 \implies d_2 = 4$$

$$a_{33} : 3.5 = d_1 l_{31}^2 + d_2 l_{32}^2 + d_3 \implies d_3 = 1,$$

Thus

$$a_{11} : 4 = d_1 \implies d_1 = 4,$$

$$a_{21} : -1 = d_1 l_{21} \implies l_{21} = -0.25$$

$$a_{31} : 1 = d_1 l_{31} \implies l_{31} = 0.25,$$

$$a_{22} : 4.25 = d_2 + d_1 l_{21}^2 \implies d_2 = 4$$

$$a_{32} : 2.75 = d_1 l_{21} l_{31} + d_2 l_{32} \implies l_{32} = 0.75, \quad a_{33} : 3.5 = d_1 l_{31}^2 + d_2 l_{32}^2 + d_3 \implies d_3 = 1,$$

and we have

$$A = LDL^t = \begin{bmatrix} 1 & 0 & 0 \\ -0.25 & 1 & 0 \\ 0.25 & 0.75 & 1 \end{bmatrix} \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -0.25 & 0.25 \\ 0 & 1 & 0.75 \\ 0 & 0 & 1 \end{bmatrix}. \quad \blacksquare$$



**ALGORITHM**  
**6.5**

### **$LDL^t$ Factorization**

To factor the positive definite  $n \times n$  matrix  $A$  into the form  $LDL^t$ , where  $L$  is a lower triangular matrix with 1s along the diagonal and  $D$  is a diagonal matrix with positive entries on the diagonal:

**INPUT** the dimension  $n$ ; entries  $a_{ij}$ , for  $1 \leq i, j \leq n$  of  $A$ .

**OUTPUT** the entries  $l_{ij}$ , for  $1 \leq j < i$  and  $1 \leq i \leq n$  of  $L$ , and  $d_i$ , for  $1 \leq i \leq n$  of  $D$ .

**Step 1** For  $i = 1, \dots, n$  do Steps 2–4.

**Step 2** For  $j = 1, \dots, i - 1$ , set  $v_j = l_{ij}d_j$ .

**Step 3** Set  $d_i = a_{ii} - \sum_{j=1}^{i-1} l_{ij}v_j$ .

**Step 4** For  $j = i + 1, \dots, n$  set  $l_{ji} = (a_{ji} - \sum_{k=1}^{i-1} l_{jk}v_k)/d_i$ .

**Step 5** **OUTPUT** ( $l_{ij}$  for  $j = 1, \dots, i - 1$  and  $i = 1, \dots, n$ );  
**OUTPUT** ( $d_i$  for  $i = 1, \dots, n$ );  
**STOP.**





## Cholesky

To factor the positive definite  $n \times n$  matrix  $A$  into  $LL'$ , where  $L$  is lower triangular:

Andre-Louis Cholesky  
(1875-1918) was a French  
military officer involved in  
geodesy and surveying in the  
early 1900s. He developed this  
factorization method to compute  
solutions to least squares  
problems.

## Cholesky

To factor the positive definite  $n \times n$  matrix  $A$  into  $LL^t$ , where  $L$  is lower triangular:

**INPUT** the dimension  $n$ ; entries  $a_{ij}$ , for  $1 \leq i, j \leq n$  of  $A$ .

**OUTPUT** the entries  $l_{ij}$ , for  $1 \leq j \leq i$  and  $1 \leq i \leq n$  of  $L$ . (The entries of  $U = L^t$  are  $u_{ij} = l_{ji}$ , for  $i \leq j \leq n$  and  $1 \leq i \leq n$ .)

**Step 1** Set  $l_{11} = \sqrt{a_{11}}$ .

**Step 2** For  $j = 2, \dots, n$ , set  $l_{j1} = a_{j1}/l_{11}$ .

**Step 3** For  $i = 2, \dots, n - 1$  do Steps 4 and 5.

**Step 4** Set  $l_{ii} = \left(a_{ii} - \sum_{k=1}^{i-1} l_{ik}^2\right)^{1/2}$ .

**Step 5** For  $j = i + 1, \dots, n$

set  $l_{ji} = \left(a_{ji} - \sum_{k=1}^{i-1} l_{jk}l_{ik}\right) / l_{ii}$ .

**Step 6** Set  $l_{nn} = \left(a_{nn} - \sum_{k=1}^{n-1} l_{nk}^2\right)^{1/2}$ .

**Step 7** OUTPUT ( $l_{ij}$  for  $j = 1, \dots, i$  and  $i = 1, \dots, n$ );  
STOP.

Determine the Cholesky  $LL^T$  factorization of the positive definite matrix

$$A = \begin{bmatrix} 4 & -1 & 1 \\ -1 & 4.25 & 2.75 \\ 1 & 2.75 & 3.5 \end{bmatrix}.$$

**Solution** The  $LL^T$  factorization does not necessarily has 1s on the diagonal of the lower triangular matrix  $L$  so we need to have

Determine the Cholesky  $LL'$  factorization of the positive definite matrix

$$A = \begin{bmatrix} 4 & -1 & 1 \\ -1 & 4.25 & 2.75 \\ 1 & 2.75 & 3.5 \end{bmatrix}.$$

$$A = \begin{bmatrix} a_{11} & a_{21} & a_{31} \\ a_{21} & a_{22} & a_{32} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} l_{11} & 0 & 0 \\ l_{21} & l_{22} & 0 \\ l_{31} & l_{32} & l_{33} \end{bmatrix} \begin{bmatrix} l_{11} & l_{21} & l_{31} \\ 0 & l_{22} & l_{32} \\ 0 & 0 & l_{33} \end{bmatrix}$$

$$A = \begin{bmatrix} a_{11} & a_{21} & a_{31} \\ a_{21} & a_{22} & a_{32} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} l_{11} & 0 & 0 \\ l_{21} & l_{22} & 0 \\ l_{31} & l_{32} & l_{33} \end{bmatrix} \begin{bmatrix} l_{11} & l_{21} & l_{31} \\ 0 & l_{22} & l_{32} \\ 0 & 0 & l_{33} \end{bmatrix}$$

$$\begin{bmatrix} 4 & -1 & 1 \\ -1 & 4.25 & 2.75 \\ 1 & 2.75 & 3.5 \end{bmatrix} = \begin{bmatrix} l_{11}^2 & l_{11}l_{21} & l_{11}l_{31} \\ l_{11}l_{21} & l_{21}^2 + l_{22}^2 & l_{21}l_{31} + l_{22}l_{32} \\ l_{11}l_{31} & l_{21}l_{31} + l_{22}l_{32} & l_{31}^2 + l_{32}^2 + l_{33}^2 \end{bmatrix}$$

$$\begin{bmatrix} 4 & -1 & 1 \\ -1 & 4.25 & 2.75 \\ 1 & 2.75 & 3.5 \end{bmatrix} = \begin{bmatrix} l_{11}^2 & l_{11}l_{21} & l_{11}l_{31} \\ l_{11}l_{21} & l_{21}^2 + l_{22}^2 & l_{21}l_{31} + l_{22}l_{32} \\ l_{11}l_{31} & l_{21}l_{31} + l_{22}l_{32} & l_{31}^2 + l_{32}^2 + l_{33}^2 \end{bmatrix}$$

Thus

$$a_{11} : 4 = l_{11}^2 \implies l_{11} = 2,$$

$$a_{21} : -1 = l_{11}l_{21} \implies l_{21} = -0.5$$

$$a_{31} : 1 = l_{11}l_{31} \implies l_{31} = 0.5,$$

$$a_{22} : 4.25 = l_{21}^2 + l_{22}^2 \implies l_{22} = 2$$

$$a_{32} : 2.75 = l_{21}l_{31} + l_{22}l_{32} \implies l_{32} = 1.5, \quad a_{33} : 3.5 = l_{31}^2 + l_{32}^2 + l_{33}^2 \implies l_{33} = 1,$$

and we have

$$A = LL^t = \begin{bmatrix} 2 & 0 & 0 \\ -0.5 & 2 & 0 \\ 0.5 & 1.5 & 1 \end{bmatrix} \begin{bmatrix} 2 & -0.5 & 0.5 \\ 0 & 2 & 1.5 \\ 0 & 0 & 1 \end{bmatrix}.$$

■

## Band Matrices

The last class of matrices considered are *band matrices*. In many applications, the band matrices are also strictly diagonally dominant or positive definite.

An  $n \times n$  matrix is called a **band matrix** if integers  $p$  and  $q$ , with  $1 < p, q < n$ , exist with the property that  $a_{ij} = 0$  whenever  $p \leq j - i$  or  $q \leq i - j$ . The **band width** of a band matrix is defined as  $w = p + q - 1$ . ■



## Band Matrices

An  $n \times n$  matrix is called a **band matrix** if integers  $p$  and  $q$ , with  $1 < p, q < n$ , exist with the property that  $a_{ij} = 0$  whenever  $p \leq j - i$  or  $q \leq i - j$ . The **band width** of a band matrix is defined as  $w = p + q - 1$ . ■

The number  $p$  describes the number of diagonals above, and including, the main diagonal on which nonzero entries may lie. The number  $q$  describes the number of diagonals below, and including, the main diagonal on which nonzero entries may lie. For example, the matrix

$$A = \begin{bmatrix} 7 & 2 & 0 \\ 3 & 5 & -1 \\ 0 & -5 & -6 \end{bmatrix}$$

is a band matrix with  $p = q = 2$  and bandwidth  $2 + 2 - 1 = 3$ .



## Tridiagonal Matrices

Matrices of **bandwidth 3** occurring when  $p = q = 2$  are called **tridiagonal** because they have the form

$$A = \begin{bmatrix} a_{11} & a_{12} & 0 & \cdots & \cdots & 0 \\ a_{21} & a_{22} & a_{23} & \ddots & & \\ 0 & a_{32} & a_{33} & a_{34} & \ddots & \\ \vdots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & a_{n,n-1} & a_{nn} \end{bmatrix}.$$

$$L = \begin{bmatrix} l_{11} & 0 & \dots & 0 \\ l_{21} & l_{22} & \dots & 0 \\ 0 & \dots & \dots & 0 \\ \vdots & \dots & \dots & \vdots \\ 0 & \dots & 0 & l_{n,n-1} & l_{nn} \end{bmatrix} \quad \text{and} \quad U = \begin{bmatrix} 1 & u_{12} & 0 & \dots & 0 \\ 0 & 1 & \dots & \dots & 0 \\ \vdots & \dots & \dots & \dots & \vdots \\ 0 & \dots & 0 & u_{n-1,n} & 1 \end{bmatrix}.$$

## Crout Factorization for Tridiagonal Linear Systems

To solve the  $n \times n$  linear system

$$\begin{aligned}
 E_1 : \quad & a_{11}x_1 + a_{12}x_2 & & = a_{1,n+1}, \\
 E_2 : \quad & a_{21}x_1 + a_{22}x_2 + a_{23}x_3 & & = a_{2,n+1}, \\
 & \vdots & \vdots & \vdots \\
 E_{n-1} : \quad & & a_{n-1,n-2}x_{n-2} + a_{n-1,n-1}x_{n-1} + a_{n-1,n}x_n & = a_{n-1,n+1}, \\
 E_n : \quad & & a_{n,n-1}x_{n-1} + a_{nn}x_n & = a_{n,n+1},
 \end{aligned}$$

which is assumed to have a unique solution:

Determine the Crout factorization of the symmetric tridiagonal matrix

$$\begin{bmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix},$$

and use this factorization to solve the linear system

$$\begin{aligned} 2x_1 - x_2 &= 1, \\ -x_1 + 2x_2 - x_3 &= 0, \\ -x_2 + 2x_3 - x_4 &= 0, \\ -x_3 + 2x_4 &= 1. \end{aligned}$$

**Solution** The  $LU$  factorization of  $A$  has the form

$$A = \begin{bmatrix} a_{11} & a_{12} & 0 & 0 \\ a_{21} & a_{22} & a_{23} & 0 \\ 0 & a_{32} & a_{33} & a_{34} \\ 0 & 0 & a_{43} & a_{44} \end{bmatrix} = \begin{bmatrix} l_{11} & 0 & 0 & 0 \\ l_{21} & l_{22} & 0 & 0 \\ 0 & l_{32} & l_{33} & 0 \\ 0 & 0 & l_{43} & l_{44} \end{bmatrix} \begin{bmatrix} 1 & u_{12} & 0 & 0 \\ 0 & 1 & u_{23} & 0 \\ 0 & 0 & 1 & u_{34} \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

**Solution** The  $LU$  factorization of  $A$  has the form

$$A = \begin{bmatrix} a_{11} & a_{12} & 0 & 0 \\ a_{21} & a_{22} & a_{23} & 0 \\ 0 & a_{32} & a_{33} & a_{34} \\ 0 & 0 & a_{43} & a_{44} \end{bmatrix} = \begin{bmatrix} l_{11} & 0 & 0 & 0 \\ l_{21} & l_{22} & 0 & 0 \\ 0 & l_{32} & l_{33} & 0 \\ 0 & 0 & l_{43} & l_{44} \end{bmatrix} \begin{bmatrix} 1 & u_{12} & 0 & 0 \\ 0 & 1 & u_{23} & 0 \\ 0 & 0 & 1 & u_{34} \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix} = \begin{bmatrix} l_{11} & l_{11}u_{12} & 0 & 0 \\ l_{21} & l_{22} + l_{21}u_{12} & l_{22}u_{23} & 0 \\ 0 & l_{32} & l_{33} + l_{32}u_{23} & l_{33}u_{34} \\ 0 & 0 & l_{43} & l_{44} + l_{43}u_{34} \end{bmatrix}.$$

Thus

$$\begin{aligned}
 a_{11} : \quad 2 &= l_{11} \implies l_{11} = 2, & a_{12} : \quad -1 &= l_{11}u_{12} \implies u_{12} = -\frac{1}{2}, \\
 a_{21} : \quad -1 &= l_{21} \implies l_{21} = -1, & a_{22} : \quad 2 &= l_{22} + l_{21}u_{12} \implies l_{22} = -\frac{3}{2}, \\
 a_{23} : \quad -1 &= l_{22}u_{23} \implies u_{23} = -\frac{2}{3}, & a_{32} : \quad -1 &= l_{32} \implies l_{32} = -1, \\
 a_{33} : \quad 2 &= l_{33} + l_{32}u_{23} \implies l_{33} = \frac{4}{3}, & a_{34} : \quad -1 &= l_{33}u_{34} \implies u_{34} = -\frac{3}{4}, \\
 a_{43} : \quad -1 &= l_{43} \implies l_{43} = -1, & a_{44} : \quad 2 &= l_{44} + l_{43}u_{34} \implies l_{44} = \frac{5}{4}.
 \end{aligned}$$

This gives the Crout factorization

$$A = \begin{bmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 & 0 \\ -1 & \frac{3}{2} & 0 & 0 \\ 0 & -1 & \frac{4}{3} & 0 \\ 0 & 0 & -1 & \frac{5}{4} \end{bmatrix} \begin{bmatrix} 1 & -\frac{1}{2} & 0 & 0 \\ 0 & 1 & -\frac{2}{3} & 0 \\ 0 & 0 & 1 & -\frac{3}{4} \\ 0 & 0 & 0 & 1 \end{bmatrix} = LU.$$

Solving the system

$$L\mathbf{z} = \begin{bmatrix} 2 & 0 & 0 & 0 \\ -1 & \frac{3}{2} & 0 & 0 \\ 0 & -1 & \frac{4}{3} & 0 \\ 0 & 0 & -1 & \frac{5}{4} \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \\ z_3 \\ z_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} \quad \text{gives} \quad \begin{bmatrix} z_1 \\ z_2 \\ z_3 \\ z_4 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ \frac{1}{3} \\ \frac{1}{4} \\ 1 \end{bmatrix},$$

and then solving

$$U\mathbf{x} = \begin{bmatrix} 1 & -\frac{1}{2} & 0 & 0 \\ 0 & 1 & -\frac{2}{3} & 0 \\ 0 & 0 & 1 & -\frac{3}{4} \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ \frac{1}{3} \\ \frac{1}{4} \\ 1 \end{bmatrix} \quad \text{gives} \quad \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}. \quad \blacksquare$$



Do the following from Book

(Ex 6.6 )

Do Q 1,3,5, 11 & 12