

33 Polyagglutination and Cryptantigens

33.1 Introduction, 515

33.2 Acquired polyagglutination and the cryptantigens involved, 515

33.3 Inherited polyagglutination, 519

33.4 Polyagglutination of undetermined status, 520

33.1 Introduction

Polyagglutination was defined by Bird [1] as, 'The agglutination of red cells, irrespective of blood group, by many sera, the abnormality being a property of the red cells and not the sera'. Microbial polyagglutination generally results from the effect of bacterial or viral glycosidases revealing concealed antigenic determinants, called cryptantigens. These enzymes catalyse the cleavage of terminal monosaccharides or acetyl groups from the oligosaccharide chains of membrane glycoproteins or glycolipids. Polyagglutination occurs because of IgM antibodies against these cryptantigens present in the serum of most people; sera from those individuals with polyagglutinable red cells usually lack the corresponding antibody. Somatic mutation resulting in incomplete biosynthesis of oligosaccharides may lead to a more persistent expression of cryptantigen. Another type of polyagglutination arises from the inheritance of very rare antigens, which are detected by antibodies in most adult human sera. A classification of polyagglutination is provided in Table 33.1. For reviews on polyagglutination see [1–7].

Lectins are non-immunoglobulin extracts from plants and occasionally animals. They have proved invaluable in the identification and classification of cryptantigens and polyagglutination. Some of the most useful lectins are listed in Table 33.2.

33.2 Acquired polyagglutination and the cryptantigens involved

33.2.1 Microbial polyagglutination

Microbial polyagglutination is associated with septicaemia, bowel or respiratory tract infections, wound infections, and conditions, such as tumours or obstructions, that compromise the integrity of the bowel wall and permit entry of microbial enzymes into the blood [1,2,5–7,17]. Occasionally microbial polyagglutination is detected in apparently healthy individuals, presumably because of subclinical infection or an undetected intestinal lesion. Before microbial enzymes can modify red cell membrane structures they must be present in sufficient quantity to neutralise plasma enzyme inhibitors. Cryptantigens are quite commonly detectable with lectins, but not with human sera, on the red cells of patients with various infections, sometimes before the infection has been diagnosed. Microbial polyagglutination is usually transient, the polyagglutinability of the cells diminishing as infection abates.

Red cells may also become polyagglutinable *in vitro*, either as a result of bacterial contamination of blood samples or by the use of enzymes isolated from micro-organisms.

33.2.1.1 T polyagglutination

The Hübener–Thomsen–Friedenreich phenomenon [18] or T activation of red cells results from microbial

sialidases catalysing the cleavage of sialic acid residues from the O-linked disialotetrasaccharides of the major red cell sialoglycoproteins, glycophorins A and B, and from other glycoproteins and glycolipids. The T determinant (CD176) is basically terminal D-galactose in β 1,3 linkage with GalNAc, the core 1 O-glycan (Table 33.3 and see Chapter 3). T exposure causes depression of M and N antigens [5] and the reduction in cell surface charge makes T-active cells agglutinable by *Glycine soja* lectin [14]. Sialidases from bacteria such as *Clostridium perfringens*, *Vibrio cholerae*, and pneumococci, and from the influenza virus, have all been responsible for *in vivo* T

cryptantigen exposure. T activation is more common in infants than in adults and is particularly associated with paediatric intestinal disorders [17,19,20].

In 1964, Bird [8] found that extracts of peanuts, *Arachis hypogaea*, made a powerful anti-T lectin. The term ‘peanut-positive polyagglutination’ now refers to the exposure of any of five cryptantigens, T, Tk, Th, Tx, or Tr, that bind *Arachis hypogaea* lectin (Table 33.2). Red cells of 52 (0.5%) of 9672 hospital patients were agglutinated by *Arachis hypogaea* lectin, but in only one patient were the cells polyagglutinable [21].

Almost all normal human sera agglutinate T-activated red cells, but only a very small proportion (2% [22]) are haemolytic *in vitro*. This haemolytic activity is usually weak and dependent on the degree of desialylation [22]. Monoclonal antibodies with T specificity have been produced following the immunisation of rodents with desialylated red cells [23–25]. Some monoclonal anti-Le^a cross-react with T-activated Le(a–) red cells [26].

T polyagglutination is occasionally accompanied by haemolysis [27]. Whether this is due to complement fixing anti-T in the patient’s serum or to an effect of the associated infection is unclear. Haemolytic-uraemic syndrome in children following pneumococcal infection may be caused by the effect of bacterial sialidase; T activation of red cells, platelets, and renal glomeruli could be responsible for the characteristic symptoms of haemolytic anaemia, thrombocytopenia, and renal failure [28–30]. Transfusion of plasma or whole blood containing anti-T has been implicated as a cause of severe haemolytic reactions in infants with necrotising enterocolitis

Table 33.1 Classification of polyagglutination.

Acquired

Microbial

Cryptantigens uncovered by microbial enzymes

T, Tk, Th, Tx, Acquired B

Adsorption of bacteria or bacterial products onto cell surface

Non-microbial (persistent)

Biosynthetic blockage resulting from somatic mutation

Tn, ?Th

Inherited

Sd(a++) (Cad), CDA II, NOR, Hyde Park

Undetermined

VA, Tr

Table 33.2 Reactions of lectins with cryptantigens and with Sd(a++) and Hyde Park polyagglutinable cells.

Lectins	T	Th	Tk	Tx	Tn	Sd(a++)	Hyde Park	Tr	References
<i>Arachis hypogaea</i>	+	+	+	+	–	–	w	+	[8]
<i>Vicia hircanica</i>	+	+	+		–	–	+		[9]
<i>Vicia cretica</i>	+	+	–	–	–	–	w		[10]
<i>Vicia villosa</i>	+	–	–		+	+	+	+	[11]
<i>Griffonia simplicifolia</i> (GSII)	–	–	+	–	–	–	+	+	[12]
<i>Medicago disciformis</i>	+	+	–	–	–		+		[13]
<i>Dolichos biflorus</i> (B and O only)	–	–	–	–	+	+	–	–	[14]
<i>Salvia sclarea</i>	–	–	–	–	+	–	–	+	[15]
<i>Salvia horminum</i>	–	–	–	–	+	+	w	+	[15]
<i>Leonurus cardiaca</i>	w	–	–	–	–	+	–		[16]
<i>Glycine soja</i>	+	–	–	–	+	+	+/–		[14]

w, weak agglutination; +/–, reactive with strong examples only.

Table 33.3 O-linked oligosaccharides of sialoglycoproteins from normal red cells and from T- and Tn-activated cells. See Table 2.4 for abbreviations.

Normal	NeuAc α 2 \rightarrow 3Gal β 1 \rightarrow 3GalNAc—Ser/Thr
	6 ↑
	NeuAc α 2
T	Gal β 1 \rightarrow 3GalNAc—Ser/Thr
(core 1)	
Tn	GalNAc—Ser/Thr
Sialyl-Tn	GalNAc—Ser/Thr
	6 ↑
	NeuAc α 2

(NEC) and T-activated red cells, though this complication is rare [17,31]. Whether patients likely to have T-activated cells, such as children with NEC, should be transfused with plasma, is controversial [17,22], and plasma exchange may be considered beneficial in extreme cases [32].

Leucocytes and platelets of individuals with T-polyagglutinable red cells also express T antigen [33].

33.2.1.2 Th polyagglutination

Th polyagglutination is distinguished from T polyagglutination by the use of lectins (Table 33.2) and probably represents a weak expression or early stage of T activation caused by sialidase from *Corynebacterium aquaticum* [34,35]. Sialidases from *Vibrio cholerae* used under conditions of particularly mild hydrolysis will also produce Th activity, *in vitro*. Release of sialic acid below a critical quantity (20 μ g per 10¹⁰ red cells) leads to Th activity, whereas liberation of greater quantities of sialic acid results in T polyagglutinability [35]. It is possible that Th activity represents the removal of only one sialic acid residue from the tetrasaccharide shown in Table 32.3, exposing terminal Gal, but not the T disaccharide (Gal β 1 \rightarrow 3GalNAc).

Red cells from 22 of 200 cord samples were Th-activated (but not polyagglutinable); in six of these the mother's red cells were also Th-active [36]. A patient with Th polyagglutination and peritonitis arising from a perforated colonic tumour died as a result of severe intravascular haemolysis [37].

It is possible that a persistent form of Th polyagglutination resulting from incomplete biosynthesis also

occurs. Persistent Th activation in the absence of infection was identified in five of seven children with congenital hypoplastic anaemia [38] and in a patient with myelodysplasia whose red cells were also Tn-active [39].

33.2.1.3 Tk polyagglutination

Tk polyagglutination [40] is most readily distinguished from T polyagglutination by the use of GSII, a Tk-specific lectin isolated from the seeds of *Griffonia simplicifolia* [12]. Expression of Tk is enhanced by papain treatment and Tk-activated cells are not sialic acid deficient [40]. *Vicia hircanica* seeds contain separable T- and Tk-specific lectins [9]. Monoclonal anti-Tk has been produced by immunising mice with endo- β -galactosidase-modified red cells [41].

Tk polyagglutination is usually associated with *Bacteroides fragilis* infection [42], but *Serratia marcescens*, *Candida albicans*, *Aspergillus niger* and, in children, invasive pneumococcal infection, have also been responsible [3,4,43]. Endo- and exo- β -galactosidases secreted by these pathogens cleave oligosaccharides at Gal β 1 \rightarrow 4GlcNAc linkages on ABH and Ii active blood group substances (Tables 2.3, 2.5, and 25.1) exposing terminal GlcNAc residues, the Tk receptor [44,45]. Tk-transformed red cells may have reduced expression of ABH and Ii antigens [46,47]. GSII lectin is inhibited by GlcNAc [12,48] and monoclonal anti-Tk was adsorbed by GlcNAc β 1 \rightarrow 3Gal β 1 \rightarrow 4Glc β -R coupled to silica beads [41]. Endo- β -galactosidases from *B. fragilis*, *Escherichia freundii*, and *Flavobacterium keratolyticus* have been used to produce Tk-activated red cells *in vitro* [41,45,49].

Tk polyagglutination is often associated with acquired B polyagglutination (Section 33.2.1.5) and with T activation in pneumococcal infections in infants [43].

33.2.1.4 Tx cryptantigen

Tx was identified on the red cells of children with pneumococcal infections and could be produced, *in vitro*, by the action on red cells of culture supernatant from pneumococci derived from these patients [50]. Tx is distinguished from T by *Vicia cretica* lectin (Table 33.2). Tx is only slightly weakened by papain treatment [50]. A persistent form of Tx polyagglutination was detected over a period of 5 years in a woman with myelodysplastic syndrome and no symptoms of infection [51].

33.2.1.5 Acquired B

Acquired B phenomenon is described in detail in Section 2.13.1. It is caused by microbial deacetylases converting GalNAc of an A determinant to galactosamine, which

resembles Gal closely enough for the cells to be agglutinated by some anti-B reagents. Acquired B is usually associated with a unique polyagglutination, but is often also associated with Tk and occasionally T or Th polyagglutination.

33.2.2 Non-microbial polyagglutination

33.2.2.1 Tn polyagglutination

In Tn polyagglutination [52] only a proportion of the red cells are polyagglutinable: between 30 and 90% are agglutinated by anti-Tn, the remainder are Tn-negative. Like T, Tn is located on O-linked oligosaccharides of red cell glycoproteins, chiefly glycophorins A and B [53]. The Tn determinant (CD175) is GalNAc attached to serine or threonine (Table 33.3) [51–55]. The GalNAc is sometimes sialylated, to form sialyl-Tn (CD175s, Table 33.3) [56].

Tn is exposed by defective oligosaccharide biosynthesis [57,58]. Tn is the biosynthetic precursor of T (core 1), which is a precursor of the disialotetrasaccharide typical of red cell sialoglycoproteins (Table 33.3). In Tn-polyagglutinable cells, conversion of the Tn monosaccharide to the T disaccharide does not occur because of deficiency of T synthase, the core 1 β -3-D-galactosyltransferase [59,60]. This enzyme deficiency arises from a somatic mutation within a pluripotent haemopoietic stem cell and subsequent clonal expansion and proliferation of populations of Tn+ red cells, platelets, granulocytes, and lymphocytes [61–63]. Tn-active red cells are not totally deficient in T synthase as some normal tetrasaccharides are present on the glycophorin molecules [64].

The somatic mutations responsible for Tn phenotype do not occur in *C1GALT1*, the gene encoding T synthase on chromosome 7p21.3, but in *C1GALT1C1*, the gene encoding Cosmc, a molecular chaperone required for effective functioning of T synthase [65–67]. Cosmc interacts with denatured, but not native, T synthase to form a non-covalent and reversible complex that results in the acquisition of T-synthase activity [68]. Four Cosmc mutations have been shown to be responsible for the Tn phenotype in different individuals: 3G>C, Met1Ile; 202C>T, Arg68stop; 454G>A, Glu152Lys; and 577T>C, Ser 193Pro. In one other Tn individual no Cosmc cDNA was detected [66,67]. Cells of the Jurkat human lymphoblastoid T-cell line express Tn and have a mutated Cosmc gene with a deletion introducing a premature stop codon [65]. Transfection of Jurkat cells with Cosmc cDNA restores T-synthase activity and ablates Tn expression, but transfection of Cosmc cDNA containing any of

the Tn mutations has no effect on Tn expression [66, 67]. The Cosmc gene is X-linked, located at Xq24 [65]. Owing to the presence of a Y-chromosome in males and X-chromosome inactivation in females, somatic cells contain only one active X-chromosome and consequently only one active Cosmc gene. If a similar process occurs in Tn phenotype as that in PNH (see Section 19.5), then cytotoxic T cells targeting the normal O-glycans of blood cells would be expected in individuals having large populations of Tn-positive red cells. Experimental ablation of the Cosmc gene in mice results in embryonic death, with Tn expressed on every observable cell of the embryo [69].

Tn-active red cells have reduced sialic acid content [53,70] and depressed M and N antigens and T cryptantigen [54,71,72]. Tn is destroyed by papain treatment of red cells [70,73].

The immunodominant monosaccharide of Tn is GalNAc, so Tn-activated red cells are agglutinated by lectins that also agglutinate group A cells, such as *Dolichos biflorus* and *Helix pomatia* [57,70,73]. *Salvia sclarea* lectin is more specific for Tn (Table 33.2) [15]. Tn cells are more strongly agglutinated by human sera containing anti-A than those lacking it [70]. Many murine monoclonal Tn antibodies have been produced, though they may cross-react with other carbohydrate epitopes [74], including sialyl-Tn [75]. An scFv–Fc fusion protein, produced by employing a simultaneous positive and negative selection strategy with Tn and A antigens, respectively, had strict Tn specificity [76]. All Tn antibodies and lectins react only with the Tn+ population of cells giving the characteristic mixed field pattern of agglutination with red cells of individuals with Tn polyagglutination.

Tn polyagglutination is often associated with haemolytic anaemia, leucopenia, and thrombocytopenia [27] and, on occasion, is detected in healthy blood donors [70,77]. Although usually persistent, Tn polyagglutination has been known to recede [58]. There are three reports of transient Tn polyagglutination in newborns, possibly resulting from late development of full T-synthase activity [78–80]. Flow cytometry with monoclonal anti-Tn revealed less than 1×10^{-6} Tn red cells in the peripheral blood of healthy donors [81].

33.2.3 T, Tn, sialyl-Tn, and Tk in malignancy

T and Tn are cryptantigens on epithelial tissues, in the carbohydrate chains of glycoproteins and glycolipids. Springer and others have shown that T, Tn, or sialyl-Tn are exposed on about 90% of primary and metastatic

carcinomas, probably as a result of incomplete biosynthesis [82]. All human tumour cells examined that express Tn and sialyl-Tn harbour mutations in the *Cosmc* gene, resulting in loss of T-synthase activity [83]. Increased density of Tn over T is an indicator of high metastatic potential of the tumour. Immunotherapy with a vaccine derived from T- and Tn-activated red cells has been successful in preventing recurrence of breast cancer and has a potential for wider application (reviewed in [84,85]). Monoclonal anti-Tk reacted with 48% of human colorectal carcinomas [41].

Reports of patients with Tn polyagglutination and concurrent acute myeloid leukaemia (AML), or who subsequently developed leukaemia, led Ness *et al.* [86] to propose that Tn polyagglutination may represent a pre-leukaemic state and that careful clinical observation of individuals with Tn polyagglutination would be circum-spect. Polyagglutination vanished during chemotherapy [58,86]. Tn polyagglutination has also been associated with myelodysplasia [39,77].

33.3 Inherited polyagglutination

33.3.1 Sd(a++) (Cad)

Sd^a is a red cell antigen with an incidence of about 91% and of variable strength. Red cells of the very rare individuals with an extra strong form of Sd^a antigen (referred to as Sd(a++) or Cad) are polyagglutinable. The immunodominant structure of Sd^a is GalNAc in β -linkage with Gal. Sd(a++) red cells are agglutinated by *Dolichos biflorus* and *Helix pomatia* lectins, but can be distinguished from Tn cells by *Salvia sclarea* and *Leonurus cardiaca* (Table 33.2). Sd^a and Sd(a++) are described in detail in Chapter 31.

33.3.2 Congenital dyserythropoietic anaemia type II (CDA II or HEMPAS)

CDA II (or hereditary erythroblastic multinuclearity with a positive acidified serum lysis test, HEMPAS) is a rare, autosomal-recessive syndrome found predominantly in southern Italy and Central Europe [87,88]. Most patients have a mild anaemia, although some become transfusion-dependent. Red cells of patients with CDA II are agglutinated at 20°C, or lysed at 37°C, by complement fixing IgM antibodies in about one third of normal sera [89,90]. CDA II red cells have elevated i, normal I, and depressed H [89–92]. They have reduced levels of *N*-glycosylation of cell surface glycoproteins, particularly band 3 and

band 4.5 (glucose transporter), which contain repeating *N*-acetylglucosamine (Gal β 1 \rightarrow 4GlcNAc) units [93], revealing a cryptantigen responsible for the polyagglutination and resulting in excess production of i-active linear *N*-acetylglucosaminylceramides. In addition, 10–35% of erythroblasts in CDA II are binucleated and some red cells have a second membrane close to the outer membrane [87].

CDA II is caused by mutations in *SEC23B*, a gene comprising 20 exons and located on chromosome 20p11.23 [94,95]. *SEC23B* encodes a component of the COPII coat protein complex [96], which is involved in protein-trafficking through membrane vesicles. Abnormalities of *SEC23B* may disturb endoplasmic reticulum-to-Golgi trafficking and interfere with sugar transporters and glycosyltransferases, affecting glycosylation pathways [97]. Homozygosity and compound heterozygosity for numerous *SEC23B* missense or inactivating mutations result in CDA II [94,95,98–100]. Homozygosity or compound heterozygosity for two inactivating mutations has been found only once [100], suggesting that complete lack of the protein could be incompatible with life; compound heterozygosity for a missense and an inactivating mutation tends to result in more severe symptoms than homozygosity or compound heterozygosity for missense mutations [98]. The most common mutations encode Glu109Lys and Arg14Trp and represent 32% and 19%, respectively, of mutations identified [100]. CDAII patients exhibit 40–60% decrease in *SEC23B* mRNA levels, and the patient with a nonsense mutation and a splice site mutation had 30% mRNA level [100].

33.3.3 NOR polyagglutination

NOR is a very rare form of polyagglutination found in only two families. It results from the expression of abnormal glycolipids resulting from a mutation in *A4GALT*, the gene encoding the enzyme responsible for biosynthesis of P1 and P antigens. Consequently, NOR antigen is P1PK4 and is described in Section 4.5.

33.3.4 Hyde Park polyagglutination

The association between a unique form of polyagglutination associated with a rare haemoglobin variant in a large South African family of mixed ethnicity remains a puzzle. Thirty-five members of the family were studied: 12 had the variant haemoglobin M-Hyde Park and polyagglutinable red cells; 23 had neither [101]. Polyagglutination had not been observed previously in association with haemoglobin M-Hyde Park.

Group O, N+ Hyde Park red cells show the following serological characteristics [101,102]:

- 1 agglutination with a minority of normal human sera (7 of 40);
- 2 enhanced agglutination with *Vicia graminea* and *Ulex europaeus* lectins and with human anti-I and -i, but normal strength agglutination with rabbit and monoclonal anti-N;
- 3 agglutination by monoclonal anti-Tn, but not by Tn-specific *Salvia sclarea* lectin;
- 4 agglutination by *Glycine soja*, *Sophora japonica*, and *Arachis hypogaea* (weakly), lectins that detect desialylated O-glycans; and
- 5 agglutination by the GlcNAc-specific lectins *Vicia hircanica* and GSII (weakly) (Table 33.2).

These serological characteristics, together with results of biochemical analyses, suggest that the polyagglutination results from two unrelated anomalies, one associated with heterogeneity of sialylation of the O-glycans of glycophorin molecules and the other associated with exposed GalNAc residues on the N-glycans of band 3 and band 4.5 [103].

33.4 Polyagglutination of undetermined status

33.4.1 VA polyagglutination

VA polyagglutination is very rare. The red cells are not agglutinated by *Arachis hypogaea* or *Dolichos biflorus* lectins, but give a mixed-field agglutination with *Helix pomatia* lectin [103,104]; they also give a characteristic stippled appearance in immunofluorescence with *H. pomatia* lectin [104]. An associated depression of H antigen has led to the suggestion that microbial α -fucosidase could be responsible for VA exposure [103, 105]. In the original case VA polyagglutination was persistent and associated with haemolytic anaemia [103,104]; in the only other reported example, VA-active red cells were also Tk active [105].

33.4.2 Tr polyagglutination

Tr polyagglutination has been found in one individual [106]. Tr red cells gave a unique pattern of reactions with a panel of lectins (Table 33.2) and reacted with monoclonal anti-T + Tn and -Tk. Results of tests with lectins, reduced periodic acid-Schiff staining of glycophorins A, B, and C, and increased electrophoretic mobility of band 3 suggested a reduction in sialylation of both N- and O-glycans, exposing $\beta 1 \rightarrow 4$ Gal and $\beta 1 \rightarrow 3$ Gal residues. It is

feasible that Tr results from a defective glycosyltransferase, probably an $\alpha 1,6$ -sialyltransferase. If so, that may also account for a bleeding defect, probably resulting from reduced platelet glycoprotein Ib levels, and for other defects suggesting a connective tissue disorder.

References

- 1 Bird GWG. Complexity of erythrocyte polyagglutinability. In: Mohn JE, Plunkett RW, Cunningham RK, Lambert RM, eds. *Human Blood Groups, 5th Int Convoc Immunol, Buffalo NY*. Basel: Karger, 1977:335–343.
- 2 Bird GWG. Clinical aspects of red blood cell polyagglutinability of microbial origin. In: Salmon C, ed. *Blood Groups and Other Red Cell Surface Markers in Health and Disease*. New York: Masson, 1982:55–64.
- 3 Judd WJ. Microbial-associated forms of polyagglutination (T, Tk and acquired-B). In: Beck ML, Judd WJ, eds. *Polyagglutination, a Technical Workshop*. Arlington: American Association of Blood Banks, 1980:23–53.
- 4 Judd WJ. Review: polyagglutination. *Immunohematology* 1992;8:58–69.
- 5 Vaith P, Uhlenbruck G. The Thomsen agglutination phenomenon: a discovery revisited 50 years later. *Z Immun Forsch* 1978;154:1–14.
- 6 Horn KD. The classification, recognition and significance of polyagglutination in transfusion medicine. *Blood Rev* 1999;13:36–44.
- 7 Beck ML. Red blood cell polyagglutination: clinical aspects. *Semin Hematol* 2000;37:186–196.
- 8 Bird GWG. Anti-T in peanuts. *Vox Sang* 1964;9:748–749.
- 9 Liew YW, Bird GWG. Separable anti-T and anti-Tk lectins from the seeds of *Vicia hircanica*. *Vox Sang* 1988;54: 226–227.
- 10 Bird GWG, Wingham J. *Vicia cretica*: a powerful lectin for T- and Th- but not Tk- or other polyagglutinable erythrocytes. *J Clin Pathol* 1981;34:69–70.
- 11 Bird GWG, Wingham J. Lectins for polyagglutinable red cells: *Cytisus scoparius*, *Spartium junceum* and *Vicia villosa*. *Clin Lab Haematol* 1980;2:21–23.
- 12 Judd WJ, Beck ML, Hicklin BL, Shankar Iyer PN, Goldstein IJ. BSII lectin: a second hemagglutinin isolated from *Ban-deiraea simplicifolia* seeds with affinity for type III polyagglutinable red cells. *Vox Sang* 1977;33:246–251.
- 13 Bird GWG, Wingham J. 'New' lectins for the identification of erythrocyte cryptantigens and the classification of erythrocyte polyagglutinability: *Medicago disciformis* and *Medicago turbinata*. *J Clin Pathol* 1983;36:195–196.
- 14 Bird GWG. Lectins in immunohematology. *Transfus Med Rev* 1989;3:55–62.
- 15 Bird GWG, Wingham J. Haemagglutinins from *Salvia*. *Vox Sang* 1974;26:163–166.

- 16 Bird GWG, Wingham J. Anti-Cad lectin from the seeds of *Leonurus cardiaca*. *Clin Lab Haematol* 1979;1:57–59.
- 17 Eder AF, Manno CS. Does red cell T activation matter? *Br J Haematol* 2001;114:25–30.
- 18 Friedenreich V. *The Thomsen Hemagglutination Phenomenon*. Copenhagen: Levin and Munksgaard, 1930.
- 19 Seger R, Joller P, Bird GWG, *et al*. Necrotising enterocolitis and neuraminidase-producing bacteria. *Helv Paediat Acta* 1980;35:121–128.
- 20 Seger RA, Kenny A, Bird GWG, *et al*. Pediatric surgical patients with severe anaerobic infection: report of 16 T-antigen positive cases and possible hazards of blood transfusion. *J Pediatr Surg* 1981;16:905–910.
- 21 Rawlinson VI, Stratton F. Incidence of T activation in a hospital population. *Vox Sang* 1984;46:306–317.
- 22 Burin des Roziers N, Bodivit G, Chadebec P, *et al*. Anti-T haemolysins: the effects of sialic acid removal and 2-aminoethylisothiuronium bromide treatment of erythrocytes on immune lysis. *Vox Sang* 2011;100:401–408.
- 23 Rahman AFR, Longenecker BM. A monoclonal antibody specific for the Thomsen–Friedenreich cryptic T antigen. *J Immunol* 1982;129:2021–2024.
- 24 Seitz R, Fischer K, Poschmann A. Differentiation of red cell membrane abnormalities causing T-polyagglutination by use of monoclonal antibodies. *Rev Franc Transfus Immuno-Hémat* 1983;26:420.
- 25 Metcalfe S, Springer GF, Svvennsen RJ, Tegtmeyer H. Monoclonal antibodies specific for human Thomsen–Friedenreich (T) and Tn blood group precursor antigens. *Prot Biol Fluids* 1985;32:765–768.
- 26 Pisacka M, Stambergová M. Activation of Thomsen–Friedenreich antigen on red cells: a possible source of errors in antigen typing with some monoclonal antibodies. *Vox Sang* 1994;66:300.
- 27 Klein HG, Anstee DJ. *Mollison's Blood Transfusion in Clinical Medicine*, 11th edn. Oxford: Blackwell Publishing, 2005.
- 28 Klein PJ, Bulla M, Newman RA, *et al*. Thomsen–Friedenreich antigen in haemolytic-uraemic syndrome. *Lancet* 1977;ii: 1024–1025.
- 29 Seger R, Joller P, Baerlocher K, *et al*. Hemolytic-uremic syndrome associated with neuraminidase-producing microorganisms: treatment by exchange transfusion. *Helv Paediat Acta* 1980;35:359–367.
- 30 Wang L-Y, Chan Y-S, Chang F-C, Wang C-L, Lin M. Thomsen–Friedenreich activation in infants with necrotizing enterocolitis in Taiwan. *Transfusion* 2011;51:1972–1976.
- 31 Engelfriet CP, Reesink HW, Strauss RG, *et al*. Blood transfusion in premature or young infants with polyagglutination and activation of the T antigen. *Vox Sang* 1999;76: 128–132.
- 32 Hopkins CK, Yuan S, Lu Q, Ziman A, Goldfinger D. A severe case of atypical hemolytic uremic syndrome associated with pneumococcal infection and T activation treated successfully with plasma exchange. *Transfusion* 2008;48: 2448–2452.
- 33 Hysell JK, Hysell JW, Nichols ME, Leonardi RG, Marsh WL. *In vivo* and *in vitro* activation of T-antigen receptors on leukocytes and platelets. *Vox Sang* 1976;31(Suppl. 1):9–15.
- 34 Bird GWG, Wingham J, Beck ML, *et al*. Th, a 'new' form of erythrocyte polyagglutination. *Lancet* 1978;i:1215–1216.
- 35 Sondag-Thull D, Levene NA, Levene C, *et al*. Characterization of a neuraminidase from *Corynebacterium aquaticum* responsible for Th polyagglutination. *Vox Sang* 1989;57: 193–198.
- 36 Wahl CM, Herman JH, Shirey RS, Kickler TS, Ness PM. Th activation of maternal and cord blood. *Transfusion* 1989;29:635–637.
- 37 Levene NA, Levene C, Gekker K, *et al*. Th polyagglutination with fatal outcome in a patient with massive intravascular hemolysis and perforated tumour of colon. *Am J Hematol* 1990;35:127–128.
- 38 Herman JH, Shirey RS, Smith B, Kickler TS, Ness PM. Th activation in congenital hypoplastic anemia. *Transfusion* 1987;27:253–256.
- 39 Janvier D, Guignier F, Reviron M, Benbunan M. Concomitant exposure of Tn and Th cryptantigens on the red cells of a patient with myelodysplasia. *Vox Sang* 1991;61:142–143.
- 40 Bird GWG, Wingham J. Tk: a new form of red cell polyagglutination. *Br J Haematol* 1972;23:759–763.
- 41 Meichenin M, Rocher J, Galanina O, *et al*. Tk, a new colon tumor-associated antigen resulting from altered O-glycosylation. *Cancer Res* 2000;60:5499–5507.
- 42 Inglis G, Bird GWG, Mitchell AAB, Milne GR, Wingham J. Effect of *Bacteroides fragilis* on the human erythrocyte membrane: pathogenesis of Tk polyagglutination. *J Clin Path* 1975;28:964–968.
- 43 Janvier D, Sellami F, Boyeldieu D, *et al*. Invasive pneumococcal infections in young children as a new cause of Tk activation. *Vox Sang* 2008;95(Suppl. 1):197–198.
- 44 Doinel C, Andreu G, Cartron J-P, Salmon C, Fukuda MN. Tk polyagglutination produced *in vitro* by an endo-beta-galactosidase. *Vox Sang* 1980;38:94–98.
- 45 Doinel C, Rufin JM, Andreu G. The Tk antigenic determinant studies of Tk-activated red blood cells with endoglycosidases. *Rev Franc Transfus Immuno-Hémat* 1981;24: 109–116.
- 46 Inglis G, Bird GWG, Mitchell AAB, Wingham J. Tk polyagglutination associated with reduced A and H activity. *Vox Sang* 1978;35:370–374.
- 47 Andreu G, Doinel C, Cartron J-P, Mativet S. Induction of Tk polyagglutination by *Bacteroides fragilis* culture supernatants. Associated modifications of ABH and Ii antigens. *Rev Franc Transfus Immuno-Hémat* 1979;22:551–561.
- 48 Iyer PNS, Wilkinson KD, Goldstein IJ. An N-acetyl-D-glucosamine binding lectin from *Bandeiraea simplicifolia* seeds. *Arch Biochem Biophys* 1976;177:330–333.

- 49 Liew YW, Bird GWG, King MJ. The human erythrocyte cryptantigen Tk: exposure by an endo-beta-galactosidase from *Flavobacterium keratolyticus*. *Rev Franc Transfus Immuno-Hémat* 1982;25:639–641.
- 50 Bird GWG, Wingham J, Seger R, Kenny AB. Tx, a 'new' red cell cryptantigen exposed by pneumococcal enzymes. *Rev Franc Transfus Immuno-Hémat* 1982;25:215–216.
- 51 Pisacka M, Karasova R, Prosicka M, Cermak J. Two cases of erythrocyte cryptantigen reactive with *Arachis hypogaea* in association with myelodysplastic syndrome. *VI Regional Eur Congr Int Soc Blood Transfus*, 1999:81 [Abstracts].
- 52 Dausset J, Moullec J, Bernard J. Acquired hemolytic anemia with polyagglutinability of red blood cells due to a new factor present in normal human serum (anti-Tn). *Blood* 1959;14:1079–1093.
- 53 Dahr W, Uhlenbruck G, Gunson HH, van der Hart M. Molecular basis of Tn-polyagglutinability. *Vox Sang* 1975;29:36–50.
- 54 Sturgeon P, Luner SJ, McQuiston DT. Permanent mixed-field polyagglutinability (PMFP). II. Hematological, biochemical and biochemical observations. *Vox Sang* 1973;25:498–512.
- 55 Dahr W, Uhlenbruck G, Bird GWG. Cryptic A-like receptor sites in human erythrocyte glycoproteins: proposed nature of Tn-antigen. *Vox Sang* 1974;27:29–42.
- 56 Kjeldsen T, Hakomori S, Springer GE, *et al.* Coexpression of sialosyl-Tn (NeuAc α 2 \rightarrow 6GalNAc α 1 \rightarrow O-Ser/Thr) and Tn (GalNAc α 1 \rightarrow O-Ser/Thr) blood group antigens on Tn erythrocytes. *Vox Sang* 1989;57:81–87.
- 57 Bird GWG, Shinton NK, Wingham J. Persistent mixed-field polyagglutination. *Br J Haematol* 1971;21:443–453.
- 58 Bird GWG, Wingham J, Pippard MJ, Hoult JG, Melikian V. Erythrocyte membrane modification in malignant diseases of myeloid and lymphoreticular tissues. *Br J Haematol* 1976;33:289–294.
- 59 Cartron J-P, Andreu G, Cartron J, *et al.* Demonstration of T-transferase deficiency in Tn-polyagglutinable blood samples. *Eur J Biochem* 1978;92:111–119.
- 60 Berger EG, Kozdrowski I. Permanent mixed-field polyagglutinable erythrocytes lack galactosyltransferase activity. *FEBS Letts* 1978;93:105–108.
- 61 Cartron J-P, Nurden AT. Galactosyltransferase and membrane glycoprotein abnormality in human platelets from Tn-syndrome donors. *Nature* 1979;282:621–623.
- 62 Vainchenker W, Testa U, Deschamps JF, *et al.* Clonal expression of the Tn antigen in erythroid and granulocyte colonies and its application to determination of the clonality of the human megakaryocyte colony assay. *J Clin Invest* 1982;69:1081–1091.
- 63 Brouet J-C, Vainchenker W, Blanchard D, Testa U, Cartron J-P. The origin of human B and T cells from multipotent stem cells. *Eur J Immunol* 1983;13:350–352.
- 64 Blumenfeld OO, Lalezari P, Khorshidi M, Puglia K, Fukuda M. O-linked oligosaccharides of glycoporphins A and B in erythrocytes of two individuals with the Tn polyagglutinability syndrome. *Blood* 1992;80:2388–2395.
- 65 Ju T, Cummings RD. A unique molecular chaperone Cosmc required for activity of the mammalian core 1 β -galactosyltransferase. *Proc Natl Acad Sci USA* 2002;99:16613–16618.
- 66 Ju T, Cummings RD. Chaperone mutation in Tn syndrome. *Nature* 2005;437:1252.
- 67 Karamatic Crew V, Singleton BK, Green C, *et al.* New mutations in *C1GALT1C1* in individuals with Tn positive phenotype. *Br J Haematol* 2008;142:657–667.
- 68 Aryal RP, Ju T, Cummings RD. Tight complex formation between Cosmc chaperone and its specific client non-native T-synthase leads to enzyme activity and client-driven dissociation. *J Biol Chem* 2012;287:15317–15329.
- 69 Wang Y, Ju T, Ding X, *et al.* Cosmc is an essential chaperone for correct protein O-glycosylation. *Proc Natl Acad Sci USA* 2010;107:9228–923.
- 70 Myllylä G, Furuholm U, Nordling S, *et al.* Persistent mixed field polyagglutinability. Electrokinetic and serological aspects. *Vox Sang* 1971;20:7–23.
- 71 Sturgeon P, McQuiston DT, Taswell HF, Allan CJ. Permanent mixed-field polyagglutinability (PMFP). I. Serological Observations. *Vox Sang* 1973;25:481–497.
- 72 Bird GWG, Wingham J. The M, N and N_v receptors of Tn-erythrocytes. *Vox Sang* 1974;26:171–175.
- 73 Gunson HH, Stratton F, Mullard GW. An example of polyagglutinability due to the Tn antigen. *Br J Haematol* 1970;18:309–316.
- 74 Manimala JC, Li Z, Jain A, VedBrat S, Gildersleeve JC. Carbohydrate array analysis of anti-Tn antibodies and lectins reveals unexpected specificities: implications for diagnostic and vaccine development. *ChemBiochem* 2005;6:2229–2241.
- 75 O'Boyle KP, Markowitz AL, Khorshidi M, *et al.* Specificity analysis of murine monoclonal antibodies reactive with Tn, sialylated Tn, T, and monosialylated (2 \rightarrow 6) T antigens. *Hybridoma* 1996;15:401–408.
- 76 Kubota T, Matsushita T, Niwa R, Kumagai I, Nakamura K. Novel anti-Tn single-chain Fv-Fc fusion proteins derived from immunized phage library and antibody Fc domain. *Anticancer Res* 2010;30:3397–3405.
- 77 Bird GWG, Wingham J, Richardson SGN. Myelofibrosis, autoimmune haemolytic anaemia and Tn-polyagglutinability. *Haematologia* 1985;18:99–103.
- 78 Wilson MJ, Cott ME, Sotus PC. Probable Tn-activation *in utero*. *Transfusion* 1980;20:622 [Abstract].
- 79 Schultz M, Fortes P, Brewer L, Miller A, Beck M. 'In utero' exposure of Tn and Th cryptantigens. *Transfusion* 1983;23:422 [Abstract].
- 80 Rose RR, Skradski KJ, Polesky HF, *et al.* Transient neonatal Tn-activation: another example. *Transfusion* 1983;23:422 [Abstract].

- 81 Bigbee WL, Langlois RG, Stanker LH, Vanderlaan M, Jensen RH. Flow cytometric analysis of erythrocyte populations in Tn syndrome blood using monoclonal antibodies to glycophorin A and the Tn antigen. *Cytometry* 1990;11:261–271.
- 82 Springer GE. T and Tn, general carcinoma autoantigens. *Science* 1984;224:1198–1206.
- 83 Ju T, Lanneau GS, Gautam G, *et al.* Human tumor antigens Tn and sialyl Tn arise from mutations in *Cosmc*. *Cancer Res* 2008;68:1636–1646.
- 84 Desai PR. Immunoreactive T and Tn antigens in malignancy: role in carcinoma diagnosis, prognosis, and immunotherapy. *Transfus Med Rev* 2000;14:312–325.
- 85 Heimburg-Molinaro J, Lum M, Vijay G, *et al.* Cancer vaccines and carbohydrate epitopes. *Vaccine* 2011;29:8802–8826.
- 86 Ness PM, Garratty G, Morel PA, Perkins HA. Tn polyagglutination preceding acute leukemia. *Blood* 1979;54:30–34.
- 87 Renella R, Wood WG. The congenital dyserythropoietic anemias. *Hematol Oncol Clin N Am* 2009;23:283–306.
- 88 Iolascon A, Russo R, Delaunay J. Congenital dyserythropoietic anemias. *Curr Opin Hematol* 2011;18:146–151.
- 89 Crookston JH, Crookston MC, Burnie KL, *et al.* Hereditary erythroblastic multinuclearity associated with a positive acidified-serum test: a type of congenital dyserythropoietic anaemia. *Br J Haematol* 1969;17:11–26.
- 90 Crookston JH, Crookston MC, Rosse WF. Red-cell abnormalities in HEMPAS (hereditary erythroblastic multinuclearity with a positive acidified-serum test). *Br J Haematol* 1972;23(Suppl.):83–91.
- 91 Bird GWG, Wingham J. The action of seed and other reagents on HEMPAS erythrocytes. *Acta Haematol* 1976;55:174–180.
- 92 Rochant H, Gerbal A. Polyagglutinabilité due à l'antigène HEMPAS. *Rev Franc Transfus Immuno-Hémat* 1976;14:239–245.
- 93 Fukuda MN, Masri KA, Dell A, Luzzatto L, Moremen KW. Incomplete synthesis of N-glycans in congenital dyserythropoietic anemia type II caused by a defect in the gene encoding α -mannosidase II. *Proc Natl Acad Sci USA* 1990;87:7443–7447.
- 94 Schwarz K, Iolascon A, Verissimo F, *et al.* Mutations affecting the secretory COPII coat component SEC23B cause congenital dyserythropoietic anemia type II. *Nature Genet* 2009;41:936–940.
- 95 Bianchi P, Fermo E, Vercellati C, *et al.* Congenital dyserythropoietic anemia type II (CDAIL) is caused by mutations in the SEC23B gene. *Hum Mutation* 2009;30:1–7.
- 96 Paccaud J-P, Reith W, Carpentier J-L, *et al.* Cloning and functional characterization of mammalian homologues of the COPII component Sec23. *Mol Biol Cell* 1996;7:1535–1546.
- 97 Hughes H, Stephens DJ. Assembly, organization, and function of the COPII coat. *Histochem Cell Biol* 2008;129:129–151.
- 98 Iolascon A, Russo R, Esposito MR, *et al.* Molecular analysis of 42 patients with congenital dyserythropoietic anemia type II: new mutations in the SEC23B gene and a search for a genotype–phenotype relationship. *Haematologica* 2010;95:708–715.
- 99 Russo R, Esposito MR, Asci R, *et al.* Mutational spectrum in congenital dyserythropoietic anemia type II: identification of 19 novel variants in SEC23B gene. *Am J Hematol* 2010;85:915–920.
- 100 Punzo F, Bertoli-Avella A, Scianguetta S, *et al.* Congenital dyserythropoietic anemia type II: molecular analysis and expression of the SEC23B gene. *Orphanet J Rare Dis* 2011;6:89.
- 101 Bird AR, Kent P, Moores PP, Elliott T. Haemoglobin M-Hyde Park associated with polyagglutinable red blood cells in a South African family. *Br J Haematol* 1988;68:459–464.
- 102 King MJ, Liew YW, Moores PP, Bird GWG. Enhanced reaction with *Vicia graminea* lectin and exposed terminal N-acetyl-D-glucosaminyl residues on a sample of human red cells with Hb M-Hyde Park. *Transfusion* 1988;28:549–555.
- 103 Graninger W, Rameis H, Fischer K, *et al.* 'VA', a new type of erythrocyte polyagglutination characterized by depressed H receptors and associated with hemolytic anemia. I. Serological and hematological observations. *Vox Sang* 1977;32:195–200.
- 104 Graninger W, Poschmann A, Fischer K, *et al.* 'VA', a new type of erythrocyte polyagglutination characterized by depressed H receptors and associated with hemolytic anemia. II. Observations by immunofluorescence, electron microscopy, cell electrophoresis and biochemistry. *Vox Sang* 1977;32:201–207.
- 105 Beck ML, Myers MA, Moulds J, *et al.* Coexistent Tk and VA polyagglutination. *Transfusion* 1978;18:680–684.
- 106 Halverson GR, Lee AH, Øyen R, *et al.* Altered glycosylation leads to Tr polyagglutination. *Transfusion* 2004;44:1588–1592.