

21 Indian Blood Group System and the AnWj Antigen

21.1 Introduction, 449
21.2 CD44 and the Indian antigens, 449
21.3 Indian antigens, 450
21.4 Effects of *In*(Lu) on CD44 and Indian antigens, 451

21.5 Indian antibodies, 452
21.6 Functional aspects of CD44, 452
21.7 AnWj (901009), 453

21.1 Introduction

In^b/*In*^a, a polymorphism in people from the Indian sub-continent and in Arabs, results from an Arg46Pro substitution in CD44. *In*^a and *In*^b are of low and high incidence, respectively. The Indian system also contains two antigens of very high frequency, IN3 and IN4 (Table 21.1). CD44 is a ubiquitous glycoprotein with a variety of functions, mostly associated with its ability to bind hyaluronan, a component of the extracellular matrix. The gene encoding CD44 is on chromosome 11p13.

The high frequency antigen AnWj (901009) has not been assigned to the Indian system, but is either located on an isoform of CD44 or is closely associated with it.

21.2 CD44 and the Indian antigens

21.2.1 CD44

CD44, which is present on cells of most tissues, is a member of the link module superfamily of proteoglycans (for review see [1]). It is a major red cell membrane component of apparent MW 80 kDa [2,3]. The *CD44* gene spans 50 kb of DNA and consists of 20 exons [4,5]. CD44 exists as multiple isoforms, arising partly from alternative splicing of 10 variant exons (Figure 21.1) and partly from variation in glycosylation [4,5]. CD44s (standard), which contains products of none of the alternatively spliced exons, is the isoform present on

haemopoietic cells. Exon 1 encodes the signal sequence. The 248 amino acid N-terminal extracellular domain can be divided into two regions:

- 1 the *N*-glycosylated link module encoded by exons 2–3, which is maintained in a folded configuration by three Cys-Cys bonds; and

- 2 an *O*-glycosylated mucin-like domain encoded by exons 4–7, which contains sites for covalent linkage of the glycosaminoglycan chondroitin sulphate.

Exon 8 encodes the 21 amino acid membrane-spanning domain and exon 9 the 72 amino acid C-terminal cytoplasmic tail (Figure 21.2) [6–8]. The cytoplasmic domain interacts with the membrane skeleton [3]: CD44 can bind protein 4.1R and ankyrin [9] of the two band 3-macrocomplexes (see Figure 10.2) and 4.1R-deficient red cells lack CD44 [10], yet protein 4.2 deficiency causes elevated CD44 expression [11]. There are an estimated 6000 to 10 000 copies of CD44 per red cell [12].

CD44 was assigned to chromosome 11 by testing a panel of somatic cell hybrids with the original anti-CD44 [13] and localised to 11p13 [14,15].

21.2.2 Indian antigens are located on CD44

Spring *et al.* [3] showed that *In*^a and *In*^b are carried on the CD44 glycoprotein. Immunoblotting of membranes from antigen-positive cells under non-reducing conditions with human anti-*In*^a and -*In*^b revealed an 80 kDa component of identical mobility to CD44; no such component was detected in membranes from *In*(a-b+) and

Table 21.1 Antigens of the Indian system.

Antigen				Molecular basis*		
No.	Name	Frequency	Antithetical antigen	Nucleotides	Exon	Amino acids
IN1	In ^a	Low	IN2	137G>C	2	Arg46Pro
IN2	In ^b	High	IN1	137G (C)	2	Arg46 (Pro)
IN3	INFI	High		255C (G)	2	His85 (Gln)
IN4	INJA	High		488C (G)	2	Thr163 (Lys)

*Molecular basis of antigen-negative phenotype in parentheses.




Figure 21.1 Organisation of *CD44* showing the 20 exons, including the 10 alternatively spliced exons, v1–v10, within the region encoding the extracellular domain (exons 1–17) and the exon encoding the transmembrane domain (TM, exon 18). The lower figure (*CD44H*) shows the exons encoding the haemopoietic form of *CD44*, *CD44s*.

In(a+b–) cells, respectively. *CD44* glycoprotein isolated from human red cell membranes by immunoprecipitation with monoclonal anti-*CD44* reacted with anti-In^b on an immunoblot. Anti-In^b was shown, by radioimmunoassay, to bind *CD44* glycoprotein isolated from human red cells and leucocytes [3,16] and positive results were obtained in MAIEA assays performed with human alloanti-In^b and mouse monoclonal anti-*CD44* [17,18].

21.3 Indian antigens

21.3.1 In^a and In^b (IN1 and IN2)

In 1973, a new antigen present on red cells of about 3% of Indians from Bombay was named In^a by Badakere *et al.* [19,20]. Two years later Giles [21] found that an antibody to a public antigen, Salis, was antithetical to anti-In^a and the antibody became anti-In^b.

The In^b/In^a polymorphism results from Arg46Pro at the N-terminal of an α 1 helix within the link module of *CD44* [22] (Table 21.1, Figure 21.1). This was confirmed

by cDNA transfection and site-directed mutagenesis experiments with Jurkat human leukaemia cell line. Other changes detected in some In(a+b–) individuals, encoding Tyr109Ser and Glu239Gly, do not appear to affect expression of the Indian antigens.

Of 1749 Bombay Indians, 51 (3%) were In(a+) [20]. This gives a frequency for *IN*^{*}A of 0.0147, and the following genotype frequencies can be deduced: *IN*^{*}A/A 0.0002; *IN*^{*}A/B 0.0290; *IN*^{*}B/B 0.9708. A higher frequency of In^a was found in some Arabs: 10.6% of Iranians and 11.8% of Arabs in Bombay were In(a+) [23]. In^a is virtually unknown in other populations. Two of 700 Indian blood donors were In(b–) [24], far in excess of the number expected from the calculated genotype frequencies given above. Of 251 members of the Asian immigrant population of northern England, two were In(a+b–), 8 In(a+b+), and 241 In(a–b+); again an excess of In(a+b–) [25].

21.3.2 IN3 (INFI) and IN4 (INJA)

IN3 and IN4 are two antigens of high frequency, shown to be located on *CD44* by immunoblotting and by MAIEA analysis [26]. Anti-IN3 was found in three pregnant Moroccan women. IN:–3,4 phenotype in the three probands was associated with His85Gln in the β 4 strand of the link module of *CD44* (*IN*^{*}02.–03). Anti-IN4 was found in two pregnant Pakistani women. IN:3,–4 phenotype in the two probands and in the father and two siblings of one of the probands was associated with Thr163Lys in the mucin-like C-terminal extension (*IN*^{*}02.–04) (Table 21.1, Figure 21.1).

IN:–3 and IN:–4 red cells were In(a–b+) but showed weakness of In^b in titration studies [26]. Observation of three-dimensional models of the link module indicates that Arg46 (IN2) and His85 (IN3) may be in close proximity on the folded protein, but that Thr163 (IN4) would




Figure 21.2 Model of CD44s. The extracellular domain consists of two regions: a membrane-proximal region (black) containing one *N*-glycosylation site (*N*), several *O*-glycosylation sites (*O*), and several Ser-Gly chondroitin sulphate linkage sites (SG); and a distal region containing five *N*-glycosylation sites and six cysteine residues (*C*), which suggests the presence of three disulphide bonds (dotted lines), the link module (red), and flanking extended lobes (blue). The position of the amino acid substitution responsible for the *In*^a/*In*^b polymorphism and IN3 and IN4 antigens are shown.

not be [27,28], suggesting that amino acid substitutions responsible for the rare IN:-3 and IN:-4 phenotypes affect the conformation of the link module and reduce binding of anti-*In*^b. Furthermore, binding of a monoclonal antibody to CD44 appeared to block binding of

anti-*In*^b, -IN3, and -IN4, suggesting either that binding of the monoclonal antibody affects the conformation of the whole link module or that *In*^b, IN3, and IN4 are in closer proximity than suggested by the three-dimensional models.

21.3.3 Antigen characteristics

Indian antigens are destroyed by the proteases papain, pronase, trypsin, and chymotrypsin, but are resistant to treatment of red cells with sialidase. They are also destroyed by the disulphide bond reducing agents AET and DTT. Higher concentrations of the reducing agents than generally used for treating red cells may be required to destroy antigen expression.

Red cells from cord samples and from pregnant women show reduced expression of *In*^a, with about 25% of the adult number of *In*^a antigen sites [29]. Red cells of pregnant women have about 38% of the normal adult number, the number of sites returning to normal 3–6 months after delivery. No weakness of *In*^b was detected on cord red cell samples by serological titration [25].

CD44 is present in serum [30,31] and *In*^b can be detected in serum by haemagglutination inhibition.

21.4 Effects of *In*(Lu) on CD44 and Indian antigens

Telen *et al.* [30,32] demonstrated by flow cytometry that a CD44 monoclonal antibody (A3D8) showed markedly reduced levels of binding with red cells of the *In*(Lu) phenotype. *In*(Lu) is associated with low expression of Lutheran antigens and some other red cell antigens, and with mutations in *EKLF* (see Section 6.8). One example of *In*(Lu) cells bound between 25 and 39% of the quantity of anti-CD44 bound by normal cells [32]. Reduced binding was not seen with *Lu*_{null} cells or with *Lu*_{mod} cells of the X-linked type (Section 6.8.3), and the *Lu*_{null} cells may even have had enhanced binding of CD44 antibodies [33]. With some CD44 antibodies the reduced binding to *In*(Lu) cells can also be detected by conventional serological techniques [3,34]. In inhibition experiments, normal human sera reduced binding of anti-CD44 to red cells by 67% whereas serum from an *In*(Lu) individual reduced binding by only 33% [30]. Immunoprecipitation and immunoblotting of membranes from *In*(Lu) cells revealed only a trace of CD44 [2,3,12,34], whereas *Lu*_{null} and X-linked *Lu*_{mod} cells had normal, or even slightly enhanced, CD44 expression [33,35].

Anti-In^b has a reduced titre with In(Lu) cells, compared with cells of normal Lutheran type and with Lu_{null} and X-linked Lu_{mod} cells [3,36], and anti-IN3 and -IN4 reacted weakly or not at all with In(Lu) cells [26]. A band of markedly reduced intensity was seen with membranes from In(Lu) cells blotted with anti-In^b [3].

Red cells of a girl with a novel form of congenital dyserythropoietic anaemia associated with heterozygosity for an *EKL*F mutation (Glu325Lys) had a gross deficiency of red cell CD44 and were In(a-b-): they were also IN:-3,-4 (as determined by immunoblotting [26]), Co(a-b-), AnWj-, and weak for LW^{ab} [37-39] (see Section 15.4 for further information).

21.5 Indian antibodies

Indian antigens appear to be good immunogens. Thirty of 39 In(a-) donors immunised for anti-D production with D+ In(a+) red cells made anti-In^a [40]. One anti-In^b and one anti-IN4 were produced in untransfused women during their first pregnancies [26,36]. Anti-In^a and -In^b often agglutinate antigen-positive red cells directly, although the strength of reaction is generally enhanced by antiglobulin [21,25,40].

Radiolabelled In(a+) red cells were eliminated from the circulation of two individuals with anti-In^a within 20 minutes, suggesting potential for a transfusion reaction [40]. There is one case of anti-In^b causing an immediate HTR after infusion of 50 ml of incompatible blood [24]. Reduced *in vivo* survival of In(b+) cells in a patient with anti-In^b was observed at 24 h [36]. Antigen-negative red cells for transfusion to patients with anti-In^b, -IN3, or -IN4, may be very difficult to obtain, though In(Lu) cells would probably be suitable for transfusion. No Indian antibody has been implicated in HDFN and it has been suggested that binding of anti-In^b to CD44 on fetal monocytes and macrophages could have a blocking effect on FcγR1 [41].

Numerous murine monoclonal antibodies to non-polymorphic epitopes on CD44 have been produced, one of which had In^b specificity [42].

21.6 Functional aspects of CD44

CD44 proteoglycan is a ubiquitous structure: very few tissues or cells lack CD44. It is present on circulating red cells, B and T lymphocytes, granulocytes, and monocytes, but not platelets [3,30,34], and is present on thymus,

central nervous system white matter, epidermis, skeletal muscle, and epithelium from stomach, intestine, liver, bladder, lung, and breast [12,43].

Many roles have been attributed to CD44, the functions being regulated by the inclusion of the products of the various alternatively spliced exons. Functions of CD44 include the following: adhesion of leucocytes to endothelial cells, stromal cells, and the extracellular matrix (ECM); participation in T and B cell activation in response to immunological stimuli; lymphocyte-endothelial cell interactions involved in the localisation of lymphocytes to the site of inflammation; modelling of the ECM during wound healing and embryonic development. CD44 has also been implicated in tumour metastasis (reviews in [44,45]). Most of these interactions involve binding of CD44 to hyaluronan, a high molecular weight glycoaminoglycan that is a major component of the ECM and is also present on cell surfaces. CD44 also binds collagen, fibronectin, and laminin; proteins of the ECM that fill the spaces between cells [1]. CD44 appears to play a regulatory role in normal haemopoiesis (review in [46]). Hyaluronan binding by CD44 may be involved in adhesion of haemopoietic progenitors, including burst-forming units-erythroid (BFU-E), to the bone marrow stroma [46,47]. Surface expression of CD44 in mice progressively decreased 30-fold in late-stage erythroblasts [48]. Consequently, CD44 on circulating red cells may be vestigial, its functions being completed during erythropoiesis.

CD44 contains three copies of a hyaluronan binding motif, BX₇B (B = arginine or lysine; X₇ = seven non-acidic amino acids), present in proteins that bind hyaluronan [49]. Experiments involving site-directed mutagenesis identified several amino acid residues that were considered important in the binding of CD44 to hyaluronan, one of which was Arg46 (the second B of a BX₇B hyaluronan-binding motif). Mutating Arg46 to Gly in CD44 constructs abolished hyaluronan binding [49]. In(a+b-) phenotype results from homozygosity for CD44 alleles encoding Pro46 (Section 21.3.1), yet the In^a Arg-46Pro substitution does not reduce hyaluronan binding to intact CD44s, *in vitro* [22]. A combined modelling and mutagenesis study has subsequently suggested that Arg41, Tyr42, Arg78, and Tyr79 are critical for hyaluronan binding, and Lys68, Asn100, Asn 101, and Tyr105 support binding [50]. All of these residues, except Lys68, come together in a three-dimensional module, as a contiguous linear patch [27]. Glycosylation of Asn25 and Asn120, which are outside the link module, also play a key role in hyaluronan binding.

21.7 AnWj (901009)

Anti-Anton [51] and anti-Wj [52] were names given to alloantibodies and autoantibodies that failed to react with In(Lu) phenotype cells and with cord cells, but did react with Lu_{null} cells. It subsequently became clear that anti-Anton and -Wj had the same specificity [53,54] and the name AnWj was given to the antigen they define.

21.7.1 Inheritance and frequency

The rare AnWj⁻ phenotype is usually acquired and may be transient, but one family demonstrated that it could also be inherited [55]. Two of seven siblings of an AnWj⁻ Arab woman with anti-AnWj were AnWj⁻. The consanguineous parents and the six children of the propositus were AnWj⁺, suggesting that the AnWj⁻ phenotype in this family results from homozygosity of a rare recessive gene. The family study demonstrated that AnWj is not controlled by *LU*, or by *ABO*, *MNS*, *RH*, *KEL*, *FY*, *JK*, *XG*, or *XK*.

Anti-AnWj screening of red cells from 2400 American donors revealed three In(Lu) samples, but no AnWj⁻ red cells with normal Lutheran antigens [56].

21.7.2 AnWj and the In(Lu) phenotype

Like In^b and several other red cell antigens outside the Lutheran system, AnWj is expressed only very weakly on red cells of individuals with the dominant gene *In(Lu)* (see Section 6.8). Usually AnWj cannot be detected on these Lu_{mod} cells by direct testing, but anti-AnWj can be adsorbed and eluted from them [55]. Lu_{null} and X-linked Lu_{mod} cells have normal AnWj expression [51,57].

AnWj differs from Lutheran and Indian system antigens in being resistant to trypsin, chymotrypsin, and the disulphide bond reducing agent AET.

21.7.3 Development of AnWj

Analysis of red cells from 36 infants revealed that the age at which conversion from AnWj⁻ to AnWj⁺ takes place varies from infant to infant, but occurs between the ages of three days and 46 days and requires less than one day to complete [58]. This rapid, 'all or nothing' phenomenon is unexpected as the red cells in the circulation are not all produced at the same time and no evidence could be found for a conversion factor in the serum. Whatever causes the change from AnWj⁻ to AnWj⁺ in infants might be reversible on rare occasions, as the adult AnWj⁻ phenotype with concurrent presence of anti-AnWj has

been found to be transient in several patients and in a healthy individual [54,58–60].

21.7.4 AnWj as a receptor for *Haemophilus influenzae*

H. influenzae is a commensal bacterium of the throat of most healthy people, but it may also cause respiratory tract infections and, more seriously, is a major cause of bacterial meningitis in young children. Some strains of *H. influenzae* express fimbriae (short, thread-like processes attached to the cell walls), which are probably involved in adherence to nasopharyngeal epithelial cells.

Fimbriae-bearing strains of *H. influenzae* isolated from patients with invasive disease and respiratory tract infections agglutinated most red cell samples from adults, including Lu_{null} cells, but did not agglutinate cord cells, In(Lu) cells, or AnWj⁻ red cells of both acquired and inherited types [61,62]. Anti-AnWj inhibited agglutination of AnWj⁺ red cells by the bacteria.

H. influenzae bound to buccal epithelial cells, including those from one individual with transient AnWj⁻ red cells, two with *In(Lu)* genes, and several neonates, but did not bind to buccal epithelial cells from three AnWj⁻ members of the Arab family with inherited AnWj⁻ phenotype [62,63]. Adherence of *H. influenzae* to epithelial cells was not inhibited by anti-AnWj [63]. Although the receptors for *H. influenzae* adherence on red cells and epithelial cells may not be identical, they appear to have a common genetic basis.

21.7.5 Anti-AnWj

Anti-AnWj may be autoantibodies or alloantibodies, or apparent alloantibodies in patients, usually with lymphoid tumours, with an acquired AnWj⁻ phenotype. In the only family with more than one AnWj⁻ member, the propositus and her AnWj⁻ sister both had anti-AnWj and both had been pregnant, but not transfused [55]. It is a little surprising that anti-AnWj can be stimulated by pregnancy considering the very low level of AnWj antigen on neonatal red cells.

Anti-AnWj has been incriminated in severe HTRs [60,64–68], although there are several cases where AnWj⁺ cells have been transfused uneventfully to patients with anti-AnWj. The haemolytic potential of anti-AnWj has been supported by *in vivo* red cell survival studies [64,69] and monocyte monolayer analysis [67,70]. In a patient with autoanti-AnWj and depressed red cell AnWj expression, radiolabelled autologous red cells survived normally, but AnWj⁺ allogeneic cells had reduced survival

[71]. If available, In(Lu) cells are most suitable for transfusing patients with anti-AnWj.

There was no indication of HDFN in any of the children of a mother with alloanti-AnWj [55].

Two monoclonal antibodies with AnWj specificity were produced from mice immunised with human T-cell lines derived from patients with acute lymphocytic leukaemia [72].

21.7.6 AnWj may be located on CD44 glycoprotein

In(a-b-) red cells of the patient with a novel form of CDA and a gross deficiency of red cell CD44 were AnWj- [37–39]. Positive results were obtained in MAIEA analyses performed with either anti-CD44 or -AnWj murine monoclonal antibodies and anti-In^b or -AnWj human antibodies [17]. The CD44-negative human leukaemia cell line, Jurkat, transfected with *CD44* cDNA reacted with human and mouse monoclonal anti-AnWj [73]. Immunoblotting with anti-AnWj revealed an 80kDa component in normal red cells and in Chinese hamster ovary cells (CHO) and murine erythroleukaemia (MEL) cells transfected with *CD44* cDNA, but not in non-transfected CHO or MEL cells, or in In(Lu) red cells. The anti-AnWj also bound a 200kDa structure in red cells and transfected CHO cells. Telen *et al.* [73] suggest that AnWj antigen is located on a trypsin-resistant region of an isoform of CD44 that is not present on red cells of the newborn. AnWj might also be on the 200kDa chondroitinated isoform of CD44.

References

- Day AJ, Prestwich GD. Hyaluronan-binding proteins: tying up the giant. *J Biol Chem* 2002;277:4585–4588.
- Telen MJ, Palker TJ, Haynes BF. Human erythrocyte antigens: II. The *In(Lu)* gene regulates expression of an antigen on an 80-Kilodalton protein of human erythrocytes. *Blood* 1984; 64:599–606.
- Spring FA, Dalchau R, Daniels GL, *et al.* The In^a and In^b blood group antigens are located on a glycoprotein of 80 000 MW (the CDw44 glycoprotein) whose expression is influenced by the *In(Lu)* gene. *Immunology* 1988;64:37–43.
- Screaton GR, Bell MV, Jackson DG, *et al.* Genomic structure of DNA encoding the lymphocyte homing receptor CD44 reveals at least 12 alternatively spliced exons. *Proc Natl Acad Sci USA* 1992;89:12160–12164.
- Tölg C, Hofmann M, Herrlich P, Ponta H. Splicing choice from ten variant exons establishes CD44 variability. *Nucleic Acid Res* 1993;21:1225–1229.
- Stamenkovic I, Amiot M, Pesando JM, Seed B. A lymphocyte molecule implicated in lymph node homing is a member of the cartilage link protein family. *Cell* 1989;56:1057–1062.
- Goldstein LA, Zhou DFH, Picker LJ, *et al.* A human lymphocyte homing receptor, the Hermes antigen, is related to cartilage proteoglycan core and link proteins. *Cell* 1989;56: 1063–1072.
- Harn H-J, Isola N, Cooper DL. The multispecific cell adhesion molecule CD44 is represented in reticulocyte cDNA. *Biochem Biophys Res Comm* 1991;178:1127–1134.
- Nunomura W, Takakuwa Y, Tokimitsu R, *et al.* Regulation of CD44-protein 4.1 interaction by Ca²⁺ and calmodulin. Implications for modulation of CD44-ankyrin interaction. *J Biol Chem* 1997;272:30322–30328.
- Jeremy KP, Plummer ZE, Head DJ, *et al.* 4.1R-deficient human red blood cells have altered phosphatidylserine exposure pathways and are deficient in CD44 and CD47 glycoproteins. *Haematologica* 2009;94:1354–1361.
- van den Akker E, Satchwell TJ, Pellegrin S, *et al.* Investigating the key membrane protein changes during in vitro erythropoiesis of protein 4.2 null cells (protein 4.2 Chartres 1 and 2). *Haematologica* 2010;95:1278–1286.
- Anstee DJ, Gardner B, Spring FA, *et al.* New monoclonal antibodies in CD44 and CD58: their use to quantify CD44 and CD58 on normal human erythrocytes and to compare the distribution of CD44 and CD58 in human tissues. *Immunology* 1991;74:197–205.
- Goodfellow PN, Banting G, Wiles MV, *et al.* The gene, MIC4, which controls expression of the antigen defined by monoclonal antibody F10.44.2, is on human chromosome 11. *Eur J Immunol* 1982;12:659–663.
- Couillin P, Azoulay M, Henry I, *et al.* Characterization of a panel of somatic cell hybrids for subregional mapping along 11p and within band 11p13. *Hum Genet* 1989;82:171–178.
- Couillin P, Azoulay M, Metezeau P, Grisard M-C, Junien C. The gene for catalase is assigned between the antigen loci MIC4 and MIC11. *Genomics* 1989;4:7–11.
- Telen MJ, Ferguson DJ. Relationship of In^b antigen to other antigens on *In(Lu)*-related p80. *Vox Sang* 1990;58:118–121.
- Rao N, Udani M, Telen MJ. Demonstration by monoclonal antibody immobilization of erythrocyte antigens and dot blot that both the In and AnWj blood group antigens reside on CD44. *Transfusion* 1994;34(Suppl.):25S [Abstract].
- Petty AC, Green CA, Daniels GL. The monoclonal antibody-specific immobilization of erythrocyte antigens assay (MAIEA) in the investigation of human red-cell antigens and their associated membrane proteins. *Transfus Med* 1997;7:179–188.
- Badakere SS, Joshi SR, Bhatia HM, *et al.* Evidence for a new blood group antigen in the Indian population (a preliminary report). *Ind J Med Res* 1973;61:563.
- Badakere SS, Parab BB, Bhatia HM. Further observations on the In^a (Indian) antigen in Indian populations. *Vox Sang* 1974;26:400–403.

- 21 Giles CM. Antithetical relationship of anti-In^a with the Salis antibody. *Vox Sang* 1975;29:73–76.
- 22 Telen MJ, Udani M, Washington MK, *et al.* A blood group-related polymorphism of CD44 abolishes a hyaluronan-binding consensus sequence without preventing hyaluronan binding. *J Biol Chem* 1996;271:7147–7153.
- 23 Badakere SS, Vasantha K, Bhatia HM, *et al.* High frequency of In^a antigen among Iranians and Arabs. *Hum Hered* 1980;30:262–263.
- 24 Joshi SR. Immediate haemolytic transfusion reaction due to anti-In^b. *Vox Sang* 1992;63:232–233.
- 25 Longster GH, Robinson EAE. Four further examples of anti-In^b detected during pregnancy. *Clin Lab Haematol* 1981;3:351–356.
- 26 Poole J, Tilley L, Warke N, *et al.* Two missense mutations in the CD44 gene encode two new antigens of the Indian blood group system. *Transfusion* 2007;47:1306–1311. Correction: *Transfusion* 2007;47:1741.
- 27 Teriete P, Banerji S, Noble M, *et al.* Structure of the regulatory hyaluronan binding domain in the inflammatory leukocyte homing receptor CD44. *Mol Cell* 2004;13:483–496.
- 28 Takeda M, Ogino S, Umemoto R. Ligand-induced structural changes of the CD44 hyaluronan-binding domain revealed by NMR. *J Biol Chem* 2006;281:40089–40095.
- 29 Dumasia AN, Gupta SC. Quantitation of In^a blood group antigens. *Ind J Med Res B* 1990;92:50–53.
- 30 Telen MJ, Eisenbarth GS, Haynes BF. Human erythrocyte antigens. Regulation of expression of a novel erythrocyte surface antigen by the inhibitor Lutheran In(Lu) gene. *J Clin Invest* 1983;71:1878–1886.
- 31 Lucas MG, Green AM, Telen MJ. Characterization of the serum In(Lu)-related antigen: identification of a serum protein related to erythrocyte p80. *Blood* 1989;73:596–600.
- 32 Telen MJ, Rogers I, Letarte M. Further characterization of erythrocyte p80 and the membrane protein defect of In(Lu) Lu(a–b–) erythrocytes. *Blood* 1987;70:1475–1481.
- 33 Telen MJ, Green AM. Human red cell antigens. V. Expression of In(Lu)-related p80 antigens by recessive-type Lu(a–b–) red cells. *Transfusion* 1988;28:430–434.
- 34 Telen MJ, Shehata H, Haynes BF. Human medullary thymocyte p80 antigen and In(Lu)-related p80 antigen reside on the same protein. *Hum Immunol* 1986;17:311–324.
- 35 Judson PA, Spring FA, Parsons SF, Anstee DJ, Mallinson G. Report on group 8 (Lutheran) antibodies. *Rev Franc Transfus Immuno-Hémat* 1988;31:433–440.
- 36 Ferguson DJ, Gaal HD. Some observations on the In^b antigen and evidence that anti-In^b causes accelerated destruction of radiolabeled red cells. *Transfusion* 1988;28:479–482.
- 37 Parsons SF, Jones J, Anstee DJ, *et al.* A novel form of congenital dyserythropoietic anemia associated with deficiency of erythroid CD44 and a unique blood group phenotype [In(a–b–) Co(a–b–)]. *Blood* 1994;83:860–868.
- 38 Singleton BK, Fairweather VSS, Lau W, *et al.* A novel EKLK mutation in a patient with dyserythropoietic anemia: the first association of EKLK with disease in man. *Blood* 2009;114:162 [Abstract].
- 39 Arnaud L, Saison C, Helias V, *et al.* A dominant mutation in the gene encoding the erythroid transcription factor KLF1 causes a congenital dyserythropoietic anemia. *Am J Hum Genet* 2010;87:721–727.
- 40 Bhatia HM, Badakere SS, Mokashi SA, Parab BB. Studies on the blood group antigen In^a. *Immunol Commun* 1980;9:203–215.
- 41 Garner SF, Devenish A. Do monocyte ADCC assays accurately predict the severity of hemolytic disease of the newborn caused by antibodies to high-frequency antigens? *Immunohematology* 1996;12:20–26.
- 42 Stoll M, Dalchau R, Schmidt RE. Cluster report: CD44. In: Knapp W, Dörken B, Gilks WR, *et al.*, eds. *Leukocyte Typing IV*. Oxford: Oxford University Press, 1989:619–622.
- 43 Flanagan BF, Dalchau R, Allen AK, Daar AS, Fabre JW. Chemical composition and tissue distribution of the human CDw44 glycoprotein. *Immunology* 1989;67:167–175.
- 44 Borland G, Ross JA, Guy K. Forms and functions of CD44. *Immunology* 1998;93:139–148.
- 45 Bajorath J. Molecular organization, structural features, and ligand binding characteristics of CD44, a highly variable cell surface glycoprotein with multiple functions. *Proteins* 2000;39:103–111.
- 46 Chan JY, Watt SM. Adhesion receptors on haematopoietic progenitor cells. *Br J Haematol* 2001;112:541–557.
- 47 Oostendorp RAJ, Spitzer E, Brandl M, Eaves CJ, Dörmer P. Evidence for differences in the mechanisms by which antibodies against CD44 promote adhesion of erythroid and granulopoietic progenitors to marrow stromal cells. *Br J Haematol* 1998;101:436–445.
- 48 Chen K, Liu J, Heck S, *et al.* Resolving the distinct stages in erythroid differentiation based on dynamic changes in membrane protein expression during erythropoiesis. *Proc Natl Acad Sci USA* 2009;106:17413–17418.
- 49 Yang B, Yang BL, Savani RC, Turley EA. Identification of a common hyaluronan binding motif in the hyaluronan binding proteins RHAMM, CD44 and link protein. *EMBO J* 1994;13:286–296.
- 50 Bajorath J, Greenfield B, Munro SB, Day AJ, Aruffo A. Identification of CD44 residues important for hyaluronan binding and delineation of the binding site. *J Biol Chem* 1998;273:338–343.
- 51 Poole J, Giles CM. Observations on the Anton antigen and antibody. *Vox Sang* 1982;43:220–222.
- 52 Marsh WL, Brown PJ, DiNapoli J, *et al.* Anti-Wj: an autoantibody that defines a high-incidence antigen modified by the In(Lu) gene. *Transfusion* 1983;23:128–130.
- 53 Poole J, Giles C. Anton and Wj, are they related? *Transfusion* 1985;25:443.
- 54 Mannessier L, Rouger P, Johnson CL, Mueller KA, Marsh WL. Acquired loss of red-cell Wj antigen in a patient with Hodgkin's disease. *Vox Sang* 1986;50:240–244.

- 55 Poole J, Levene C, Bennett M, *et al.* A family showing inheritance of the Anton blood group antigen AnWj and independence of AnWj from Lutheran. *Transfus Med* 1991;1: 245–251.
- 56 Lukasavage T. Donor screening with anti-AnWj. *Immunohematology* 1993;9:112.
- 57 Norman PC, Tippet P, Beal RW. An Lu(a–b–) phenotype caused by an X-linked recessive gene. *Vox Sang* 1986;51: 49–52.
- 58 Poole J, Van Alphen L. *Haemophilus influenzae* receptor and the AnWj antigen. *Transfusion* 1988;28:289.
- 59 Harris T, Steiert S, Marsh WL, Berman LB. A Wj-negative patient with anti-Wj. *Transfusion* 1986;26:117.
- 60 Magrin G, Harrison C. One hour ⁵¹Cr survival in a patient with anti-AnWj. *20th Congr Int Soc Blood Transfus*, 1988:228 [Abstracts].
- 61 van Alphen L, Poole J, Overbeeke M. The Anton blood group antigen is the erythrocyte receptor for *Haemophilus influenzae*. *FEMS Microbiol Letts* 1986;37:69–71.
- 62 van Alphen L, Levene C, Geelan-van den Broek L, *et al.* Combined inheritance of epithelial and erythrocyte receptors for *Haemophilus influenzae*. *Infect Immun* 1990;58: 3807–3809.
- 63 van Alphen L, Poole J, Geelen L, Zanen HC. The erythrocyte and epithelial cell receptors for *Haemophilus influenzae* are expressed independently. *Infect Immun* 1987;55:2355–2358.
- 64 de Man AJM, van Dijk BA, Daniels GL. An example of anti-AnWj causing haemolytic transfusion reaction. *Vox Sang* 1992;63:238.
- 65 Bishop D, Powell T, Banks J, Massey E, Copplestone A. An example of haemolytic transfusion reaction caused by anti-AnWj. *Transfus Med* 2005;15(Suppl. 1):49 [Abstract].
- 66 Long S, Moulds J, Moulds J, Poole J. Transfusion reaction due to anti-AnWj. *Transfusion* 2008;48(Suppl.):181A [Abstract].
- 67 Grigoriadis G, Condon J, Green K, *et al.* Persistent complement-dependent anti-AnWj in a lymphoproliferative disorder: a case study and review. *Immunohematology* 2011;27:83–88.
- 68 Xu Z, Duffett L, Tokessy M, *et al.* Anti-AnWj causing acute hemolytic transfusion reaction in a patient with aplastic anemia. *Transfusion* 2012;52:1476–1481.
- 69 van Gaalen FA, Zanin DEA, Brand A. Erythrocyte survival tests in cases of anti-AnWj antibodies. *Vox Sang* 2009;97: 275–276.
- 70 Stowers RE, Richa EM, Stubbs JR, Moore SB. Red blood cell transfusion in a patient with anti-AnWj: a case report. *Immunohematology* 2007;23:55–58.
- 71 Whitsett CF, Hare VW, Oxendine SM, Pierce JA. Autologous and allogeneic red cell survival studies in the presence of autoanti-AnWj. *Transfusion* 1993;33:845–847.
- 72 Knowles RW, Bai Y, Lomas C, Green C, Tippet P. Two monoclonal antibodies detecting high frequency antigens absent from red cells of the dominant type of Lu(a–b–) Lu:–3. *J Immunogenet* 1982;9:353–357.
- 73 Telen MJ, Rao N, Udani M, Liao H-X, Haynes BF. Relationship of the AnWj blood group antigen to expression of CD44. *Transfusion* 1993;33(Suppl.):48S [Abstract].