

# 7 Kell and Kx Blood Group Systems

- |  |  |
|--|--|
| <ul style="list-style-type: none"><li>7.1 Introduction, 278</li><li>7.2 The Kell glycoprotein and the gene that encodes it, 280</li><li>7.3 K and k (KEL1 and KEL2), 281</li><li>7.4 Kp<sup>a</sup>, Kp<sup>b</sup>, and Kp<sup>c</sup> (KEL3, KEL4, and KEL21), 285</li><li>7.5 Js<sup>a</sup> and Js<sup>b</sup> (KEL6 and KEL7), 286</li><li>7.6 Other Kell-system antigens, 287</li><li>7.7 The Kell-null and Kell-mod phenotypes and anti-Ku (-KEL5), 290</li></ul> | <ul style="list-style-type: none"><li>7.8 Kell depression in Gerbich-negative phenotypes, 291</li><li>7.9 Acquired and transient depressed Kell phenotypes, 292</li><li>7.10 Effects of enzymes and reducing agents on Kell antigens, 292</li><li>7.11 Kell antigens on other cells and in other species, 293</li><li>7.12 Functional aspects, 293</li><li>7.13 The Kx blood group system, 293</li></ul> |
|--|--|

## 7.1 Introduction

Kell was the first of many blood group systems disclosed by the antiglobulin test [1]. When Allen *et al.* [2] described the fourth Kell system antigen, Kp<sup>b</sup>, they concluded prophetically, 'There is, probably, much still to be learned about the Kell blood group system'. There are now 35 antigens in the Kell system (Table 7.1) and Kell is closely related to the Kx system.

There are seven sets of antigens in the Kell system with allelic relationships: K and k; Kp<sup>a</sup>, Kp<sup>b</sup>, and Kp<sup>c</sup>; Js<sup>a</sup> and Js<sup>b</sup>; K11 and K17 (Wk<sup>a</sup>); KEL14 and KEL24; KEL25 and KEL28; KEL31 and KEL38. There are an additional 17 high frequency antigens and three low frequency antigens. All have been shown to be associated with nucleotide changes in *KEL*, except KEL13, which has been shown to be on the Kell-glycoprotein, and Km, which requires the presence of Xk (see below). Recombination as a result of crossing-over has never been observed within *KEL*.

None of the Kell antigens are expressed on cells of the Kell-null phenotype, K<sub>o</sub>, which arises from homozygosity for *KEL* inactivating mutations. Ku antigen is present on all cells save those of the K<sub>o</sub> phenotype. In K<sub>mod</sub>, which also arises from *KEL* mutations, all antigens of the Kell system are expressed weakly (Section 7.7).

Several rare phenotypes occur in which all or most of the high frequency Kell antigens are expressed only weakly. Some are due to epistasis, such as the McLeod phenotype and depressed Kell associated with some Gerbich negative phenotypes, and some arise from interactions within the *KEL* gene. In patients with Kell-related autoantibodies, the depressed Kell phenotype may be acquired and transient (Section 7.9).

The Kell antigens are located on CD238, a red cell transmembrane glycoprotein of apparent MW 93 kDa, a metalloendopeptidase that processes endothelin-3 (Section 7.2).

The *KEL* gene is situated on chromosome 7q33 and consists of 19 exons.

McLeod syndrome is a form of neuroacanthocytosis, which includes an abnormal Kell red cell phenotype. McLeod phenotype red cells have depressed Kell antigens and lack the high frequency antigen Kx. The inheritance of Kx is controlled by an X-borne gene, XK, and represents a blood group system (the Kx system) independent of Kell. The Xk protein and Kell glycoprotein are linked by a disulphide bond. The Kx system is described in this chapter (Section 7.13) because of its phenotypic and biochemical associations with Kell.

The numerical antigen notation is generally used in this chapter, except for those antigens more commonly known by their traditional symbols: K, k, Kp<sup>a</sup>, Kp<sup>b</sup>, Kp<sup>c</sup>, Ku, Js<sup>a</sup>, Js<sup>b</sup>, U1<sup>a</sup>, and Km.

**Table 7.1** Antigens of the Kell system.

Antigen				Molecular basis*		
No.	Name	Frequency	Antithetical	Nucleotides†	Exon	Amino acids
KEL1	K	Polymorphic	k	578C>T	6	Thr193Met
KEL2	k	High	K	578C (T)	6	Thr193 (Met)
KEL3	Kp <sup>a</sup>	Polymorphic	Kp <sup>b</sup> (Kp <sup>c</sup> )	841C>T	8	Arg281Trp
KEL4	Kp <sup>b</sup>	High	Kp <sup>a</sup>	1. 841C (T)	8	Arg281 (Trp)
			Kp <sup>c</sup>	2. 842G (A)	8	Arg281 (Gln)
KEL5	Ku	High		Various		Various
KEL6	Js <sup>a</sup>	Polymorphic	Js <sup>b</sup>	1790T>C	17	Leu597Pro
KEL7	Js <sup>b</sup>	High	Js <sup>a</sup>	1790T(C)	17	Leu597 (Pro)
KEL10	U1 <sup>a</sup>	Low		1481A>T	13	Glu494Val
KEL11	K11 (Côté)	High	KEL17	905T (C)	8	Val302 (Ala)
KEL12	K12 (Boc)	High		1523A (G)	15	His548 (Arg)
KEL13	K13	High		986T (C)	9	Leu329 (Pro)
KEL14	K14 (San)	High	KEL24	1. 539G (C)	6	Arg180 (Pro)
				2. 538C (T)	6	Arg180 (Cys)
				3. 539G (A)	6	Arg180 (His)
KEL16	'k-like'	High		Not known		
KEL17	K17 (Wk <sup>a</sup> )	Low	KEL11	905T>C	8	Val302Ala
KEL18	K18	High		1. 388C (T)	4	Arg130 (Trp)
				2. 389G (A)		Arg130 (Gln)
KEL19	K19 (Sub)	High		1475G (A)	13	Arg492 (Gln)
KEL20	Km	High				Absence of Xk
KEL21	Kp <sup>c</sup>	Low	Kp <sup>b</sup> (Kp <sup>a</sup> )	842G>A	8	Arg281Gln
KEL22	K22	High		965C (T)	9	Ala322 (Val)
KEL23	K23	Low		1145A>G	10	Gln382Arg
KEL24	K24 (Cls)	Low	KEL14	539G>C	6	Arg180Pro
KEL25	VLAN	Low	KEL28	743G>A	8	Arg248Gln
KEL26	TOU	High		1217G (A)	11	Arg406 (Gln)
KEL27	RAZ	High		745G (A)	8	Glu249 (Lys)
KEL28	VONG	Low	KEL25	742C>T	8	Arg248Trp
KEL29	KALT	High		1868G (A)	17	Arg623 (Lys)
KEL30	KTIM	High		913G (A)	8	Asp305 (Asn)
KEL31	KYO	Low	KEL38	875G>A	8	Arg292Gln
KEL32	KUCI	High		1271C (T)	11	Ala424 (Val)
KEL33	KANT	High		1283G (T)	11	Arg428 (Leu)
KEL34	KASH	High		758A (G)	8	Tyr253 (Cys)
KEL35	KELP	High		708G (T), 2024G (A)	8,18	Leu260 (Phe), Arg675 (Gln)
KEL36	KETI	High		1391C (T)	12	Thr464 (Ile)
KEL37	KUHL	High		877C>T	8	Arg293 (Trp)
KEL38	KYOR	High	KEL31	875G (A)	8	Arg292 (Gln)

\*Molecular basis of antigen-negative phenotype in parentheses.

†1 is the first nucleotide of the translation-initiating codon, which is 120 bp downstream of the traditional position for the first nucleotide in early reports.

Obsolete: KEL8, previously Kw; KEL9, previously KL; KEL15, previously Kx (now XK1).


## 7.2 The Kell glycoprotein and the gene that encodes it

### 7.2.1 The Kell glycoprotein

A glycoprotein of apparent MW 93 kDa was isolated from detergent-solubilised red cell membranes in the 1980s by immunoprecipitation with antibodies to Kell-system antigens: anti-K, -k, -Js<sup>b</sup>, or -KEL22 [3–5]. Treatment of the Kell glycoprotein with an *N*-glycanase reduced the apparent MW by about 15 kDa, whereas O-glycanase had little effect. The Kell glycoprotein is phosphorylated, but not palmitoylated [6,7]. Kell antibodies do not generally react with isolated Kell glycoprotein by immunoblotting, though mouse monoclonal and rabbit antibodies, produced by immunising animals with purified Kell glycoprotein, detect the 93 kDa Kell glycoprotein on immunoblots [3,8]. No Kell glycoprotein was detected on blots of K<sub>0</sub> cells or isolated from K<sub>0</sub> cells by immunoprecipitation with a variety of polyclonal and monoclonal Kell antibodies.

Based on the amino acid sequence of a tryptic peptide, primers were synthesised and a specific oligonucleotide probe prepared by the polymerase chain reaction. Lee *et al.* [9] used this probe to screen a human bone marrow cDNA library and a clone was isolated with an open reading frame encoding a 732 amino acid polypeptide. Rabbit antibody prepared to a synthetic 30 amino acid peptide derived from the cDNA sequence bound to Kell glycoprotein on an immunoblot.

Hydropathy analysis indicated a type II membrane protein with a single hydrophobic membrane-spanning region, a highly hydrophilic N-terminal cytoplasmic domain of 47 amino acids (or 28 amino acids if the codon for Met 20 is used for translation initiation), and a large, 665-amino acid, C-terminal extracellular domain (Figure 7.1). The N-terminal methionine residue is probably cleaved from the mature protein. The extracellular domain has six Asn-Xxx-Ser/Thr putative *N*-glycosylation sites (positions 94, 115, 191, 345, 627, and 724), though Asn724 is unlikely to be glycosylated as residue 725 is proline, which usually inhibits glycosylation. There are 15 extracellular cysteine residues, suggesting the presence of seven intramolecular disulphide bonds, resulting in extensive folding of the molecule. The Kell protein has structural and sequence homology with a family of zinc-binding endopeptidases (for functional aspects see Section 7.12) and has been modelled, based on the crystal structure of the external domains of neutral endopeptidase 24.11 (NEP) and endothelin-converting enzyme 1 (ECE-1) [10,11]. The Kell-glycoprotein has two globular



**Figure 7.1** Homology model for Kell ectodomain, with the highly conserved peptidase domain in green and the variable distal membrane domain in blue. The linker region is the site of attachment to the transmembrane domain. Positions associated with Kell antigens are labelled. Modified from [11].

extracellular domains, consisting mostly of  $\alpha$ -helical segments. The domain closest to the membrane contains the N- and C-terminal sequences and the enzyme-active site; the outer domain contains almost all of the amino acid sites responsible for Kell-system alloantigenicity [10].

Kell glycoprotein is closely associated in the membrane with the Xk protein and a 120 kDa heterodimer can be isolated by immunoprecipitation under non-reducing conditions [12]. The two proteins are linked by disulphide bonding between Cys72 of Kell and Cys347 of Xk [13] (Section 7.13.2). The Kell-Xk heterodimer is part of the 'junctional' or 4.1R red cell membrane complex that contains band 3, Rh proteins, and glycophorin C, and is linked to the spectrin-actin junction of the cytoskeleton through protein 4.1R and p55 [14] (see Section 10.7 and Figure 10.2).

With Fab fragments of three monoclonal antibodies directed at epitopes on the Kell glycoprotein, figures of 4000–8000 sites per red cell were obtained, but Fab fragments from a fourth antibody gave a figure of 18 000 sites per cell [15]. The number of K antigen sites per red cell has been estimated as 4000–6200 on K+ k- cells and 2500–3500 on K+ k+ cells by use of radioiodinated polyclonal and monoclonal anti-K [16,17].

### 7.2.2 Organisation of the *KEL* gene

*KEL* spans about 21.5 kb organised into 19 exons of coding sequence [18] (Table 7.2). Exon 1 encodes a possible translation initiating methionine residue and Sp1 and GATA-1 binding sites. The exon 1 region is involved

**Table 7.2** Exon/intron organisation of *KEL*.

Exon	Codons	3' intron size kb	Comments
1	5'UT Met1	0.34	
2	2–27	0.29	Cytoplasmic
3	28–74	0.26	Transmembrane
4	75–133	~2.6	KEL18
5	134–175	0.33	
6	176–224	~3.2	KEL14/KEL24, K/k
7	225–245	0.093	
8	246–308	0.23	KEL25/KEL28, KEL27, KEL34, KEL35, Kp <sup>a</sup> /Kp <sup>b</sup> /Kp <sup>c</sup> , KEL31, KUHL, KEL11/KEL17, KEL30
9	309–358	~1.3	KEL22, KEL13
10	359–401	~6	KEL23
11	402–438	~1.6	KEL26, KEL32, KEL33
12	439–471	0.24	KEL36
13	472–497	0.44	KEL19, UI <sup>a</sup>
14	498–531	0.19	
15	532–568	0.15	KEL12
16	569–590	0.23	HELLH
17	591–647	0.35	Js <sup>a</sup> /Js <sup>b</sup> , KEL29
18	648–679	~1.3	KEL35
19	680–732 3'UT		

UT, untranslated; HELLH, consensus sequence for zinc neutral endopeptidases.

in negative regulation of the promoter in non-erythroid tissue [19]. Exon 2 encodes the cytoplasmic domain and a second possible translation initiation site at Met20, exon 3 the membrane-spanning domain, and exons 4–19 the large extracellular domain. The 5' flanking region to nucleotide –176 contains two GATA-1 binding sites and a CACCC box [18].

### 7.2.3 Linkage and chromosome location

*KEL* was located on 7q33 through linkage to *PIP*, the gene for prolactin-inducible protein [20], indirect linkage to the cystic fibrosis gene (*CFTR*) [21], and *in situ* hybridisation with cDNA encoding the Kell protein [22,23]. An analysis of 31 families informative for segregation of Yt blood group gene and *KEL* revealed loose linkage between these loci, with maximum likelihood of a recombination fraction of 0.26 [24].

## 7.3 K and k (KEL1 and KEL2)

In 1946, in the first report on the applications of the direct antiglobulin test, Coombs, Mourant, and Race [1]

described an antibody of new specificity. This antibody, originally called anti-Kell and subsequently anti-K or anti-KEL1, reacted with the red cells of the husband and two children of the antibody producer and with about 7% of random blood samples [25].

Three years later, Levine *et al.* [26] described anti-Cellano, an antibody antithetical to anti-K. As *k* had already been used to represent the common allele of *K*, the symbol *k* was subsequently adopted for the product of that gene, despite *K* and *k* being products of co-dominant alleles.

Kell antigens are well developed at birth. *K* was found in fetuses of 10–11 weeks gestation and *k* at 6–7 weeks [27].

### 7.3.1 The molecular basis of the K/k polymorphism

The *k/K* polymorphism results from a C578T transition within exon 6 of the *KEL* gene, which gives rise to an amino acid substitution in the Kell glycoprotein: Met193 in *K* and Thr193 in *k* [28,29]. A *BsmI* (*BsaMI*) restriction site is present in *KEL*\*01 (*K*), but not *KEL*\*02 (*k*) [28] (Table 7.3). In the *KEL*\*02 product, Asn-Arg-Thr193 is a

**Table 7.3** Sequences from nucleotides 571–582 of *KEL*, encoding amino acids 191–194, in four alleles encoding k, K, and two variants.

Allele		Antigen			
<i>k</i>	<i>KEL*02</i>	AAC	CGA	<b>578</b> ACG	CTG
		Asn <sup>N</sup>	Arg	Thr	Leu
<i>K</i>	<i>KEL*02</i>	AAC	<u>CGA</u>	<u>ATG</u>	<u>CTG</u>
		Asn	Arg	Met	Leu
	<i>KEL*01.2</i> [30]	AAC	CGA	TCG	CTG
		Asn <sup>N*</sup>	Arg	Ser	Leu
	<i>KEL*01M.01</i> [31]	AAC	CGA	AGG	CTG
		Asn*	Arg	Arg	Leu
				<b>193</b>	

N, *N*-glycosylation. N\*, probable *N*-glycosylation. \*, probable no *N*-glycosylation. Underlining in *K* allele sequence shows *BsmI* restriction site.

consensus sequence for *N*-glycosylation of Asn191, whereas Asn-Arg-Met in the product of *KEL\*01* is not. Immunoblotting revealed that the K and k proteins are of apparent MW 110 and 115 kDa, respectively, supporting the suggestion that Asn191 is glycosylated in the latter, but not in the former [32,33]. Enzymatic deglycosylation of the Kell glycoprotein on red cells did not affect binding of either anti-K or -k to cells of the appropriate phenotype, demonstrating that k expression is not dependent on *N*-glycosylation of Asn191 [33].

7.3.2 Frequencies of K and k

In tests on nearly 10 000 English blood donors (mostly white), 9.02% were K+ [34]. From this figure the following gene and genotype frequencies have been calculated: *K* 0.0462; *k* 0.9538; *K/K* 0.0021; *K/k* 0.0881; *k/k* 0.9097 (assuming *k* is the only allele of *K*). *K* is much less common in Africans and extremely rare in eastern Asia and in Native Americans [35] (Table 7.4). *K* achieves its highest level among people of the Arabian and Sinai peninsulas, where up to 25% may be K+.

The k antigen has a high incidence in all populations. From the gene frequencies given above it can be estimated that the incidence of K+ k– would be 1 in 476. The incidence of k– was found to be one in 549 London blood donors [50].

Table 7.5 shows genotype frequencies in Austrian blood donors obtained by genotyping with allele-specific primers [51].

7.3.3 K/k genotyping

A variety of methods, involving SNP testing in exon 6 of *KEL*, are available for predicting K/k phenotypes for DNA. All these tests may give a false prediction if a Kell null or mod allele or the *K* variant allele *KEL\*01.2* is present (Sections 7.3.4 and 7.7).

Anti-K is a relatively common cause of severe HDFN (Section 7.3.5.2), so in pregnant women with anti-K it is advantageous to predict fetal K phenotype. This can be done from fetal DNA in maternal plasma, thus avoiding invasive procedures such as amniocentesis or chorionic villus sampling (see Section 5.7.1). Finning *et al.* [52] achieved 98.6% accuracy by real-time quantitative PCR, involving the application of locked nucleic acids to prevent mispriming of the *KEL\*01*-specific primer on the *KEL\*02* allele. This technology is now employed in England to provide a routine service to pregnant women with anti-K, with a level of accuracy of around 99.6% [53]. A method for fetal *K* detection incorporating matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) gave 94% accuracy [54].

K genotyping has also been applied to preimplantation genetic diagnosis (PGD). In K– women with a K+ k+ partner, half of the embryos derived from *in vitro* fertilisation (IVF) will have a paternal *KEL\*01* allele and half will not. K genotyping of DNA from individual blastomeres obtained by IVF was used for PGD to ensure implantation of a K– embryo in two women with anti-K,

**Table 7.4** Frequency of some Kell system antigens and deduced gene frequencies.

Antigen	Population	No. tested	Positive (%)	Gene frequency	References
K	English	9875	9.02	0.0462	[34]
	Parisians	81 962	8.55	0.0437	[36]
	Finns	5000	4.10	0.0207	[37]
	African Americans	4079	1.50	0.0075	[38]
	Japanese	14 541	0.02	0.0001	[39]
Kp <sup>a</sup>	White people	18 934	2.28	0.0114	[34,40–42]
Kp <sup>c</sup>	Japanese (Osaka)	4442	0.32	0.0016	[*]
	Japanese (Miyagi)	5974	0.18	0.0009	[43]
Js <sup>a</sup>	African Americans	1298	15.87	0.0828	[38,44,45]
	Black Africans	593	15.68	0.0818	[46]
Ul <sup>a</sup>	Finns	2620	2.6	0.0131	[37]
	English	5000	0	0.0000	[37]
	Swedes	501	0.2	0.0011	[37]
	Chinese	12	1 pos.		[37]
	Japanese	8000	0.46	0.0023	[47]
KEL17	English	11 044	0.29	0.0015	[48]
KEL31	Japanese	400	1.5	0.0075	[49]

\*H. Yamaguchi, Y. Okubo, T. Seno, unpublished observations.

**Table 7.5** Results of genotyping tests on about 11 000 Austrian blood donors [51].

	<i>KEL*02</i> <i>k</i>	<i>KEL*04</i> <i>Kp<sup>b</sup></i>	<i>KEL*07</i> <i>Js<sup>b</sup></i>	<i>KEL*11</i> <i>K<sup>11</sup></i>
Homozygous positive (%)	91.98	98.31	99.99	99.61
Heterozygous positive (%)	7.94	1.67	0.01	0.39
Negative (%)	0.08	0.02	0	0

both of whom had previously lost babies as a result of fatal HDFN [55].

### 7.3.4 Unusual K and k expression

Red cells of two unrelated Swiss-German blood donors, and the mother of one of them, reacted with most polyclonal and monoclonal anti-K, albeit slightly less strongly than with normal K+ cells, but gave significantly weak or negative reactions with some anti-K reagents [30]. Like *KEL\*02* homozygotes, they were homozygous for 578C, but were also heterozygous for 577A>T, encoding Ser193

(*KEL\*01.02*) (Table 7.3). Ser193 would be expected to support *N*-glycosylation of Asn191. It is somewhat surprising that Ser193 would be responsible for K expression, particularly as flow cytometric analyses suggested a homozygous dose of *k*. Consequently the *KEL\*01.02* allele appears to produce both variant K and *k* antigens. Poole *et al.* [30] suggest that anti-*k* detect an epitope that is not substantially altered by a Thr193Ser substitution, whereas anti-K recognise a distinct conformational epitope created by any substitution of Thr193. A third unrelated example of this variant was found in an American Caucasian [56]. Despite encoding K, *KEL\*01.02* would be recognised as a *KEL\*02* allele by most genotyping systems.

Weak K is produced by the very rare *KEL\*01* alleles that also produce Kp<sup>a</sup> [57] (Section 7.4). Whereas this K could be missed by some serological methods, it would be predicted by genotyping.

The term K<sub>mod</sub> was used for four individuals with very weak K that could only be detected by adsorption and elution, no *k*, and weak expression of high frequency Kell-system antigens [31]. All four were homozygous for a *KEL* 578C>G encoding Thr193Arg (*KEL\*01M.01*) (Table 7.3). Weakness of other Kell antigens was due to reduced quantity of Kell glycoprotein. Heterozygosity for



other  $K_{mod}$  alleles on a  $KEL^*01$  background has also been responsible for weakened expression of K.

A K+ woman had an antibody resembling anti-K in the serum. Her red cells and those of her daughter expressed a weak K antigen, which did not react with her K-like antibody [58]. Weak expression of K has also occurred in the McLeod [59] and Gerbich-negative [60] phenotypes (Sections 7.10 and 7.13).

During a terminal episode of sepsis, red cells of a patient previously known to be K- k+ became K+, as did K- transfused cells [61]. Post-mortem blood samples contained a Gram-positive organism, *Streptococcus faecium*. K- red cells incubated with a culture containing disrupted *S. faecium* were converted to K+.

Weakness of k in K+ k+ individuals, associated with a missense mutation in the *KEL* gene, is likely to result from heterozygosity for a  $K_{mod}$  allele and  $KEL^*01$ , the weakness of the other high frequency Kell antigens being masked by normal strength antigens being produced by the  $KEL^*01$  allele (Section 7.7.2).

### 7.3.5 Anti-K

#### 7.3.5.1 Alloanti-K

Anti-K is the most common immune red cell antibody outside of the ABO and Rh systems; about two-thirds of non-Rh red cell immune antibodies are anti-K [62]. Giblett [63] estimated the relative potency of antigens in stimulating the formation of antibodies and, excluding ABO and D, K attained the highest score with a relative potency of twice that for c, about 20 times that for  $Fy^a$ , and over 100 times that for S. Anti-K is often found in sera containing antibodies to high incidence Kell system antigens. Seventy-five percent of people with IgG autoantibodies related to the Kell system also have alloanti-K in their serum [64]. Conflicting results were obtained in two studies comparing HLA Class II genotypes with anti-K production. In one, anti-K alloimmunisation was not restricted by HLA-DRB1 genotype [65], whereas in the other study frequencies of *HLA-DRB1\*11* and *HLA-DRB1\*13* were significantly higher in patients with anti-K, than in those without [66].

Anti-K, like other Kell system antibodies, are generally IgG, and predominantly IgG1 [67]. Although IgG anti-K may occasionally agglutinate K+ red cells directly, the antiglobulin test is usually the method of choice. Anti-K often react poorly in low ionic strength solutions (LISS) [68,69] and fewer molecules of anti-K bind to red cells in LISS than in normal strength saline [70]. Problems in detecting anti-K have also been encountered in automated systems [71].

A few examples of apparent non-red cell immune anti-K have been described in untransfused, healthy, male blood donors [72,73]. Microbial infection has been associated with the presence of IgM or IgA anti-K [74–78]. *Escherichia coli*, *Enterococcus faecalis*, *Morganella morganii*, and mycobacterium, responsible for pulmonary tuberculosis, have been implicated. In some cases cell-free preparations from these stool cultures inhibited IgM anti-K and K antigens were detected on the bacterial cells [74,79].

Human monoclonal anti-K are generally used as grouping reagents. Murine monoclonal anti-K have been produced by immunising mice with plasmids encoding K, followed by a boost injection of plasmid-transfected cells [80].

#### 7.3.5.2 Clinical significance of anti-K

All Kell-system antibodies must be considered potentially clinically significant and, where possible, antigen-negative red cells should be selected for transfusion. Anti-K can be responsible for severe and fatal HTRs.

Anti-K can cause severe HDFN [62,81]. In one series of tests [82], maternal anti-K was detected in 127 of 127076 pregnancies (0.1%); 13 of the pregnancies with maternal anti-K produced a K+ baby, five (38%) of whom were severely affected with HDFN. K immunisation usually results from transfusion. In a Dutch survey, 83% of women with anti-K had a history of red cell transfusion [83]. It is common practice, therefore, for girls and premenopausal women to be transfused only with K- red cells. In addition, first trimester screening for red cell antibodies in the Netherlands resulted in an improvement from 61 to 100% survival of severely affected fetuses in K alloimmunised pregnancies [84].

Unlike RhD (Section 5.18.1.4), no prophylaxis is available for the prevention of K alloimmunisation during pregnancy and at delivery. A 15-residue peptide representing the K protein, with Met179 at the C-terminus, was identified as the major helper T-cell epitope in the alloresponse to K [85]. Administration of this peptide via a suppressive route, such as the nasal mucosa, may have the potential to reduce or prevent K alloantibody production in susceptible women.

The pathogenesis of anti-K HDFN differs from that caused by anti-D. Severity of the anti-K disease is harder to predict than the anti-D disease. There is very little correlation between anti-K titre and severity of disease [62], though severe HDFN due to anti-K of titre less than 32 is extremely rare [86]. Anti-K HDFN is associated with

lower concentrations of amniotic fluid bilirubin than in anti-D HDFN of equivalent severity and post-natal hyperbilirubinaemia is not prominent in babies with anaemia caused by anti-K [82,87,88]. There is also reduced reticulocytosis and erythroblastosis in the anti-K disease, compared with anti-D HDFN. These symptoms suggest that anti-K HDFN is associated with a lower degree of haemolysis and the fetal anaemia appears to result predominantly from a suppression of erythropoiesis [87,88]. Kell glycoprotein appears on erythroid progenitors very early in erythropoiesis, whereas the Rh proteins are late to appear [89,90]. Vaughan *et al.* [91] found that *in vitro* growth of K+ erythroid blast-forming units (BFU-E) and colony-forming units (CFU-E) was specifically inhibited by monoclonal and polyclonal anti-K. They speculated that the Kell glycoprotein, an endopeptidase (Section 7.13), might be involved in regulating the growth and differentiation of erythroid progenitors, possibly by modulating peptide growth factors on the cell surface. Consequently, binding of anti-K to the Kell glycoprotein might impede its enzymatic activity and suppress erythropoiesis. Unfortunately, this theory does not take into account the K<sub>0</sub> phenotype, in which no Kell glycoprotein is present on the surface of erythroid cells, yet erythropoiesis is apparently normal. It is more likely, therefore, that anti-K suppresses erythropoiesis through the immune destruction of early erythroid progenitors. Early erythroid progenitors cultured from CD34<sup>+</sup> cells derived from K+ neonates expressed K and elicited a strong response from monocytes in a functional assay in the presence of anti-K; no response was obtained with anti-D because Rh antigens do not appear on erythroid cells until much later, when they have become haemoglobinised erythroblasts [92]. Anti-k and anti-Kp<sup>b</sup> also inhibit BFU-E growth *in vitro* [93].

In addition to inhibiting erythropoiesis, Kell antibodies also inhibit *in vitro* proliferation of granulocyte-monocyte and megakaryocyte progenitors (CFU-GM and CFU-MK) [75,76]. Pronounced thrombocytopenia, leukopenia, and neutropenia have been detected in cases of HDFN caused by anti-K [94,95].

### 7.3.5.3 'Mimicking' autoanti-K

Autoantibodies that appear to have K specificity, but which can be adsorbed and eluted from K- cells, have been detected in the serum and red cell eluates of K- patients [96–98]. These antibodies caused strong DATs and were not associated with any weakening of high frequency Kell-system antigens.

### 7.3.6 Anti-k

Less than 2 in 1000 people are k- and capable of making anti-k, yet many examples of this rare antibody have been described [99]. Most anti-k are IgG (often IgG1 [67]) and work best by the antiglobulin test, but cold agglutinating IgM anti-k are known [100,101]. Anti-k has been responsible for HTRs [99,102] and HDFN [99,102–104]. The characteristics of fetal anaemia (reticulocytopenia and normal bilirubin levels) caused by anti-k are similar to those due to anti-K, suggestive of suppression of erythropoiesis [104] (Section 7.3.5.2)

IgG1 and IgG2a monoclonal anti-k, which could not be adsorbed and eluted from k- red cells, have been raised in mice [105–107]. One antibody which behaved like anti-k, but did not react with k+ KEL:-22 red cells [108] (k and KEL22 are probably spatially related [109], Figure 7.1). Some murine monoclonal antibodies react with red cells of all Kell phenotypes except K<sub>0</sub>, but react more strongly with K- k+ and K+ k+ cells than with K+ k- cells and may behave as anti-k at an appropriate dilution [80,107,110].

## 7.4 Kp<sup>a</sup>, Kp<sup>b</sup>, and Kp<sup>c</sup> (KEL3, KEL4, and KEL21)

In 1957, Allen and Lewis [40] described anti-Kp<sup>a</sup> and its probable antithetical antibody anti-Kp<sup>b</sup>. Kell became a complex blood group system in the following year when Kp<sup>a</sup> and Kp<sup>b</sup> alleles were shown to be linked to K and k [2]. Family evidence confirmed this very close linkage; K+ Kp(a+) people never receive both K and Kp<sup>a</sup> from the same parent and never pass them on to the same child. Despite numerous studies of families with K+ Kp(a+) *propositi*, the KKp<sup>a</sup> allele was never found. However, in 2009 gene sequencing demonstrated that two K+ k+ Kp(a+b+) unrelated Caucasians with very weak K antigens were heterozygous for KKp<sup>a</sup> (KEL\*01.03) and kKp<sup>b</sup> (KEL\*02.04) [57]. The suppressive effect of Kp<sup>a</sup> on other Kell antigens expressed on the same molecule is described in Section 7.7.3.

Tests with anti-Kp<sup>a</sup> on just under 19000 white people from Europe and North America [34,40–42], showed 2.28% to be Kp(a+), a gene frequency of 0.0114 for Kp<sup>a</sup> (Table 7.4). A Kp<sup>a</sup> gene frequency of 0.0086 in Austrian blood donors was obtained by molecular genotyping [51] (Table 7.5). Only 1.21% of K+ people are Kp(a+) [34]. Although about 9% of white people are K+, only 2.7% of Kp(a+) mostly white people from Boston were K+ [40].



Kp<sup>a</sup> appears to be extremely rare in other ethnic groups. Kp<sup>b</sup> is a public antigen in all populations.

In 1979, Yamaguchi *et al.* [111] found that the red cells of a Japanese blood donor were Kp(a–b–) with otherwise unremarkable Kell antigens. Her red cells reacted with the serum containing anti-Levay [112], an antibody to the first inherited private red cell antigen, originally reported in 1945 [113,114]. Study of the informative family of the Japanese propositus proved that Levay is the product of Kp<sup>c</sup>, a third Kp allele. In one Japanese family, two Kp(a–b–c+) members appeared to be heterozygous for Kp<sup>c</sup> and a Kell-null gene K<sup>o</sup> [43]. The incidence of Kp<sup>c</sup> in Japan is shown in Table 7.4. Other than the original Levay-positive propositus and her family, only one Kp(c+) individual has been found outside Japan: a Kp(a+b–c+) Spanish American with anti-Kp<sup>b</sup>.

Kp<sup>a</sup> and Kp<sup>c</sup> differ from the common allele, Kp<sup>b</sup>, by single nucleotide changes at adjacent sites within the same codon in exon 8 [115]. Kp<sup>b</sup> has CGG encoding Arg281; Kp<sup>a</sup> has TGG, Trp281 and Kp<sup>c</sup> has CAG, Gln281. The Kp<sup>a</sup> and Kp<sup>c</sup> mutations introduce *Nla*III and *Pvu*II restriction sites, respectively. Site-directed mutagenesis experiments confirmed that the single base change is responsible for the Kp<sup>a</sup>/Kp<sup>b</sup> polymorphism [29].

#### 7.4.1 Anti-Kp<sup>a</sup>

The first (Penney) appeared to be ‘naturally occurring’ but, as with most anti-Kp<sup>a</sup>, reacted best by the antiglobulin test [34,40]. Anti-Kp<sup>a</sup> can cause delayed HTRs; only one case is reported as severe [116]. Anti-Kp<sup>a</sup> very rarely causes severe HDFN, but there are reports of a requirement for neonatal transfusion [117,118], one case of hydrops fetalis [119], and one neonate who presented with purpura, respiratory failure, severe liver dysfunction, hyperbilirubinaemia, and anaemia [120]. In one case, symptoms are described as consistent with suppression of erythropoiesis in addition to immune red cell destruction [118]. Owing to the relative rarity of Kp<sup>a</sup>, anti-Kp<sup>a</sup>, and serious clinical sequelae of incompatible transfusion, Kp(a+) red cells are not required in antibody screening panels [121].

Murine monoclonal anti-Kp<sup>a</sup> have been produced by immunising mice with plasmid DNA followed by a boost injection of plasmid-transfected cells [80].

#### 7.4.2 Anti-Kp<sup>b</sup>

The first anti-Kp<sup>b</sup> (Rautenberg) was found during routine crossmatching [2]; the serum also contained anti-K, as do some other examples [34]. Although anti-Kp<sup>b</sup> is usually

IgG (IgG1 + IgG4 [67]), two anti-Kp<sup>b</sup> that appear to be ‘naturally occurring’ and did not react by the antiglobulin test have been mentioned [34].

Serious HDFN due to anti-Kp<sup>b</sup> is very rare, but two cases are reported where obstetric intervention and, in one case, transfusion, were required [122,123]. Both mothers had been transfused during childhood. *In vivo* red cell survival studies and monocyte monolayer assays predict that anti-Kp<sup>b</sup> has the potential to cause reduced survival of transfused Kp(b+) cells [124,125]. Anti-Kp<sup>b</sup> has been responsible for a delayed HTR [124], although Kp(b+) units of blood have been administered to patients with anti-Kp<sup>b</sup>, with no indications of transfusion reaction or reduced red cell survival [126,127].

Autoanti-Kp<sup>b</sup> has been responsible for AIHA [128–130], in one case in a 12-week-old infant [129]. One autoanti-Kp<sup>b</sup> was pure IgM [131]. Autoantibodies to Kell-system antigens are often associated with weakened expression of Kell (Section 7.9).

A murine monoclonal antibody (BRIC 203) defined an epitope shared by Kp<sup>b</sup> and Kp<sup>c</sup>, but not Kp<sup>a</sup> [15]. A human single-chain Fv (scFv) antibody fragment specific for Kp<sup>b</sup> has been isolated from a V gene phage-display library derived from non-immunised donors [132].

#### 7.4.3 Anti-Kp<sup>c</sup>

The first anti-Kp<sup>c</sup>, which was called anti-Levay for 34 years, was made in response to transfusion in a patient who also made the first examples of anti-Lu<sup>a</sup>, -C<sup>w</sup>, and human anti-N [113,114]. Several more anti-Kp<sup>c</sup> have been found since, all immune and all in Japanese.

### 7.5 Js<sup>a</sup> and Js<sup>b</sup> (KEL6 and KEL7)

Js<sup>a</sup>, a new antigen present on the red cells of about 20% of African Americans, but in none of 500 white people, was first described in 1958 [44,133]. Anti-Js<sup>b</sup> was found in 1963 in the serum of a Js(a+) black woman with four Js(a+) children [134,135]. This antibody failed to react with the red cells of 13 of 1269 black donors. Twelve of the 13 were tested with anti-Js<sup>a</sup> and all were positive. The antibody did not react with the Js(a+) red cells of two sisters, believed to be homozygous for Js<sup>a</sup> because all of their 10 children were Js(a+).

The first hint that Js<sup>a</sup> and Js<sup>b</sup> might belong to the Kell system came from the observation that cells of the Kell-null phenotype (K<sub>o</sub>) were Js(a–b–) [38]. A search of 4000 black donors revealed six K+ Js(a+) propoiti and the

subsequent family studies suggested control of  $J_s^a$  and  $J_s^b$  at the *KEL* locus. This was confirmed by four large Brazilian families with K+  $J_s(a+)$  probands [136].

$J_s^a$  is almost completely confined to people of African origin [35]. The incidence of  $J_s^a$  among African Americans is about 16%, giving a frequency of 8% for the  $J_s^a$  gene (Table 7.4).  $J_s^a$  is very rare in white people. Of 11 000 African Americans tested with anti- $J_s^b$ , 34 were  $J_s(b-)$  [137]. The  $J_s(a+b-)$  phenotype has not been reported in a person of non-African origin. Genetic testing revealed  $J_s^a$  (*KEL\*06*) frequencies of 8.18% and 11.68% in Afro-Caribbean donors and from donors originating from Ngazidja (an island off the east coast of Africa), respectively [138].

An 1790T>C SNP in exon 17 encoding a single amino acid substitution in the Kell glycoprotein is responsible for the  $J_s^b/J_s^a$  polymorphism: Leu597 for  $J_s^b$ ; Pro597 for  $J_s^a$  [139] (Table 7.1). This has been confirmed by site-directed mutagenesis experiments [29]. An *MnII* restriction site is eliminated in the  $J_s^a$  allele. The Leu597Pro substitution is between two cysteine residues and could affect disulphide bonding and, consequently, folding of the molecule. In a A>G1899 synonymous SNP at Leu633 codon, about 80% of  $J_s^b$  alleles have 1899A and 20% G1899 [140].

### 7.5.1 Anti- $J_s^a$

Anti- $J_s^a$  generally react best by the antiglobulin test and are red cell immune in origin. An apparently 'naturally occurring' IgM anti- $J_s^a$  in a Japanese woman directly agglutinated  $J_s(a+)$  cells [141].

Anti- $J_s^a$  has been responsible for HDFN, including hydrops in one case [142–144]. Two anti- $J_s^a$ , barely detectable by routine serological tests, caused delayed HTRs [145,146].

### 7.5.2 Anti- $J_s^b$

All examples of anti- $J_s^b$  have been found in black people. They generally work best by the antiglobulin test. Anti- $J_s^b$  has caused severe HDFN resulting in fatal hydrops fetalis [147,148]. The poor predictive value of anti- $J_s^b$  titres is typical of Kell-system antibodies [149]. The mother of a hydropic baby received a transfusion of 275 ml  $J_s(b+)$  red cells and suffered no symptoms of an HTR, although the survival of the  $J_s(b+)$  cells was substantially reduced [147]. Anti- $J_s^b$  has been responsible for a delayed HTR [150]; multiple transfusions of  $J_s(b+)$  red cells to transfusion-dependent patients with anti- $J_s^b$  have resulted in no adverse reactions [151].

Autoanti- $J_s^b$ , enhanced by polyethylene glycol, was detected in the serum of a  $J_s(a-b+)$  renal patient whose red cells gave a weakly positive DAT [152].

A potent murine monoclonal IgG anti- $J_s^b$  was raised by immunising a mouse with a murine erythroleukaemia (MEL) cell line expressing recombinant human Kell glycoprotein [153]. Human-mouse chimeric IgM antibodies have been engineered from this clone, in order to produce a directly agglutinating reagent [154].

## 7.6 Other Kell-system antigens

In addition to the Kell polymorphisms – K/k,  $Kp^a/Kp^b/Kp^c$ , and  $J_s^a/J_s^b$  – a number of other Kell-system antigens are known, all of either high frequency or low frequency (Table 7.1). They are absent from  $K_0$  cells and expressed either weakly or not at all on McLeod phenotype cells. All have been allocated to the Kell system through family evidence, location of the antigen on the Kell glycoprotein by immunochemical means, and/or association of antigen presence or absence with a sequence change in the *KEL* gene. Ku and Km are described in Sections 7.7 and 7.13.3, respectively.

### 7.6.1 $U_l^a$ (KEL10)

An incompatible crossmatch revealed an antibody, named anti- $U_l^a$ , that reacted with the red cells of 2.6% of Helsinki blood donors, but is rare in most other populations [37]. Three families with K+  $U_l(a+)$  members had demonstrated that  $U_l^a$  belongs to the Kell system [155]. An antibody antithetical to anti- $U_l^a$  has not been found.  $U_l^a$  is often considered a predominantly Finnish characteristic, but 0.46% of Japanese [47] and one of 12 Chinese [37] were  $U_l(a+)$  (Table 7.4).

$U_l^a$  results from a mutation encoding a Glu494Val substitution and acquiring an *AccI* restriction site [115].

Anti- $U_l^a$  is very rare. No anti- $U_l^a$  was detected in the serum of 19  $U_l(a-)$  mothers of  $U_l(a+)$  children [37]. One case of HDFN caused by anti- $U_l^a$  is reported [156].

### 7.6.2 KEL11 and KEL17 ( $Wk^a$ )

The original anti-KEL11, found, in the serum of a French Canadian woman (Mrs Côté), reacted with all red cells tested except for her own, those of two siblings, and  $K_0$  phenotype cells, and reacted extremely weakly with McLeod phenotype cells [157]. Thus Côté serum appeared to contain an antibody recognising a new high frequency antigen related to the Kell system.

Anti-Wk<sup>a</sup> reacted with red cells of 0.3% of English blood donors (Table 7.4), but with those of only 0.1% of K+ donors [48]. None of 1000 Kp(a+) donors was Wk(a+). The families of five K+ Wk(a+) donors showed that Wk<sup>a</sup> was always inherited with k; there was no recombinant and 13 non-recombinants. KEL:–11 red cells were found to be Wk(a+) and the allelic status of *KEL11* and Wk<sup>a</sup> was confirmed by family studies [48,158]. As KEL11 has never been called Wk<sup>b</sup>, the numerical notation for Wk<sup>a</sup>, KEL17, will now be used here.

*KEL\*11* and *KEL\*17* differ by a single nucleotide, creating an *MscI* restriction site in the *KEL\*17* allele. *KEL\*11* encodes Val302; *KEL\*17* encodes Ala302 [115].

Anti-KEL11 and -KEL17 are rare antibodies. A patient with anti-KEL11 was transfused with 11 units of KEL:11 red cells with no adverse clinical outcome [159]. <sup>51</sup>Cr-labelled KEL:11 cells survived normally and there was no increase in reactive monocytes in a monocyte monolayer assay. One example of anti-KEL11 was implicated in severe fetal anaemia, despite results in a chemiluminescent functional assay suggesting that the antibody was not clinically significant [160].

### 7.6.3 KEL12

Five examples of anti-KEL12 and four KEL:–12 propiati are reported [161–164]. All are white (although one was originally described as black [165]). Two of the propiati each had a KEL:–12 sibling [163,164]. One of the KEL:–12 propiati and her KEL:–12 sister had both been transfused and both had anti-KEL12 [163]. Two of the propiati were transfused with KEL:12 blood with no evidence of *in vivo* destruction [163,164].

Two unrelated KEL:–12 individuals had a mutation encoding His548Arg and abolishing an *NlaIII* restriction site [32].

### 7.6.4 KEL13

The only reported KEL:–13 propiatus was a much transfused man of Italian parentage [166]. His red cells and those of his KEL:–13 sister displayed weakened expression of k, Kp<sup>b</sup>, Js<sup>b</sup>, Ku, and KEL12 and gave an enhanced score with anti-Kx (typical of cells from a K<sup>o</sup> heterozygote), suggesting that the KEL:–13 siblings have a K<sup>o</sup> allele. This was confirmed by molecular testing, which revealed heterozygosity for one allele encoding Leu329Pro and the other containing a nonsense mutation, Gln532stop [167]. KEL:–13 can be considered a K<sub>mod</sub> phenotype (*KEL\*02M.03*). In transfected human embryonic kidney cells, fewer Leu329Pro mutant Kell proteins were

transported to the cell surface, compared with control cells [167].

### 7.6.5 KEL14 and KEL24

The original anti-KEL14 was found in the serum of a white woman [161,168], and KEL14 was shown to be an inherited character retrospectively when Dp, a previously described public antigen [169], was found to be KEL14 [170]. The KEL:–14 propiatus, a white woman with consanguineous parents, had four KEL:14 and two KEL:–14 siblings [169].

IgG and IgM murine and IgG human monoclonal anti-KEL14 have been produced [105,108].

An antibody in the serum of a white woman, which reacted with the red cells of her baby and several of the baby's relatives, appeared to be antithetical to anti-KEL14 and was numbered anti-KEL24 [171]. Anti-KEL24 reacted with all three KEL:–14 samples tested, but with none of 700 other red cell samples, and gave a higher titre with KEL:–14,24 cells than with KEL:14,24 cells.

Two unrelated KEL:–14,24 individuals had a mutation introducing a *HaeIII* restriction site [172]: KEL14 represents Arg180; KEL24, Pro180. Two other mutations in two unrelated KEL:–14 Japanese encoded Arg180Cys and Arg180His [173], and the Arg180Cys mutation was also found in a KEL:–14 patient of Middle Eastern descent with anti-KEL14 [174] (Table 7.1).

### 7.6.6 KEL18

The only three reported KEL:–18 individuals were white and had anti-KEL18 [175–177]. Despite being serologically identical, the first two unrelated propiati had different single base mutations in the same codon, encoding different amino acid substitutions: Arg130Trp and Arg130Gln [32]. The two mutations created *Eco571* and *TaqII* restriction sites, respectively. No example of KEL:–18 was revealed by tests on 54450 blood donors [178].

*In vivo* survival studies and mononuclear phagocyte assays predicted that the original anti-KEL18 would not cause an acute HTR, but that transfusion therapy with KEL:18 red cells would be ineffective in all but an emergency [178]. One anti-KEL18 caused mild HDFN, necessitating phototherapy for hyperbilirubinaemia [177].

### 7.6.7 KEL19

The first anti-KEL19 was found in a KEL:–19 woman with a KEL:–19 brother and two KEL:19 sisters [179]. The second anti-KEL19, identified in the serum of a

black man, caused a delayed HTR, eliminating four units of incompatible blood [180]. None of 10757 donors tested with anti-KEL19 was KEL:–19 [180]. Two unrelated KEL:–19 individuals had a mutation encoding Arg492Gln [32].

### 7.6.8 KEL22

Two examples of anti-KEL22 were found in the sera of unrelated Israeli women of Iranian Jewish origin, with a total of three KEL:–22 siblings [181,182]. Three unrelated KEL:–22 individuals had a mutation encoding Ala322Val [32].

Anti-KEL22 in the second KEL:–22 proband was responsible for mild HDFN in her fourth and fifth children and severe HDFN in her sixth child, requiring exchange transfusion with the mother's washed red cells [182,183]. The IgG isotype was IgG1 during the fourth and fifth pregnancies and IgG1 plus IgG3 during the sixth.

### 7.6.9 KEL23

An antibody in the serum of a white woman of Italian ancestry reacted with red cells of her two children, her husband, and his mother, but with none of 2100 reference samples [184]. The antibody precipitated Kell glycoprotein from the husband's red cells and the antigen was designated KEL23. Red cells lacking high frequency Kell antigens were all KEL:–23. Two KEL:23 family members were heterozygous for a mutation encoding Gln382Arg and creating a *BcnI* restriction site [32].

Anti-KEL23 caused a strongly positive DAT on the red cells of the third baby of the proband, but did not cause HDFN [184].

### 7.6.10 KEL25 (VLAN) and KEL28 (VONG)

KEL25 and KEL28 are low frequency antigens representing mutations in the same codon encoding Arg248Gln and Arg248Trp, respectively [185,186]. Both antigens were initially shown to be located on the Kell glycoprotein by MAIEA analyses [186,187]. KEL25 was detected on the red cells of a Dutch blood donor when they were crossmatched with the serum of a patient of unknown transfusion history [187]. Two sisters and a niece of the donor were also KEL:25. Anti-KEL25 consisted of IgG1 and IgG2 isotypes and directly agglutinated KEL:25 red cells. None of 1068 donors was KEL:25. Anti-KEL28 was responsible for fetal anaemia, suggestive of suppressed erythropoiesis, in a family of ethnic Chinese from Timor [186].

### 7.6.11 KEL26 (TOU)

KEL26 is an antigen of high frequency absent from K<sub>0</sub> cells and shown to be located on the Kell glycoprotein by a MAIEA analysis [188]. Two examples of anti-KEL26 have been identified, one in a Native American man and the other in a Latino woman. Neither had been transfused, but the woman had been pregnant twice. A mutation was detected, encoding Arg406Gln in three KEL:–26 samples from two families [32]. A monocyte monolayer assay suggested that the original anti-KEL26 was not clinically significant [188].

### 7.6.12 KEL27 (RAZ)

KEL27 is a high frequency antigen, not present on K<sub>0</sub> cells, expressed weakly on McLeod phenotype cells, and located on the Kell glycoprotein as determined by a MAIEA assay [189]. Anti-KEL27 was found in a Kenyan Indian woman, the only KEL:–27 person known. She is homozygous for a mutation encoding Glu249Lys [185].

### 7.6.13 KEL29 (KALT)

Anti-KEL29 was found in a Mexican with a history of pregnancies, but no transfusion. She was homozygous for a mutation encoding Arg623Lys and deleting a *TfI* restriction site. KEL29 is the most C-terminal of the Kell antigens on the Kell glycoprotein and is unique for a Kell-system antigen because it is destroyed by trypsin treatment of intact red cells [190]. Consequently, the most N-terminal trypsin cleavage site must be between Leu597 (Js<sup>b</sup>, trypsin-resistant) and Arg623.

Red cells of the baby of the proband gave a positive DAT, but no treatment for HDFN was required.

### 7.6.14 KEL30 (KTIM)

Anti-KEL30 was made by a white American, with a history of pregnancies and transfusion. KEL:–30 resulted from homozygosity for a mutation encoding Asp305Asn eliminating a *TaqI* recognition site [190].

### 7.6.15 KEL31 (KYO), KEL38 (KYOR), and KEL37 (KUHL)

Anti-KEL31 was found through routine antibody screening of Japanese blood donors, but no further example of anti-KEL31 was found in 100 000 donors [49]. Six of 400 Japanese donors were KEL:31. KEL31 was located on the Kell glycoprotein by a MAIEA analysis and results from an Arg292Gln substitution. Two KEL31 Japanese patients who were homozygous for the mutation encoding Arg-292Gln had antibodies to KEL38, the antigen antithetical to KEL31 [191].

The only two known KEL:–37 individuals are an Asian woman and her sister with apparently normal Kell phenotypes, except that the red cells of the sister were KEL:–31 [192]. Both sisters are homozygous for a mutation encoding Arg293Trp, in the codon adjacent to that responsible for KEL:–31. Despite the propositus being Kp(a–b+), the antibody initially presented as an alloanti-Kp<sup>b</sup>, but 4 years later reacted strongly with Kp(b–) red cells, but not with K<sub>0</sub> cells.

#### 7.6.16 KEL32 (KUCI) and KEL33 (KANT)

KEL32 and KEL33 are serologically related Kell antigens of high frequency. Anti-KEL32 was produced by a previously transfused African American woman with a KEL:–32 sibling [193]. Anti-KEL33 was produced by an individual of European origin. KEL:–32 red cells are also KEL:–33, but KEL:–33 red cells appear to be KEL:32 [193]. KEL:–32 results from homozygosity for a mutation encoding Ala424Val. The KEL:–33 individual was heterozygous for a mutation encoding Arg428Leu and a Kell-null mutation encoding Arg406stop. The close proximity of the amino acid changes responsible for the KEL:–32 and KEL:–33 phenotypes probably explains the partial serological compatibility.

#### 7.6.17 KEL34 (KASH)

The antibody defining KEL34 is non-reactive with the red cells of the antibody maker, a woman of Pakistani origin, and with those of her brother, both of whom have a K<sub>mod</sub> phenotype and are homozygous for a *KEL* allele encoding a Tyr253Cys substitution [194]. The additional cysteine could disrupt disulphide bonding, causing misfolding in the Kell glycoprotein and the K<sub>mod</sub> phenotype.

#### 7.6.18 KEL35 (KELP)

A pregnant woman with an antibody to a high frequency Kell-system antigen was homozygous for two *KEL* mutations, encoding two amino acid substitutions: Leu260Phe, which is surface exposed and most likely the cause of the KEL:–35 phenotype, and Tyr253Cys, which is buried a short distance from the surface (6Å) and could affect the conformation of a nearby surface-exposed loop [195]. KEL:–35 red cells are also KEL:–12, but no explanation is provided by the protein model.

#### 7.6.19 KEL36 (KETI)

Anti-KEL36 was found in a British patient homozygous for a *KEL* allele encoding Thr464Ile [196]. Anti-KEL36

was non-reactive with two examples of K<sub>0</sub> red cells and with K<sub>mod</sub> KEL:–34 cells.

## 7.7 The Kell-null and Kell-mod phenotypes and anti-Ku (-KEL5)

### 7.7.1 K<sub>0</sub>, the Kell null phenotype

In the same year as the discovery of Kp<sup>a</sup> and Kp<sup>b</sup>, Chown *et al.* [197] found a new Kell phenotype, K– k– Kp(a–b–), in two sisters. The consanguineous parents and two other sisters were of the common Kell phenotype K– k+ Kp(a–b+). The propositus had made an antibody that reacted with all but K– k– Kp(a–b–) cells. This antibody was used to search for another example of the new phenotype [198,199]: the 3122nd blood tested did not react and was also K– k– Kp(a–b–), and the rare phenotype was named K<sub>0</sub> [2]. Family studies showed that K<sub>0</sub> results from apparent homozygosity for an amorph gene at the *KEL* locus [34]. In several families heterozygosity for a silent gene producing no K or k explains abnormal inheritance.

K<sub>0</sub> phenotype results from homozygosity or compound heterozygosity for a variety of mutations, including numerous nonsense mutations, single nucleotide insertions or deletions, splice site mutations, and several missense mutations [200–205]. The presence of null alleles has explained unexpected *KEL*\*01/02 genotyping results in individuals with K+ k– red cells and occasionally individuals with K– k+ cells. Many null and mod alleles are listed in the dbRBC and ISBT databases [206,207]. The Kell-glycoprotein has a large, extracellular C-terminal domain, so many of the mutations that cause early termination of mRNA translation might be expected to produce a truncated protein that could be detected by immunoblotting with Kell antibodies, but there is no evidence for this [3,8,201]. One possible explanation involves nonsense-mediated mRNA decay, which clears eukaryotic cells mRNA molecules containing premature termination codons [203,205]. Ser363Asn and Ser676Asn mutants, expressed in human embryonic kidney cells, were retained in a pre-Golgi compartment and not transported to the cell surface [201]. Arg128stop mutations, homozygous in two African Americans, were present in *J*<sup>s</sup> (*KEL*\*06) alleles [201].

K<sub>0</sub> red cells lack expression of all Kell antigens, including, by definition, Ku and Km. The strength of Kx antigen detected on the surface of intact K<sub>0</sub> red cells is reported to be enhanced [208], yet the quantity of Kx protein is reduced [209] (Section 7.13.2). K<sub>0</sub> red cells demonstrate no morphological abnormality [210] or unusual



expression of antigens belonging to other blood group systems, except Kx.

Only one  $K_0$  was found from testing 16518 white donors with the serum of the original  $K_0$  proband [42]. Several studies provided only one example of  $K_0$  from 24953 white people [34]. These results suggest a frequency of about 0.007 for  $K^0$  alleles in white people. One  $K_0$  was found among 14541 Japanese, suggesting a similar frequency [39].

### 7.7.2 $K_{mod}$ phenotype

Marsh and Redman [211] introduced  $K_{mod}$  as an umbrella term to describe phenotypes in which Kell antigens are expressed very weakly, often requiring adsorption/elution tests for detection, and in which Kx antigen expression is elevated.  $K_{mod}$  cells have reduced quantity of the Kell glycoprotein. Because some  $K_{mod}$  red cells have very weak expression of Kell antigens, the distinction between  $K_0$  and  $K_{mod}$  in some cases, may be dependent on the serological reagents and methods used.

$K_{mod}$  usually arises from homozygosity for a missense mutation in the *KEL* gene, or heterozygosity with another such mutation or a null mutation [167,204,206,207]. *KEL*:−13 (Leu329Pro) [167] and *KEL*:−34 (Tyr253Cys) [194] (Section 7.6.17) could also be considered  $K_{mod}$  phenotypes. A synonymous mutation within the Gly573 codon (*KEL*\*02M.04), 16 bp downstream of the 3' splice site of exon 16, caused exon 16 skipping and introduction of a premature splice site [212]. The weak expression of Kell antigens probably results from some normal splicing events. Homozygosity for a Ser363Asn mutation (*KEL*\*02M.01) resulted in a  $K_{mod}$  phenotype [167], whereas heterozygosity for the same mutation and for a null allele (Arg192stop or intron 3 splice site mutations) gave rise to  $K_0$  phenotype [201]. Based on transfection studies with HEH cells, the majority of the Ser363Asn and Tyr677Cys (*KEL*\*02M.02) mutant proteins are degraded intracellularly and not transported to the cell surface, whereas Leu329Pro (*KEL*\*02M.03) and Gly703Arg (*KEL*\*02M.04) proteins are degraded to a lesser extent, but more so than in controls [167,201].

In Austria, genotyping of 401 apparent K+ k− samples revealed *KEL*\*01/02 (578C/T) heterozygosity in 14 (3.5%) cases: nine were genuinely k− and six of these had null splice site or nonsense mutations, whereas in the other three no *KEL* mutation was detected; in four of the remaining five, k could be detected by adsorption and elution and all four had mod mutations; and the other one had weak k and was heterozygous for a  $Kp^a$  allele (Section 7.3) [204].

### 7.7.3 The $Kp^a$ effect

In the original description of  $Kp^a$ , Allen and Lewis [40] noted some difficulty in k typing some K+  $Kp(a+)$  family members, now known to result from weakening of k and other Kell antigens owing to a reduced quantity of Kell-glycoprotein with the Arg281Trp substitution responsible for  $Kp^a$  expression [29]. The  $Kp^a$  effect is only recognised under certain conditions:

- 1 when an alternative allele, such as K, is present on the opposite chromosome;
- 2 when there is a  $K^0$  gene *in trans* [213,214]; or
- 3 with difficulty, when there is homozygosity for  $Kp^a$ .  $Kp^c$ , a low incidence allele of  $Kp^a$ , does not appear to produce a similar effect when *in trans* with  $K^0$  [43].

Expression of cDNA constructs in human embryonic kidney cells showed that the  $Kp^a$  mutation causes retention of most of the Kell glycoprotein in a pre-Golgi compartment owing to differential processing, suggesting aberrant transport of the Kell glycoprotein to the cell surface [29].

### 7.7.4 Anti-Ku (-KEL5)

Anti-Ku is the typical antibody of immunized  $K_0$  individuals and detects an antigen present on all red cells apart from those of the  $K_0$  phenotype. It appears to be a single specificity and cannot be separated into components of other Kell specificity by adsorption and elution [215]. Exceptional  $K_0$  individuals with anti- $Kp^b$  or -k have been reported [34,216].

Anti-Ku has been responsible for severe and fatal HTRs [197,217,218] and for HDFN characterised by fetal anaemia [219]. Monocyte monolayer assays on 11 examples of anti-Ku suggested a high potential for causing HTRs and HDFN [220].

Some  $K_{mod}$  individuals make an antibody that resembles anti-Ku, but differs from anti-Ku by being non-reactive with the weakly Ku+  $K_{mod}$  cells of the antibody maker. Antibodies made by different  $K_{mod}$  individuals are often not mutually compatible, because  $K_{mod}$  arises from a variety of different amino acid substitutions [167]. Anti-KEL13, -KEL34, and even - $Kp^b$  could be considered as Ku-like antibodies in individuals with  $K_{mod}$  phenotypes.

## 7.8 Kell depression in Gerbich-negative phenotypes

The phenomenon of Kell depression associated with some Gerbich-negative phenotypes was first recognised



in a K+ woman and her brother with the rare Ge:–2,–3 phenotype [60]. Both showed weakened expression of K, k, and Kp<sup>b</sup>, with about half the number of K antigen sites of K+ k+ Ge:2,3 cells [16]. Nine of 11 red cell samples from Ge:–2,–3 people showed at least some degree of weakening of Kell antigens [221], as did red cells of the Ge:–2,–3,–4 Leach phenotype [222,223]. All six Ge:–2,3 samples had normal expression of Kell antigens [221]. Red cells with the K<sub>o</sub>, K<sub>mod</sub>, and McLeod phenotypes have normal expression of Gerbich antigens. The biochemical nature of the phenotypic association between Gerbich and Kell is not understood, but studies on mice suggest that the Kell glycoprotein belongs to the same membrane complex as glycophorin C, which expresses Gerbich antigens [14].

## 7.9 Acquired and transient depressed Kell phenotypes

In 1972, Seyfried *et al.* [128] described the case of a boy with severe AIHA whose red cells gave a weakly positive DAT and had weak expression of k, Kp<sup>b</sup>, Js<sup>b</sup>, and Ku. His serum contained a potent anti-Kp<sup>b</sup> responsible for an HTR. Within 16 weeks of the start of the investigation, his positive DAT had virtually disappeared, there was no sign of the anti-Kp<sup>b</sup>, and his Kell antigens were back to normal strength. Similar examples of Kell-related autoantibodies associated with weak Kell antigens have been described since [224–228].

Anti-Kp<sup>b</sup> was responsible for a positive DAT on the cells of a patient who was genetically Kp(a+b–) [226]. Her k and Js<sup>b</sup> antigens were weakly expressed, but she had strong Kp<sup>a</sup>. Nine months later the DAT was negative, the anti-Kp<sup>b</sup> undetectable, her k and Js<sup>b</sup> back to the strength expected for Kp(a+b–) cells, and own anti-Kp<sup>b</sup> from the initial study no longer reacted with her cells.

A patient with autoimmune idiopathic thrombocytopenic purpura (AITP) had a potent antibody to a high frequency Kell antigen. His red cells gave a negative DAT and displayed profound depression of Kell system antigens. Transfused cells also lost their Kell antigens. Five months later the antibody had disappeared and the patient's Kell antigens had returned to normal. An environmental agent, possibly of microbial origin, may have been responsible [229]. Another similar case in an AITP patient is reported [230]; in remission his Kell antibody disappeared and his Kell antigens returned to normal, but

his red cells lost their Lutheran antigens during a subsequent relapse (see Section 6.9).

Another patient with AITP and with IgM anti-Kp<sup>b</sup> was typed as K– k– Kp(b–), but Js(b+), with a positive DAT with anti-IgM, but not with anti-IgG [131]. When the anti-Kp<sup>b</sup> disappeared, his red cells were K– k+ Kp(b+) Js(b+), with no DAT. Boscoe *et al.* [131] suggest that there was no reduction in expression of Kell glycoprotein in this patient, but that his apparent Kp(b–) phenotype resulted from blocking of the Kp<sup>b</sup> epitopes by the IgM autoantibody, and that the large IgM molecules also blocked k epitopes by steric hindrance. This is consistent with the relatively close proximity of k and Kp<sup>b</sup> as predicted by a three-dimensional model of the Kell glycoprotein [10] and by a competitive MAIEA assay [109].

The possibility that Kell glycoprotein is expressed on megakaryocytes [94] might provide an explanation for the ITP in these patients.

## 7.10 Effects of enzymes and reducing agents on Kell antigens

Treatment of red cells with the proteases papain, ficin, or trypsin does not reduce expression of Kell antigens (with the exception of KEL29 [190] and KEL38 [191], Sections 7.6.13 and 7.6.15, which are trypsin-sensitive); the effects of  $\alpha$ -chymotrypsin and pronase are variable [231]. Treatment of red cells with a mixture of trypsin and chymotrypsin, with trypsin followed by chymotrypsin, or *vice versa*, abolishes activity of Kell antigens [231,232], but some Kell-related monoclonal antibodies continue to agglutinate red cells treated in this way [15].

The Kell glycoprotein has 15 cysteine residues in its extracellular domain and thiol reducing agents, which dissociate disulphide bonds between cysteine residues, destroy Kell antigens on intact red cells. Kell antigens are conformational and are destroyed by 100–200 mM DTT and by 6% AET at pH 8 [233,234]. Js<sup>a</sup> and Js<sup>b</sup> are inactivated by substantially lower concentrations of DTT (<2 mM) [233]. Two cysteine residues flank the position of the amino acid substitution associated with the Js<sup>a</sup>/Js<sup>b</sup> polymorphism (residue 597) and probably form part of the Js<sup>a</sup> and Js<sup>b</sup> epitopes [139]. Like natural K<sub>o</sub> cells, artificial K<sub>o</sub> cells produced by AET treatment of red cells have enhanced expression of Kx antigen [234]. AET treatment of red cells is not a reliable way of identifying Kell-system antibodies as AET destroys many other red cell antigens,

including the Lutheran, Yt, Dombrock, LW, Knops, and JMh antigens.

## 7.11 Kell antigens on other cells and in other species

### 7.11.1 Other cells and other tissues

No Kell-system antigens or Kell glycoprotein has been detected on human lymphocytes, granulocytes, monocytes, or platelets, by flow cytometry with several Kell-related monoclonal antibodies [235] or by immunoblotting with a monoclonal antibody to purified Kell glycoprotein [8]. Cells of the human erythroleukaemic line, K562, do not express Kell antigens, unless induced to synthesise haemoglobin by hemin [235,236]. Kell mRNA transcripts were detected in haemopoietic tissue, bone marrow, and fetal liver, and in peripheral blood leucocytes [237]. There is also indirect evidence that Kell glycoprotein could be present on progenitors of granulocytes, monocytes, and megakaryocytes [93,94] (Section 7.3.5.2).

*KEL* mRNA transcripts were found to be about equally abundant in erythroid tissues and testis, and were detected in lesser amounts in lymph node, brain, colon, spleen, skeletal muscle, and several other tissues. Immunoblotting and immunohistochemistry revealed Kell glycoprotein in testis, lymphoid tissues, and skeletal muscle [19,237]. Kell glycoprotein was co-isolated with Kx protein in skeletal muscle [237].

### 7.11.2 Evolutionary aspects

Chimpanzees (*Pan troglodyte*) have the Kell phenotype that is common in humans, except that they are Js(a+b−) [238]. Antigens k, Kp<sup>b</sup>, and Js<sup>a</sup> are present on the red cells of gorilla and gibbon, K and Js<sup>a</sup> on those of old world monkeys, and k on those of new world monkeys [239].

*KEL* and its mouse homologue, *Kel*, share 80% sequence identity; their glycoprotein shares 74% amino acid sequence identity [240]. The mouse Kell glycoprotein is disulphide-linked to Xk protein.

## 7.12 Functional aspects

Kell protein shares a pentameric zinc-binding motif, His-Glu-Xxx-Xxx-His (HEXXH, HELH in Kell, Figure 7.1), with zinc-dependent endopeptidases [9]. Closest homology is with the neprilysin (M13) family, which includes neutral endopeptidases 24.11 (NEP or CD10) and two

endothelium-converting enzymes (ECE-1 and ECE-2) [241]. These enzymes process a variety of biologically active peptides. ECE-1 and ECE-2 cleave big endothelin (ET)-1, big ET-2, and big ET-3, inactive peptides of about 40 amino acids, to create 21-amino-acid peptides with vasoconstrictor activity, ET-1, ET-2, and ET-3. A *KEL* cDNA construct lacking the regions encoding the cytoplasmic and membrane-spanning domains was expressed in insect (sf9) cells and a truncated Kell glycoprotein was secreted. This secreted glycoprotein cleaved big ET-3 at Trp21-Ile22 to produce ET-3. It could also process ET-1 and ET-2 from big ET-1 and big ET-2, but to a much lesser extent [242]. Secreted Kell glycoprotein was inactivated by mutating the HELH motif, essential for catalytic activity, to HGLH. Red cells of normal Kell phenotype, but not those of the K<sub>o</sub> phenotype, were also capable of processing ET-3. Clapéron *et al.* [33] claimed that Kell-glycoprotein expressing K is enzymatically inactive, but this is disputed by Sha *et al.* [243], who found that the proteins expressing either K or Js<sup>a</sup> had similar activity to the common form. ET-3 is a biologically active peptide with multiple roles, so the function of the Kell glycoprotein remains unclear. *Kel* knockout mice, like K<sub>o</sub> humans, have no obvious abnormalities, but minor effects suggested possible physiological functions of Kell glycoprotein in heart, red cell ion transport, neovascularisation in tumours, and motor function [244].

Kell glycoprotein appears on erythroid progenitor cells at an early stage of erythropoiesis, before glycophorin A or band 3 [89,90]. When K562 erythroleukaemia cell line is stimulated with hemin to develop as an erythroid line it expresses Kell but no NEP, but when stimulated into megakaryocyte development with phorbol ester it expresses NEP but no Kell [245]. Kell may, therefore, play a role in erythropoiesis (but see Section 7.3.5.2).

## 7.13 The Kx blood group system

The Kx system consists of one antigen, Kx (XK1 or 019001), located on the Xk protein, which is encoded by an X-linked gene, XK. Absence of Kx from red cells results in severe reduction in expression of Kell antigens, the McLeod phenotype.


### 7.13.1 The Xk protein and the gene that encodes it

Xk, a red cell membrane protein of apparent MW 37 kDa, was isolated from red cells by immunoprecipitation with

human alloanti-Kx [246]. Xk protein is not glycosylated [12,247]; it is phosphorylated and palmitoylated [6,7].

Patients with McLeod syndrome (MLS) lack Kx and occasionally have interstitial deletions of chromosome Xp21. Ho *et al.* [247] assembled a cosmid contig of 360 kb covering the region between genes flanking XK. They then hybridised the cosmids with genomic DNA from patients with MLS and identified the breakpoints of a deletion of about 50kb. Genomic fragments that spanned deletion endpoints were used to screen human cDNA libraries and a consensus full-length transcript for a candidate XK gene was derived from seven cDNA clones. The open reading frame predicted a 444 amino acid polypeptide with no *N*-glycosylation site and a calculated MW of 50913Da. Rabbit antibodies raised to synthetic peptides with sequences corresponding to the XK cDNA clone bound to the 37kDa Xk protein on immunoblots of membrane proteins derived from red cells of common phenotype, but not to blots of proteins from McLeod phenotype red cells [209]. The N-terminal 22 amino acids of the protein were in accordance with the corresponding nucleotide sequence of the cDNA clone [12].

Hydropathy analysis of the amino acid sequence of Xk suggested a protein that spans the membrane 10 times, with internal N- and C-termini [247] (Figure 7.2). The predicted topographical arrangement is similar to that of members of a family of proteins that co-transport a neurotransmitter together with Na<sup>+</sup> and Cl<sup>-</sup> ions [248], the amino acid sequence bearing closest resemblance to an Na<sup>+</sup>-dependent glutamate transporter [247]. Xk also



**Figure 7.2** Diagrammatic representation of the topology of the Xk protein in the red cell membrane, showing cytoplasmic termini, 10 membrane-spanning domains, and (•) Cys347, which is linked by a disulphide bond with Cys72 of the Kell glycoprotein.

shares similarity with ced-8, a protein of the nematode *Caenorhabditis elegans*, which is involved in the regulation of apoptosis [249].

XK is organised into three exons, encoding amino acids 1–82, 83–168, and 169–444, respectively [249]. XK mRNA showed widespread distribution, with high levels of expression detected in fetal liver and in adult skeletal muscle, brain, and heart [249]. Immunohistochemical analysis showed Xk in human type II muscle fibres [250] and, in the rat, Xk is widespread in the brain, but its neuronal location is cytoplasmic and not at the cell surface [251]. Kx was not detected on neutrophils [252].

A mouse homologue of XK encodes a protein with 82% identity to human Xk protein with 10 predicted membrane-spanning domains [253].

### 7.13.2 Association of Xk protein and Kell glycoprotein

When immunoprecipitation experiments were performed under non-reducing conditions with monoclonal anti-K and K+ red cells, a disulphide-bonded heterodimer comprising the 93 kDa Kell glycoprotein and the 37 kDa Xk protein was detected [12]. Rabbit antibody raised to the purified complex reacted with both components on immunoblots and precipitated the Xk protein from K<sub>o</sub> red cells, but not from McLeod phenotype cells.

Precipitation with a rabbit antibody raised to a peptide representing the second extracellular loop of Xk (Figure 7.2) isolated the Kell-Xk complex from COS-1 mammalian cells co-transfected with *KEL* and *XK* cDNA. Conversion of cysteine residues to serine by site-directed mutagenesis demonstrated that Cys347 on the fifth extracellular loop of Xk is linked by a single disulphide bond to Cys72 of the Kell glycoprotein [13]. Time-course studies on the *KEL* and *XK* transfected COS-1 cells demonstrated that the Kell-Xk complex is assembled in the endoplasmic reticulum and transported to the cell surface [254].

The relative quantity of Xk protein in K<sub>o</sub> is lower than in cells of common Kell type [201,209]. In contrast, Kx antigen, as determined by serological techniques with alloanti-Kx, is expressed more strongly on K<sub>o</sub> cells than on cells of common Kell type [208], suggesting that the presence of the Kell glycoprotein may impair access of the anti-Kx to the Xk protein in intact cells. Unlike Kell system antigens, Kx is not denatured by the disulphide-bond reducing agents DTT and AET. On the contrary, red cells treated with the appropriate concentrations of these chemicals resemble K<sub>o</sub> cells and have enhanced serological expression of Kx [234,255].

Both *KEL* and *XK* transcripts were detected in early stages of human erythropoiesis, but in mice *Kel*, but not *Xk* transcripts were detected in erythroid progenitors [256].

### 7.13.3 McLeod phenotype and McLeod syndrome (MLS)

Routine tests on medical students led Allen *et al.* [257] to recognise that one of the students, Mr McLeod, had an unusual Kell phenotype. In the McLeod red cell phenotype all high frequency Kell antigens are expressed weakly, the degree of depression of these antigens varying in different individuals. K is also weakly expressed when present [59]. McLeod phenotype red cells lack the Kx (XK1) and Km (KEL20) antigens (Section 7.13.5).

McLeod phenotype is very rare and no frequency estimate has been published. Two unrelated men with the McLeod phenotype were found as a result of testing, with anti-k, red cells from many thousands of donors from South East England [258].

McLeod phenotype is only one of a number of characteristics that collectively make up a wider phenomenon known as McLeod syndrome (MLS), a form of neuroacanthocytosis [259]. Originally described as a 'benign X-linked myopathy with acanthocytes' [258], MLS is now recognised as a multisystem degenerative disorder sharing many features with Huntington's disease [260]. McLeod cells are acanthocytic [258,259,261] with decreased whole cell deformability [262] and reduced *in vivo* survival [263]. A variety of muscular, neurological, and psychiatric defects have been associated with MLS, with an average age of onset of around 40 years [260, 264,265]. Symptoms include muscle wasting, diminished deep tendon reflex, choreiform movements, and cardiomyopathy, sometimes leading to chronic heart failure and death, personality disorders, and generalised seizures. Elevated serum creatine phosphokinase is almost a constant feature. The reason for the association between Xk deficiency and neuroacanthocytosis and muscular defects is unknown, but the relationship of Xk protein to mammalian neurotransmitter transporters could provide a clue.

MLS red cells appear biochemically and physiologically relatively normal, despite their grossly abnormal morphology and the virtual absence of Kell system antigens. The protein profile as determined by SDS PAGE is normal, suggesting no defect of the membrane skeleton [246,266,267]. Phospholipid content is essentially normal [268,269], but there is evidence for abnormalities in the composition of the membrane lipid bilayers [270] and

enhanced transbilayer mobility of phosphatidylcholine [269]. Electrolyte transport in McLeod cells is normal, but osmotic water permeability is reduced [268].

### 7.13.4 Inheritance and molecular genetics of Kx and McLeod syndrome

Expression of Kx antigen, present on all red cells save those of the McLeod phenotype, is controlled by an X-borne gene. MLS is, therefore, inherited as an X-linked recessive disease. With only rare exceptions [271], McLeod has always been found in males and the rare gene is inherited from the mother.

MLS has resulted from hemizygosity for a large variety of inactivating mutations including deletions of the whole gene, exon deletions, deletion of one or a few nucleotides introducing frameshifts, nonsense mutations, and splice site mutations (listed in [206,207,260, 264,272,273]). Two missense mutations, encoding Arg-222Gly (*XK\*N.27*) [272,273] and Glu327Lys (*XK\*N.29*) [274], have been found in two pairs of brothers manifesting no or minimal symptoms of MLS, despite all being over 50 years of age. Xk with Arg222Gly did not travel to the surface of transfected COS-1 cells [272]. A deletion of exon 2 of *XK* (*XK\*N.04*) was detected in a man with MLS and in his grandson, predicting that the boy will be afflicted with the disease in the future [275]. Studies of sets of brothers with identical *XK* mutations revealed significant variation in clinical presentation, including causes of morbidity and mortality [276]. A nonsense mutation in *XK* exon 3 (*XK\*N.21*) of a man with MLS was probably a new mutation as it was not present in his mother or sister [277].

A mutation in the fifth nucleotide of the 5' donor splice site of intron 2 of *XK* (*XK\*N.25*) was found in a man with almost no Kell antigens on his red cells and in his two daughters [278]. He did not have neuroacanthocytosis or muscle defects, possibly because of some degree of normal *XK* RNA splicing. The extreme reduction in Kell antigen was attributed to the combined effects of homozygosity for a *Kp<sup>a</sup>* allele and the Kx deficiency (Section 7.7.3).

Chronic granulomatous disease (CGD) is an inherited immunodeficiency that may be X-linked (about 60% of cases) or autosomal. X-linked CGD result from deletion of *CYBB*, an X-linked gene that encodes the NADPH oxidase NOX2, or from mutations within that gene [279]. A small minority of CGD patients, all of the X-linked type, have MLS. The locus for X-linked CGD and the *XK* locus are discrete and the association of MLS with CGD results from a deletion of part of the X-chromosome that

encompasses both genes. Some large deletions also include genes for Duchenne Muscular Dystrophy (*DMD*), retinitis pigmentosa (*RPGR*), and ornithine transcarbamylase deficiency (*OTC*), causing a condition known as contiguous gene deletion syndrome [280–284].

7.13.4.1 X-chromosome inactivation

*XK* is subject to the phenomenon of X-chromosome inactivation, in which all somatic cells in female mammals have one active X-chromosome and one inactive X-chromosome (described in Section 12.7.1). Mixed populations of Kx+ and Kx– red cells, or of red cells with strong and weak Kell antigen expression, have been recognised in many female carriers of genes responsible for MLS. The proportion of McLeod phenotype red cells in female McLeod carriers usually varies from 5 to 85% [211]. This dual population of red cells is often difficult to detect serologically, especially if Kell antibodies and not anti-Kx are used, but flow cytometry permits an accurate estimation of the two red cell populations [211,285].

The only female described with MLS was heterozygous for a single basepair deletion in exon 2 of *XK* (*XK*\*N.08) [271]. Her severe neurological and muscular defects and weakening of her Kell antigens were attributed to extreme skewing of inactivation of her X-chromosome carrying the normal *XK* gene.

7.13.5 Anti-Kx and -Km (-KEL20)

A five-year-old boy afflicted with recurrent infections (later presumed to be CGD), the second example of the McLeod phenotype, had anti-Kx and -Km (initially called anti-KL) and suffered an HTR. The antibody reacted with all cells tested, except for his own and those of Mr McLeod [286,287]. Anti-Kx reacts strongly with K<sub>0</sub> cells, weakly with red cells of common Kell phenotype, and not at all with McLeod phenotype cells [208] (Table 7.6).

**Table 7.6** Expression of Kx, Km, and Ku on red cells of common, K<sub>0</sub>, McLeod, and K<sub>mod</sub> phenotypes.

Phenotype	Kx	Km	Ku*
Common	Weak	Strong	Strong
K <sub>0</sub>	Strong	Negative	Negative
McLeod	Negative	Negative	Weak
K <sub>mod</sub>	Strong	Not tested	Weak

\* Ku represents all high frequency Kell antigens.

Adsorption of anti-Kx + -Km serum with K<sub>0</sub> cells removes anti-Kx and isolates anti-Km. The anti-Kx can be recovered by elution. Unfortunately, the separation of anti-Kx from anti-Km is often difficult to achieve and sera containing these antibodies are in very short supply. Anti-Km reacts with red cells of common Kell phenotype, but not with K<sub>0</sub> or McLeod phenotype red cells [208] (Table 7.6). The Kx determinant is located on the Xk protein whereas Km is probably a discontinuous antigen, the product of interaction between Kell glycoprotein and Kx protein. Although Km has been numbered KEL20, it could belong to the Kx system.

Anti-Kx + -Km is the typical immune response of McLeod phenotype patients with CGD following transfusion [208,286,288], and has been responsible for an HTR [263,286]. An untransfused McLeod phenotype CGD patient made anti-Km during septic shock [289]. Two transfused McLeod individuals without CGD made anti-Km, but no anti-Kx [290,291]. When one of these patients required further transfusion, a monocyte monolayer assay was strongly positive and no radiolabelled red cells of common Kell phenotype survived, *in vivo*, 24 hours after injection. Consequently, the patient received four units of K<sub>0</sub> and one unit of K<sub>mod</sub> blood. The transfusion was successful and, despite receiving strongly Kx-positive red cells, the patient did not make anti-Kx [291]. However, some patients with MLS, but no CGD, can, make anti-Kx [292,293] or anti-Kx + -Km [294] in response to transfusion. Whether the molecular background to MLS affects the immune response is not clear.

An IgG autoanti-Kx in a man with common Kell phenotype did not cause haemolysis of autologous or transfused Kx+ cells [295].

References

- Coombs RRA, Mourant AE, Race RR. *In-vivo* isosensitisation of red cells in babies with haemolytic disease. *Lancet* 1946;i:264–266.
- Allen FH, Lewis SJ, Fudenburg H. Studies of anti-Kp<sup>b</sup>, a new antibody in the Kell blood group system. *Vox Sang* 1958;3:1–13.
- Redman CM, Avellino G, Pfeffer SR, *et al*. Kell blood group antigens are part of a 93000-Dalton red cell membrane protein. *J Biol Chem* 1986;261:9521–9525.
- Redman CM, Marsh WL, Mueller KA, Avellino GP, Johnson CL. Isolation of Kell-active protein from the red cell membrane. *Transfusion* 1984;24:176–177.
- Wallas C, Simon R, Sharpe MA, Byler C. Isolation of a Kell-reactive protein from red cell membranes. *Transfusion* 1986; 26:173–176.



- 6 Carbonnet F, Hattab C, Cartron J-P, Bertrand O. Kell and Kx, two disulphide-linked proteins of the human erythrocyte membrane are phosphorylated *in vivo*. *Biochem Biophys Res Comm* 1998;247:569–575.
- 7 Carbonnet F, Hattab C, Callebaut I, *et al*. Kx, a quantitatively minor protein from human erythrocytes, is palmitoylated *in vivo*. *Biochem Biophys Res Comm* 1998;250:569–574.
- 8 Jaber A, Loirat M-J, Willem C, *et al*. Characterization of murine monoclonal antibodies directed against the Kell blood group glycoprotein. *Br J Haematol* 1991;79:311–315.
- 9 Lee S, Zambas ED, Marsh WL, Redman CM. Molecular cloning and primary structure of Kell blood group protein. *Proc Natl Acad Sci USA* 1991;88:6353–6357.
- 10 Lee S, Debnath AK, Redman CM. Active amino acids of the Kell blood group protein and model of the ectodomain based on the structure of neutral endopeptidase 24.11. *Blood* 2003;102:3028–3034.
- 11 Burton NM, Daniels G. Structural modelling of red cell surface proteins. *Vox Sang* 2011;100:129–139.
- 12 Khamlichi S, Bailly P, Blanchard D, *et al*. Purification and partial characterization of the erythrocyte Kx protein deficient in McLeod patients. *Eur J Biochem* 1995;228:931–934.
- 13 Russo D, Redman C, Lee S. Association of XK and Kell blood group proteins. *J Biol Chem* 1998;273:13950–13956.
- 14 Salomao M, Zhang X, Yang Y, *et al*. Protein 4.1R-dependent multiprotein complex: new insights into the structural organization of the red blood cell membrane. *Proc Natl Acad Sci USA* 2008;105:8026–8031.
- 15 Parsons SF, Gardner B, Anstee DJ. Monoclonal antibodies against Kell glycoprotein: serology, immunochemistry and quantitation of antigen sites. *Transfus Med* 1993;3:137–142.
- 16 Jaber A, Blanchard D, Goossens D, *et al*. Characterization of the blood group Kell (K1) antigen with a human monoclonal antibody. *Blood* 1989;73:1597–1602.
- 17 Hughes-Jones NC, Gardner B. The Kell system studied with radioactively labelled anti-K. *Vox Sang* 1971;21:154–158.
- 18 Lee S, Zambas E, Green ED, Redman C. Organization of the gene encoding the human Kell blood group protein. *Blood* 1995;85:1364–1370.
- 19 Camara-Clayette V, Rahuel C, Lopez C, *et al*. Transcriptional regulation of the *KEL* gene and Kell protein expression in erythroid and non-erythroid cells. *Biochem J* 2001;356:171–180.
- 20 Zelinski T, Coghlan G, Myal Y, *et al*. Genetic linkage between the Kell blood group system and prolactin-inducible protein loci: provisional assignment of *KEL* to chromosome 7. *Ann Hum Genet* 1991;55:137–140.
- 21 Purohit KR, Weber JL, Ward LJ, Keats BJB. The Kell blood group locus is close to the cystic fibrosis locus on chromosome 7. *Hum Genet* 1992;89:457–458.
- 22 Lee S, Zambas ED, Marsh WL, Redman CM. The human Kell blood group gene maps to chromosome 7q33 and its expression is restricted to erythroid cells. *Blood* 1993;81:2804–2809.
- 23 Murphy MT, Morrison N, Miles JS, *et al*. Regional chromosomal assignment of the Kell blood group locus (*KEL*) to chromosome 7q33–q35 by fluorescence in situ hybridization: evidence for the polypeptide nature of antigenic variation. *Hum Genet* 1993;91:585–588.
- 24 Coghlan G, Kaita H, Belcher E, Philipps S, Lewis M. Evidence for genetic linkage between the *KEL* and *YT* blood group loci. *Vox Sang* 1989;57:88–89.
- 25 Race RR. A summary of present knowledge of human blood-groups, with special reference to serological incompatibility as a cause of congenital disease. *Br Med Bull* 1946;4:188–193.
- 26 Levine P, Backer M, Wigod M, Ponder R. A new human hereditary blood property (Cellano) present in 99.8% of all bloods. *Science* 1949;109:464–466.
- 27 Toivanen P, Hirvonen T. Antigens Duffy, Kell, Kidd, Lutheran and Xg<sup>a</sup> on fetal red cells. *Vox Sang* 1973;24:372–376.
- 28 Lee S, Wu X, Reid M, Zelinski T, Redman C. Molecular basis of the Kell (K1) phenotype. *Blood* 1995;85:912–916.
- 29 Yazdanbakhsh K, Lee S, Lu Q, Reid ME. Identification in a defect in the intracellular trafficking of a Kell blood group variant. *Blood* 1999;94:310–318.
- 30 Poole J, Warke N, Hustinx H, *et al*. A *KEL* gene encoding serine at position 193 of the Kell glycoprotein results in expression of KEL1 antigen. *Transfusion* 2006;46:1879–1885.
- 31 Uchikawa M, Onodera T, Tsuneyama H, *et al*. Molecular basis of unusual K<sub>mod</sub> phenotype with K+<sup>a</sup>k-. *Vox Sang* 2000;78(Suppl. 1):abstract O011.
- 32 Lee S. Molecular basis of Kell blood group phenotypes. *Vox Sang* 1997;73:1–11 and *Vox Sang* 1998;74:58.
- 33 Clapéron A, Rose C, Gane P, *et al*. The Kell protein of the common K2 phenotype is a catalytically active metalloprotease, whereas the rare Kell K1 antigen is inactive. *J Biol Chem* 2005;280:21272–21283.
- 34 Race RR, Sanger R. *Blood Groups in Man*, 6th edn. Oxford: Blackwell Scientific Publications, 1975.
- 35 Mourant AE, Kopec AC, Domaniewska-Sobczak K. *The Distribution of the Human Blood Groups and Other Polymorphisms*, 2nd edn. London: Oxford University Press, 1976.
- 36 Garretta M, Gener J, Muller A. Analyse de 280 000 déterminations du facteur Kell sur les équipements Groupamatic. *Rev Franc Transfus Immuno-Hémat* 1978;21:379–386.
- 37 Furuholm U, Nevanlinna HR, Nurkka R, *et al*. The blood group antigen U<sup>1</sup> (Karhula). *Vox Sang* 1968;15:118–124.
- 38 Stroup M, MacIlroy M, Walker R, Aydelotte JV. Evidence that Sutter belongs to the Kell blood group system. *Transfusion* 1965;5:309–314.



- 39 Hamilton HB, Nakahara Y. The rare Kell blood group phenotype K<sup>o</sup> in a Japanese family. *Vox Sang* 1971;20:24–28.
- 40 Allen FH, Lewis SJ. Kp<sup>a</sup> (Penney), a new antigen in the Kell blood group system. *Vox Sang* 1957;2:81–87.
- 41 Dichupa PJ, Anderson C, Chown B. A further search for hypothetical K<sup>p</sup> of the Kell-system. *Vox Sang* 1969;17:1–4.
- 42 Chown B, Lewis M, Kaita H, Philipps S. Some blood group frequencies in a Caucasian population. *Vox Sang* 1963;8:378–381.
- 43 Kikuchi M, Endo N, Seno T, Okubo Y, Yamaguchi H. A Japanese family with two Kp(a–b–c+) members, presumed genotype Kp<sup>f</sup>/K<sup>o</sup>. *Transfusion* 1983;23:254–255.
- 44 Giblett ER, Chase J. Js<sup>a</sup>, a ‘new’ red-cell antigen found in Negroes; evidence for an eleventh blood group system. *Br J Haematol* 1959;5:319–326.
- 45 Jarkowski TL, Hinshaw CP, Beattie KM, Silberberg B. Another example of anti-Js<sup>a</sup>. *Transfusion* 1962;2:423–424.
- 46 Spielmann W, Teixidor D, Renninger W, Matznetter T. Blutgruppen und Lepra bei moçambiquanischen Völkern. *Humangenetik* 1970;10:304–317.
- 47 Okubo Y, Yamaguchi H, Seno T, *et al.* The first example of anti-UI<sup>a</sup> and UI(a+) red cells found in Japan. *Transfusion* 1986;26:215.
- 48 Strange JJ, Kenworthy RJ, Webb AJ, Giles CM. Wk<sup>a</sup> (Weeks), a new antigen in the Kell blood group system. *Vox Sang* 1974;27:81–86.
- 49 Uchikawa M, Onodera T, Ogasawara K, *et al.* Molecular basis for a novel low-frequency antigen in the Kell blood group system, KYO. *Vox Sang* 2006;91(Suppl. 3):137 [Abstract].
- 50 Ball M, Cant B, Garwood P, Rogers KL, Tills D, Sonneborn H-H. Evaluation of a murine monoclonal anti-k (BS45). *Biotest Bull* 1989;4:15–18.
- 51 Jungbauer C. Molecular bases and genotyping for rare blood groups. *Transfus Med Hemother* 2009;36:213–218.
- 52 Finning K, Martin P, Summers J, Daniels G. Fetal genotyping for the K (Kell) and Rh C, c, and E blood groups on cell-free fetal DNA from maternal plasma. *Transfusion* 2007;47:2126–2133.
- 53 Daniels G, Finning K, Martin P. Non-invasive fetal blood grouping: present and future. *Clin Biol Sci* 2010;30:431–442.
- 54 Li Y, Finning K, Daniels G, Hahn S, Zhong X, Holzgreve W. Noninvasive genotyping fetal Kell blood group (KEL1) using cell-free fetal DNA in maternal plasma by MALDI-TOF mass spectrometry. *Prenat Diagn* 2008;28:203–208.
- 55 Verlinsky Y, Rechitsky S, Ozen S, *et al.* Preimplantation genetic diagnosis for the Kell genotype. *Fertil Steril* 2003;80:1047–1051.
- 56 Lee-Stroka H, Slezak SL, Adams S, *et al.* Another example of KEL1 variant red cell phenotype due to a threonine to serine change at position 193 of Kell glycoprotein. *Transfusion* 2008;48:925–929.
- 57 Körmöcz G, Scharberg EA, Gassner C. A novel KEL\*1,3 allele with weak Kell antigen expression confirming the cis-modifier effect of KEL3. *Transfusion* 2009;49:733–739.
- 58 McDowell MA, Mann JM, Milakovic K. Kell-like antibody in a Kell positive patient. *Transfusion* 1978;18:389 [Abstract].
- 59 Marsh WL, Schnipper EF, Johnson CL, Mueller KA, Schwartz SA. An individual with McLeod syndrome and the Kell blood group antigen K (K1). *Transfusion* 1983;23:336–338.
- 60 Muller A, André-Liardet J, Garretta M, Brocteur J, Moullec J. Observations sur un anticorps rare: l’anti-Gerbich. *Rev Franc Transfus* 1973;16:251–257.
- 61 McGinniss MH, MacLowry JD, Holland PV. Acquisition of K:1-like antigen during terminal sepsis. *Transfusion* 1984;24:28–30.
- 62 Klein HG, Anstee DJ. *Mollison’s Blood Transfusion in Clinical Medicine*, 11th edn. Oxford: Blackwell Publishing, 2005.
- 63 Giblett ER. A critique of the theoretical hazard of inter vs. intra-racial transfusion. *Transfusion* 1961;1:233–238.
- 64 Marsh WL, Mueller KA, Johnson CL. Use of AET-treated cells in the investigation of Kell related auto-immunity. *Transfusion* 1982;22:419 [Abstract].
- 65 Noizat-Pirenne F, Tournamille C, Bierling P, *et al.* Relative immunogenicity of Fy<sup>a</sup> and K antigens in a Caucasian population, based on HLA class II restriction analysis. *Transfusion* 2006;46:1328–1333.
- 66 Chiaroni J, Dettori I, Ferrera V, *et al.* HLA-DRB1 polymorphism is associated with Kell immunisation. *Br J Haematol* 2005;132:374–378.
- 67 Hardman JT, Beck ML. Hemagglutination in capillaries: correlation with blood group specificity and IgG subclass. *Transfusion* 1981;21:343–346.
- 68 Molthan L, Strohm PL. Hemolytic transfusion reaction due to anti-Kell undetectable in low-ionic-strength solutions. *Am J Clin Path* 1981;75:629–631.
- 69 Voak D, Downie M, Haigh T, Cook N. Improved antiglobulin tests to detect difficult antibodies: detection of anti-Kell by LISS. *Med Lab Sci* 1982;39:363–370.
- 70 Merry AH, Thomson EE, Lagar J, *et al.* Quantitation of antibody binding to erythrocytes in LISS. *Vox Sang* 1984;47:125–132.
- 71 West NC, Jenkins JA, Johnston BR, Modi N. Interdonor incompatibility due to anti-Kell antibody undetectable by automated antibody screening. *Vox Sang* 1986;50:174–176.
- 72 Morgan P, Bossom EL. ‘Naturally occurring’ anti-Kell (K1): two examples. *Transfusion* 1963;3:397–398.
- 73 Clark A, Monaghan WP, Martin CA. Two additional examples of non-transfusion-stimulated anti-Kell. *Am J Med Tech* 1981;47:983–984.
- 74 Marsh WL, Nichols ME, Øyen R, *et al.* Naturally occurring anti-Kell stimulated by *E. Coli* enterocolitis in a 20-day-old child. *Transfusion* 1978;18:149–154.

- 75 Tegoli J, Sausais L, Issitt PD. Another example of a 'naturally-occurring' anti-K1. *Vox Sang* 1967;12:305–307.
- 76 Kanel GC, Davis I, Bowman JE. 'Naturally occurring' anti-K1: possible association with mycobacterium infection. *Transfusion* 1978;18:472–473.
- 77 Doelman CJA, Westermann WF, van Voorst tot Voorst E, Miedema K. An anti-K apparently induced by *Enterococcus faecalis* in a 30-year-old man. *Transfusion* 1992;32:790.
- 78 Pereira A, Monteagudo J, Rovira M, et al. Anti-K1 of the IgA class associated with *Morganella morganii* infection. *Transfusion* 1989;29:549–551.
- 79 Savalonis JM, Kalish RI, Cummings EA, Ryan RW, Aloisi R. Kell blood group activity of gram-negative bacteria. *Transfusion* 1988;28:229–232.
- 80 Tearina Chu T-H, Halverson GR, Yazdanbakhsh K, Øyen R, Reid ME. A DNA-based immunization protocol to produce monoclonal antibodies to blood group antigens. *Br J Haematol* 2001;113:32–36.
- 81 Poole J, Daniels G. Blood group antibodies and their significance in transfusion medicine. *Transfus Med Rev* 2007;21:58–71.
- 82 Caine ME, Mueller-Heubach E. Kell sensitization in pregnancy. *Am J Obstet Gynecol* 1986;154:85–90.
- 83 Koelwijn JM, Vrijkotte TGM, de Haas M, van der Schoot CE, Bonsel GJ. Risk factors for the presence of non-rhesus D red blood cell antibodies in pregnancy. *BJOG* 2009;116:655–664.
- 84 Kamphuis MM, Lindenburg I, van Kamp IL, et al. Implementation of routine screening for Kell antibodies: does it improve perinatal survival? *Transfusion* 2008;48:953–957.
- 85 Stephen J, Cairns LS, Pickard WJ, et al. Identification, immunomodulatory activity and immunogenicity of the major helper T cell epitope on the K blood group antigen. *Blood* 2012;119:5563–5574.
- 86 McKenna DS, Nagaraja HN, O'Shaughnessy R. Management of pregnancies complicated by anti-Kell isoimmunization. *Obstet Gynecol* 1999;93:667–673.
- 87 Vaughan JI, Warwick R, Letsky E, et al. Erythropoietic suppression in fetal anemia because of Kell alloimmunization. *Am J Obstet Gynecol* 1994;171:247–252.
- 88 Weiner CP, Widness JA. Decreased fetal erythropoiesis and hemolysis in Kell hemolytic anemia. *Am J Obstet Gynecol* 1996;174:547–551.
- 89 Southcott MJG, Tanner MJA, Anstee DJ. The expression of human blood group antigens during erythropoiesis in a cell culture system. *Blood* 1999;93:4425–4435.
- 90 Bony V, Gane P, Bailly P, Cartron J-P. Time-course expression of polypeptides carrying blood group antigens during human erythroid differentiation. *Br J Haematol* 1999;107:263–274.
- 91 Vaughan JI, Manning M, Warwick RM, et al. Inhibition of erythroid progenitor cells by anti-Kell antibodies in fetal alloimmune anemia. *New Engl J Med* 1998;338:798–803.
- 92 Daniels G, Hadley A, Green CA. Causes of fetal anaemia in haemolytic disease due to anti-K. *Transfusion* 2003;43:115–116.
- 93 Wagner T, Berer A, Lanzer G, Geissler K. Kell is not restricted to erythropoietic lineage but is also expressed on myeloid progenitor cells. *Br J Haematol* 2000;110:409–411.
- 94 Wagner T, Bernaschek G, Geissler K. Inhibition of megakaryopoiesis by Kell-related antibodies. *New Engl J Med* 2000;343:72.
- 95 Wagner T, Resch B, Reiterer F, Gassner C, Lanzer G. Pancytopenia due to suppressed hematopoiesis in a case of fatal hemolytic disease of the newborn associated with anti-K supported by molecular K1 typing. *J Pediatr Hematol Oncol* 2004;26:13–15.
- 96 Garratty G, Sattler MS, Petz LD, Flannery EP. Immune hemolytic anemia associated with anti-Kell and a carrier state for chronic granulomatous disease. *Rev Franc Transfus Immuno-Hémat* 1979;22:529–549.
- 97 Hare V, Wilson MJ, Wilkinson S, Issitt PD. A Kell system antibody with highly unusual characteristics. *Transfusion* 1981;21:613 [Abstract].
- 98 Viggiano E, Clary NL, Ballas SK. Autoanti-K antibody mimicking an alloantibody. *Transfusion* 1982;22:329–332.
- 99 Kluge A, Jungfer H. Anti-K2 (Cellano) blood group antibodies. Typing as IgG and IgA with a review of their clinical significance. *Blut* 1970;21:357–365.
- 100 Thomas MJ, Konugres AA. An anti-K2 (Cellano) serum with unusual properties. *Vox Sang* 1966;11:227–229.
- 101 Dinning G, Doughty RW, Collins AK. A further example of IgM anti-K2 (Cellano). *Vox Sang* 1985;48:317–318.
- 102 Schultz MH. Serology and clinical significance of Kell blood group system antibodies. In: Laird-Fryer B, Daniels G, Levitt J, eds. *Blood Group Systems: Kell*. Arlington: American Association of Blood Banks, 1990:37–68.
- 103 Bowman JM, Harman FA, Manning CR, Pollock JM. Erythroblastosis fetalis produced by anti-k. *Vox Sang* 1989;56:187–189 and *Vox Sang* 1990;58:139.
- 104 Win N, Amess P, Needs M, Hewitt PE. Use of red cells preserved in extended storage media for exchange transfusion in anti-k haemolytic disease of the newborn. *Transfus Med* 2005;15:157–160.
- 105 Nichols ME, Rosenfield RE, Rubenstein P. Monoclonal anti-K14 and anti-K2. *Vox Sang* 1987;52:231–235.
- 106 Chester MA, Johnson U, Lundblad A, et al. *Proc 2nd Int Workshop on Monoclonal Antibodies Against Human Red Blood Cell and Related Antigens*, 1990:126–138.
- 107 Rouger P, Anstee D, Salmon C, eds. *1st International Workshop on Monoclonal Antibodies Against Human Red Blood Cell and Related Antigens*. *Rev Franc Transfus Immuno-Hémat* 1988;31:381–418.
- 108 Daniels G. Section 4: antibodies to other blood group antigens. Coordinator's report. *Transfus Clin Biol* 2002;9:75–80.

- 109 Petty AC, Green CA, Daniels GL. The monoclonal antibody-specific immobilisation of erythrocyte antigens assay (MAIEA) in the investigation of human red cell antigens and their associated membrane proteins. *Transfus Med* 1997;7:179–188.
- 110 Inglis G, Fraser RH, McTaggart S, *et al.* Monoclonal antibodies to high incidence Kell epitopes: characterization and application in automated screening of donor samples. *Transfus Med* 1994;4:209–212.
- 111 Yamaguchi H, Okubo Y, Seno T, Matsushita K, Daniels GL. A 'new' allele, *Kp<sup>c</sup>*, at the *Kell* complex locus. *Vox Sang* 1979;36:29–30.
- 112 Gavin J, Daniels GL, Yamaguchi H, Okubo Y, Seno T. The red cell antigen once called Levay is the antigen *Kp<sup>c</sup>* of the *Kell* system. *Vox Sang* 1979;36:31–33.
- 113 Callender S, Race RR, Paykoc ZV. Hypersensitivity to transfused blood. *Br Med J* 1945;ii:83.
- 114 Callender ST, Race RR. A serological and genetical study of multiple antibodies formed in response to blood transfusion by a patient with lupus erythematosus diffusum. *Ann Eugen* 1946;13:102–117.
- 115 Lee S, Wu X, Son S, *et al.* Point mutations characterize *KEL10*, the *KEL3*, *KEL4*, and *KEL21* alleles, and the *KEL17* and *KEL11* alleles. *Transfusion* 1996;36:490–494.
- 116 Koshy R, Patel B, Harrison JS. Anti-*Kp<sup>a</sup>*-induced severe delayed haemolytic transfusion reaction. *Immunohematology* 2009;25:44–47.
- 117 Costamagna L, Barbarini M, Viarengo GL, *et al.* A case of hemolytic disease of the newborn due to anti-*Kp<sup>a</sup>*. *Immunohematology* 1997;13:61–62.
- 118 Tuson M, Hue-Roye K, Koval K, *et al.* Possible suppression of fetal erythropoiesis by the *Kell* blood group antibody anti-*Kp<sup>a</sup>*. *Immunohematology* 2011;27:58–60.
- 119 Smoleniek J, Anderson N, Poole GD. Hydrops fetalis caused by anti-*Kp<sup>a</sup>*, an antibody not usually detected in routine screening. *Transfus Med* 1994;4(Suppl. 1):48 [Abstract].
- 120 Brumbaugh JE, Morgan S, Beck JC, *et al.* Blueberry muffin rash, hyperbilirubinemia, and hypoglycemia: A case of hemolytic disease of the fetus and newborn due to anti-*Kp<sup>a</sup>*. *J Perinatol* 2011;31:373–376.
- 121 Garratty G. Screening for RBC antibodies – what should we expect from antibody detection RBCs. *Immunohematology* 2002;18:71–77.
- 122 Dacus JV, Spinnato JA. Severe erythroblastosis fetalis secondary to anti-*Kp<sup>b</sup>* sensitization. *Am J Obstet Gynecol* 1984;150:888–889.
- 123 Gorlin JB, Kelly L. Alloimmunisation via previous transfusion places female *Kp<sup>b</sup>*-negative recipients at risk for having children with clinically significant hemolytic disease of the newborn. *Vox Sang* 1994;66:46–48.
- 124 Wren MR, Issitt PD. The monocyte monolayer assay and *in vivo* antibody activity. *Transfusion* 1986;26:548 [Abstract].
- 125 Eby CS, Cowan JL, Ramos RR, Chaplin H. *In-vivo* and *in-vitro* studies of anti-*Kp<sup>b</sup>* allo-antibody. *Joint Congr Int Soc Blood Transfus and Am Assoc Blood Banks*, 1990:156 [Abstracts].
- 126 Kohan AI, Reybaud JF, Salamone HJ, *et al.* Management of a severe transfusional problem in a patient with alloantibody to *Kp<sup>b</sup>* (*K4*). *Vox Sang* 1990;59:216–217.
- 127 Mazzara R, Lozano M, Salmerón JM, *et al.* Transfusion of incompatible RBCs to a patient with alloanti-*Kp<sup>b</sup>*. *Transfusion* 2001;41:611–614.
- 128 Seyfried H, Górská B, Maj S, *et al.* Apparent depression of antigens of the *Kell* blood group system associated with autoimmune acquired haemolytic anaemia. *Vox Sang* 1972;23:528–536.
- 129 Win N, Kaye T, Mir N, Damain-Willems C, Chatfield C. Autoimmune haemolytic anaemia in infancy with anti-*Kp<sup>b</sup>* specificity. *Vox Sang* 1996;71:187–188.
- 130 Lee E, Burgess, Win N. Autoimmune hemolytic anemia and a further example of autoanti-*Kp<sup>b</sup>*. *Immunohematology* 2005;21:119–121.
- 131 Bosco A, Xenocostas A, Kinney J, Cadwell CM, Zimring JC. An autoanti-*Kp<sup>b</sup>* immunoglobulin M that stimulated antigen suppression. *Transfusion* 2009;49:750–756.
- 132 Hughes-Jones NC, Gorick BD, Bye JM, *et al.* Characterization of human blood group scFv antibodies derived from a V gene phage-display library. *Br J Haematol* 1994;88:180–186.
- 133 Giblett ER. Js, a 'new' blood group antigen found in Negroes. *Nature* 1958;181:1221–1222.
- 134 Walker RH, Argall CI, Steane EA, Sasaki TT, Greenwalt TJ. Anti-*Js<sup>b</sup>*, the expected antithetical antibody of the Sutter blood group system. *Nature* 1963;197:295–296.
- 135 Walker RH, Argall CI, Steane EA, Sasaki TT, Greenwalt TJ. *Js<sup>b</sup>* of the Sutter blood group system. *Transfusion* 1963;3:94–99.
- 136 Morton NE, Krieger H, Steinberg AG, Rosenfield RE. Genetic evidence confirming the localization of Sutter in the *Kell* blood-group system. *Vox Sang* 1965;10:608–613.
- 137 Beattie KM, Shafer AW, Sigmund K, Cisco S. Mass screening of American Black donors to identify high incidence antigen-negative bloods. *19th Congr Int Soc Blood Transfus*, 1986:312 [Abstracts].
- 138 Silvy M, Di Cristofaro J, Beley S, *et al.* Identification of *RHCE* and *KEL* alleles in large cohorts of Afro-Caribbean and Comorian donors by multiplex SNaPshot and fragment assays: a transfusion support for sickle cell disease patients. *Br J Haematol* 2011;154:260–270.
- 139 Lee S, Wu X, Reid M, Redman C. Molecular basis of the *K:6,-7* [*Js(a+b-)*] phenotype in the *Kell* blood groups system. *Transfusion* 1995;35:822–825.
- 140 Hansen BJ, Robbins F-M, Adams S, *et al.* Identification of a *KEL7* subtype: implications for genotyping red blood cell *Js<sup>a</sup>* and *Js<sup>b</sup>* antigens. *Transfus Med* 2006;16:445–446.
- 141 Ito K, Mukumoto Y, Konishi H. An example of 'naturally occurring' anti-*Js<sup>a</sup>* (*K6*) in a Japanese female. *Vox Sang* 1979;37:350–351.

- 142 Donovan LM, Tripp KL, Zuckerman JE, Konugres AA. Hemolytic disease of the newborn due to anti-Js<sup>a</sup>. *Transfusion* 1973;13:153.
- 143 Levene C, Rudolphson Y, Shechter Y. A second case of hemolytic disease of the newborn due to anti-Js<sup>a</sup>. *Transfusion* 1980;20:714–715.
- 144 Ringressi A, Periti E, Magrini A, Falchi L, Gentilini G. First case of severe HDN with foetal hydropsy and intra uterine transfusion therapy, by anti-Js<sup>a</sup> antibody. *VIII Europ Congr Int Soc Blood Transfus, Istanbul*, 2003;93 [Abstracts].
- 145 Taddie SJ, Barrasso C, Ness PM. A delayed transfusion reaction caused by anti-K6. *Transfusion* 1982;22:68–69.
- 146 Anderson RR, Sosler SD, Kovach J, DeChristopher PJ. Delayed hemolytic transfusion reaction due to anti-Js<sup>a</sup> in an alloimmunized patient with a sickle cell syndrome. *Am J Clin Pathol* 1997;108:658–661.
- 147 Ratcliff D, Fiorenza S, Culotta E, Arndt P, Garratty G. Hydrops fetalis (HF) and a maternal hemolytic transfusion reaction associated with anti-Js<sup>b</sup>. *Transfusion* 1987;27:534 [Abstract].
- 148 Gordon MC, Kennedy MS, O'Shaughnessy RW, Waheed A. Severe hemolytic disease of the newborn due to anti-Js<sup>b</sup>. *Vox Sang* 1995;69:140–141.
- 149 Stanworth S, Fleetwood P, de Silva. Severe haemolytic disease of the newborn due to anti-Js<sup>b</sup>. *Vox Sang* 2001;81:134–135.
- 150 Waheed A, Kennedy MS. Delayed hemolytic transfusion reaction caused by anti-Js<sup>b</sup> in a Js(a+b+) patient. *Transfusion* 1982;22:161–162.
- 151 Yuan S, Ewing NP, Bailey D, Salvador M, Wang S. Transfusion of multiple units of Js(b+) red blood cells in the presence of anti-Js<sup>b</sup> in a patient with sickle  $\beta$ -thalassemia disease and a review of the literature. *Immunohematology* 2007;23:75–80.
- 152 Eveland D. Autoanti-Js<sup>b</sup> enhanced by polyethylene glycol. *Joint Congr Int Soc Blood Transfus and Am Assoc Blood Banks*, 1990:156 [Abstracts].
- 153 Chu T-H T, Yazdanbakhsh K, Øyen R, Smart E, Reid ME. Production and characterization of anti-Kell monoclonal antibodies using transfected cells as the immunogen. *Br J Haematol* 1999;106:817–823.
- 154 Huang TJ, Reid ME, Halverson GR, Yazdanbakhsh K. Production of recombinant murine-human chimeric IgM and IgG anti-Js<sup>b</sup> for use in the clinical laboratory. *Transfusion* 2003;43:758–764.
- 155 Furuhjelm U, Nevanlinna HR, Nurkka R, Gavin J, Sanger R. Evidence that the antigen U1<sup>a</sup> is controlled from the Kell complex locus. *Vox Sang* 1969;16:496–499.
- 156 Sakuma K, Suzuki H, Ohto H, Tsuneyama H, Uchikawa M. First case of hemolytic disease of the newborn due to anti-U1<sup>a</sup> antibodies. *Vox Sang* 1994;66:293–294.
- 157 Guévin RM, Taliano V, Waldmann O. The Côté serum (anti-K11), an antibody defining a new variant in the Kell system. *Vox Sang* 1976;31(Suppl. 1):96–100.
- 158 Sabo B, McCreary J, Gellerman M, *et al.* Confirmation of K<sup>11</sup> and K<sup>17</sup> as alleles in the Kell blood group system. *Vox Sang* 1975;29:450–455.
- 159 Kelley CM, Karwal MW, Schlueter AJ, Olson JD. Outcome of transfusion of K:11 erythrocytes in a patient with anti-K11 antibody. *Vox Sang* 1998;74:205–208.
- 160 Flommersfeld S, Geipel A, Heep A, Oldenburg J, Hoch J. K11-incompatibility: rare causes of severe fetal anaemia. *Transfus Med Hemother* 2007;34(Suppl. 1):52 [Abstract].
- 161 Heistø H, Guévin R-M, Taliano V, *et al.* Three further antigen-antibody specificities associated with the Kell blood group system. *Vox Sang* 1973;24:179–180.
- 162 Marsh WL, Stroup M, Macilroy M, *et al.* A new antibody, anti-K12, associated with the Kell blood group system. *Vox Sang* 1973;24:200–205.
- 163 Beattie KM, Heinz B, Korol S, Øyen R, Marsh WL. Anti-K12 in the serum of two brothers: inheritance of the K:-12 phenotype. *Rev Franc Transfus Immuno-Hémat* 1982;25:611–618.
- 164 Reid ME, Øyen R, Redman CM, *et al.* K12 is located on the Kell blood group protein in proximity to K/k and Js<sup>a</sup>/Js<sup>b</sup>. *Vox Sang* 1995;68:40–45.
- 165 Taylor HL. Anti-K12 in a Black? *Transfusion* 1979;19:787–788.
- 166 Marsh WL, Jensen L, Øyen R, *et al.* Anti-K13 and the K:-13 phenotype. *Vox Sang* 1974;26:34–40.
- 167 Lee S, Russo DCW, Reid ME, Redman CM. Mutations that diminish expression of Kell surface protein and lead to the K<sub>mod</sub> RBC phenotype. *Transfusion* 2003;43:1121–1125.
- 168 Wallace ME, Bouysou C, de Jongh DS, *et al.* Anti-K14: an antibody specificity associated with the Kell blood group system. *Vox Sang* 1976;30:300–304.
- 169 Frank S, Schmidt RP, Baugh M. Three new antibodies to high-incidence antigenic determinants (anti-E1, anti-Dp, and anti-So). *Transfusion* 1970;10:254–257.
- 170 Sabo B, McCreary J, Harris P. Anti-Dp is anti-K14. *Vox Sang* 1982;43:56.
- 171 Eicher C, Kirkley K, Porter M, Kao Y. A new low frequency antigen in the Kell system: K24 (Cl<sub>s</sub>). *Transfusion* 1985;25:448 [Abstract].
- 172 Lee S, Naime, Reid M, Redman C. The KEL24 and KEL14 alleles of the Kell blood group system. *Transfusion* 1997;37:1035–1038.
- 173 Uchikawa M, Onodera T, Tsuneyama H, *et al.* Different point mutations in the same codon of KEL:-14 phenotype. *Transfusion* 1999;39(Suppl.):50S [Abstract].
- 174 Thomas N, Vege S, Mills J, Nance SJ, Westhoff CM. A new allele in the Kell blood group system maps to the K14/K24 polymorphism. *Transfusion* 2006;46 (Suppl.):138A–139A [Abstract].
- 175 Barrasso C, Eska P, Grindon AJ, Øyen R, Marsh WL. Anti-K18: an antibody defining another high-frequency antigen related to the Kell blood group system. *Vox Sang* 1975;29:124–127.



- 176 Pehta JC, Valinsky J, Redman C, Marsh WL. Biochemical and flow cytometric analysis of the second example of anti-K18. *Joint Congr Int Soc Blood Transfus and Am Assoc Blood Banks*, 1990;57 [Abstracts].
- 177 O'Leary MF, Larimore KS, Hue-Roye K, *et al.* Anti-K18 causing hemolytic disease of the fetus and newborn. *Transfusion* 2011;51(Suppl.):156A [Abstract].
- 178 Barrasso C, Baldwin ML, Drew H, Ness PM. *In vivo* survival of K:18 red cells in a recipient with anti-K18. *Transfusion* 1983;23:258–259.
- 179 Sabo B, McCreary J, Stroup M, Smith DE, Weidner JG. Another Kell-related antibody, anti-K19. *Vox Sang* 1979;36:97–102.
- 180 Marsh WL, DiNapoli J, Øyen R, *et al.* Delayed hemolytic transfusion reaction caused by the second example of anti-K19. *Transfusion* 1979;19:604–608.
- 181 Bar Shany S, Ben Porath D, Levene C, Sela R, Daniels GL. K22, a 'new' para-Kell antigen of high frequency. *Vox Sang* 1982;42:87–90.
- 182 Manny N, Levene C, Harel N, *et al.* The second example of anti-K22 and a family genetically informative for K and K<sup>22</sup>. *Vox Sang* 1985;49:135–137.
- 183 Levene C, Sela R, DaCosta Y, Manny N. Clinical significant of anti-K22. *20th Congr Int Soc Blood Transfus*, 1988:295 [Abstracts].
- 184 Marsh WL, Redman CM, Kessler LA, *et al.* A low-incidence antigen in the Kell blood group system identified by biochemical characterization. *Transfusion* 1987;27:36–40.
- 185 Lee S, Reid ME, Redman CM. Point mutations in KEL exon 8 determine a high-incidence (RAZ) and a low-incidence (KEL25, VLAN) antigen of the Kell blood group system. *Vox Sang* 2001;81:258–63.
- 186 Grey D, Poole J, Martin P, *et al.* Haemolytic disease of the newborn caused by a new Kell antigen. *Transfus Med* 2003;13(Suppl. 1):30 [Abstract].
- 187 Jongerius JM, Daniels GL, Overbeeke MAM, *et al.* A new low-incidence antigen in the Kell blood group system: VLAN (KEL25). *Vox Sang* 1996;71:43–47.
- 188 Jones J, Reid ME, Øyen R, *et al.* A novel common Kell antigen, TOU, and its spatial relationship to other Kell antigens. *Vox Sang* 1995;69:53–60.
- 189 Daniels GL, Petty A, Reid M, *et al.* Demonstration by the MAIEA assay that a new red cell antigen belongs to the Kell blood group system. *Transfusion* 1994;34:818–820.
- 190 Lee S, Debnath AK, Wu X, *et al.* Molecular basis of two novel high prevalence antigens in the Kell blood group system, KALT and KTIM. *Transfusion* 2006;46:1323–1327.
- 191 Fuchisawa A, Lomas-Francis C, Uchikawa M, *et al.* A new high prevalence Kell antigen, KYOR, is antithetical to the low-prevalence antigen KYO and the second trypsin-sensitive Kell system antigen. *Transfusion* 2012;52(Suppl. 3):158A [Abstract].
- 192 Vege S, Lomas-Francis C, Velliquette RW, *et al.* A new high prevalence antigen (KUHL) in the Kell blood group system. *Transfusion* 2011;51(Suppl.):25A [Abstract].
- 193 Velliquette RW, Sausais L, Lomas-Francis C, *et al.* Two novel and related high-prevalence antigens in the Kell blood group system. *Transfusion* 2007;47(Suppl.):164A [Abstract].
- 194 Karamatic Crew V, Poole J, Watson T, *et al.* KASH (KEL34): a novel high incidence antigen in the Kell blood group system. *Vox Sang* 2010;99(Suppl. 1):357 [Abstract].
- 195 Karamatic Crew V, Poole J, Bullock T, *et al.* KELP (KEL35): a new high incidence antigen in the Kell blood group system defined by two homozygous missense mutations in KEL. *Transfus Med* (2010);20(Suppl. 1):30 [Abstract].
- 196 Karamatic Crew V, Poole J, Bullock T, *et al.* KETI, a novel high incidence antigen in the Kell blood group system: a serological and molecular study. *Vox Sang* 2011;101(Suppl. 1):19 [Abstract].
- 197 Chown B, Lewis M, Kaita K. A 'new' Kell blood-group phenotype. *Nature* 1957;180:711.
- 198 Kaita H, Lewis M, Chown B, Gard E. A further example of the Kell blood group phenotype K<sup>–</sup>,k<sup>–</sup>,Kp(a<sup>–</sup>b<sup>–</sup>). *Nature* 1959;183:1586.
- 199 Chown B, Lewis M, Kaita H, Nevanlinna HR, Soltan HC. The pedigrees of two people already reported as of phenotype K<sup>–</sup>, k<sup>–</sup>, Kp(a<sup>–</sup>b<sup>–</sup>). *Vox Sang* 1961;6:620–623.
- 200 Yu L-C, Twu Y-C, Chang C-Y, Lin M. Molecular basis of the Kell-null phenotype. A mutation at the splice site of human *KEL* gene abolishes the expression of Kell blood group antigens. *J Biol Chem* 2001;276:10247–10252.
- 201 Lee S, Russo DCW, Reiner AP, *et al.* Molecular defects underlying the Kell null phenotype. *J Biol Chem* 2001;276:27281–27289.
- 202 Koda Y, Soejima M, Tsuneoka M, *et al.* Heterozygosity for two novel null alleles of the *KEL* gene causes the Kell-null phenotype in a Japanese woman. *Br J Haematol* 2002;117:220–225.
- 203 Wester ES, Storry JR, Schneider K, *et al.* Genetic basis of the K<sub>0</sub> phenotype in the Swedish population. *Transfusion* 2005;45:545–549.
- 204 Körmöcz GE, Wagner T, Jungbauer C, *et al.* Genetic diversity of KELnull and KELel: a nationwide Austrian survey. *Transfusion* 2007;47:703–714.
- 205 Yang Y, Wang LL, Wang C, *et al.* Two novel null alleles of the *KEL* gene detected in two Chinese women with the K<sub>null</sub> phenotype. *Transfus Med* 2009;19:235–244.
- 206 Blood Group Antigen Gene Mutation Database (dbRBC). <http://www.ncbi.nlm.nih.gov/projects/gv/mhc/xslcgi.cgi?cmd=bgmut/home> (last accessed 16 October 2012).
- 207 The International Society of Blood Transfusion Red Cell Immunogenetics and Blood Group Terminology Working Party. <http://www.isbtweb.org/working-parties/red-cell-immunogenetics-and-blood-group-terminology/> (last accessed 16 October 2012).

- 208 Marsh WL, Øyen R, Nichols ME, Allen FH. Chronic granulomatous disease and the Kell blood groups. *Br J Haematol* 1975;29:247–262.
- 209 Carbonnet F, Hattab C, Collec E, *et al.* Immunochemical analysis of the Kx protein from human red cells of different Kell phenotypes using antibodies raised against synthetic peptides. *Br J Haematol* 1997;96:857–863.
- 210 Marsh WL, Øyen R, Nichols ME. Kx antigen, the McLeod phenotype, and chronic granulomatous disease: further studies. *Vox Sang* 1976;31:356–362.
- 211 Marsh WL, Redman CM. Recent developments in the Kell blood group system. *Transfus Med Rev* 1987;1:4–20.
- 212 Karamatic Crew V, Poole J, Halverson G, Daniels G. The first example of a silent (synonymous) mutation causing aberrant splicing of KEL and silencing of the KEL gene. *Vox Sang* 2010;99(Suppl. 1):377 [Abstract].
- 213 Walsh TJ, Daniels GL, Tippet P. A family with unusual Kell genotypes. *Forens Sci Int* 1981;18:161–163.
- 214 Ford DS, Knight AE, Smith F. A further example of Kp<sup>a</sup>/K<sup>o</sup> exhibiting depression of some Kell group antigens. *Vox Sang* 1977;32:220–223.
- 215 Corcoran PA, Allen FH, Lewis M, Chown B. A new antibody, anti-Ku (anti-Peltz), in the Kell blood group system. *Transfusion* 1961;1:181–183.
- 216 Marsh WL, Redman CM. The Kell blood group system: a review. *Transfusion* 1990;30:158–167.
- 217 Nunn HD, Giles CM, Dormandy KM. A second example of anti-Ku in a patient who has the rare Kell phenotype, K<sup>o</sup>. *Vox Sang* 1966;11:611–619.
- 218 Lin M, Wang CL, Chen FS, Ho LH. Fatal hemolytic transfusion reaction due to anti-Ku in a K<sub>null</sub> patient. *Immunohematology* 2003;19:19–21.
- 219 Manoura A, Korakaki E, Hatzidaki E, *et al.* Use of recombinant erythropoietin for the management of severe hemolytic disease of the newborn of a K0 phenotype mother. *Pediatr Hematol Oncol* 2007;24:69–73.
- 220 Noumsi GT, Billingsley K, Moulds JM, Moulds JJ. Evaluation of the rare anti-Ku antibody using monocyte monolayer assay (MMA). *Transfusion* 2011;51(Suppl.):135A [Abstract].
- 221 Daniels GL. Studies on Gerbich negative phenotypes and Gerbich antibodies. *Transfusion* 1982;22:405 [Abstract].
- 222 Anstee DJ, Ridgwell K, Tanner MJA, Daniels GL, Parsons SF. Individuals lacking the Gerbich blood-group antigen have alterations in the human erythrocyte membrane sialoglycoproteins β and γ. *Biochem J* 1984;221:97–104.
- 223 Daniels GL, Shaw M-A, Judson PA, *et al.* A family demonstrating inheritance of the Leach phenotype: a Gerbich-negative phenotype associated with elliptocytosis. *Vox Sang* 1986;50:117–121.
- 224 Beck ML, Marsh WL, Pierce SR, *et al.* Auto-anti-Kp<sup>b</sup> associated with weakened antigenicity in the Kell blood group system: a second example. *Transfusion* 1979;19:197–202.
- 225 Marsh WL, Øyen R, Alicea E, Linter M, Horton S. Autoimmune hemolytic anemia and the Kell blood groups. *Am J Hematol* 1979;7:155–162.
- 226 Manny N, Levene C, Sela R, *et al.* Autoimmunity and the Kell blood groups: auto-anti-Kp<sup>b</sup> in a Kp(a+b-) patient. *Vox Sang* 1983;45:252–256.
- 227 Puig N, Carbonell F, Marty ML. Another example of mimicking anti-Kp<sup>b</sup> in a Kp(a+b-) patient. *Vox Sang* 1986;51:57–59.
- 228 Rowe GP, Tozzo GG, Poole J, Liew YW. The elucidation of a Kell-related autoantibody using ZZAP-treated red cells. *Immunohematology* 1989;5:79–82.
- 229 Vengelen-Tyler V, Gonzalez B, Garratty G, *et al.* Acquired loss of red cell Kell antigens. *Br J Haematol* 1987;65:231–234.
- 230 Williamson LM, Poole J, Redman C, *et al.* Transient loss of proteins carrying Kell and Lutheran red cell antigens during consecutive relapses of autoimmune thrombocytopenia. *Br J Haematol* 1994;87:805–812.
- 231 Daniels G. Kell related antibodies. *Rev Franc Transfus Immuno-Hémat* 1988;31:395–405.
- 232 Judson PA, Anstee DJ. Comparative effect of trypsin and chymotrypsin on blood group antigens. *Med Lab Sci* 1977;34:1–6.
- 233 Branch DR, Muensch HA, SySiok Hian AL, Petz LD. Disulfide bonds are a requirement for Kell and Cartwright (Yt<sup>a</sup>) blood group antigen integrity. *Br J Haematol* 1983;54:573–578.
- 234 Advani H, Zamor J, Judd WJ, Johnson CL, Marsh WL. Inactivation of Kell blood group antigens by 2-aminoethylisothiouonium bromide. *Br J Haematol* 1982;51:107–115.
- 235 Parsons SF, Judson PA, Spring FA, Mallinson G, Anstee DJ. Antibodies with specificities related to the Kell blood group system. *Rev Franc Transfus Immuno-Hémat* 1988;31:401–405.
- 236 McGinniss MH, Dean A. Expression of red cell antigens by K562 human leukemia cells before and after induction of hemoglobin synthesis by hemin. *Transfusion* 1985;25:105–109.
- 237 Russo D, Wu X, Redman CM, Lee S. Expression of Kell blood group protein in non-erythroid tissues. *Blood* 2000;96:340–346.
- 238 Redman CM, Lee S, Ten Bokkel Huinink D, *et al.* Comparison of human and chimpanzee Kell blood group systems. *Transfusion* 1989;29:486–490.
- 239 Blancher A, Reid ME, Socha WW. Cross-reactivity of antibodies to human and primate red cell antigens. *Transfus Med Rev* 2000;14:161–179.
- 240 Lee S, Russo DCW, Pu J, Ho M, Redman CM. The mouse Kell blood group gene (*Kel*): cDNA sequence, genomic organization, expression, and enzymatic function. *Immunogenetics* 2000;52:53–62.
- 241 Turner AJ, Isaac RE, Coates D. The neprilysin (NEP) family of zinc metalloproteases: genomics and function. *Bioessays* 2001;23:261–269.



- 242 Lee S, Lin M, Mele A, *et al.* Proteolytic processing of big endothelin-3 by the Kell blood group protein. *Blood* 1999;94:1440–1450.
- 243 Sha Q, Redman CM, Lee S. Endothelin-3-converting enzyme activity of the KEL1 and KEL6 phenotypes of the Kell blood group system. *J Biol Chem* 2006;281:7180–7182
- 244 Zhu X, Rivera A, Golub MS, *et al.* Changes in red cell ion transport, reduced intramural neovascularisation, and some mild motor function abnormalities accompany targeted disruption of the mouse Kell gene (*Kel*). *Am J Hematol* 2009;84:492–498.
- 245 Belhacène N, Maulon L, Guérin S, *et al.* Differential expression of the Kell blood group and CD10 antigens: two related membrane metallopeptidases during differentiation of K562 cells by phorbol ester and hemin. *FASEB J* 1998;12:531–539.
- 246 Redman CM, Marsh WL, Scarborough A, *et al.* Biochemical studies on McLeod phenotype red cells and isolation of Kx antigen. *Br J Haematol* 1988;68:131–136.
- 247 Ho M, Chelly J, Carter N, *et al.* Isolation of the gene for McLeod syndrome that encodes a novel membrane transport protein. *Cell* 1994;77:869–880.
- 248 Gether U, Andersen PH, Larsson OM, Schousboe A. Neurotransmitter transporters: molecular function of important drug targets. *Trends Pharmacol Sci* 2006;27: 375–383.
- 249 Stanfield GM, Horvitz HR. The *ced-8* gene controls the timing of programmed cell deaths in *C. elegans*. *Mol Cell* 2000;5:423–433.
- 250 Jung HH, Russo D, Redman C, Brandner S. Kell and XK immunohistochemistry in McLeod myopathy. *Muscle Nerve* 2001;24:1346–1351.
- 251 Clapéron A, Hattab C, Armand V, *et al.* The Kell and Xk proteins of the Kell blood group are not co-expressed in the central nervous system. *Brain Res* 2007;1147:12–24.
- 252 Branch DR, Gaidulis L, Lazar GS. Human granulocytes lack red cell Kx antigen. *Br J Haematol* 1986;62:747–755.
- 253 Collec E, Colin Y, Carbonnet F, *et al.* Structure and expression of the mouse homologue of the XK gene. *Immunogenetics* 1999;50:16–21.
- 254 Russo D, Lee S, Redman C. Intracellular assembly of Kell and XK blood group proteins. *Biochim Biophys Acta* 1999;1461:10–18.
- 255 Branch DR, Sy Siok Hian AL, Petz LD. Unmasking of Kx antigen by reduction of disulphide bonds on normal and McLeod red cells. *Br J Haematol* 1985;59:505–512.
- 256 Pu JJ, Redman CM, Visser JWM, Lee S. Onset of expression of the components of the Kell blood group system. *Transfusion* 2005;45:969–975.
- 257 Allen FH, Krabbe SMR, Corcoran PA. A new phenotype (McLeod) in the Kell blood-group system. *Vox Sang* 1961; 6:555–560.
- 258 Swash M, Schwartz MS, Carter ND, *et al.* Benign X-linked myopathy with acanthocytes (McLeod syndrome). Its relationship to X-linked muscular dystrophy. *Brain* 1983;106: 717–733.
- 259 Jung HH, Danek A, Walker RH. Neuroacanthocytosis syndromes. *Orphanet J Rare Dis* 2011;6:68.
- 260 Hewer E, Danek A, Schoser BG, *et al.* McLeod myopathy revisited: more neurogenic and less benign. *Brain* 2007; 130:3285–3296.
- 261 Wimer BM, Marsh WL, Taswell HF, Galey WR. Haematological changes associated with the McLeod phenotype of the Kell blood group system. *Br J Haematol* 1977;36: 219–224.
- 262 Ballas SK, Bator SM, Aubuchon JP, *et al.* Abnormal membrane physical properties of red cells in McLeod syndrome. *Transfusion* 1990;30:722–727.
- 263 Brzica SM, Rhodes KH, Pineda AA, Taswell HF. Chronic granulomatous disease and the McLeod phenotype. Successful treatment of infection with granulocyte transfusions resulting in subsequent hemolytic transfusion reaction. *Mayo Clin Proc* 1977;52:153–156.
- 264 Danek A, Rubio JP, Rampoldi L, *et al.* McLeod neuroacanthocytosis: genotype and phenotype. *Ann Neurol* 2001;50: 755–764.
- 265 Jung HH, Danek A, Frey BM. McLeod syndrome: a neurohaematological disorder. *Vox Sang* 2007;93:112–121.
- 266 Tang LL, Redman CM, Williams D, Marsh WL. Biochemical studies on McLeod phenotype erythrocytes. *Vox Sang* 1981;40:17–26.
- 267 Glaubenskleee CS, Evan AP, Galey WR. Structural and biochemical analysis of the McLeod erythrocyte membrane. I. Freeze fracture and discontinuous polyacrylamide gel electrophoresis analysis. *Vox Sang* 1982;42:262–271.
- 268 Galey WR, Evan AP, Van Nice PS, *et al.* Morphology and physiology of the McLeod erythrocyte. I. Scanning electron microscopy and electrolyte and water transport properties. *Vox Sang* 1978;34:152–161.
- 269 Kuypers FA, van Linde-Sibenius Trip M, Roelofsen B, *et al.* The phospholipid organisation in the membranes of McLeod and Leach phenotype erythrocytes. *FEBS Lett* 1985;184:20–24.
- 270 Redman CM, Huima T, Robbins E, Lee S, Marsh WL. Effect of phosphatidylserine on the shape of McLeod red cell acanthocytes. *Blood* 1989;74:1826–1835.
- 271 Ho ME, Chalmers RM, Davis MB, Harding AE, Monaco AP. A novel point mutation in the McLeod syndrome gene in neuroacanthocytosis. *Ann Neurol* 1996;39:672–675.
- 272 Russo DCW, Lee S, Reid ME, Redman CM. Point mutations causing the McLeod phenotype. *Transfusion* 2002;42: 287–293.
- 273 Walker RH, Danek A, Uttner I, *et al.* McLeod phenotype without McLeod syndrome. *Transfusion* 2007;47:299–305.
- 274 Jung HH, Hergersberg M, Vogt M, *et al.* McLeod phenotype associated with a XK missense mutation without hematology, neuromuscular, or cerebral involvement. *Transfusion* 2003;43:928–938.

- 275 Singleton BK, Green CA, Renaud S, *et al.* McLeod syndrome resulting from a novel XK mutation. *Transfus Med* 2003; 122:682–685.
- 276 Walker RH, Jung HH, Tison F, Lee S, Danek A. Phenotypic variation among brothers with the McLeod neuroacanthocytosis syndrome. *Movement Dis* 2007;22: 244–248.
- 277 Supple SG, Iland HJ, Barnett MH, Pollard JD. A spontaneous novel XK gene mutation in a patient with McLeod syndrome. *Br J Haematol* 2001;115:369–372.
- 278 Daniels GL, Weinauer F, Stone C, *et al.* A combination of effects of rare genotypes at the XK and KEL blood group loci results in absence of Kell system antigens from the red blood cells. *Blood* 1996;88:4045–4050.
- 279 Stasia MJ, Li XJ. Genetics and immunopathology of chronic granulomatous disease. *Semin Immunopathol* 2008;30: 209–235.
- 280 Francke U, Ochs HD, de Martinville B, *et al.* Minor Xp21 chromosome deletion in a male associated with expression of Duchenne muscular dystrophy, chronic granulomatous disease, retinitis pigmentosa, and McLeod syndrome. *Am J Hum Genet* 1985;37:250–267.
- 281 Frey D, Mächler M, Seger R, Schmid W, Orkin SH. Gene deletion in a patient with chronic granulomatous disease and McLeod syndrome: fine mapping of the Xk gene locus. *Blood* 1988;71:252–255.
- 282 de Saint-Basile G, Bohler MC, Fischer A, *et al.* Xp21 DNA microdeletion in a patient with chronic granulomatous disease, retinitis pigmentosa, and McLeod phenotype. *Hum Genet* 1988;80:85–89.
- 283 Bertelson CJ, Pogo AO, Chaudhuri A, *et al.* Localization of the McLeod locus (XK) within XP21 by deletion analysis. *Am J Hum Genet* 1988;42:703–711.
- 284 Peng J, Redman C, Wu X, *et al.* Insights into extensive deletions around the XK locus associated with McLeod phenotype and characterization of two novel cases. *Gene* 2007;392:142–150.
- 285 Øyen R, Reid ME, Rubenstein P, Ralph H. A method to detect McLeod phenotype red blood cells. *Immunohematology* 1996;12:160–163.
- 286 van der Hart M, Szaloky A, van Loghem JJ. A ‘new’ antibody associated with the Kell blood group system. *Vox Sang* 1968;15:456–458.
- 287 Marsh WL. Letter. *Vox Sang* 1979;36:375–376.
- 288 Giblett ER, Klebanoff SJ, Pincus SH, *et al.* Kell phenotypes in chronic granulomatous disease: a potential transfusion hazard. *Lancet* 1971;i:1235–1236.
- 289 Carstairs K, Rutledge Harding S, Lue K, Johnson CL, Marsh WL. Anti-K9 in an untransfused McLeod individual. *Transfusion* 1988;28(Suppl.):20S [Abstract].
- 290 White W, Washington ED, Sabo BH, *et al.* Anti-Km in a transfused man with McLeod syndrome. *Rev Franc Transfus Immuno-Hémat* 1980;23:305–317.
- 291 Sharp D, Rogers S, Dickstein B, Johnson C, Toy E. Successful transfusion of K<sub>0</sub> blood to a Km–Kx– patient with anti-Km. *Transfusion* 1988;28(Suppl.):37S [Abstract].
- 292 Russo DC, Øyen R, Powell VI, *et al.* First example of anti-Kx in a person with the McLeod phenotype and without chronic granulomatous disease. *Transfusion* 2000;40: 1371–1375.
- 293 Mathur G, Sutor LJ, Kresie L, *et al.* Rare anti-Kx in a McLeod phenotype patient without chronic granulomatous disease. *Transfusion* 2011;51(Suppl.):135A [Abstract].
- 294 Bansal I, Jeon H-R, Hui SR, *et al.* Transfusion support for a patient with McLeod phenotype without chronic granulomatous disease and with antibodies to Kx and Km. *Vox Sang* 2008;94:216–220.
- 295 Sullivan CM, Kline WE, Rabin BI, Johnson CL, Marsh WL. The first example of autoanti-Kx. *Transfusion* 1987;27: 322–324.