

# 31 Sid Antigen

31.1 Introduction, 505  
31.2 Sd<sup>a</sup> and Cad, 505  
31.3 Sid antibodies and lectins, 507

31.4 Biochemistry, 507  
31.5 Sd<sup>a</sup> and gastrointestinal cancer, 509  
31.6 Malaria, 509

## 31.1 Introduction

Numerous antibodies of identical specificity were studied in several laboratories for at least 10 years before Macvie *et al.* [1] and Renton *et al.* [2] simultaneously reported the specificity in 1967 and named it anti-Sd<sup>a</sup>. About 91% of white people have Sd(a+) red cells and so Sid qualifies for inclusion in the 901 series of high frequency antigens and has the number 901011. As Sid differs so significantly, both in frequency and characteristics, from the other 901 series antigens, it has been given a separate chapter. Although the gene encoding the  $\beta$ 1,4-*N*-acetylgalactosa minyltransferase responsible for Sd<sup>a</sup> expression has been isolated, Sid has not been made a blood group system because the genetic basis of the Sid polymorphism has not been elucidated.

## 31.2 Sd<sup>a</sup> and Cad

### 31.2.1 Variation in strength of Sd<sup>a</sup>: the Sd(a++) or Cad phenotype

The strength of expression of Sd<sup>a</sup> on Sd(a+) cells varies from barely recognisable to extremely strong. A 'mixed field' reaction, with small agglutinates among a sea of unagglutinated cells, is a characteristic of reactions with anti-Sd<sup>a</sup>. If the unagglutinated cells are separated and retested with anti-Sd<sup>a</sup>, a 'mixed field' picture is again apparent [2]. In haemagglutination tests with a strong anti-Sd<sup>a</sup>, about 1% of individuals have strong agglutination with big clumps of cells, about 80% have moderate sized agglutinates consisting of 10–20 cells, about 10%

have only occasional tiny agglutinates, and about 9% have no agglutination [1]. Weak examples of anti-Sd<sup>a</sup> react with only a few Sd(a+) samples and may be mistaken for antibodies to a low frequency antigen.

In 1968, Cazal *et al.* [3] described a new private antigen, Cad, in a Mauritian family. Cells of the Cad+ members were polyagglutinable and, despite being group O or B, were agglutinated by *Dolichos biflorus* seed extract, a lectin that generally only agglutinates A<sub>1</sub> cells (Chapter 2). After demonstrating that red cells with very strong Sd<sup>a</sup> expression are also agglutinated by *Dolichos* lectin and that Cad cells are very strongly Sd(a+), Sanger *et al.* [4] concluded that Cad represents the high end of a continuous curve of Sd<sup>a</sup> antigen strength. There is considerable variation in agglutination strength with *Dolichos* and with human anti-Sd<sup>a</sup> of these 'super-Sid' or Sd(a++) cells between individuals and within families [5–11]. As Cad probably represents the expression of an extra strong Sd<sup>a</sup> gene, the term Sd(a++) will be used here to describe all phenotypes in which the red cells have sufficiently strong Sd<sup>a</sup> to be agglutinated by *Dolichos* lectin. The term Cad will be restricted to describe the rare phenotype of members of the Cad family, still probably the strongest Sd(a++) cells encountered.

Sd(a++) cells of the Cad family are polyagglutinable [3]; that is, they are agglutinated by most human sera. Although other Sd(a++) cells are generally polyagglutinable, this is usually less obvious and selected sera and sensitive methods are required to detect the agglutination [9,10]. Sd(a++) polyagglutination differs from all other forms of polyagglutination (Chapter 33).

Weak anti-Sd<sup>a</sup> appears to be present in the sera of most people and responsible for the polyagglutination

**Table 31.1** Frequencies of Sd(a++) in various populations as detected by different preparations of *Dolichos biflorus* lectin.

Population	No. tested	No. positive	Frequency	References
French, Montpellier	250 000	0		[3]
French (four centres)	78 526	56	0.0007	[10]
Canadian, Winnipeg	1425	2	0.0014	[8]
Canadian, Toronto	2191	1	0.0005	[5]
Japanese	51 420	15	0.0003	[7]
Hong Kong Chinese	36 037	110	0.0031	[18]
Thai	14 261	37	0.0026	[12,19]

of Sd(a++) cells. Polyagglutinable Sd(a++) cells are not agglutinated by sera from people with Sd(a++) cells and the stronger the Sd<sup>a</sup> antigen on Sd(a++) cells, the greater the number of ‘normal’ sera that would agglutinate them [12].

**31.2.2 Sd<sup>a</sup> in body fluids and other tissues**

Sd<sup>a</sup> can be detected, by haemagglutination inhibition, in the saliva of individuals with Sd(a+) red cells, with greater quantities present in neonatal than adult saliva [1,13]. Sd<sup>a</sup> can also be detected in human serum and milk, but by far the greatest abundance is found in meconium and urine [13]. Inhibition of anti-Sd<sup>a</sup> with urine is considered the most reliable method of determining Sd<sup>a</sup> phenotype.

Sd<sup>a</sup> has been found, in variable quantities, in the urine of 12 species of mammals, but not in birds [13]. Guinea pig urine is an extremely concentrated source of Sd<sup>a</sup> and a useful tool in blood group antibody identification.

Of the human tissues, kidney is a very rich source of Sd<sup>a</sup> [13], with activity localised to the distal convoluted tubules and collecting ducts [14]. Sd<sup>a</sup> is also present in colon and stomach, but was not detected in small intestine, muscle, liver, spleen, or brain [15,16].

**31.2.3 Frequency and inheritance**

The frequency of Sd<sup>a</sup>, as determined by testing red cells, is about 90%. Three surveys gave the following results: 290 English blood donors, 91.4% Sd(a+) [1]; 131 English donors, 91.0% Sd(a+) [2]; 1307 Italian donors, 89.3% Sd(a+) [17]. A higher incidence of Sd(a+) is obtained when urine is tested by haemagglutination inhibition. Sd<sup>a</sup> was detected in the urine of 96.1% of English blood donors [13] and 93.4% of north Italians [17]. The gene

and genotype frequencies in English donors determined by urine inhibition are: Sd<sup>a</sup> 0.8030; Sd 0.1970; Sd<sup>a</sup>/Sd<sup>a</sup> 0.6448; Sd<sup>a</sup>/Sd 0.3164; Sd/Sd 0.0388.

Sd(a++) is very rare in Europeans, but may be less infrequent in the Far East. Table 31.1 shows some frequency determinations on group O and B donors tested with *Dolichos* lectin. These frequencies are somewhat arbitrary as they are partly dependent on the potency of the lectin preparation used.

Tests with anti-Sd<sup>a</sup> on 55 families with a total of 168 children, demonstrated that Sd<sup>a</sup> is inherited as a dominant character [1]. Sd(a++), as detected by *Dolichos* lectin, is also inherited in a dominant manner [3–8,10–12,20]. Family studies have shown that Sd<sup>a</sup> is not part of the ABO, MNS, P1PK, Rh, Lutheran, Kell, Duffy, Kidd, Xg, Dombrock, or XK systems [1,5,8,10]. One Canadian family provided a suggestion of linkage between the genes for Sd(a++) and Wr<sup>a</sup> (*SLC4A1* or *DI*), with six non-recombinants and no recombinants, providing a lod score of 1.806 at a recombination fraction (θ) of zero [8]. Both *SLC4A1* and *B4GALNT2*, the Sd<sup>a</sup> gene (Section 31.4.3) are on the long arm of chromosome 17.

**31.2.4 Sd<sup>a</sup> in babies and pregnant women**

Red cells of fetuses and newborns are Sd(a–) and become Sd(a+) about 10 weeks post-natally [1,2,15]. Red cells of the son of an Sd(a++) father were non-reactive with *Dolichos* lectin at birth, but were agglutinated by the lectin 6 months later [10]. Saliva, urine, and meconium from neonates generally contain abundant Sd<sup>a</sup> substance [13].

An increased frequency of Sd(a–) red cell phenotype is found among pregnant women [1,15,21]. In different studies, 36% and 75% were Sd(a–) at term, and 25% six weeks post-partum had Sd(a–) red cells [15,21]. Sd<sup>a</sup> is present in the urine of most Sd(a–) pregnant women [13].

### 31.3 Sid antibodies and lectins

#### 31.3.1 Anti-Sd<sup>a</sup>

Anti-Sd<sup>a</sup> is disclosed in about 1% of donor sera when Sd(a+) cells are used for detection [1,2]. The incidence of anti-Sd<sup>a</sup> is increased when Sd(a++) cells are used [21], and the exceptionally strong Sd(a++) Cad cells are agglutinated by most sera. Sd<sup>a</sup> antibodies are usually IgM and active at 20°C [1,2,15], although IgG anti-Sd<sup>a</sup> has been identified [21,22].

Anti-Sd<sup>a</sup> are not generally considered a transfusion hazard [1,15,21,23], though transfusion of Sd(a+) red cells to patients with anti-Sd<sup>a</sup> may evoke a significant increase in IgG antibody titre [21]. Most survival studies with <sup>51</sup>Cr-labelled cells in patients with anti-Sd<sup>a</sup> have shown insignificant reduction in red cell survival, even when Sd(a++) cells were used [24,25]. A couple of HTRs, allegedly caused by transfusion of Sd(a++) cells to a patient with IgM anti-Sd<sup>a</sup>, are reported [24,25].

#### 31.3.2 Lectins and other non-human sources of anti-Sd<sup>a</sup>

A single lectin in *Dolichos biflorus* extracts specific for terminal GalNAc agglutinates A<sub>1</sub>, Tn-positive, and Sd(a++) red cells [26]. These reactions can be inhibited by GalNAc [4]. Sd(a++) cells express more determinants than A<sub>1</sub> or Tn cells, so an apparently specific anti-Sd(a++) can be produced by dilution of the lectin or by partial adsorption with A<sub>1</sub> or Tn cells; further adsorptions, or adsorption with Sd(a++) cells, removes all activity.

The seed lectins *Salvia horminum*, *S. farinacea*, and *Leonurus cardiaca* [27–29], and the snail lectins *Helix pomatia* and *H. aspersa* [6,30–33] agglutinate Sd(a++) cells and these reactions are inhibited by GalNAc. Sd(a++) and Tn specificity in the *Salvia* lectins can be separated by adsorption [27]. The anti-A<sub>1</sub> lectins *Phaseolus lunatus* and *P. limensis* do not agglutinate even the strongest of Sd(a++) cells [30–32].

Chicken serum contains a ‘naturally-occurring’ antibody reactive with Sd(a++) cells [34]. Separable anti-Tn and an antibody that agglutinates Sd(a++) cells were identified in the serum of the snake *Python sebae* [35].

### 31.4 Biochemistry

#### 31.4.1 Sd<sup>a</sup> in urine







In 1970, Morton and Terry [36] partially separated Sd<sup>a</sup>-active material from human urine by ethanol precipita-

tion. Ten years later, Soh *et al.* [37] showed that Sd<sup>a</sup> activity is associated with Tamm–Horsfall protein, now usually called uromodulin, the most abundant glycoprotein in urine [38]. Uromodulin is secreted by the kidney and protects against urinary tract infection (UTI). It is composed of 70% protein and 30% carbohydrate, with eight potential *N*-glycosylation sites. Uromodulin in urine is a key anti-adherence factor for *Escherichia coli*, the principal cause of UTI. Mutations in *UMOD*, the gene encoding the uromodulin protein, lead to rare autosomal dominant diseases, referred to as uromodulin-associated kidney disease, and uromodulin is also a risk factor for chronic kidney disease [38].

Sd(a+) uromodulin contains between 1–2% GalNAc, whereas that from Sd(a–) individuals contains virtually none [37,39]. Sd<sup>a</sup>-active uromodulin can be precipitated by *Dolichos* and *Helix pomatia* lectins and by human anti-Sd<sup>a</sup> [39,40]. A pentasaccharide isolated from Sd<sup>a</sup>-active uromodulin is shown in Table 31.2b [41,42]. This pentasaccharide strongly inhibits the reactivity of anti-Sd<sup>a</sup> with Sd(a+) cells and of *Dolichos* lectin with Sd(a++) cells. The pentasaccharide is not present on uromodulin from Sd(a–) urine, whereas a tetrasaccharide lacking the GalNAc in  $\beta$ -linkage with Gal (Table 31.2a), is present on both Sd<sup>a</sup>-active and -inactive glycoproteins [43]. Tetrasaccharides prepared by the cleavage of GalNAc or sialic acid residues from the Sd<sup>a</sup>-active pentasaccharides lack Sd<sup>a</sup> activity [44]. A mucin with high GalNAc content isolated from urine with *H. pomatia* lectin inhibited anti-Sd<sup>a</sup> as effectively as uromodulin [45], so at least two substances in human urine carry the Sd<sup>a</sup> determinant.

#### 31.4.2 Sd<sup>a</sup> on red cells

Most biochemical analysis of Sd<sup>a</sup> on red cells has been carried out on the Sd(a++) cells of members of the Cad family, which have exceptionally high levels of Sd<sup>a</sup> activity. SDS PAGE revealed that glycophorin A (GPA) and glycophorin B (GPB) from Cad red cells have an increase in apparent MW of 3 and 2 kDa, respectively, compared with normal red cells [46]. This results from the addition of a  $\beta$ -linked GalNAc residue to most of the disialotetrasaccharides characteristic of these glycoproteins [47] (Table 31.2c and d, and Chapter 3). The resulting pentasaccharide (Table 31.2d) inhibited *Dolichos* lectin and anti-Sd<sup>a</sup>, and the altered GPA molecules bound to the lectin in affinity columns [46–48]. Red cells from the original Cad propositus had approximately 12 pentasaccharides per GPA molecule, cells from two other Sd(a++) individuals had 2–3 pentasaccharides per GPA molecule, and Sd(a+) and Sd(a–) cells had no pentasaccharides

**Table 31.2** Some structures associated with different Sd<sup>a</sup> phenotypes (for abbreviations see Table 2.4).**Tetrasaccharide and pentasaccharide isolated from uromodulin**(a) Sd(a<sup>-</sup>) and Sd(a<sup>+</sup>)(b) Sd(a<sup>+</sup>)**Tetrasaccharide and pentasaccharide from glycophorin A**(c) Sd(a<sup>-</sup>), Sd(a<sup>+</sup>), and Sd(a<sup>++</sup>)(d) Sd(a<sup>++</sup>)**Sialosylparagloboside and Sd<sup>a</sup> ganglioside**(e) Sd(a<sup>-</sup>), Sd(a<sup>+</sup>), and Sd(a<sup>++</sup>)(f) Sd(a<sup>+</sup>), and Sd(a<sup>++</sup>)

of a new ganglioside of lower mobility [51]. The unusual ganglioside, which binds *H. pomatia* lectin and inhibits anti-Sd<sup>a</sup>, represents sialosylparagloboside with an additional terminal  $\beta$ -GalNAc residue (Table 31.2f) [51,52]. The structure of the terminal pentasaccharide of this ganglioside is identical to that of Sd<sup>a</sup>-active uromodulin shown in Table 31.2b [52]. Small quantities of this ganglioside were detected in membranes from Sd(a<sup>+</sup>), but not Sd(a<sup>-</sup>) red cells, and might represent the major Sd<sup>a</sup>-active structure in Sd(a<sup>+</sup>) red cells [51].

**31.4.3 The Sd<sup>a</sup> glycosyltransferase and the gene that encodes it**

The product of the *Sd<sup>a</sup>* gene is a  $\beta$ 1,4-*N*-acetylgalactosaminyltransferase that catalyses the transfer of GalNAc in  $\beta$ -linkage from a nucleotide carrier to an appropriate tetrasaccharide acceptor substrate molecule. Such enzymes have been detected in human urine from Sd(a<sup>+</sup>) but not Sd(a<sup>-</sup>) individuals [43,53], in human kidney [54], in human large intestine [55], and in guinea pig kidney [56,57]. Sd(a<sup>-</sup>) uromodulin was the best acceptor for the human urine enzyme, but the disialotetrasaccharides of GPA were also suitable acceptors [53]. Preparations from human kidney catalysed the transfer of GalNAc to sialosylparagloboside, but not to native GPA, although tryptic peptides of GPA were good acceptor substrates [51,54].

In 2003, two groups exploited the sequence of a mouse  $\beta$ 1,4-*N*-acetylgalactosaminyltransferase gene to isolate human Sd<sup>a</sup> cDNA from an expressed sequence tag database and from colon carcinoma (Caco2) cells [58,59]. The Sd<sup>a</sup> gene, *B4GALNT2*, is on chromosome 17q23.1 and consists of at least 12 exons encoding two proteins of 506 and 566 amino acids, identical in their sequence with the exception of their cytoplasmic tail. Through alternative splicing of one of two exons 1 most of either a very long (exon 1<sub>L</sub>) or very short (exon 1<sub>S</sub>) cytoplasmic N-terminal domain is encoded; exon 2 encodes the membrane-spanning domain and the stem region; exons 3–11 the putative catalytic site. Transient transfection of Cos-7 cells with the catalytic domain of the enzyme produced a soluble form of the protein that catalysed the transfer of GalNAc to  $\alpha$ 2,3-sialylated acceptor substrates to form the GalNAc $\beta$ 1-4(Neu5Ac $\alpha$ 2-3)Gal $\beta$ 1-R trisaccharide common to Sd<sup>a</sup> antigens [58]. The short form of the enzyme is associated with a higher level of activity [60]. Northern blot analysis indicated that five transcripts were highly expressed in colon and to a lesser extent in kidney, stomach, ileum, and rectum [58].

[49]. Increased apparent MW of decay-accelerating factor (CD55) in Cad cells was also assumed to result from the presence of additional GalNAc residues [50].

Thin layer chromatography of Cad red cell membranes demonstrated an unusual profile characterised by reduction in the content of sialosylparagloboside (Table 31.2e), the major ganglioside in normal cells, and the presence

### 31.5 Sd<sup>a</sup> and gastrointestinal cancer

Sd<sup>a</sup> antigen is expressed abundantly in normal gastrointestinal mucosae [14,61], but neither Sd<sup>a</sup> antigen nor the Sd<sup>a</sup>-transferase is detected in gastrointestinal cancer cells [14,55]. Transfection of colonic or gastric cell lines with the Sd<sup>a</sup>-transferase cDNA resulted in increased cell surface Sd<sup>a</sup> expression with concomitant loss of the E-selectin ligands sialyl-Le<sup>x</sup> (sLe<sup>x</sup>) and sialyl-Le<sup>a</sup> (sLe<sup>a</sup>) (Section 2.3.4), and significant decreased adhesion to activated human umbilical vein endothelial cells [62]. Kawamura *et al.* [62] propose that the Sd<sup>a</sup>-transferase competes with fucosyltransferases, blocking expression of sLe<sup>x</sup> and sLe<sup>a</sup> biosynthesis and eliminating metastasis. From experiments on colorectal cancer specimens, Malagolini *et al.* [60] found that Sd<sup>a</sup> and sLe<sup>x</sup> are expressed on different proteins and consider that the effect described by Kawamura *et al.* [62] is an artefact of the experimental model and not applicable to colon cancer tissues.

### 31.6 Malaria

Sd(a++) cells of the Cad family are relatively resistant to invasion by the malarial parasite *Plasmodium falciparum* [63] (see Section 3.21.1). Sialic acid on glycoprotein molecules is required by *P. falciparum* merozoites for invasion of the red cells and this sialic acid might not be available on the O-glycans of Cad cells because of chemical interaction with the acetyl group of the extra GalNAc residues [63].

### References

- Macvie SI, Morton JA, Pickles MM. The reactions and inheritance of a new blood group antigen, Sd<sup>a</sup>. *Vox Sang* 1967;13:485–492.
- Renton PH, Howell P, Ikin EW, Giles CM, Goldsmith KLG. Anti-Sd<sup>a</sup>, a new blood group antibody. *Vox Sang* 1967;13:493–501.
- Cazal P, Monis M, Caubel J, Brives J. Polyagglutinabilité héréditaire dominante: antigène privé (Cad) correspondant à un anticorps public et à une lectine de *Dolichos biflorus*. *Rev Franc Transfus* 1968;11:209–221.
- Sanger R, Gavin J, Tippet P, Teesdale P, Eldon K. Plant agglutinin for another human blood-group. *Lancet* 1971;ii:1130.
- Race RR, Sanger R. *Blood Groups in Man*, 6th edn. Oxford: Blackwell Scientific Publications, 1975.
- Cazal P, Monis M, Bizot M. Les antigènes Cad et leurs rapports avec les antigènes A. *Rev Franc Transfus* 1971;14:321–334.
- Yamaguchi H, Okubo Y, Ogawa Y, Tanaka M. Japanese families with group O and B red cells agglutinable by *Dolichos biflorus* extract. *Vox Sang* 1973;25:361–369.
- Lewis M, Kaita H, Chown B, *et al.* A family with the rare red cell antigens Wr<sup>a</sup> and 'super' Sd<sup>a</sup>. *Vox Sang* 1973;25:336–340.
- Lopez M, Gerbal A, Bony V, Salmon C. Cad antigen: comparative study of 50 samples. *Vox Sang* 1975;28:305–313.
- Gerbal A, Lopez M, Chassaigne M, *et al.* L'antigène Cad dans la population française. *Rev Franc Transfus Immuno-Hémat* 1976;14:415–429.
- Cazal P, Monis M, Bizot M. Les antigènes Cad en 1976. *Rev Franc Transfus Immuno-Hémat* 1977;20:165–173.
- Sringarm S, Chiewsilp P, Tubrod J. Cad receptor in Thai blood donors. *Vox Sang* 1974;26:462–466.
- Morton JA, Pickles MM, Terry AM. The Sd<sup>a</sup> blood group antigen in tissues and body fluids. *Vox Sang* 1970;19:472–482.
- Morton JA, Pickles MM, Vanhegan RI. The Sd<sup>a</sup> antigen in the human kidney and colon. *Immunol Invest* 1988;17:217–224.
- Pickles MM, Morton JA. The Sd<sup>a</sup> blood group. In: *Human Blood Groups, 5th Int Convoc Immunol*, Buffalo NY. Basel: Karger, 1977:277–286.
- Piller F, Cartron J-P, Tuppy H. Increase of blood group A and loss of blood group Sd<sup>a</sup> activity in the mucus from human neoplastic colon. *Rev Franc Transfus Immuno-Hémat* 1980;23:599–611.
- Conte R, Serafini-Cessi F. Comparison between the erythrocyte and urinary Sd<sup>a</sup> antigen distribution in a large number of individuals from Emilia-Romagna, a region of northern Italy. *Transfus Med* 1991;1:47–49.
- Mak KH, Leong S, Chan NK. Incidence of Cad antigen among Chinese donors in Hong Kong. *20th Congr Int Soc Blood Transfus*, 1988:303 [Abstracts].
- Sringarm S, Chupungart C, Giles CM. The use of *Ulex europaeus* and *Dolichos biflorus* extracts in routine ABO grouping of blood donors in Thailand. Some unexpected findings. *Vox Sang* 1972;23:537–545.
- Lopez M, Gerbal A, Girard-Debord M, Salmon C. Trois sujets A<sub>end</sub> Cad dans une famille française. *Rev Franc Transfus Immuno-Hémat* 1977;20:457–466.
- Spitalnik S, Cox MT, Spennacchio J, Guenther R, Blumberg N. The serology of Sd<sup>a</sup> effects of transfusion and pregnancy. *Vox Sang* 1982;42:308–312.
- Silvergleid AJ, Wells RF, Hafleigh EB, *et al.* Compatibility test using <sup>51</sup>chromium-labeled red blood cells in crossmatch positive patients. *Transfusion* 1978;18:8–14.
- Colledge KI, Kaplan HS, Marsh WL. Massive transfusion of Sd(a+) blood to a recipient with anti-Sd<sup>a</sup>, without clinical complication. *Transfusion* 1973;13:340 [Abstract].



- 24 Peetermans ME, Cole-Dergent J. Haemolytic transfusion reaction due to anti-Sd<sup>a</sup>. *Vox Sang* 1970;18:67–70.
- 25 Reznicek MJ, Cordle DG, Strauss RG. A hemolytic reaction implicating Sd<sup>a</sup> antibody missed by immediate spin cross-match. *Vox Sang* 1992;62:173–175.
- 26 Bird GWG, Wingham J. Cad (super Sd<sup>a</sup>) in a British family with Eastern connections: a note on the specificity of the *Dolichos biflorus* lectin. *J Immunogenet* 1976;3:297–302.
- 27 Bird GWG, Wingham J. Haemagglutinins from *Salvia*. *Vox Sang* 1974;26:163–166.
- 28 Moore BPL, Marsh S. Identification of strong Sd(a+) and Sd(a++) red cells by hemagglutinins from *Salvia horminum*. *Transfusion* 1975;15:132–134.
- 29 Bird GWG, Wingham J. Anti-Cad lectin from the seeds of *Leonurus cardiaca*. *Clin Lab Haematol* 1979;1:57–59.
- 30 Myllylä G, Furuhjelm U, Nordling S, *et al.* Persistent mixed field polyagglutinability. Electrokinetic and serological aspects. *Vox Sang* 1971;20:7–23.
- 31 Bird GWG, Wingham J. Some serological properties of the Cad receptor. *Vox Sang* 1971;20:55–61.
- 32 Uhlenbruck G, Sprenger I, Heggen M, Leseney AM. Diagnosis of the 'Cad' blood group with agglutinins from snails and plants. *Z Immun Forsch* 1971;141:290–291.
- 33 Bizot M. Comportement de quelques extraits de gastéropodes terrestres vis-à-vis des substances de spécificité Sd<sup>a</sup>. *Rev Franc Transfus* 1972;15:371–375.
- 34 Bizot M, Cayla J-P. Hétéro-anticorps anti-Cad du poulet. *Rev Franc Transfus* 1972;15:195–202.
- 35 Lockyer WJ, Gold ER, Bird GWG. Anti-Tn and anti-Cad in the serum of the non-poisonous snake *Python sebae*. *Med Lab Sci* 1977;34:311–317.
- 36 Morton JA, Terry AM. The Sd<sup>a</sup> blood group antigen. Biochemical properties of urinary Sd<sup>a</sup>. *Vox Sang* 1970;19:151–161.
- 37 Soh CPC, Morgan WTJ, Watkins WM, Donald ASR. The relationship between the *N*-acetylgalactosamine content and the blood group Sd<sup>a</sup> activity of Tamm and Horsfall urinary glycoprotein. *Biochem Biophys Res Comm* 1980;93:1132–1139.
- 38 Rampoldi L, Scolari F, Amoroso A, Ghiggeri G, Devuyt O. The rediscovery of uromodulin (Tamm–Horsfall protein): from tubulointerstitial nephropathy to chronic kidney disease. *Kidney Int* 2011;80:338–347.
- 39 Morgan WTJ, Soh CPC, Donald ASR, Watkins WM. Observations on the blood group Sd<sup>a</sup> activity of Tamm–Horsfall urinary glycoprotein. *Rev Franc Transfus Immuno-Hémat* 1981;24:37–51.
- 40 Serafini-Cessi F, Conte R. Precipitin reaction between Sd<sup>a</sup>-active human Tamm–Horsfall glycoprotein and anti-Sd<sup>a</sup> serum. *Vox Sang* 1982;42:141–144.
- 41 Donald ASR, Yates AD, Soh CPC, Morgan WTJ, Watkins WM. A blood group Sd<sup>a</sup>-active pentasaccharide isolated from Tamm–Horsfall urinary glycoprotein. *Biochem Biophys Res Comm* 1983;115:625–631.
- 42 Donald ASR, Feeney J. Oligosaccharides obtained from a blood-group-Sd<sup>a+</sup> Tamm–Horsfall glycoprotein. *Biochem J* 1986;236:821–828.
- 43 Donald ASR, Soh CPC, Yates AD, *et al.* Structure, biosynthesis and genetics of the Sd<sup>a</sup> antigen. *Biochem Soc Trans* 1987;15:606–608.
- 44 Donald ASR, Yates AD, Soh CPC, Morgan WTJ, Watkins WM. The human blood-group-Sd<sup>a</sup> determinant: a terminal non-reducing carbohydrate structure in *N*-linked and mucin-type glycoproteins. *Biochem Soc Trans* 1984;12:596–599.
- 45 Cartron J-P, Kornprobst M, Lemonnier M, *et al.* Isolation from human urines of a mucin with blood group Sd<sup>a</sup> activity. *Biochem Biophys Res Comm* 1982;106:331–337.
- 46 Cartron J-P, Blanchard D. Association of human erythrocyte membrane glycoproteins with blood-group Cad specificity. *Biochem J* 1982;207:497–504.
- 47 Blanchard D, Cartron J-P, Fournet B, *et al.* Primary structure of the oligosaccharide determinant of blood group Cad specificity. *J Biol Chem* 1983;258:7691–7695.
- 48 Herkt F, Parente JP, Leroy Y, *et al.* Structure determination of oligosaccharides isolated from Cad erythrocyte membranes by permethylation analysis and 500-MHz <sup>1</sup>H-NMR spectroscopy. *Eur J Biochem* 1985;146:125–129.
- 49 Blanchard D, Capon C, Leroy Y, Cartron J-P, Fournet B. Comparative study of glycophorin A derived O-glycans from human Cad, Sd(a+) and Sd(a-) erythrocytes. *Biochem J* 1985;232:813–818.
- 50 Spring FA, Judson PA, Daniels GL, *et al.* A human cell-surface glycoprotein that carries Cromer-related blood group antigens on erythrocytes and is also expressed on leucocytes and platelets. *Immunology* 1987;62:307–313.
- 51 Blanchard D, Piller F, Gillard B, Marcus D, Cartron J-P. Identification of a novel ganglioside on erythrocytes with blood group Cad specificity. *J Biol Chem* 1985;260:7813–7816.
- 52 Gillard BK, Blanchard D, Bouhours J-F, *et al.* Structure of a ganglioside with Cad blood group antigen activity. *Biochemistry* 1988;27:4601–4606.
- 53 Serafini-Cessi F, Malagolini N, Dall'Olio F. Characterization and partial purification of β-*N*-acetylgalactosaminyltransferase from urine of Sd<sup>a+</sup> individuals. *Arch Biochem Biophys* 1988;266:573–582.
- 54 Piller F, Blanchard D, Huet M, Cartron J-P. Identification of a α-NeuAc-(2→3)-β-D-galactopyranosyl *N*-acetyl-β-D-galactosaminyltransferase in human kidney. *Carbohydrate Res* 1986;149:171–184.
- 55 Malagolini N, Dall'Olio F, Di Stefano G, *et al.* Expression of UDP-GalNAc:NeuAcα2,3Galβ-R β1,4(GalNAc to Gal) *N*-acetylgalactosaminyltransferase involved in the synthesis of Sd<sup>a</sup> antigen in human large intestine and colorectal carcinomas. *Cancer Res* 1989;49:6466–6470.
- 56 Serafini-Cessi F, Dall'Olio F. Guinea-pig kidney β-*N*-acetylgalactosaminyltransferase towards Tamm–Horsfall glyco-

- protein. Requirement of sialic acid in the acceptor for transferase activity. *Biochem J* 1983;215:483–489.
- 57 Soh CPC, Donald ASR, Feeney J, Morgan WTJ, Watkins WM. Enzymic synthesis, chemical characterisation and Sd<sup>a</sup> activity of GalNAc $\beta$ 1-4[NeuAc $\alpha$ 2-3]Gal $\beta$ 1-4GlcNAc and GalNAc $\beta$ 1-4[NeuAc $\alpha$ 2-3]Gal $\beta$ 1-4Glc. *Glycoconjugate J* 1989;6:319–332.
  - 58 Montiel M-D, Krzewinski-Recchi M-A, Delannoy P, Harduin-Lepers A. Molecular cloning, gene organisation and expression of human UDP-GalNAc:Neu5Ac $\alpha$ 2-3Gal $\beta$ -R  $\beta$ 1,4-N-acetylgalactosaminyltransferase responsible for the biosynthesis of the blood group Sd<sup>a</sup>/Cad antigen: evidence for an unusual extended cytoplasmic domain. *Biochem J* 2003;373:369–379.
  - 59 Presti LL, Cabury E, Chiricolo M, Dall'Ollio F. Molecular cloning of the human  $\beta$ 1,4N-acetylgalactosaminyltransferase responsible for the biosynthesis of the Sd<sup>a</sup> histo-blood group antigen: the sequence predicts a very long cytoplasmic domain. *J Biochem* 2003;134:675–682.
  - 60 Malagolini N, Santini D, Chiricolo M, Dall'Ollio F. Biosynthesis and expression of the Sd<sup>a</sup> and sialyl Lewis x antigens in normal and cancer colon. *Glycobiology* 2007;17:688–697.
  - 61 Capon C, Maes E, Michelski J-C, Leffler H, Kim YS. Sd<sup>a</sup>-antigen-like structures carried on core 3 are prominent features of glycans from the mucin of normal human descending colon. *Bioch J* 2001;358:657–664.
  - 62 Kawamura YI, Kawashima R, Fukunaga R, *et al.* Introduction of Sd<sup>a</sup> carbohydrate antigen in gastrointestinal cancer cells eliminates selectin ligands and inhibits metastasis. *Cancer Res* 2005;65:6220–6227.
  - 63 Cartron J-P, Prou O, Luilier M, Soulier JP. Susceptibility to invasion by *Plasmodium falciparum* of some human erythrocytes carrying rare blood group antigens. *Br J Haematol* 1983;55:639–647.