

Gap Junctional Signaling in Pattern Regulation: physiological network connectivity instructs growth and form

Juanita Mathews and Michael Levin^{*}

Department of Biology, and
Tufts Center for Regenerative and Developmental Biology
Tufts University

^{*} Author for correspondence:

200 Boston Ave.
Suite 4600
Medford, MA 02155
Tel. (617) 627-6161
Email: michael.levin@tufts.edu

Running title: Gap junctional signals control patterning

Keywords: gap junctions, networks, bioelectric, patterning, morphogenesis

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version record](#). Please cite this article as [doi:10.1002/dneu.22405](https://doi.org/10.1002/dneu.22405).

Abstract

Gap junctions are aqueous channels that allow cells to communicate via physiological signals directly. The role of gap junctional connectivity in determining single-cell functions has long been recognized. However, gap junctions have another important role: the regulation of large-scale anatomical pattern. Gap junctions are not only versatile computational elements that allow cells to control which small molecule signals they receive and emit, but also establish connectivity patterns within large groups of cells. By dynamically regulating the topology of bioelectric networks in vivo, gap junctions underlie the ability of many tissues to implement complex morphogenesis. Here, we review recent data on patterning roles of gap junctions in growth of the zebrafish fin, the establishment of left-right patterning, the developmental dysregulation known as cancer, and the control of large-scale head-tail polarity and head shape in planarian regeneration. We propose a perspective in which gap junctions are not only molecular features functioning in single cells, but enable global neural-like dynamics in non-neural somatic tissues. This view suggests a rich program of future work which capitalizes on the rapid advances in the biophysics of gap junctions to exploit gap junction-mediated global dynamics for applications in birth defects, regenerative medicine, and morphogenetic bioengineering.

Introduction

Gap junctions (GJs), formed by connexin or innexins proteins, form aqueous channels directly connecting the internal cytoplasmic space of nearby cells for the transfer of ions and other small molecules (Phelan, 2005; Scemes et al., 2007). The molecular biology and physiology of gap junctions, as well as their many roles in the nervous system, have recently been expertly reviewed (Baker and Macagno, 2014; Pereda et al., 2013b; Sohl and Willecke, 2004; Steyn-Ross et al., 2007). Here, we focus not on the cellular-level signaling mechanics of GJs, but on their role in determining large-scale pattern formation. Tissue and organ patterning includes embryonic morphogenesis, regeneration of limbs and other organs in a range of model species, and continuous tumor suppression in adult metazoan organisms. Coordinating cell behaviors in vivo toward the creation and repair of complex structures depends on a tight coordination among cells. This coordination is implemented by a rich system of information-bearing signals that propagate throughout the body. Because GJs enable small molecule physiological signals (e.g., ion flow) to pass among cells, they serve as an important modality for cell-cell communication (Trosko, 2007). In this overview, we focus on true channel functions of GJs, and do not discuss permeability-independent functions of connexin proteins (Clasadonte and Haydon, 2014; Dang et al., 2003; Dang et al., 2006; Vinken et al., 2012).

GJ-mediated intercellular signaling functions in parallel with the better-understood secreted gradients of extracellular signaling molecules (Ben-Zvi et al., 2011; Hironaka and Morishita, 2012; Rogers and Schier, 2011). GJ channels are restricted to small molecules (unlike extracellular signaling, which is often mediated by sizeable

proteins). However, GJ signaling networks are extremely versatile because they allow cells to control coupling at the transcriptional, translational, and physiological levels (Solan and Lampe, 2014). GJs can be gated by a range of regulatory inputs (phosphorylation, voltage, pH, etc.) (Peracchia, 2004) and depending on their specific Connexin composition, can exercise considerable selectivity over permeant molecules (Goldberg et al., 2004). Because of this, GJ-coupled cell networks can set up very rich patterns of connectivity in vivo – potentially, with far greater complexity than is possible for diffusing chemicals. Thus, it is no surprise that now classical (Chuang-Tseng et al., 1982; Guthrie and Gilula, 1989; Lo, 1996; Warner, 1985; Weir and Lo, 1982) and recent data have implicated gap junctional communication (GJC) in a range of patterning events in both vertebrate and invertebrate model systems (Ahir and Pratten, 2014; Elias and Kriegstein, 2008b; Sutor and Hagerty, 2005) (Ahir and Pratten, 2014; Elias and Kriegstein, 2008a; Irion et al., 2014; Merrifield and Laird, 2015; Sutor and Hagerty, 2005).

GJs are a powerful mechanism for coordinating long-range signaling during pattern regulation, which is only beginning to be understood. Cells often up-regulate gap junctional communication when they need to share ions with their neighbors (Aslanidi et al., 1991; Larre et al., 2006). Thus, GJC is a perfect conduit for information flow during development, which depends on the ability of cells and tissues to communicate. The converse however is also paramount - embryos contain independent compartments (Rela and Szczupak, 2004; Sutor and Hagerty, 2005) that must remain isolated for proper morphology to result. Mutations in GJ genes are now known to be involved in a wide spectrum of patterning defects (Table 1); however, the true impact of GJC will be

realized not only through ever finer-resolution analysis of the protein dynamics in single cells, but also through an appreciation of their contribution to network dynamics. Here, we review several non-neural contexts in which GJ-mediated signals instruct growth and form. We also present a novel perspective on this field, proposing that GJs may underlie information processing during somatic remodeling that is very similar to the role of these electrical synapses (Bennett, 1997) in implementing plasticity and memory in the brain. This hypothesis sheds new light on emerging data in this exciting field and suggests a research program focused on understanding the global dynamics of GJ-mediated physiological networks.

Gap junctions in cellular regulation

The first patterning tasks required of any animal is embryogenesis – the remarkable self-assembly of a complex body from the descendants of a single egg cell. This process necessarily involves partitioning the embryo into diverse components with different fates and functions.

Endogenous expression and GJC-dependent domains

Embryos of frog, fruit fly, fish, and nematodes have long been known to possess compartments – discrete regions of GJ-coupled cells with distinct selectivity of permeable signals (Blennerhassett and Caveney, 1984; Bozhkova, 1998; Lo and Gilula, 1979; Pitts et al., 1988; Weir and Lo, 1982, 1984). For example, the wing imaginal disk in *Drosophila* is subdivided into a number of communication compartments during differentiation (Weir and Lo, 1984). Such developmental domains are cell groups with

distinct physiological properties, which thus could underlie the diversification and patterning of different embryonic regions. Early mollusk development includes a number of specific GJC patterns that functionally determine cell determination (de Laat et al., 1980). Chick limbs exhibit a gradient of GJC along the AP axis (Coelho and Kosher, 1991). The highest GJC was observed in cells adjacent to the zone of polarizing activity, while no GJC was present at the opposite end of the limb bud. Mesenchymal tissues in the middle of the limb had an intermediate level of dye coupling and it has been hypothesized that polarizing region cells communicate to anterior mesenchyme cells via gap junctions (Allen et al., 1990).

The potential complexity of GJ-based regionalization is augmented by the fact that gap junctions composed of different Connexin subunits provide a degree of functional compensation (Vink et al., 2004), but also confer the ability to sense different signals upon cells that express them (Elfgang et al., 1995). For example, homomeric Cx32 channels are permeable to both cAMP and cGMP, whereas heteromeric Cx32/Cx26 channels retain permeability to cAMP but prevent transfer of cGMP (Bevans et al., 1998). Indeed, substitution studies in tissues such as lens showed that distinct connexins play different roles in development, forming signaling conduits that are not interchangeable (White, 2002). In addition to molecular selectivity, gap junctions also enable rectification and unidirectional transfer (Bruzzone and Giaume, 1999; Fan et al., 2005; Palacios-Prado et al., 2014; Robinson et al., 1993; Zhang et al., 2003), enabling embryogenesis to take advantage of a very rich and dynamic set of signaling paths. For example, unidirectional junctions are thought to form between Cx32 and Cx43

(Robinson et al., 1993; Xin and Bloomfield, 1997), potentially allowing an embryo to establish one-way signaling paths.

GJC and stem cells

Pattern formation is the result of a tight spatio-temporal regulation of cell proliferation, differentiation, migration, and programmed cell death. Recent data have shown that GJ-mediated signaling is a powerful regulator of these properties, especially in stem cells (Oviedo and Levin, 2007; Wong et al., 2008). GJs are involved in regulating multipotency (Dyce et al., 2014), differentiation (Araya et al., 2005; Araya et al., 2003; Bani-Yaghoub et al., 1999; Gu et al., 2003b; Hirschi et al., 2003; Li et al., 2015; Zhang et al., 2002), self-renewal (Hitomi et al., 2015), proliferation (Paraguassu-Braga et al., 2003; Pearson et al., 2005; Starich et al., 2014), and motility (Huang et al., 1998a; Kotini and Mayor, 2015; Marins et al., 2009; Xu et al., 2001; Zahler et al., 2003). neural stem cells exhibit a unique signature based on GJC and ion transporters, and GJC based on Cx43 and Cx45 is essential for their survival and proliferation (Cai et al., 2004). For example, the gap junction protein Zero Population Growth (zpg) is required for germ cell differentiation in *Drosophila* ovary. In the absence of ZPG, the stem cell daughter destined to differentiate instead dies (Gilboa et al., 2003). Germ line stem cells differentiate upon losing contact with their niche, which likely involves GJC; as an example, it has been proposed that gap junction-mediated cAMP signaling between blastomeres and somatic cells results in changes in somatic cell gene expression (Burnside and Collas, 2002). Many instances of GJC-dependent regulation is bi-directional, with several cell types exchanging instructive information, such as occurs in

the communication between stem cells and their neighbors in spermatogenesis during germline differentiation (Smendziuk et al., 2015).

Pigmentation control networks

One of the exciting recent developments is the elegant merging of genetic evidence for GJC roles with a mathematical analysis of the resulting pattern formation that has taken place in the context of zebrafish pigment patterning. In 1953 Alan Turing wrote a paper entitled 'The chemical basis of morphogenesis'. In it, Turing describes a mathematical analysis of reaction-diffusion systems that could account for the formation of stable periodic patterns in organisms. His system consisted of two substances with differing diffusion rates. One of those unknown compounds was an activator that could enhance its own formation and that of the other unknown compound, the inhibitor. The inhibitor not only was capable of inhibiting the formation of the activator, but it also possessed the faster diffusion rate of the two (Turing, 1953). The model predicted self-organization of spatial patterns, which would change as the size of the organism changed, but the chemical identity of the morphogens was unknown (Schiffmann, 2005). Confirmation came in 1995 when Kondo and Asai found that pigmentation patterns on angelfish changed as the fish grew, and showed a traveling wave that behaved similar to the computer simulations of Turing's reaction-diffusion system (Kondo and Asai, 1995). In order to elucidate the molecules that played the role of the activator and inhibitor, research began to focus on zebrafish pigmentation patterns due to the increasing amount molecular tools available for experimentation in that model

organism (Asai et al., 1999a; Iwashita et al., 2006a; Kondo, 2002; Watanabe et al., 2006a; Watanabe and Kondo, 2015; Yamaguchi et al., 2007).

It was found that the zebrafish *leopard* mutation, which confers spots instead of the typical stripes, gives rise to distinct pigmentation changes that suggested a reaction-diffusion wave as the mechanism controlling pigment pattern (Asai et al., 1999b). Interestingly, it was then discovered that the *leopard* mutation is in *connexin41.8* (Watanabe et al., 2006b), suggesting that in this case of pigment system morphogenesis, the relevant signaling molecules may be moving through gap junctions. The zebrafish mutant, *jaguar*, also showed changes in pigmentation pattern. This mutant had irregular spacing of its stripes that was caused by a mutation in the gene for the inward rectifying potassium channel 7.1 (Kir7.1) (Iwashita et al., 2006b). Since Kir7.1 is responsible for maintaining transmembrane voltage potential and is required in the black pigment cells (melanophores), experiments were done to determine how membrane potential may be involved in pigment patterning changes. These experiments found that melanophore membrane potential was more depolarized in the *jaguar* mutants and that this interfered with the transient depolarization signal conferred by contact with xanthophores, the yellow pigment cells. Loss of this transient signal resulted in melanophores that stayed in close contact with xanthophores, rather than moving away from each other as was the case in the wildtype (Inaba et al., 2012).

Further research by Kondo et al. showed that interactions between the two main pigment cells, xanthophores and melanophores, also involved Delta/Notch signals. These signals were activated in the cells by long processes that connect the different chromatophores. The xanthophore processes is short and presents the Delta ligand

that binds to the Notch receptor on the membrane of the melanophore and promotes its survival. If the melanophore is near a xanophore then it can bind to the Delta ligand and survive. This relationship would satisfy the requirement for an activator with a short range. In addition, the melanophores have a long process that reach out to xanophores, but after a certain distance they can no longer reach, resulting in long-range inactivation (Hamada et al., 2014; Watanabe and Kondo, 2014). However, it is possible that Delta/Notch works in conjunction with a molecule that still depends on diffusion through gap junctions (Hamada et al., 2014). Indeed, recent work on mutated connexins 41.8 and 39.4 show that heterotypic gap junctions do play a part in zebrafish pigmentation patterning. These experiments also show that a third class of chromatophore called an iridophore is involved in patterning on the trunk of the animal but not on the tail, and that this relationship is possibly mediated by molecules that are diffused by gap junctions (Bullara and Decker, 2015; Irion et al., 2014; Kondo, 2002; Takagi and Kaneko, 2005). Together, these studies form an elegant body of work on the relationship of voltage and gap junctional signaling in pattern regulation.

The theoretical discussion as to the identity of these diffusible molecules has suggested the GJC-permeable molecules cAMP and ATP may be the relevant Turing couple. Whereby cAMP is the short range activator and ATP a long range inhibitor (Schiffmann, 1991). However, the small molecule serotonin may also play a role in the reaction-diffusion model of zebrafish coloring pattern formation, as it does in other examples of GJC-regulated patterning (Blackiston et al., 2015; Fukumoto et al., 2005b; Gairhe et al., 2012). Recently it was found that the zebrafish *leopard* mutation was correlated with behavioral abnormalities that pointed to lowered levels of serotonin

transport and that treatments with serotonergic drugs resulted in altered pigmentation patterns (Maximino et al., 2013; Stewart et al., 2013).

GJC and nerve sprouting control

In addition to development, GJs are also important for physiological maintenance (Maeda and Tsukihara, 2011), regeneration (Hoptak-Solga et al., 2008; Umino and Saito, 2002), and remodeling. A key cell type involved in these processes is the neuron. Axonal processes must interact with surrounding tissues and make decisions about direction and magnitude of sprouting, in order to correctly pattern and maintain innervation to target organs. The signaling underlying this, and the involvement of GJs, was recently revealed by using a transplantation assay in which eye primordial from a fluorescently-labeled donor embryo were transplanted onto the flank of a host (Vandenberg et al., 2014). Normally, the ectopic eye produced one major nerve bundle, which extended to the host's spinal cord. However, when the host cells were depolarized, the eye instead generated a huge overproliferation of nerve which spread throughout the tissue. Testing a number of transduction mechanisms for this effect on nerve growth, the authors showed that inhibiting GJC in the host abolished the ability of depolarization to cause hyperinnervation from the implanted organ. The authors showed that the effect only applied to ectopic innervation (not the host's native nerve pattern), and that serotonergic signaling was both required for the hyperinnervation and could rescue the blockade of GJC. A model was advanced in which ectopic nerve, searching for guidance cues, was making proliferation and extension decisions based on serotonin molecules arriving via GJs. As will be seen in the discussion of left-right patterning

below, the movement of serotonin via GJs, under an electrophoretic force, is a conserved theme among a number of patterning systems.

Gap junctions in cancer

Another example in which GJs play an important role is the occasional defection of cells away from the normal anatomical plan towards carcinogenesis (Mesnil et al., 2005; Omori et al., 1998; Trosko, 2005) and subsequent metastasis (Defamie et al., 2014). In a sense, cancer is the derangement of developmental patterning – a disease of geometry (Chernet and Levin, 2013a; Maffini et al., 2005; Rubin, 1985; Soto and Sonnenschein, 2011; Tarin, 2012). Cancer cells abandon correct morphogenesis in favor of unrestrained growth reminiscent of unicellular organisms prior to the appearance of multicellularity and to the GJ-mediated coupling of cells to other cells within a larger somatic context (Loch-Caruso and Trosko, 1985). Reduction of GJ-dependent communication with other somatic tissues is an important early step in the process in which cells begin to treat the rest of the body as an “external environment” within which they must survive by any means necessary.

Studies have long noticed that lowered GJC was associated with induction of tumorigenesis (Loch-Caruso and Trosko, 1985; Potter, 1980; Yamasaki et al., 1984); normal tissue generally possesses a much higher degree of GJC than tumor tissue, and a loss of GJC accompanies early steps in neoplastic transformation (Pitts et al., 1988). A neoplastic phenotype can be induced in cell culture by ectopic closing of gap junctions using pharmacological agents or dominant-negative constructs (Omori and Yamasaki, 1998). Most interestingly, neoplastic characteristics can be suppressed by

ectopic induction of GJC in tumor tissue (Hellmann et al., 1999; Mehta et al., 1991; Rose et al., 1993). Connexin32-deficient mice have a 25-fold increased incidence of spontaneous liver tumors (Temme et al., 1997), and Connexin32 is also an anti-invasive agent in renal cell carcinoma (Yano et al., 2006). Thus, gap junctional isolation is known to be a tumor-promoting agent (Leithe et al., 2006; Loewenstein, 1969, 1979, 1980; Loewenstein and Kanno, 1966; Mesnil et al., 2005; Rose et al., 1993; Yamasaki et al., 1995). This is consistent with a view of GJs as mediating morphogenetic cues that could be keeping cells under differentiation and growth limitation consistent with adult morphostasis. It should be noted however that a few studies have indicated the opposite (Stoletov et al., 2013), for example the finding that Connexins 26 and 43 mediate metastasis in melanoma and breast cancer (Stoletov et al., 2013), and a role for GJs in promoting metastasis by transfer of cGAMP (Chen et al., 2016).

The fact that some Connexins are pro-, and others anti-oncogenic, or even can exert both types of effects in different contexts (El-Saghir et al., 2011; Sin et al., 2012), is less puzzling if GJs are considered not as typical proteins that induce or suppress a particular phenotype, but rather as conduits for cell-cell communication. The outcome is expected to be a function of what signals arrive via the GJs (Lim et al., 2011; Mendoza-Naranjo et al., 2007), and are thus hard to predict from knowledge of the connexin type, because it's partially dependent on the neighboring cells, and physiological parameters that dictate the open/closed state of the GJs and what permeant molecule might be traversing it. This has implications for therapies targeting GJs in cancer and other diseases (Cogliati et al., 2016; Grek et al., 2014; Kandouz and Batist, 2010), because it

shifts the focus onto the signaling molecules and their transfer profiles, in addition to targeting Connexin gene products themselves.

In *Xenopus*, cells expressing KRAS mutations make tumor structures; the incidence of tumors can be significantly reduced, despite the strong presence of the oncogenic mutation, by hyperpolarizing cells in vivo by misexpression of hyperpolarizing ion channels (Chernet and Levin, 2013b). Remarkably however, this also works if cells at the opposite end of the animal are targeted (Chernet and Levin, 2014). How can such long-range communication of physiological state occur? Functional experiments implicated the movement of serotonin and butyrate in bioelectric control of tumorigenesis (Blackiston et al., 2011; Chernet and Levin, 2014; Lobikin et al., 2012), and a recent study looked at the role of GJs in this process. Chernet et al. showed that loss of GJC taking place within nascent tumor sites, within remote host tissues, or between the host and the prospective tumor region significantly lowered the incidence of tumorigenesis, with the most pronounced suppression taking place when GJC inhibition occurred far from the oncogene-expressing cells (Chernet et al., 2015). In contrast, overexpression of wild-type Connexin26 increased tumor incidence. The authors presented a mechanistic model, based on an oscillating signal that propagates through GJs and controls proliferation, that quantitatively explained these puzzling data (and made an unexpected, and subsequently validated, prediction about the role of the left-right axis in this process).

From single cell properties to multiscale pattern

Active GJ communication allows cells to make sophisticated decisions comparing relative levels of specific compounds between themselves and their neighbors (Moreno, 2008) and thus can underlie the transmission of physiological patterning signals (Dobrowolski et al., 2009; Jin et al., 2008; Lecanda et al., 2000b; Levin, 2007; Levin and Mercola, 1998b; Levin and Mercola, 1999; Levin and Mercola, 2000; Oviedo et al., 2010; Pizard et al., 2005; Schiffmann, 2008; Warner, 1999). Cells utilize GJs to communicate directly with their neighbors, or with more distant cells via tunneling nanotubes – extended cell processes with gap junctions at their end (Antanaviciute et al., 2014; Rimkute et al., 2016; Sherer, 2013; Wang et al., 2010; Wittig et al., 2012).

Some of the earliest data implicating GJs in development came from studies with functional antibodies. Introduction of antibodies raised to specific portions of connexin proteins in mouse embryos resulted in developmental defects (Becker et al., 1995). In *Xenopus*, microinjection of antibodies has been reported to disrupt axial patterning (Warner et al., 1984); the treated embryos contained differentiated mesodermal derivatives such as notochord and muscle tissue and Warner *et al.* argued that GJC has a role in pattern formation *per se*, rather than in the induction of specific cell types.

In invertebrate systems, *C. elegans* and *Drosophila* have been paramount in revealing GJC roles via genetic approaches. Numerous defects were observed in a *C. elegans* innexin-3 mutant, including rupture of hypodermis, failure of elongation, and failure of pharynx to attach to anterior (Starich et al., 2003). Likewise, epithelial organization and polarity of the embryonic epidermis is dependent on heteromerization of innexins 2 and 3 in *Drosophila* (Bauer et al., 2004; Lehmann et al., 2006). Establishment of the proventriculus requires Innexin-2, which is a target of Wingless

signaling (Bauer et al., 2002; Bauer et al., 2001). This work is particularly interesting because it links GJC to a canonical signaling pathway. Wnt signaling is likewise upstream of GJC patterns in vertebrates (Ai et al., 2000; Olson et al., 1991a; Olson and Moon, 1992; van der Heyden et al., 1998), suggesting possible conservation of signaling modules involving GJC. Interestingly, overexpression of Cx43 provides a partial recovery of the formation of the cerebellum in Wnt-1 knock-out mice (Melloy et al., 2005), strengthening the link to the Wnt pathway and suggesting that establishing ectopic domains of cell:cell communication may be a useful modality to control certain patterning defects. Gene targeting studies of connexins in mice have provided evidence that gap junctions are important for cardiac septation (Kirchhoff et al., 2000) and morphogenesis of the outflow tract (Gu et al., 2003a) and endocardial cushions (Kumai et al., 2000). Many of these effects appear mediated by effects on the behavior of neural crest (Ewart et al., 1997; Huang et al., 1998b; Huang et al., 1998c; Li et al., 2002; Lo et al., 1997; Lo et al., 1999; Sullivan et al., 1998; Waldo et al., 1999; Waller et al., 2000; Xu et al., 2001).

We next discuss a number of specific recent examples in which GJC mediates instructive morphogenetic cues.

Gap junctional communication in growth and patterning of the zebrafish fin

Connexins and joint formation

Zebrafish fin patterning is an excellent model of joint formation due to the ability of the fin to regenerate once amputated, and the continuation of fin growth in adult

animals. The caudal fin is made up of 16-18 fin rays, which are made up of multiple bone segments separated by multiple joints (Sims et al., 2009a). Gap junctions have been shown to regulate the morphology of bone segments and joints in the zebrafish fins. The fin length mutant *short fin (sof)* was found to be caused by mutations in the expression of Connexin43 (Cx43) protein (Iovine et al., 2005a). Analysis showed that reduced GJC caused by lower expression of Cx43 or missense alleles reduces the total number of dividing cells that form the fin ray segments, resulting in shorter segments. However, Cx43 could also be allowing the diffusion of signaling molecules from proximal regions to distal regions acting as a biological ruler that could inform the formation of new segments that extend the length of the fin (Hoptak-Solga et al., 2007). The new bone segments in the fin are dependent on new joint formation that occurs in a proximal to distal manner. Cx43 was found to take part in joint formation by first determining the location of the future joint in the mesenchymal compartment. Later, the Cx43 proteins polarize themselves towards the medial surface of the newly differentiating cells that will form the new joint. Lowered levels of Cx43 resulted in premature joint formation and small fins, while increased expression resulted in joint failure and large fins (Hoptak-Solga et al., 2007).

Recently, microarray investigation into the mechanism by which Cx43 exerts its effect on fin size found that expression of the gene encoding the signaling molecule Semaphorin3d (Sema3d) was perturbed in Cx43 mutants. Semaphorins are involved in the patterning of a variety of tissues and organs. The semaphorin receptors Neuropilin2a (Nrp2a) and PlexinA3 (PlxnA3), were found to be involved in the changes in cell proliferation and joint forming phenotypes associated with Cx43 expression in fin

formation, respectively (Ton and Iovine, 2012). Cx43 involvement in suppressing joint formation was also found to influence the timing of expression of the homeobox gene that encodes the transcription factor Even Skipped 1 (Evx1). Overexpression of Cx43 resulted in loss of evx-1 and decreased expression resulted in evx-1 expression that was more distal than wildtype and led to the premature expression of the genes involved in joint formation (Ton and Iovine, 2013).

The other gene identified in the microarray investigation into downstream effects of changes in Cx43 expression, was hyaluronan and “proteoglycan link protein 1a” (hapln1a). Hapln1a is a protein that links the carbohydrate polymer, hyaluronic acid (HA) with proteoglycans in the extracellular matrix (ECM). This link stabilizes HA creating the right ECM environment that works in parallel with the Sema3d pathway for correct fin formation (Govindan and Iovine, 2014). Further studies on Cx43 expression in zebrafish and how it affects fin formation have found that the gene called “establishment of cohesion1 homolog 2” (esco2) is responsible for regulating Cx43 levels. ESCO2 is an acetyltransferase that modifies cohesion proteins that are responsible for joining sister chromatids during mitosis. Mutations in esco2 cause disruptions in mitosis that increase cell apoptosis and decrease cell proliferation. Together, these data reveal elements both upstream and downstream of connexin function. However, the perturbation in mitosis caused by the mutant form of esco2 in humans could not completely explain the phenotype seen in the resulting syndrome. This syndrome, called Robert’s syndrome is characterized by craniofacial abnormalities and limb malformation that could be a more severe form of oculodentodigital dysplasia (OCDD) that arises from mutations in the human Cx43 (Banerji et al., 2016).

Gap junctions and left-right patterning

Establishing vertebrate laterality

All vertebrates, and many invertebrates, exhibit a fundamental asymmetry of the bilateral bodyplan, with some internal organs (the heart, viscera, and brain) consistently biased to one side of the midline (Speder et al., 2007; Vandenberg and Levin, 2009). Early discoveries of left- and right-specific transcriptional signaling cascades had suggested that each side develops independently (Levin, 1998; Ramsdell and Yost, 1998). However, surgical experiments in the chick showed that in the absence of the right side, the left-sided (and not directly manipulated) tissue became confused, sometimes failing to turn on left-sided markers such as *Sonic hedgehog* and *Nodal* (Levin and Mercola, 1999). Seeking a mechanistic understanding of why distant tissue is required for laterality decisions, gap junctional communication was tested as a mechanism to coordinate sidedness decisions across the whole early embryo in frog and chick models (Levin and Mercola, 1998b). The data showed that either universal blockade of GJs with dominant negative Connexins, or universal expression of constitutively-open Connexins, randomized both the expression of laterality marker genes and the sidedness of the internal organs. Targeted misexpression experiments in the frog showed that normal asymmetry requires open GJC across the embryos' dorsal side, while lack of GJC is required across the ventral midline. These endogenous differences in GJC had been shown to be set up by Wnt pathway signaling as part of dorso-ventral axis determination (Olson et al., 1991b; Olson and Moon, 1992).

A model was proposed, in which signals traversed the blastoderm in a chiral (left to right) manner, propagating through GJs and accumulating on one side of a midline zone of junctional isolation (Levin and Mercola, 1998a; Levin and Nascone, 1997). This scheme was suggested by the molecular functional data, although LR asymmetric GJ transfer was actually present in earlier data examining connectivity among frog blastomeres (Guthrie, 1984; Guthrie et al., 1988; Nagajski et al., 1989; Turin and Warner, 1980). Subsequent work identified the neurotransmitter serotonin (Adams et al., 2006; Fukumoto et al., 2005a; Fukumoto et al., 2005b; Vandenberg et al., 2012) as one left-right morphogen that moves through GJs (long before the nervous system appears) to control downstream gene expression via HDAC1-dependent chromatin modification (Carneiro et al., 2011) and several proton pumps and potassium channels that provide the electrophoretic force for unidirectional accumulation of charged morphogens (Adams et al., 2006; Aw et al., 2008; Aw et al., 2010; Levin et al., 2002; Morokuma et al., 2008). These mechanisms and the role of GJs in left-right axial patterning appear to be implemented very similarly in chick and frog, despite quite different early embryonic architectures. In chick, the relevant connexin appears to be Cx43, which is also important for laterality in zebrafish, where it is required for morphogenesis of the Kupffer's Vesicle (Hatler et al., 2009). Together, these data have also led to quantitative models of the distribution of serotonin through long-range GJ paths, and a study of the main system-level behaviors of this electrophoretic system (Esser et al., 2006; Zhang and Levin, 2009). A number of open questions remain, including possible roles of unidirectional transfer (by distinct Connexins expressed in neighboring cells).

Innexin-based GJs also pattern the nematode LR axis

Interestingly, despite a completely different bodyplan and long evolutionary distance, gap junctions are also involved in LR patterning in the nematode *C. elegans*. An elegant set of studies (Chuang et al., 2007b) focused on neural lateralization showed that establishing left-right asymmetry in *C. elegans* olfactory neurons involves Ca^{++} signaling, tight junctions, and communication through gap junctions. The central nervous system of *C. elegans* contains two bilaterally symmetrical odor sensory neurons (AWC). During late embryogenesis only one of these neurons will express the *str-2* G-protein coupled olfactory receptor. The determination of which neuron is going to express *str-2* (AWC^{ON}) is stochastic, and the remaining neuron acquires the AWC^{OFF} phenotype. Chuang et al. identified *nsy-5*, a gene encoding a member of the innexin/pannexin gap junction family of proteins as *nsy-5* mutants are unable to sense odorants normally processed by AWC^{ON} neurons. Reduction of *nsy-5* function resulted in both neurons having an AWC^{OFF} phenotype. NSY-5 can form functional hemichannels and provide electrical coupling between cells. Genetic approaches showed that *nsy-5* has specific site of actions within different neuronal cell bodies. Strikingly, *nsy-5* can act autonomously to induce the future AWC^{ON} neuron based on a feedback mechanism between both AWC neurons. Subsequent work showed that calcium levels in non-AWC sensory neurons, determined the AWC^{ON} bias, demonstrating that a network of neurons communicate with AWC via signaling dependent on *nsy-5* to determine asymmetric expression of the *str-2* gene. It was also found that the modulation and propagation of the calcium signals was not mediated by

serotonin or inositol triphosphate (IP3) (Schumacher et al., 2012). It is important to note that NSY-5 mediated connectivity also works in parallel with the NSY-4 claudin, which is related to the gamma transmembrane protein subunit of voltage-activated calcium channels and genetically down-regulates calcium channels which inhibits the calcium signaling pathway in the AWC^{ON} cell (Chuang et al., 2007a; VanHoven et al., 2006). Inhibition of the calcium signaling pathway, calcium-calmodulin-dependent kinase II (CaMKII)-MAP kinase, is also achieved by an unknown *nsy-5* mechanism and also by the *mir-71* miRNA that is stabilized by *nsy-4* and *nsy-5* (Alqadah et al., 2013). Once AWC asymmetry is achieved by *nsy-5* and *nsy-4*, maintenance of that asymmetry depends on olfactory signaling, transcriptional regulation including the zinc finger transcription factor *die-1*, and TGF- β signaling (Cochella et al., 2014; Hsieh et al., 2014).

The data revealed some striking similarities between how gap junctions are used in invertebrates to their functions in vertebrate development (Levin, 2002). In chick and frog embryos, the left and right sides of the embryos communicate with each other to correctly assign left-right identity before the onset of asymmetric gene expression. In both *Xenopus* and *C. elegans*, both over- and under-expression of gap junction proteins lead to defects in laterality. The native pattern of communication mediated by gap junctions in both vertebrates and nematodes is between the Left and Right sides and involves a zone of junctional isolation (in this case, provided by an extracellular matrix layer). Moreover, the experiments by Chuang et al. revealed an asymmetry in how the left and right AWCs respond to *nsy-5* expression, mirroring the differences in gap-junctional permeability that has been described on the left and right sides of the early

frog embryo (Guthrie et al., 1988). Crucially, the discovery of this intrinsic bias in *C. elegans* shows that as in frog and chick, the communication via gap junctions does not initiate left-right information but functions as an intermediate step of the pathway.

Gap junction-mediated signaling in head patterning

Long-range control of brain formation

The developing brain must coordinate its growth and morphogenesis. In *Xenopus* embryos, it was recently shown that brain size and patterning is in part regulated by endogenous gradients of transmembrane resting potential (Pai et al., 2015a). Experimentally modulating resting potential (by expressing hyperpolarizing or depolarizing channels) in neural cells, or even in distant (ventral) cells, could control cell proliferation and overall patterning in the nascent brain. This effect was mediated by gap junctional communication: blocking GJC either genetically or pharmacologically rescued hyperpolarization-triggered brain malformations, and prevented ventral hyperpolarization from inducing proliferative effects in the brain. The authors suggest that GJC is involved in coordinating growth across long distances in the embryo, allowing the developing brain to match its proliferative profile to the rest of the body.

Moreover, induction of specific voltage gradients via ion channel misexpression was shown to induce ectopic brain tissue in posterior regions (Pai et al., 2015a). Interestingly, brain markers are also induced in *neighbors* of cells that misexpress the brain-inducing channels. However, with time, these neighboring cells turn off the ectopic brain markers, leaving only the cells whose V_{mem} is continuously set to a brain-specific

pattern to be able maintain brain fate. The authors speculate that the brain-inducing bioelectric state is spread non-cell-autonomously via GJs, but with progressive restriction of GJ communication as development proceeds and individual tissue compartments refine, cells that do not drive abnormal resting potentials with their own channels go back to their normal fates. This normalization also works in the opposite direction, often enabling neighbors to suppress aberrant functions in cells with unique resting potentials that can otherwise induce foci of ectopic eyes (Pai et al., 2012) or tumors (Chernet and Levin, 2013b).

Species-specific head morphology during planarian regeneration

Planarian flatworms are complex creatures, with a true brain, bilateral symmetry, a complex behavioral repertoire, and many body organs (Gentile et al., 2011; Sarnat, 1985). They also have the remarkable ability to regenerate completely from partial body fragments (Lobo et al., 2012; Reddien and Sanchez Alvarado, 2004; Salo et al., 2009). Their capacity for self-repair serves as a paradigm case of dynamic morphostasis and continuous remodeling towards a specific target morphology. While many details of molecular pathways required for correct stem cell differentiation and regenerative capacity are becoming clear (Aboobaker, 2011; Handberg-Thorsager et al., 2008), the mechanisms that determine the correct shape of the head, and cease growth and remodeling when that shape is achieved, are almost completely unknown (Lobo et al., 2012; Lobo et al., 2014). Examining physiological inputs into this process, we recently asked 2 questions: 1) could modification of overall bioelectric network connectivity give rise to coherent patterning changes during regeneration, and 2) is it possible to obtain

evolutionarily-relevant patterns. The results suggested that shifting among different regions of planarian morphospace (Stone, 1997) is possible by physiological perturbation alone (Emmons-Bell et al., 2015).

When GJC was systemically disrupted in fragments of *Girardia dorotocephala* (GD), it induced the expected finer-grain V_{mem} regionalization among the endogenous bioelectric network (since GJC can normally act to establish isopotential cell fields). Remarkably, the resulting worms regenerated heads with an altered shape morphology that quantitatively resembles that of multiple other flatworm species (Emmons-Bell et al., 2015). This resemblance was more than skin deep: not only was the external shape of the head converted to the shapes of distant planarian relatives, but they also manifested different species-specific brain morphologies and stem cell distributions. The exact same treatment of a cohort of GD worms produced 4 types of worm heads in characteristic frequencies (proportional to their evolutionary distance from GD). It is not yet known whether this stochastic property is a consequence of the still relatively crude method of network topology perturbation (soaking in GJ blocker such as octanol), or whether it is an intrinsic aspect of the dynamics of this system. The ability to induce a different species' head shape in a genetically wild-type worm suggests that the bioelectric network is a profound regulator of species-specific morphology. It remains to be seen whether changes in the dynamics of bioelectrical circuits have been widely exploited by evolution to explore variations of anatomical structure.

How to infer the large-scale outcomes (which kind of head, how many heads, etc.) from cell-level properties and signals? This question has been addressed for gene regulatory networks using dynamical state spaces built to describe transcriptional

circuits have been used to map complex system behavior (Halley et al., 2012; Huang, 2011; Huang et al., 2009). More directly relevant to physiological networks, similar approaches have been used to understand global behaviors of electrical activity in neural networks during decision-making. Importantly however, in planarian regeneration, as in the brain, circuit dynamics are not directly revealed from ion channel expression data but are complex and nonlinear. Such dynamics must be modeled quantitatively to understand their emergent properties (Cervera et al., 2015; Law and Levin, 2015). One possibility is that different anatomical outcomes correspond to specific attractors in the dynamical state space of the bioelectric network formed by the planarian body; in this paradigm, bioelectric perturbations can shift the system from a default (genome-specified) attractor to another nearby one (Fig. 5). We are currently working to computationally model this process, to quantitatively map stable attractors to underlying physiological details, and thus gain more control over the resulting shapes.

Gap junctional signaling underlies anatomical pattern memory

In order for a bisected worm to regenerate as two normal worms, the cells at the posterior-facing edge of the wound must make a tail, while those at the anterior-facing edge must make a head. This simple fact reveals that the adult stem cells which implement regeneration in each blastema are not locally controlled: those cells were direct neighbors until the scalpel separated them, and thus have identical positional information, and yet they form completely different structures. Local position is not sufficient for blastema cells to know what shape to build; rather, they must communicate with the remaining tissue to decide what anatomical structures must be formed.

Searching for the mechanism of this long-range coordination, (Nogi and Levin, 2005) found that inhibition of GJC in the species *D. japonica* for just 2 days resulted in subsequent regeneration of worm fragments forming heads at both ends (Nogi and Levin, 2005; Oviedo et al., 2010). These bipolar forms were viable, and could also be phenocopied by RNAi targeting 3 Innexins. By itself, this supported a role for GJ communication in anterior-posterior patterning in this species of flatworm. However, the real surprise came when the 2-headed worms were analyzed long after the GJ blocker (octanol) was gone (as shown by HPLC).

Weeks later, when these 2-headed animals had their heads and tails amputated again (in plain water, with no further perturbation), the same 2-headed phenotype resulted, and this was repeated upon subsequent rounds of amputations (Oviedo et al., 2010). These data showed that a transient perturbation of physiological cell:cell communication via gap junctions could stably change the pattern to which the animal regenerates upon damage – its target morphology. This represents a clear example of physiological change permanently rewriting major anatomical features, despite a normal genome – a novel aspect of epigenetics in the original sense of the word (Noble, 2015). While chromatin modification processes may be involved, they are not a sufficient explanation for the effect, since the ectopic heads (tissue which might be suggested to have been epigenetically reprogrammed into a head state from its original tail identity) are discarded at each generation of cutting. What remains is a normal gut fragment, which has been reprogrammed to form 2 heads, not 1, upon future regeneration; the information about basic anatomical polarity and body organization must be stored in a distributed form throughout the animal.

GJs as electrical synapses in bioelectric networks: functional implications

Synapses among cells are a key component of multicellularity, enabling cell-cell communication and information processing necessary for complex patterning and adaptive (regulative) physiology (Baluska and Mancuso, 2014). GJs are an especially important type of synapse because of the ability to regulate current based on voltage – this makes them similar to transistors or memristors (Jo et al., 2010; Pershin and Di Ventra, 2010), which are a basic unit of computation or logic gates. The central nervous system utilizes both chemical and electrical synaptic transmission. Chemical synapses use extracellular bursts of neurotransmitter release to relay digital information between adjacent cells. Electrical synapses use gap junctions to directly connect the interiors of the cells to one another, which results in a local and long-range relay of analog information through small molecules or changes in resting membrane potential (Pereda, 2014). These electrical synapses have been found to be particularly important in the imprinting of emotional memories, such as fear conditioning (Bissiere et al., 2011).

The main connexin in the mammalian brain is Connexin36, but there appear to be other connexins involved as well (Baker and Macagno, 2014). The regulation of connexin expression and the post-translational gating of the channels affects the strength of electrical synapses as does the nature of the connection formed by distinct connexin hemichannels. Gap junctions that are formed by heterotypic channels comprised of different connexins have different conductivities than homotypic channels comprised of identical connexins (Palacios-Prado et al., 2014). Electrical synapses are bidirectional, but not necessarily equally conductive in both directions (Marder, 2009;

Palacios-Prado, 2009). They are also not dependent on action potentials; therefore they are capable of sensing the simultaneous sub threshold depolarizations of a population of connected cells. In neurons, this detection can result in synchronous firing or asynchronous firing due to changes in neuronal thresholds and frequency of firing (Gutierrez and Marder, 2013; Saraga et al., 2006). The synaptic strength of a neuron can be determined by these thresholds and is modulated by the “coupling potential” that is imparted on the neurons by the gap junction conductance but also by the capacitance and resistance of the connected neurons (Pereda et al., 2013a). One of the most important functional features implemented by these versatile electrical gating elements is that of plasticity: history-dependent conductive processes.

Plasticity of electrical synapses occurs when the strength of the electrical synapse is changed by any one of a number factors related to prior electrical activity (Postma et al., 2011). Neighboring inhibitory chemical synapses can change the coupling potential between cells by shunting away excitatory currents (O'Brien, 2014). Ion channel expression and activation in neurons can change the resistance of connected cells (Curti and Pereda, 2004). Neurotransmitters can also change the conductivity of the gap junctions themselves (Piccolino and Neyton, 1984). Dopamine is able to increase cAMP, activating cAMP-dependent protein kinase A (PKA) that phosphorylates sites on Cx36 and decreases its conductance (Urschel et al., 2006). Other neurotransmitters including nitric oxide, histamine, and noradrenaline also have a regulatory effect on gap junction conductance (Hatton and Yang, 1996; Rörig and Sutor, 1996; Zsiros and Maccaferri, 2008). Glutamate from local glutamatergic synapses has a potentiating role on electrical synapses by increasing the availability of calcium that

activates CaMKII, which then phosphorylates sites on Cx36 that are both unique to and the same as PKA (Alev et al., 2008). This binding further affects the plasticity of the gap junction conductivity, usually resulting in an increase of coupling (Pereda et al., 2013a). In this way, memory and computational circuits can be formed as physiological state changes alter GJ-dependent connectivity shaping subsequent intercellular signaling dynamics, resulting in feedback loops that can stabilize transient bioelectric states into stable cascades of downstream signaling pathways.

An important source of neuronal activity modulation in the brain is the astrocyte. These are specialized cells that support neuronal metabolism and housekeeping, provide a structural framework, and are capable of releasing a variety of signaling molecules called gliotransmitters including glutamate, D-serine, and ATP to affect neuronal spiking thresholds (Hamilton and Attwell, 2010). Astrocytes have many processes and each one can connect to a large number of neurons. Although astrocytes do not form gap junctions with neurons, they do form gap junctions between themselves. These gap junctions allow the passage of calcium waves that are induced in response to patterns of neuronal activity, which are conveyed to astrocytes via glutamate (Wade et al., 2011). The propagation of these calcium waves through GJs is important for the coordination neuronal groups that are not directly linked to one another, as these waves are capable of influencing any neuron that is coupled to the astrocytic network through which it is propagating (Pereira and Furlan, 2010). The coordination of information processing in astrocytes via GJ network plasticity is thought to underlie important aspects of cognitive information-processing in the brain (Stehberg et al., 2012; Verwey and Edwards, 2010). Since many of the same mechanisms

(bioelectric signaling, calcium waves, cAMP, etc.) occur in numerous other somatic cell types, it is possible that some of the patterning plasticity observed during regulative development and regeneration could be implemented by brain-like circuits of GJs that stabilize altered downstream signaling events as a function of prior physiological state.

Conclusion

What are the GJ-permeable signals?

What sorts of signals endogenously traverse gap junctions during patterning? In general, a variety of small molecules and metabolites are thought to permeate GJs, including cAMP (Bedner et al., 2003; Burnside and Collas, 2002; Webb et al., 2002), ATP (Bao et al., 2004; Pearson et al., 2005), cGAMP (Chen et al., 2016), Ca^{++} (Blomstrand et al., 1999; Paemeleire et al., 2000; Toyofuku et al., 1998), serotonin (Fukumoto et al., 2005b; Gairhe et al., 2012; Hou et al., 2013; Wolszon et al., 1994), and histamine (Chaturvedi et al., 2014). These are in general impossible to watch traversing GJ in situ, as these molecules are too small to be labeled (fluorescently) without radically altering their properties. However in addition to small molecule metabolites, several recent reports have demonstrated that molecules much larger than the normal ≈ 1 kDa size cutoff can penetrate through gap junctions under some circumstances (Brooks and Woodruff, 2004; Valiunas et al., 2005). These include siRNA's (Katakowski et al., 2010; Lim et al., 2011; Rimkute et al., 2016; Wolvetang et al., 2007) and even the protein calmodulin (Brooks and Woodruff, 2004; Woodruff, 2005); one possibility is that the crucial parameter is shape, not overall size, and that

long, thin molecules may be able to traverse gap junction channels. The alignment of such molecules to facilitate GJC-mediated transfer could be provided by an endogenous electric field (Woodruff, 2005). Understanding the full range of endogenous permeant signals is an important area for future investigation.

Junctional selectivity can result in radically-different permeabilities of gap junctions to different types of molecules (Bevans et al., 1998; Goldberg et al., 1999; Goldberg et al., 2002; Nicholson et al., 2000). This may have important consequences for the morphogenetic system, and means that studies of this process *in vivo* may be strongly dependent upon the experimental probe used. For example, at 7.5 days, the mouse embryo was found to be subdivided into at least nine GJC compartments with respect to Lucifer Yellow (LY) transfer, but only two domains with respect to ionic coupling (Kalimi and Lo, 1989). In the loach (*Misgurnus fossilis*), LY, fluorescein, and DAPI showed consistent differences in their ability to transfer between tissues during mesoderm induction and patterning (Bozhkova and Rozanova, 2000). Moreover, the chemical selectivity of the gap junctions connecting early embryonic cells changes appreciably during loach development (Bozhkova, 1998), suggesting that embryonic patterning can utilize regulation of not only the amount of GJC but also of the various types of molecules being passed through the gap junctions. Significant advances can be expected to result from future development of versatile fluorescent methods to reveal patterns of distinct GJ paths *in vivo*.

GJC and bioelectricity: two key components of computational networks

In addition to chemical metabolites, one of the most important signals propagated through GJ paths is current. This allows GJs to demarcate isopotential compartments of cells with similar V_{mem} (Sherman and Rinzel, 1991). This is thought to underlie normalization of aberrant founder cells in the production of ectopic brain tissue (Pai et al., 2015a) or tumors (Chernet et al., 2015). GJs and resting potential are bi-directionally linked, and can thus support physiological feedback loops since V_{mem} can regulate GJ opening, while GJs regulate the sharing of V_{mem} among neighboring cells. While functionally linked, voltage and GJ state are not interchangeable. Genetic crosses have revealed that in the zebrafish tail, Connexins and ion channels have distinct roles (Hoptak-Solga et al., 2008; Perathoner et al., 2014). And in the case of voltage control of nerve sprouting (Blackiston et al., 2015), depolarization cannot induce the hyperinnervation phenotype if GJ is shut down. Indeed, GJs allow multiple cells to act as one in response to electric fields (Cooper, 1984; Tsutsui et al., 2014) and alter their bioelectric dynamics (Baigent et al., 1997; Donnell et al., 2009). One of the best recent linkages between physiology and transcriptional regulation was characterized in the highly GJ-coupled heart tissue, in which Wnt1 controls the bioelectrical gradient via regulation of the L-type calcium channel (Panakova et al., 2010).

Thus, one mechanism by which GJ activity regulates pattern formation is by shaping endogenous bioelectric gradients. The importance of these gradients have been confirmed by numerous functional studies implicating spatiotemporal V_{mem} differences as an instructive parameter for cell behavior and large-scale patterning (Funk, 2013; Jaffe, 1981; Levin, 2014; McCaig et al., 2005; Nuccitelli, 2003). Ion channel-mediated changes in V_{mem} not only affects individual cell behaviors such as

proliferation, differentiation, apoptosis, and migration (Sundelacruz et al., 2009), but also determines large-scale parameters such as organ size, shape, and axial patterning of the entire body (Beane et al., 2011; Perathoner et al., 2014). In a range of vertebrate and invertebrate model systems, V_{mem} regulates the formation of the brain, eye, wing, and face, and controls patterning along the anterior-posterior and left-right axes during embryonic development (Dahal et al., 2012; Levin et al., 2002; Pai et al., 2015a). Moreover, experimental control of bioelectric gradients has enabled induction of regenerative ability in non-regenerative contexts (Leppik et al., 2015; Tseng et al., 2010), induced reprogramming of gut tissue into complete eyes (Pai et al., 2012), and normalized tumors (Chernet and Levin, 2013b). Electrical synapses (gap junctions, GJs) and neurotransmitters like serotonin are a key component of several patterning systems, having been implicated in embryonic left-right asymmetry, bone patterning, tumor suppression, and brain size control (Chernet et al., 2015; Iovine et al., 2005b; Levin, 2007; Levin and Mercola, 1998b; Pai et al., 2015a). As in the brain, these elements often work together, such as the bioelectrically-controlled movement of serotonin through GJs during left-right patterning and control of nerve growth (Blackiston et al., 2015; Levin et al., 2006).

In addition to the known downstream targets (Pai et al., 2015b) of the bioelectric signals propagated and limited by GJ paths, specific molecular endpoints for GJ include NFATc1 downstream of Cx37 signaling (Kanady et al., 2015), semaphorin downstream of Cx43 function (Ton and Kathryn Iovine, 2012), and numerous targets of SP1 and SP3 transcription factors (Stains and Civitelli, 2004, 2005; Stains et al., 2003). Because so many downstream targets are impacted by GJC, it is a kind of master regulator node,

like Ca^{++} (Slusarski and Pelegri, 2007) signaling or the RAS gene (Jindal et al., 2015). Thus, one of the exciting aspects of GJC function, as discussed above in the context of selecting alternate head morphologies in planaria, is the ability to regulate large-scale system-wide properties by targeting a single element GJC (or even individual Connexins). An interesting recent example is the demonstration that decreasing Connexin36 gap junction coupling compensates for overactive KATP channels to restore insulin secretion and prevent hyperglycemia in a mouse model of neonatal diabetes (Nguyen et al., 2014). Gap junctions will surely play an increasing role in the understanding and management of circuit disorders, in the fields of birth defects and cancer (Huang et al., 2009).

Next steps: finer-scale reductive studies

Confocal microscopy uncaging and FRET methodologies will greatly facilitate the study of endogenous GJC paths in embryos (Bedner et al., 2003; Braet et al., 2003; Cannell et al., 2004; Dakin et al., 2005; Di et al., 2005). Key future research areas include the characterization of factors which set up patterns of differential GJC in various embryonic tissues (at the transcriptional level, as well as at the level of controlling GJC states, such as by endogenous patterns of pH and voltage gradients (Calero et al., 1998; Ek-Vitorin et al., 1996; Francis et al., 1999; Gu et al., 2000; Morley et al., 1997), and the mapping of paths which exist in and between different tissues to molecules of various charges and sizes.

Experiments in mouse (and other systems) must, of necessity, be performed in the context of multiple connexin knock-outs, since it is becoming increasingly clear that

due to compensation and redundancy, single loss of gap junction genes can obscure important phenotypes (Simon et al., 2004). Indeed, knock-in of dominant negative mutants with different specificities for endogenous connexin families is likely to reveal many important and novel roles (Beahm et al., 2006; Fiorini et al., 2004; Paul et al., 1995). Likewise, more sophisticated technologies allowing expression of dominant negative mutants restricted in space and time during development will allow the circumvention of embryonic lethal phenotypes and likely to lead to the discovery of novel patterning mechanisms (Bakirtzis et al., 2003; Becker et al., 1992). While we now have some understanding of downstream transcriptional changes that occur when gap junctional signaling is perturbed (Kim et al., 2005), next generation sequencing analysis of GJC-inhibited tissues may lead to the discovery of proximal early-response genes to GJC-permeable morphogens. Perhaps the most impactful tool would be the ability to selectively close or open GJ channels, in the way that optogenetics currently allows for ion channel function (Bernstein et al., 2012).

Next steps: systems-level integration

Perhaps even more important than increasing the molecular resolution with which we understand GJ function at the cellular level, is the converse task of synthesizing reductive data into a systems view of cellular networks coupled by dynamic GJC paths. Long-range GJ-mediated signaling, and feedback loops such as the one implemented by calmodulin, which regulates GJC (Burr et al., 2005; Dodd et al., 2008; Peracchia et al., 1983) but is itself a GJ-permeable signal (Brooks and Woodruff, 2004; Woodruff, 2005), ensure regulatory modes that cannot be captured via simple pathway

models. Especially important is the investigation of GJs in reducing the impact of physiological variability and noise in individual cells to facilitate robustness (Hallett, 1989), and their converse roles in sharpening cell responses to weak stimuli (Cooper, 1984). Because of the inherent complexity and nonlinearity of the behavior of such networks, the first task is systems-level modeling. Theoretical analyses of GJC signaling have begun via mathematical models (Cervera et al., 2015; Cooper, 1984; Cooper et al., 1989; Fortier and Bagna, 2006; Hofer et al., 2002; Nygren and Giles, 2000; Vogel and Weingart, 1998, 2002), and in future work it will be very important to apply these to the understanding of specific patterning processes in embryonic and regenerative morphogenesis (Esser et al., 2006; Zhang and Levin, 2009). Such models, especially as they become integrated with bioelectric and transcriptional networks, will become an essential tool for understanding endogenous pattern formation and developing interventions to control complex remodeling events for regenerative medicine and bioengineering applications.

A speculative hypothesis: somatic memories

Beyond modeling the biophysics of GJ-coupled cells, the field will truly mature when it also develops conceptual tools to explain information and computation propagating through GJ-coupled cell networks (Friston et al., 2015; Pezzulo and Levin, 2015). GJs are known to be important for memory and computation in the brain (Dere and Zlomuzica, 2012; Wang and Belousov, 2011; Wu et al., 2011), and the ability of most general anesthetics to serve as GJ uncoupling agents have motivated models of cognition and consciousness based on GJ-mediated integration across the brain

(Juszczak and Swiergiel, 2009; Mantz, 1992). Interestingly, recent work has applied the same types of Hebbian plasticity concepts to understand the function of the heart (Chakravarthy and Ghosh, 1997a; Zoghi, 2004a) and pancreas (Goel and Mehta, 2013b). We have recently extended this idea by proposing that target morphologies for regenerating systems are encoded via a memory-like mechanism within somatic tissues (Friston et al., 2015; Pezzulo and Levin, 2015). Because of the similarity in signaling of GJs in the brain and in non-neural patterning tissues, it is possible that circuits that guide self-limiting, flexible remodeling and regeneration programs are implementing true memories. The central involvement of electrical synapses and the holographic-like nature of patterning information that can be stably re-written (e.g., permanently 2-headed planaria) suggest models in which the target morphology is actually stored (encoded) within the real-time current dynamics, perhaps akin to storage of spatial memory in neural networks or similar proposed processes of memory in non-neural tissues (Baluska and Mancuso, 2012; Levin, 2011, 2012; McConnell et al., 1959; Shomrat and Levin, 2013; Turner et al., 2002; Zoghi, 2004a). This view makes two major predictions, currently being addressed in our lab. First, it suggests that models taken from cognitive neuroscience for understanding the system-wide dynamics of neural networks (Balduzzi and Tononi, 2009; Ebner and Hameroff, 2011; Friston, 2010, 2013; Hoel et al., 2013; Pezzulo et al., 2015) may be appropriate formalisms for efficient prediction and control of morphogenesis.

Second, it suggests that patterning could be controlled top-down, in a complementary approach to today's focus on bottom-up (molecular-level) interventions. If GJ-coupled tissues are indeed information processing agents as they are in the brain,

then it may be possible to efficiently exploit the plasticity and Hebbian-like learning capabilities facilitated by Connexins in non-neural tissues by literally training them to desired outcomes: providing positive and negative reinforcement for collections of cells undergoing morphogenesis to encourage specific types of growth patterns. As with training in cognitive animal models, rewarding for final outcome (as opposed to directly regulating molecular signaling at each point in the tissue) may allow bioengineers to capitalize on the inherent modularity and plasticity of pattern formation. This may enable the induction of desired anatomical patterning changes without facing head-on the complexity explosion that stymies attempts to construct complex organs such as limbs directly.

Training non-neural tissues and treating them like “neural” networks with plasticity and memory could be feasible as shown by recent studies that heart and pancreas cells are capable of memory. The repolarization of the ventricles in the heart, referred to as the T-wave, can be inverted using atrial pacing. Depending on the duration and frequency of that pacing, the inversion strength and duration of inversions after pacing are increased, displaying a Hebbian-like memory (Chakravarthy and Ghosh, 1997b; Rosenbaum et al., 1982; Zoghi, 2004b). Networks of pancreatic islet beta-cells release insulin due to calcium waves that travel through GJs, and display a loss in GJC in response to heightened levels of glucose that can be modeled using learning theories also based on Hebbian-like memory (Benninger et al., 2008; Goel and Mehta, 2013a). These data suggest that gap junction-based plasticity outside the brain could be exploited to offload the computational complexity of micromanaging pattern of physiology onto the cells themselves (as occurs during reward circuit shaping in

behavioral training). A whole new paradigm for medicine that incorporates tissue training could arise if these two important tissues can be physiologically entrained to correct certain heart arrhythmias and insulin secretion disorders.

Looking to neuroscience for examples of how neural networks are trained can inform new strategies to train non-neural cellular networks for desired GJC. A number of neuronal plasticity studies grow neurons on microelectrode arrays (MEAs) that can then be used to monitor the patterns of firing neurons and to also deliver electrical impulses that can stimulate the firing of certain neurons in the network (DeMarse and Dockendorf, 2005; Franke et al., 2012; le Feber et al., 2008; Pimashkin et al., 2013). Cardiomyocytes, pancreatic islet cells, and fibroblasts can all be monitored for electrical activity using MEAs (Bornat et al., 2010; Parak et al., 1999; Pfeiffer et al., 2011; Schönecker et al., 2014; Spira and Hai, 2013). However, the appropriate method for training gap junction connectivity in non-neural networks might differ from neuronal networks.

Adaptive learning in neural networks *in vitro* requires an appropriate stimulation and feedback mechanism. In two studies, which utilized extracellular microelectrode arrays, low frequency stimulation was delivered to a culture of dissociated rat cortical neurons or murine hippocampal neurons via a stimulation electrode, and the spiking frequency was read on the other recording electrodes. Whenever the output at a specific recording electrode was over a certain threshold that considers previous output (response-to-stimulus ratio), then the next recurrence of the low frequency stimulation at the stimulation electrode was delayed (Pimashkin et al., 2013; Shahaf and Marom, 2001). This strategy for training neuronal networks has shown more promise than open-

looped strategies, since it results in more space exploration and greater changes in connectivity (le Feber et al., 2010). Indeed, closing the loop by using embodied networks that are capable of interpreting spiking outputs as instructions for robotic movement and in turn take sensory information from those robots and translate them into changes in stimulation patterns is showing great promise as way to demonstrate neuronal plasticity (Bakkum et al., 2008; Bakkum et al., 2004; Chao et al., 2008). Linking spiking outputs to flight simulator software controls with one stimulating electrode for pitch and the other stimulating electrode for roll can be used to train neuronal networks to fly a plane in the simulator and be able to correct for changes in headwind (DeMarse and Dockendorf, 2005). New optogenetic stimulation of action potentials allows for even better time resolution of output spiking due to the lack of an artifact that occurs when using electrode stimulation (Dranias et al., 2013).

Adapting these neuronal training techniques to non-neural gap junctions can be achieved using electrodes and optogenetics to change membrane potentials and thus affect the GJC between those and surrounding cells. While the details could be connexin- and cell-specific, one class of training strategies could deliver dopamine or glutamate as ways to provide GJ-modulatory feedback to the network. Microfluidics can be used to implement stimulation and also to directly measure the GJC between cells (Bathany et al., 2012). Real-time monitoring of calcium wave patterns using transgenic cell lines could also provide better measurement of output (Shigetomi et al., 2010).

Thus, one way to control the patterning outcome of GJ-coupled cell networks may be to modify their information content, whether by rewriting shape memories directly (via optogenetic-like approaches), or providing stimuli that exploit learning.

Whether the neural computation analogy turns out to be the best way of understanding GJ dynamics, it is clear that GJs are a central hub for cellular information. Cells rely on GJ-permeable signals to regulate their individual activities toward large-scale construction, remodeling, and repair of complex anatomical structures. Thus, gaining predictive control over GJ dynamics is likely to be at the forefront of future efforts to exploit the ability of biological systems to regulate not just gene expression and physiology, but 3-dimensional structure. The implications will be not only in fields like birth defects, regenerative medicine, and cancer, but are likely to impact evolutionary biology, unconventional computation, and the design of robust hybrid technologies. The technological revolution precipitated by the introduction of transistors in electronics was driven not only by their physical properties, but also by the appreciation of their unique computational capabilities. GJs support similar functions at the cellular level, with similar implications for computational capabilities of GJ-coupled networks (Scarle, 2008; Simon, 2009). Thus, it may not be unreasonable to predict that fully understanding and making use of GJ signaling dynamics in biological tissues may lead to the same scale of revolutionary advances in basic biology and applied biomedicine.

Acknowledgements

We thank Alexis Pietak, Alberto Pereda, Jose Ek Vitorin, and the members of the Levin lab for many useful discussions about the role of gap junctions in pattern regulation. We gratefully acknowledge support of the NIH (HD081326, AR055993, AR061988), NSF (CBET-0939511), The G. Harold and Leila Y. Mathers Charitable Foundation, The W. M. KECK Foundation, and the Templeton World Charity Foundation

(TWCF0089/AB55). The authors declare no conflicts or competing interests.

Accepted Article

References

- Aboobaker, A.A., 2011. Planarian stem cells: a simple paradigm for regeneration. *Trends Cell Biol* 21: 304-311.
- Adams, D.S., Robinson, K.R., Fukumoto, T., Yuan, S., Albertson, R.C., Yelick, P., Kuo, L., McSweeney, M., Levin, M., 2006. Early, H⁺-V-ATPase-dependent proton flux is necessary for consistent left-right patterning of non-mammalian vertebrates. *Development* 133: 1657-1671.
- Ahir, B.K., Pratten, M.K., 2014. Structure and function of gap junction proteins: role of gap junction proteins in embryonic heart development. *The International journal of developmental biology* 58: 649-662.
- Ai, Z., Fischer, A., Spray, D.C., Brown, A.M., Fishman, G.I., 2000. Wnt-1 regulation of connexin43 in cardiac myocytes. *J Clin Invest* 105: 161-171.
- Alev, C., Urschel, S., Sonntag, S., Zoidl, G., 2008. The neuronal connexin36 interacts with and is phosphorylated by CaMKII in a way similar to CaMKII interaction with glutamate receptors. *Proceedings of the*
- Allen, F., Tickle, C., Warner, A., 1990. The role of gap junctions in patterning of the chick limb bud. *Development* 108: 623-634.
- Alqadah, A., Hsieh, Y.-W., Chuang, C.-F., 2013. microRNA function in left-right neuronal asymmetry: perspectives from *C. elegans*. *Frontiers in cellular neuroscience* 7.
- Antanaviciute, I., Rysevaite, K., Liutkevicius, V., Marandykina, A., Rimkute, L., Sveikatiene, R., Uloza, V., Skeberdis, V.A., 2014. Long-distance communication between laryngeal carcinoma cells. *PloS one* 9: e99196.
- Araya, R., Eckardt, D., Maxeiner, S., Kruger, O., Theis, M., Willecke, K., Saez, J.C., 2005. Expression of connexins during differentiation and regeneration of skeletal muscle: functional relevance of connexin43. *J Cell Sci* 118: 27-37.
- Araya, R., Eckardt, D., Riquelme, M.A., Willecke, K., Saez, J.C., 2003. Presence and importance of connexin43 during myogenesis. *Cell Commun Adhes* 10: 451-456.
- Asai, R., Taguchi, E., Kume, Y., Saito, M., Kondo, S., 1999a. Zebrafish leopard gene as a component of the putative reaction-diffusion system. *Mechanisms of development* 89: 87-92.
- Asai, R., Taguchi, E., Kume, Y., Saito, M., Kondo, S., 1999b. Zebrafish leopard gene as a component of the putative reaction-diffusion system. *Mech Dev* 89: 87-92.
- Aslanidi, K.B., Boitsova, L., Chailakhyan, L.M., Kublik, L.N., Marachova, II, Potapova, T.V., Vinogradova, T.A., 1991. Energetic cooperation via ion-permeable junctions in mixed animal cell cultures. *FEBS Lett* 283: 295-297.
- Aw, S., Adams, D.S., Qiu, D., Levin, M., 2008. H,K-ATPase protein localization and Kir4.1 function reveal concordance of three axes during early determination of left-right asymmetry. *Mech Dev* 125: 353-372.
- Aw, S., Koster, J., Pearson, W., Nichols, C., Shi, N.Q., Carneiro, K., Levin, M., 2010. The ATP-sensitive K(+) -channel (K(ATP)) controls early left-right patterning in *Xenopus* and chick embryos. *Dev Biol* 346: 39-53.
- Baigent, S., Stark, J., Warner, A., 1997. Modelling the effect of gap junction nonlinearities in systems of coupled cells. *J Theor Biol* 186: 223-239.

- Baker, M.W., Macagno, E.R., 2014. Control of neuronal morphology and connectivity: Emerging developmental roles for gap junctional proteins. *FEBS Lett.*
- Bakirtzis, G., Jamieson, S., Aasen, T., Bryson, S., Forrow, S., Tetley, L., Finbow, M., Greenhalgh, D., Hodgins, M., 2003. The effects of a mutant connexin 26 on epidermal differentiation. *Cell Commun Adhes* 10: 359-364.
- Bakkum, D.J., Chao, Z.C., Potter, S.M., 2008. Spatio-temporal electrical stimuli shape behavior of an embodied cortical network in a goal-directed learning task. *Journal of neural engineering* 5: 310.
- Bakkum, D.J., Shkolnik, A.C., Ben-Ary, G., 2004. Removing some 'A' from AI: embodied cultured networks. *Embodied artificial*
- Balduzzi, D., Tononi, G., 2009. Qualia: the geometry of integrated information. *PLoS computational biology* 5: e1000462.
- Baluska, F., Mancuso, S., 2012. Ion channels in plants: From bioelectricity, via signaling, to behavioral actions. *Plant signaling & behavior* 8.
- Baluska, F., Mancuso, S., 2014. Synaptic view of eukaryotic cell. *Int. J. Gen. Syst.* 43: 740-756.
- Banerji, R., Eble, D.M., Iovine, K.M., Skibbens, R.V., 2016. Esco2 regulates cx43 expression during skeletal regeneration in the zebrafish fin. *Developmental Dynamics* 245: 7-21.
- Bani-Yaghoub, M., Underhill, T.M., Naus, C.C., 1999. Gap junction blockage interferes with neuronal and astroglial differentiation of mouse P19 embryonal carcinoma cells. *Dev Genet* 24: 69-81.
- Bao, L., Locovei, S., Dahl, G., 2004. Pannexin membrane channels are mechanosensitive conduits for ATP. *FEBS Lett* 572: 65-68.
- Bathany, C., Beahm, D.L., Besch, S., Sachs, F., Hua, S.Z., 2012. A microfluidic platform for measuring electrical activity across cells. *Biomicrofluidics* 6: 34121.
- Bauer, R., Lehmann, C., Fuss, B., Eckardt, F., Hoch, M., 2002. The *Drosophila* gap junction channel gene innexin 2 controls foregut development in response to Wingless signalling. *J Cell Sci* 115: 1859-1867.
- Bauer, R., Lehmann, C., Hoch, M., 2001. Gastrointestinal development in the *Drosophila* embryo requires the activity of innexin gap junction channel proteins. *Cell Commun Adhes* 8: 307-310.
- Bauer, R., Lehmann, C., Martini, J., Eckardt, F., Hoch, M., 2004. Gap junction channel protein innexin 2 is essential for epithelial morphogenesis in the *Drosophila* embryo. *Mol Biol Cell* 15: 2992-3004.
- Beahm, D.L., Oshima, A., Gaietta, G.M., Hand, G.M., Smock, A.E., Zucker, S.N., Toloue, M.M., Chandrasekhar, A., Nicholson, B.J., Sosinsky, G.E., 2006. Mutation of a conserved threonine in the third transmembrane helix of alpha- and beta-connexins creates a dominant-negative closed gap junction channel. *J Biol Chem* 281: 7994-8009.
- Beane, W.S., Morokuma, J., Adams, D.S., Levin, M., 2011. A Chemical genetics approach reveals H,K-ATPase-mediated membrane voltage is required for planarian head regeneration. *Chemistry & Biology* 18: 77-89.
- Becker, D.L., Evans, W.H., Green, C.R., Warner, A., 1995. Functional analysis of amino acid sequences in connexin43 involved in intercellular communication through gap junctions. *J Cell Sci* 108: 1455-1467.

- Becker, D.L., Leclerc-David, C., Warner, A., 1992. The relationship of gap junctions and compaction in the preimplantation mouse embryo. *Dev Suppl* 113-118.
- Becker, D.L., McGonnell, I., Makarenkova, H.P., Patel, K., Tickle, C., Lorimer, J., Green, C.R., 1999. Roles for alpha 1 connexin in morphogenesis of chick embryos revealed using a novel antisense approach. *Dev Gen* 24: 33-42.
- Bedner, P., Niessen, H., Odermatt, B., Willecke, K., Harz, H., 2003. A method to determine the relative cAMP permeability of connexin channels. *Exp Cell Res* 291: 25-35.
- Ben-Zvi, D., Shilo, B.Z., Barkai, N., 2011. Scaling of morphogen gradients. *Curr Opin Genet Dev* 21: 704-710.
- Bennett, M.V., 1997. Gap junctions as electrical synapses. *J Neurocytol* 26: 349-366.
- Benninger, R.K.P., Zhang, M., Head, W.S., Satin, L.S., 2008. Gap junction coupling and calcium waves in the pancreatic islet. *Biophysical journal*.
- Bernstein, J.G., Garrity, P.A., Boyden, E.S., 2012. Optogenetics and thermogenetics: technologies for controlling the activity of targeted cells within intact neural circuits. *Curr Opin Neurobiol* 22: 61-71.
- Bevans, C.G., Kordel, M., Rhee, S.K., Harris, A.L., 1998. Isoform composition of connexin channels determines selectivity among second messengers and uncharged molecules. *J Biol Chem* 273: 2808-2816.
- Bissiere, S., Zelikowsky, M., Ponnusamy, R., Jacobs, N.S., 2011. Electrical synapses control hippocampal contributions to fear learning and memory. *Science*.
- Blackiston, D., Adams, D.S., Lemire, J.M., Lobikin, M., Levin, M., 2011. Transmembrane potential of GlyCl-expressing instructor cells induces a neoplastic-like conversion of melanocytes via a serotonergic pathway. *Disease models & mechanisms* 4: 67-85.
- Blackiston, D.J., Anderson, G.M., Rahman, N., Bieck, C., Levin, M., 2015. A novel method for inducing nerve growth via modulation of host resting potential: gap junction-mediated and serotonergic signaling mechanisms. *Neurotherapeutics* 12: 170-184.
- Blennerhassett, M.G., Caveney, S., 1984. Separation of Developmental Compartments by a Cell Type with Reduced Junctional Permeability. *Nature* 309: 361-364.
- Blomstrand, F., Aberg, N.D., Eriksson, P.S., Hansson, E., Ronnback, L., 1999. Extent of intercellular calcium wave propagation is related to gap junction permeability and level of connexin-43 expression in astrocytes in primary cultures from four brain regions. *Neuroscience* 92: 255-265.
- Bornat, Y., Raoux, M., Boutaib, Y., Morin, F.O., 2010. Detection of electrical activity of pancreatic beta-cells using micro-electrode arrays. ... Design.
- Bozhkova, V.P., 1998. The specificity of gap junctional channels in early fish embryos and its significance for pattern formation in development. *Membr Cell Biol* 11: 803-815.
- Bozhkova, V.P., Rozanova, N.V., 2000. Local changes in gap junctional permeability accompany regionalization of the mesoderm in early fish (loach, *Misgurnus fossilis*) embryos. *Membr Cell Biol* 14: 189-198.
- Braet, K., Vandamme, W., Martin, P.E., Evans, W.H., Leybaert, L., 2003. Photoliberating inositol-1,4,5-trisphosphate triggers ATP release that is blocked by the connexin mimetic peptide gap 26. *Cell Calcium* 33: 37-48.

- Britz-Cunningham, S., Shah, M., Zuppan, C., Fletcher, W., 1995. Mutations of the connexin-43 gap-junction gene in patients with heart malformations and defects of laterality. *N Engl J Med* 332: 1323-1329.
- Brooks, R.A., Woodruff, R.I., 2004. Calmodulin transmitted through gap junctions stimulates endocytic incorporation of yolk precursors in insect oocytes. *Dev Biol* 271: 339-349.
- Bruzzone, R., Giaume, C., 1999. Connexins and information transfer through glia. *Adv Exp Med Biol* 468: 321-337.
- Bullara, D., Decker, D.Y., 2015. Pigment cell movement is not required for generation of Turing patterns in zebrafish skin. *Nature Communications* 6.
- Burnside, A.S., Collas, P., 2002. Induction of Oct-3/4 expression in somatic cells by gap junction-mediated cAMP signaling from blastomeres. *Eur J Cell Biol* 81: 585-591.
- Burr, G.S., Mitchell, C.K., Keflemariam, Y.J., Heidelberger, R., O'Brien, J., 2005. Calcium-dependent binding of calmodulin to neuronal gap junction proteins. *Biochem Biophys Res Commun* 335: 1191-1198.
- Cai, J., Cheng, A., Luo, Y., Lu, C., Mattson, M.P., Rao, M.S., Furukawa, K., 2004. Membrane properties of rat embryonic multipotent neural stem cells. *J Neurochem* 88: 212-226.
- Calero, G., Kanemitsu, M., Taffet, S.M., Lau, A.F., Delmar, M., 1998. A 17mer peptide interferes with acidification-induced uncoupling of connexin43. *Circ Res* 82: 929-935.
- Cannell, M.B., Jacobs, M.D., Donaldson, P.J., Soeller, C., 2004. Probing microscopic diffusion by 2-photon flash photolysis: measurement of isotropic and anisotropic diffusion in lens fiber cells. *Microsc Res Tech* 63: 50-57.
- Carneiro, K., Donnet, C., Rejtar, T., Karger, B.L., Barisone, G.A., Diaz, E., Kortagere, S., Lemire, J.M., Levin, M., 2011. Histone deacetylase activity is necessary for left-right patterning during vertebrate development. *BMC Dev Biol* 11: 29.
- Cervera, J., Manzanares, J.A., Mafe, S., 2015. Electrical coupling in ensembles of nonexcitable cells: modeling the spatial map of single cell potentials. *J Phys Chem B* 119: 2968-2978.
- Chakravarthy, S.V., Ghosh, J., 1997a. On Hebbian-like adaptation in heart muscle: a proposal for 'cardiac memory'. *Biol Cybern* 76: 207-215.
- Chakravarthy, S.V., Ghosh, J., 1997b. On Hebbian-like adaptation in heart muscle: a proposal for 'cardiac memory'. *Biological cybernetics*.
- Chang, Q., Tang, W., Kim, Y., Lin, X., 2014. Timed conditional null of connexin26 in mice reveals temporary requirements of connexin26 in key cochlear developmental events before the onset of hearing. *Neurobiology of disease*.
- Chao, Z.C., Bakkum, D.J., Potter, S.M., 2008. Shaping embodied neural networks for adaptive goal-directed behavior. *PLoS Comput. Biol.*
- Chaturvedi, R., Reddig, K., Li, H.S., 2014. Long-distance mechanism of neurotransmitter recycling mediated by glial network facilitates visual function in *Drosophila*. *Proceedings of the National Academy of Sciences of the United States of America* 111: 2812-2817.

- Chen, Q., Boire, A., Jin, X., Valiente, M., Er, E.E., Lopez-Soto, A., S. Jacob, L., Patwa, R., Shah, H., Xu, K., Cross, J.R., Massagué, J., 2016. Carcinoma–astrocyte gap junctions promote brain metastasis by cGAMP transfer. *Nature* 533: 493-498.
- Chernet, B., Levin, M., 2013a. Endogenous Voltage Potentials and the Microenvironment: Bioelectric Signals that Reveal, Induce and Normalize Cancer. *J Clin Exp Oncol Suppl* 1.
- Chernet, B.T., Fields, C., Levin, M., 2015. Long-range gap junctional signaling controls oncogene-mediated tumorigenesis in *Xenopus laevis* embryos. *Front Physiol* 5: 519.
- Chernet, B.T., Levin, M., 2013b. Transmembrane voltage potential is an essential cellular parameter for the detection and control of tumor development in a *Xenopus* model. *Disease models & mechanisms* 6: 595-607.
- Chernet, B.T., Levin, M., 2014. Transmembrane voltage potential of somatic cells controls oncogene-mediated tumorigenesis at long-range. *Oncotarget* 5: 3287-3306.
- Chuang, C.F., VanHoven, M.K., Fetter, R.D., Verselis, V.K., 2007a. An innexin-dependent cell network establishes left-right neuronal asymmetry in *C. elegans*. *Cell*.
- Chuang, C.F., Vanhoven, M.K., Fetter, R.D., Verselis, V.K., Bargmann, C.I., 2007b. An innexin-dependent cell network establishes left-right neuronal asymmetry in *C. elegans*. *Cell* 129: 787-799.
- Chuang-Tseng, M.P., Chuang, H.H., Sandri, C., Akert, K., 1982. Gap junctions and impulse propagation in embryonic epithelium of Amphibia. A freeze-etching study. *Cell Tissue Res*. 225: 249-258.
- Civitelli, R., 2008. Cell-cell communication in the osteoblast/osteocyte lineage. *Arch. Biochem. Biophys.* 473: 188-192.
- Clasadonte, J., Haydon, P.G., 2014. Connexin 30 controls the extension of astrocytic processes into the synaptic cleft through an unconventional non-channel function. *Neurosci Bull* 30: 1045-1048.
- Cochella, L., Tursun, B., Hsieh, Y.W., Galindo, S., 2014. Two distinct types of neuronal asymmetries are controlled by the *Caenorhabditis elegans* zinc finger transcription factor *die-1*. *Genes & ...*
- Coelho, C.N., Kosher, R.A., 1991. A gradient of gap junctional communication along the anterior-posterior axis of the developing chick limb bud. *Developmental Biology* 148: 529-535.
- Cogliati, B., Menecier, G., Willebrords, J., Da Silva, T.C., Maes, M., Pereira, I.V., Crespo Yanguas, S., Hernandez-Blazquez, F.J., Dagli, M.L., Vinken, M., 2016. Connexins, Pannexins, and Their Channels in Fibroproliferative Diseases. *J Membr Biol*.
- Cooper, M.S., 1984. Gap junctions increase the sensitivity of tissue cells to exogenous electric fields. *J Theor Biol* 111: 123-130.
- Cooper, M.S., Miller, J.P., Fraser, S.E., 1989. Electrophoretic repatterning of charged cytoplasmic molecules within tissues coupled by gap junctions by externally applied electric fields. *Developmental Biology* 132: 179-188.
- Curti, S., Pereda, A.E., 2004. Voltage-dependent enhancement of electrical coupling by a subthreshold sodium current. *The Journal of neuroscience*.

- Dahal, G.R., Rawson, J., Gassaway, B., Kwok, B., Tong, Y., Ptacek, L.J., Bates, E., 2012. An inwardly rectifying K⁺ channel is required for patterning. *Development* 139: 3653-3664.
- Dakin, K., Zhao, Y., Li, W.H., 2005. LAMP, a new imaging assay of gap junctional communication unveils that Ca²⁺ influx inhibits cell coupling. *Nat Methods* 2: 55-62.
- Dang, X., Doble, B.W., Kardami, E., 2003. The carboxy-tail of connexin-43 localizes to the nucleus and inhibits cell growth. *Mol Cell Biochem* 242: 35-38.
- Dang, X., Jeyaraman, M., Kardami, E., 2006. Regulation of connexin-43-mediated growth inhibition by a phosphorylatable amino-acid is independent of gap junction-forming ability. *Mol Cell Biochem* 289: 201-207.
- Davy, A., Bush, J.O., Soriano, P., 2006. Inhibition of gap junction communication at ectopic Eph/ephrin boundaries underlies craniofrontonasal syndrome. *PLoS Biol* 4: e315.
- de Laat, S.W., Tertoolen, L.G., Dorresteijn, A.W., van den Biggelaar, J.A., 1980. Intercellular communication patterns are involved in cell determination in early molluscan development. *Nature* 287: 546-548.
- Debeer, P., Van Esch, H., Huysmans, C., Pijkels, E., De Smet, L., Van de Ven, W., Devriendt, K., Fryns, J.P., 2005. Novel GJA1 mutations in patients with oculo-dento-digital dysplasia (ODDD). *European journal of medical genetics* 48: 377-387.
- Defamie, N., Chepied, A., Mesnil, M., 2014. Connexins, gap junctions and tissue invasion. *FEBS Lett* 588: 1331-1338.
- DeMarse, T.B., Dockendorf, K.P., 2005. Adaptive flight control with living neuronal networks on microelectrode arrays. *Neural Networks*.
- Dere, E., Zlomuzica, A., 2012. The role of gap junctions in the brain in health and disease. *Neurosci Biobehav Rev* 36: 206-217.
- Di, W.L., Gu, Y., Common, J.E., Aasen, T., O'Toole, E.A., Kelsell, D.P., Zicha, D., 2005. Connexin interaction patterns in keratinocytes revealed morphologically and by FRET analysis. *J Cell Sci* 118: 1505-1514.
- Dobrowolski, R., Hertig, G., Lechner, H., Worsdorfer, P., Wulf, V., Dicke, N., Eckert, D., Bauer, R., Schorle, H., Willecke, K., 2009. Loss of connexin43-mediated gap junctional coupling in the mesenchyme of limb buds leads to altered expression of morphogens in mice. *Hum Mol Genet* 18: 2899-2911.
- Dodd, R., Peracchia, C., Stolady, D., Torok, K., 2008. Calmodulin association with connexin32-derived peptides suggests trans-domain interaction in chemical gating of gap junction channels. *J Biol Chem* 283: 26911-26920.
- Donnell, P., Baigent, S.A., Banaji, M., 2009. Monotone dynamics of two cells dynamically coupled by a voltage-dependent gap junction. *J Theor Biol* 261: 120-125.
- Dranias, M.R., Ju, H., Rajaram, E., 2013. Short-term memory in networks of dissociated cortical neurons. *The Journal of*
- Dyce, P.W., Li, D., Barr, K.J., Kidder, G.M., 2014. Connexin43 is required for the maintenance of multipotency in skin-derived stem cells. *Stem cells and development* 23: 1636-1646.

- Ebner, M., Hameroff, S., 2011. Lateral information processing by spiking neurons: a theoretical model of the neural correlate of consciousness. *Comput Intell Neurosci* 2011: 247879.
- Ek-Vitorin, J.F., Calero, G., Morley, G.E., Coombs, W., Taffet, S.M., Delmar, M., 1996. pH regulation of connexin43: molecular analysis of the gating particle. *Biophys J* 71: 1273-1284.
- El-Saghir, J.A., El-Habre, E.T., El-Sabban, M.E., Talhouk, R.S., 2011. Connexins: a junctional crossroad to breast cancer. *Int J Dev Biol* 55: 773-780.
- Elfgang, C., Eckert, R., Lichtenberg-Frate, H., Butterweck, A., Traub, O., Klein, R., Hulser, D., Willecke, K., 1995. Specific permeability and selective formation of gap junction channels in connexin-transfected HeLa cells. *J Cell Biol* 129: 805-817.
- Elias, L., Kriegstein, A.R., 2008a. Gap junctions: multifaceted regulators of embryonic cortical development. *Trends in Neurosciences* 31: 243-250.
- Elias, L.A., Kriegstein, A.R., 2008b. Gap junctions: multifaceted regulators of embryonic cortical development. *Trends Neurosci* 31: 243-250.
- Emmons-Bell, M., Durant, F., Hammelman, J., Bessonov, N., Volpert, V., Morokuma, J., Pinet, K., Adams, D.S., Pietak, A., Lobo, D., Levin, M., 2015. Gap Junctional Blockade Stochastically Induces Different Species-Specific Head Anatomies in Genetically Wild-Type *Girardia dorotocephala* Flatworms. *Int J Mol Sci* 16: 27865-27896.
- Esser, A.T., Smith, K.C., Weaver, J.C., Levin, M., 2006. Mathematical model of morphogen electrophoresis through gap junctions. *Dev Dyn* 235: 2144-2159.
- Ewart, J.L., Cohen, M.F., Meyer, R.A., Huang, G.Y., Wessels, A., Gourdie, R.G., Chin, A.J., Park, S.M., Lazatin, B.O., Villabon, S., Lo, C.W., 1997. Heart and neural tube defects in transgenic mice overexpressing the Cx43 gap junction gene. *Development* 124: 1281-1292.
- Fan, R.J., Marin-Burgin, A., French, K.A., Otto Friesen, W., 2005. A dye mixture (Neurobiotin and Alexa 488) reveals extensive dye-coupling among neurons in leeches; physiology confirms the connections. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 191: 1157-1171.
- Fiorini, C., Mograbi, B., Cronier, L., Bourget, I., Decrouy, X., Nebout, M., Ferrua, B., Malassine, A., Samson, M., Fenichel, P., Segretain, D., Pointis, G., 2004. Dominant negative effect of connexin33 on gap junctional communication is mediated by connexin43 sequestration. *J Cell Sci Pt*.
- Fortier, P.A., Bagna, M., 2006. Estimating conductances of dual-recorded neurons within a network of coupled cells. *J Theor Biol* 240: 501-510.
- Francis, D., Stergiopoulos, K., Ek-Vitorin, J.F., Cao, F.L., Taffet, S.M., Delmar, M., 1999. Connexin diversity and gap junction regulation by pH. *Dev Gen* 24: 123-136.
- Franke, F., Jäckel, D., Dragas, J., Müller, J., 2012. High-density microelectrode array recordings and real-time spike sorting for closed-loop experiments: an emerging technology to study neural plasticity. *Frontiers in neural*
- Friston, K., 2010. Is the free-energy principle neurocentric? *Nature reviews. Neuroscience*.

- Friston, K., 2013. Active inference and free energy. *The Behavioral and brain sciences* 36: 212-213.
- Friston, K., Levin, M., Sengupta, B., Pezzulo, G., 2015. Knowing one's place: a free-energy approach to pattern regulation. *Journal of the Royal Society, Interface / the Royal Society* 12.
- Fukumoto, T., Blakely, R., Levin, M., 2005a. Serotonin transporter function is an early step in left-right patterning in chick and frog embryos. *Dev Neurosci* 27: 349-363.
- Fukumoto, T., Kema, I.P., Levin, M., 2005b. Serotonin signaling is a very early step in patterning of the left-right axis in chick and frog embryos. *Curr Biol* 15: 794-803.
- Funk, R., 2013. Ion Gradients in Tissue and Organ Biology. *Biological Systems*.
- Gairhe, S., Bauer, N.N., Gebb, S.A., McMurtry, I.F., 2012. Serotonin passes through myoendothelial gap junctions to promote pulmonary arterial smooth muscle cell differentiation. *American journal of physiology. Lung cellular and molecular physiology* 303: L767-777.
- Gentile, L., Cebria, F., Bartscherer, K., 2011. The planarian flatworm: an in vivo model for stem cell biology and nervous system regeneration. *Disease models & mechanisms* 4: 12-19.
- Gilboa, L., Forbes, A., Tazuke, S.I., Fuller, M.T., Lehmann, R., 2003. Germ line stem cell differentiation in *Drosophila* requires gap junctions and proceeds via an intermediate state. *Development* 130: 6625-6634.
- Goel, P., Mehta, A., 2013a. Learning Theories Reveal Loss of Pancreatic Electrical Connectivity in Diabetes as an Adaptive Response. *PLoS ONE* 8.
- Goel, P., Mehta, A., 2013b. Learning theories reveal loss of pancreatic electrical connectivity in diabetes as an adaptive response. *PLoS One* 8: e70366.
- Goldberg, G.S., Lampe, P.D., Nicholson, B.J., 1999. Selective transfer of endogenous metabolites through gap junctions composed of different connexins. *Nat Cell Biol* 1: 457-459.
- Goldberg, G.S., Moreno, A.P., Lampe, P.D., 2002. Gap junctions between cells expressing connexin 43 or 32 show inverse permselectivity to adenosine and ATP. *J Biol Chem* 277: 36725-36730.
- Goldberg, G.S., Valiunas, V., Brink, P.R., 2004. Selective permeability of gap junction channels. *Biochim Biophys Acta* 1662: 96-101.
- Govindan, J., Iovine, K.M., 2014. Hapln1a Is Required for Connexin43-Dependent Growth and Patterning in the Regenerating Fin Skeleton. *PLoS ONE* 9.
- Grek, C.L., Rhett, J.M., Ghatnekar, G.S., 2014. Cardiac to cancer: connecting connexins to clinical opportunity. *FEBS Lett* 588: 1349-1364.
- Gu, H., Ek-Vitorin, J.F., Taffet, S.M., Delmar, M., 2000. Coexpression of connexins 40 and 43 enhances the pH sensitivity of gap junctions: a model for synergistic interactions among connexins. *Circulation Research (Online)* 86: E98-E103.
- Gu, H., Smith, F.C., Taffet, S.M., Delmar, M., 2003a. High incidence of cardiac malformations in connexin40-deficient mice. *Circ Res* 93: 201-206.
- Gu, S., Yu, X.S., Yin, X., Jiang, J.X., 2003b. Stimulation of lens cell differentiation by gap junction protein connexin 45.6. *Invest. Ophthalmol. Vis. Sci.* 44: 2103-2111.
- Guthrie, S., 1984. Patterns of junctional communication in the early amphibian embryo. *Nature* 311: 149-151.

- Guthrie, S., Gilula, N., 1989. Gap junctional communication and development. *Trends in Neurosciences* 12: 12-16.
- Guthrie, S., Turin, L., Warner, A., 1988. Patterns of junctional communication during development of the early amphibian embryo. *Development* 103: 769-783.
- Gutierrez, G.J., Marder, E., 2013. Rectifying electrical synapses can affect the influence of synaptic modulation on output pattern robustness. *The Journal of Neuroscience*.
- Hallett, M.B., 1989. The unpredictability of cellular behavior: trivial or of fundamental importance to cell biology? *Perspect. Biol. Med.* 33: 110-119.
- Halley, J.D., Smith-Miles, K., Winkler, D.A., Kalkan, T., Huang, S., Smith, A., 2012. Self-organizing circuitry and emergent computation in mouse embryonic stem cells. *Stem cell research* 8: 324-333.
- Hamada, H., Watanabe, M., Lau, H.E., Nishida, T., Hasegawa, T., Parichy, D.M., Kondo, S., 2014. Involvement of Delta/Notch signaling in zebrafish adult pigment stripe patterning. *Development* 141: 1418-1418.
- Hamilton, N.B., Attwell, D., 2010. Do astrocytes really exocytose neurotransmitters? *Nature Reviews Neuroscience*.
- Handberg-Thorsager, M., Fernandez, E., Salo, E., 2008. Stem cells and regeneration in planarians. *Front Biosci* 13: 6374-6394.
- Hatler, J.M., Essner, J.J., Johnson, R.G., 2009. A gap junction connexin is required in the vertebrate left-right organizer. *Developmental biology* 336: 183-191.
- Hatton, G.I., Yang, Q.Z., 1996. Synaptically released histamine increases dye coupling among vasopressinergic neurons of the supraoptic nucleus: mediation by H1 receptors and cyclic nucleotides. *The Journal of neuroscience*.
- Hellmann, P., Grummer, R., Schirmacher, K., Rook, M., Traub, O., Winterhager, E., 1999. Transfection with different connexin genes alters growth and differentiation of human choriocarcinoma cells. *Exp Cell Res* 246: 480-490.
- Hironaka, K., Morishita, Y., 2012. Encoding and decoding of positional information in morphogen-dependent patterning. *Curr Opin Genet Dev* 22: 553-561.
- Hirschi, K.K., Burt, J.M., Hirschi, K.D., Dai, C., 2003. Gap junction communication mediates transforming growth factor-beta activation and endothelial-induced mural cell differentiation. *Circ Res* 93: 429-437.
- Hitomi, M., Deleyrolle, L.P., Mulkearns-Hubert, E.E., Jarrar, A., Li, M., Sinyuk, M., Otvos, B., Brunet, S., Flavahan, W.A., Hubert, C.G., Goan, W., Hale, J.S., Alvarado, A.G., Zhang, A., Rohaus, M., Oli, M., Vedam-Mai, V., Fortin, J.M., Futch, H.S., Griffith, B., Wu, Q., Xia, C.H., Gong, X., Ahluwalia, M.S., Rich, J.N., Reynolds, B.A., Lathia, J.D., 2015. Differential connexin function enhances self-renewal in glioblastoma. *Cell Rep* 11: 1031-1042.
- Hoel, E.P., Albantakis, L., Tononi, G., 2013. Quantifying causal emergence shows that macro can beat micro. *Proceedings of the National Academy of Sciences of the United States of America* 110: 19790-19795.
- Hofer, T., Venance, L., Giaume, C., 2002. Control and plasticity of intercellular calcium waves in astrocytes: a modeling approach. *J Neurosci* 22: 4850-4859.
- Hoptak-Solga, A.D., Klein, K.A., DeRosa, A.M., White, T.W., Iovine, K.M., 2007. Zebrafish short fin mutations in connexin43 lead to aberrant gap junctional intercellular communication. *FEBS Letters* 581: 3297-3302.

- Hoptak-Solga, A.D., Nielsen, S., Jain, I., Thummel, R., Hyde, D.R., Iovine, M.K., 2008. Connexin43 (GJA1) is required in the population of dividing cells during fin regeneration. *Dev Biol* 317: 541-548.
- Hou, M., Li, Y., Paul, D.L., 2013. A novel, highly sensitive method for assessing gap junctional coupling. *Journal of neuroscience methods* 220: 18-23.
- Hsieh, Y.W., Alqadah, A., Chuang, C.F., 2014. Asymmetric neural development in the *Caenorhabditis elegans* olfactory system. *genesis*.
- Huang, G.Y., Cooper, E.S., Waldo, K., Kirby, M.L., Gilula, N.B., Lo, C.W., 1998a. Gap junction-mediated cell-cell communication modulates mouse neural crest migration. *J Cell Biol* 143: 1725-1734.
- Huang, G.Y., Cooper, E.S., Waldo, K., Kirby, M.L., Gilula, N.B., Lo, C.W., 1998b. Gap junction-mediated cell-cell communication modulates mouse neural crest migration. *J Cell Biol* 143: 1725-1734.
- Huang, G.Y., Wessels, A., Smith, B.R., Linask, K.K., Ewart, J.L., Lo, C.W., 1998c. Alteration in connexin 43 gap junction gene dosage impairs conotruncal heart development. *Dev Biol* 198: 32-44.
- Huang, S., 2011. On the intrinsic inevitability of cancer: from foetal to fatal attraction. *Semin. Cancer Biol.* 21: 183-199.
- Huang, S., Ernberg, I., Kauffman, S., 2009. Cancer attractors: a systems view of tumors from a gene network dynamics and developmental perspective. *Semin Cell Dev Biol* 20: 869-876.
- Inaba, M., Yamanaka, H., Kondo, S., 2012. Pigment pattern formation by contact-dependent depolarization. *Science* 335: 677.
- Iovine, K.M., Higgins, E.P., Hindes, A., Coblitz, B., Johnson, S.L., 2005a. Mutations in connexin43 (GJA1) perturb bone growth in zebrafish fins. *Developmental Biology* 278: 208-219.
- Iovine, M.K., Higgins, E.P., Hindes, A., Coblitz, B., Johnson, S.L., 2005b. Mutations in connexin43 (GJA1) perturb bone growth in zebrafish fins. *Dev Biol* 278: 208-219.
- Irion, U., Frohnhofer, H., Krauss, J., Champollion, T., Maischein, H.-M., Geiger-Rudolph, S., Weiler, C., Nüsslein-Volhard, C., 2014. Gap junctions composed of connexins 41.8 and 39.4 are essential for colour pattern formation in zebrafish. *eLife* 3.
- Iwashita, M., Watanabe, M., Ishii, M., Chen, T., Johnson, S.L., Kurachi, Y., Okada, N., Kondo, S., 2006a. Pigment Pattern in jaguar/obelix Zebrafish Is Caused by a Kir7.1 Mutation: Implications for the Regulation of Melanosome Movement. *PLoS Genet* 2: e197.
- Iwashita, M., Watanabe, M., Ishii, M., Chen, T., Johnson, S.L., Kurachi, Y., Okada, N., Kondo, S., 2006b. Pigment pattern in jaguar/obelix zebrafish is caused by a Kir7.1 mutation: implications for the regulation of melanosome movement. *PLoS genetics* 2.
- Jaffe, L.F., 1981. The role of ionic currents in establishing developmental pattern. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 295: 553-566.

- Jin, E.J., Lee, S.Y., Jung, J.C., Bang, O.S., Kang, S.S., 2008. TGF-beta3 inhibits chondrogenesis of cultured chick leg bud mesenchymal cells via downregulation of connexin 43 and integrin beta4. *J Cell Physiol* 214: 345-353.
- Jindal, G.A., Goyal, Y., Burdine, R.D., Rauen, K.A., Shvartsman, S.Y., 2015. RASopathies: unraveling mechanisms with animal models. *Disease models & mechanisms* 8: 1167.
- Jo, S.H., Chang, T., Ebong, I., Bhadviya, B.B., Mazumder, P., Lu, W., 2010. Nanoscale memristor device as synapse in neuromorphic systems. *Nano letters* 10: 1297-1301.
- Juszczak, G.R., Swiergiel, A.H., 2009. Properties of gap junction blockers and their behavioural, cognitive and electrophysiological effects: animal and human studies. *Prog Neuropsychopharmacol Biol Psychiatry* 33: 181-198.
- Kalimi, G.H., Lo, C.W., 1989. Gap junctional communication in the extraembryonic tissues of the gastrulating mouse embryo. *J Cell Biol* 109: 3015-3026.
- Kanady, J.D., Dellinger, M.T., Munger, S.J., Witte, M.H., Simon, A.M., 2011. Connexin37 and Connexin43 deficiencies in mice disrupt lymphatic valve development and result in lymphatic disorders including lymphedema and chylothorax. *Developmental biology* 354: 253-266.
- Kanady, J.D., Munger, S.J., Witte, M.H., Simon, A.M., 2015. Combining *Foxc2* and Connexin37 deletions in mice leads to severe defects in lymphatic vascular growth and remodeling. *Developmental biology* 405: 33-46.
- Kandouz, M., Batist, G., 2010. Gap junctions and connexins as therapeutic targets in cancer. *Expert Opin Ther Targets* 14: 681-692.
- Katakowski, M., Buller, B., Wang, X., Rogers, T., Chopp, M., 2010. Functional microRNA is transferred between glioma cells. *Cancer Res* 70: 8259-8263.
- Kim, J.Y., Cho, S.W., Lee, M.J., Hwang, H.J., Lee, J.M., Lee, S.I., Muramatsu, T., Shimono, M., Jung, H.S., 2005. Inhibition of connexin 43 alters *Shh* and *Bmp-2* expression patterns in embryonic mouse tongue. *Cell Tissue Res* 320: 409-415.
- Kirchhoff, S., Kim, J.S., Hagendorff, A., Thonnissen, E., Kruger, O., Lamers, W.H., Willecke, K., 2000. Abnormal cardiac conduction and morphogenesis in connexin40 and connexin43 double-deficient mice. *Circ Res* 87: 399-405.
- Kondo, S., 2002. The reaction-diffusion system: a mechanism for autonomous pattern formation in the animal skin. *Genes Cells* 7: 535-541.
- Kondo, S., Asai, R., 1995. A reaction-diffusion wave on the skin of the marine angelfish *Pomacanthus*. *Nature* 376: 765-768.
- Kotini, M., Mayor, R., 2015. Connexins in migration during development and cancer. *Dev Biol* 401: 143-151.
- Kumai, M., Nishii, K., Nakamura, K., Takeda, N., Suzuki, M., Shibata, Y., 2000. Loss of connexin45 causes a cushion defect in early cardiogenesis. *Development* 127: 3501-3512.
- Larre, I., Ponce, A., Fiorentino, R., Shoshani, L., Contreras, R.G., Cereijido, M., 2006. Contacts and cooperation between cells depend on the hormone ouabain. *Proc Natl Acad Sci U S A* 103: 10911-10916.
- Law, R., Levin, M., 2015. Bioelectric memory: modeling resting potential bistability in amphibian embryos and mammalian cells. *Theor Biol Med Model* 12: 22.

- le Feber, J., Stegenga, J., Rutten, W., 2008. Do external stimuli, applied to train cultured cortical networks, disturb the balance between activity and connectivity? Conference proceedings : ... Annual International Conference of the IEEE Engineering in Medicine and Biology Society. IEEE Engineering in Medicine and Biology Society. Annual Conference 2008: 5081-5084.
- le Feber, J., Stegenga, J., Rutten, W.L.C., 2010. The Effect of Slow Electrical Stimuli to Achieve Learning in Cultured Networks of Rat Cortical Neurons. PLoS ONE 5.
- Lecanda, F., Warlow, P.M., Sheikh, S., Furlan, F., Steinberg, T.H., Civitelli, R., 2000a. Connexin43 deficiency causes delayed ossification, craniofacial abnormalities, and osteoblast dysfunction. The Journal of cell biology 151: 931-944.
- Lecanda, F., Warlow, P.M., Sheikh, S., Furlan, F., Steinberg, T.H., Civitelli, R., 2000b. Connexin43 deficiency causes delayed ossification, craniofacial abnormalities, and osteoblast dysfunction. J Cell Biol 151: 931-944.
- Lehmann, C., Lechner, H., Loer, B., Knieps, M., Herrmann, S., Famulok, M., Bauer, R., Hoch, M., 2006. Heteromerization of Innexin Gap Junction Proteins Regulates Epithelial Tissue Organization in Drosophila. Mol Biol Cell.
- Leithe, E., Sirnes, S., Omori, Y., Rivedal, E., 2006. Downregulation of gap junctions in cancer cells. Crit Rev Oncog 12: 225-256.
- Leppik, L.P., Froemel, D., Slavici, A., Ovadia, Z.N., Hudak, L., Henrich, D., Marzi, I., Barker, J.H., 2015. Effects of electrical stimulation on rat limb regeneration, a new look at an old model. Sci Rep 5: 18353.
- Levin, M., 1998. Left-right asymmetry and the chick embryo. Semin Cell Dev Biol 9: 67-76.
- Levin, M., 2002. Isolation and community: a review of the role of gap-junctional communication in embryonic patterning. J Membr Biol 185: 177-192.
- Levin, M., 2007. Gap junctional communication in morphogenesis. Prog. Biophys. Mol. Biol. 94: 186-206.
- Levin, M., 2011. The wisdom of the body: future techniques and approaches to morphogenetic fields in regenerative medicine, developmental biology and cancer. Regenerative medicine 6: 667-673.
- Levin, M., 2012. Morphogenetic fields in embryogenesis, regeneration, and cancer: non-local control of complex patterning. Bio Systems 109: 243-261.
- Levin, M., 2014. Molecular bioelectricity: how endogenous voltage potentials control cell behavior and instruct pattern regulation in vivo. Mol Biol Cell 25: 3835-3850.
- Levin, M., Buznikov, G.A., Lauder, J.M., 2006. Of minds and embryos: left-right asymmetry and the serotonergic controls of pre-neural morphogenesis. Dev Neurosci 28: 171-185.
- Levin, M., Mercola, M., 1998a. The compulsion of chirality: toward an understanding of left-right asymmetry. Genes Dev 12: 763-769.
- Levin, M., Mercola, M., 1998b. Gap junctions are involved in the early generation of left-right asymmetry. Dev Biol 203: 90-105.
- Levin, M., Mercola, M., 1999. Gap junction-mediated transfer of left-right patterning signals in the early chick blastoderm is upstream of Shh asymmetry in the node. Development 126: 4703-4714.
- Levin, M., Mercola, M., 2000. Expression of connexin 30 in Xenopus embryos and its involvement in hatching gland function. Developmental Dynamics 219: 96-101.

- Levin, M., Nascone, N., 1997. Two molecular models of initial left-right asymmetry generation. *Med Hypotheses* 49: 429-435.
- Levin, M., Thorlin, T., Robinson, K.R., Nogi, T., Mercola, M., 2002. Asymmetries in H⁺/K⁺-ATPase and cell membrane potentials comprise a very early step in left-right patterning. *Cell* 111: 77-89.
- Li, S., He, H., Zhang, G., Wang, F., Zhang, P., Tan, Y., 2015. Connexin43-containing gap junctions potentiate extracellular Ca-induced odontoblastic differentiation of human dental pulp stem cells via Erk1/2. *Exp Cell Res*.
- Li, W.E., Waldo, K., Linask, K.L., Chen, T., Wessels, A., Parmacek, M.S., Kirby, M.L., Lo, C.W., 2002. An essential role for connexin43 gap junctions in mouse coronary artery development. *Development* 129: 2031-2042.
- Lim, P.K., Bliss, S.A., Patel, S.A., Taborga, M., Dave, M.A., Gregory, L.A., Greco, S.J., Bryan, M., Patel, P.S., Rameshwar, P., 2011. Gap junction-mediated import of microRNA from bone marrow stromal cells can elicit cell cycle quiescence in breast cancer cells. *Cancer Res* 71: 1550-1560.
- Lo, C.W., 1996. The role of gap junction membrane channels in development. *J Bioenerg Biomembr* 28: 379-385.
- Lo, C.W., Cohen, M.F., Huang, G.Y., Lazatin, B.O., Patel, N., Sullivan, R., Pauken, C., Park, S.M., 1997. Cx43 gap junction gene expression and gap junctional communication in mouse neural crest cells. *Dev Gen* 20: 119-132.
- Lo, C.W., Gilula, N.B., 1979. Gap junctional communication in the post-implantation mouse embryo. *Cell* 18: 411-422.
- Lo, C.W., Waldo, K.L., Kirby, M.L., 1999. Gap junction communication and the modulation of cardiac neural crest cells. *Trends in Cardiovascular Medicine* 9: 63-69.
- Lobikin, M., Chernet, B., Lobo, D., Levin, M., 2012. Resting potential, oncogene-induced tumorigenesis, and metastasis: the bioelectric basis of cancer in vivo. *Physical biology* 9: 065002.
- Lobo, D., Beane, W.S., Levin, M., 2012. Modeling planarian regeneration: a primer for reverse-engineering the worm. *PLoS computational biology* 8: e1002481.
- Lobo, D., Solano, M., Bubenik, G.A., Levin, M., 2014. A linear-encoding model explains the variability of the target morphology in regeneration. *Journal of the Royal Society, Interface / the Royal Society* 11: 20130918.
- Loch-Caruso, R., Trosko, J.E., 1985. Inhibited intercellular communication as a mechanistic link between teratogenesis and carcinogenesis. *Crit. Rev. Toxicol.* 16: 157-183.
- Loewenstein, W.R., 1969. Transfer of information through cell junctions and growth control. *Proc Can Cancer Conf* 8: 162-170.
- Loewenstein, W.R., 1979. Junctional intercellular communication and the control of growth. *Biochim Biophys Acta* 560: 1-65.
- Loewenstein, W.R., 1980. Junctional cell-to-cell communication and growth control. *Ann N Y Acad Sci* 339: 39-45.
- Loewenstein, W.R., Kanno, Y., 1966. Intercellular communication and the control of tissue growth: lack of communication between cancer cells. *Nature* 209: 1248-1249.

- Maeda, S., Tsukihara, T., 2011. Structure of the gap junction channel and its implications for its biological functions. *Cell Mol Life Sci* 68: 1115-1129.
- Maffini, M.V., Calabro, J.M., Soto, A.M., Sonnenschein, C., 2005. Stromal regulation of neoplastic development: age-dependent normalization of neoplastic mammary cells by mammary stroma. *The American journal of pathology* 167: 1405-1410.
- Mantz, J., 1992. Effects of intravenous anesthetics on neurons of the central nervous system: mechanisms of cellular and molecular action. *Annales Francaises d Anesthesie et de Reanimation* 11: 540-557.
- Marder, E., 2009. Electrical synapses: rectification demystified. *Current Biology*.
- Marins, M., Xavier, A.L., Viana, N.B., Fortes, F.S., Froes, M.M., Menezes, J.R., 2009. Gap junctions are involved in cell migration in the early postnatal subventricular zone. *Developmental neurobiology* 69: 715-730.
- Maximino, C., Puty, B., Oliveira, M.K.R., Herculano, A.M., 2013. Behavioral and neurochemical changes in the zebrafish leopard strain. *Genes, Brain and Behavior* 12: 576-582.
- McCaig, C.D., Rajnicek, A.M., Song, B., Zhao, M., 2005. Controlling cell behavior electrically: current views and future potential. *Physiol Rev* 85: 943-978.
- McConnell, J.V., Jacobson, A.L., Kimble, D.P., 1959. The effects of regeneration upon retention of a conditioned response in the planarian. *Journal of Comparative Physiology and Psychology* 52: 1-5.
- Mehta, P.P., Hotz-Wagenblatt, A., Rose, B., Shalloway, D., Loewenstein, W.R., 1991. Incorporation of the gene for a cell-cell channel protein into transformed cells leads to normalization of growth. *J Membr Biol* 124: 207-225.
- Melloy, P.G., Kusnierczyk, M.K., Meyer, R.A., Lo, C.W., Desmond, M.E., 2005. Overexpression of connexin43 alters the mutant phenotype of midgestational wnt-1 null mice resulting in recovery of the midbrain and cerebellum. *Anat Rec A Discov Mol Cell Evol Biol* 283: 224-238.
- Mendoza-Naranjo, A., Saez, P.J., Johansson, C.C., Ramirez, M., Mandakovic, D., Pereda, C., Lopez, M.N., Kiessling, R., Saez, J.C., Salazar-Onfray, F., 2007. Functional gap junctions facilitate melanoma antigen transfer and cross-presentation between human dendritic cells. *J. Immunol.* 178: 6949-6957.
- Merrifield, P.A., Laird, D.W., 2015. Connexins in skeletal muscle development and disease. *Seminars in cell & developmental biology*.
- Mesnil, M., Crespín, S., Avanzo, J.L., Zaidan-Dagli, M.L., 2005. Defective gap junctional intercellular communication in the carcinogenic process. *Biochim Biophys Acta* 1719: 125-145.
- Moreno, E., 2008. Is cell competition relevant to cancer? *Nature reviews. Cancer* 8: 141-147.
- Morley, G.E., Ek-Vitorin, J.F., Taffet, S.M., Delmar, M., 1997. Structure of connexin43 and its regulation by pHi. *J Cardiovasc Electrophysiol* 8: 939-951.
- Morokuma, J., Blackiston, D., Levin, M., 2008. KCNQ1 and KCNE1 K⁺ channel components are involved in early left-right patterning in *Xenopus laevis* embryos. *Cell Physiol Biochem* 21: 357-372.
- Nagajski, D., Guthrie, S., Ford, C., Warner, A., 1989. The correlation between patterns of dye transfer through gap junctions and future developmental fate in *Xenopus*. *Development* 105: 747-752.

- Nguyen, L.M., Pozzoli, M., Hraha, T.H., Benninger, R.K., 2014. Decreasing cx36 gap junction coupling compensates for overactive KATP channels to restore insulin secretion and prevent hyperglycemia in a mouse model of neonatal diabetes. *Diabetes* 63: 1685-1697.
- Nicholson, B.J., Weber, P.A., Cao, F., Chang, H., Lampe, P., Goldberg, G., 2000. The molecular basis of selective permeability of connexins is complex and includes both size and charge. *Braz J Med Biol Res* 33: 369-378.
- Nishii, K., Kumai, M., Shibata, Y., 2001. Regulation of the epithelial-mesenchymal transformation through gap junction channels in heart development. *Trends in cardiovascular medicine* 11: 213-218.
- Noble, D., 2015. Conrad Waddington and the origin of epigenetics. *J Exp Biol* 218: 816-818.
- Nogi, T., Levin, M., 2005. Characterization of innexin gene expression and functional roles of gap-junctional communication in planarian regeneration. *Dev Biol* 287: 314-335.
- Nuccitelli, R., 2003. Endogenous electric fields in embryos during development, regeneration and wound healing. *Radiat Prot Dosimetry* 106: 375-383.
- Nygren, A., Giles, W.R., 2000. Mathematical simulation of slowing of cardiac conduction velocity by elevated extracellular. *Ann Biomed Eng* 28: 951-957.
- O'Brien, J., 2014. The ever-changing electrical synapse. *Current opinion in neurobiology*.
- Oh, S.K., Shin, J.O., Baek, J.I., Lee, J., Bae, J.W., Ankamerddy, H., Kim, M.J., Huh, T.L., Ryoo, Z.Y., Kim, U.K., Bok, J., Lee, K.Y., 2015. Pannexin 3 is required for normal progression of skeletal development in vertebrates. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*.
- Olson, D.J., Christian, J.L., Moon, R.T., 1991a. Effect of wnt-1 and related proteins on gap junctional communication in *Xenopus* embryos. *Science* 252: 1173-1176.
- Olson, D.J., Christian, J.L., Moon, R.T., 1991b. Effect of wnt-1 and related proteins on gap junctional communication in *Xenopus* embryos. *Science* 252: 1173-1176.
- Olson, D.J., Moon, R.T., 1992. Distinct effects of ectopic expression of Wnt-1, activin B, and bFGF on gap junctional permeability in 32-cell *Xenopus* embryos. *Developmental Biology* 151: 204-212.
- Omori, Y., Duflot-Dancer, A., Mesnil, M., Yamasaki, H., 1998. Role of connexin (gap junction) genes in cell growth control. *Toxicol Lett* 96-97: 105-110.
- Omori, Y., Yamasaki, H., 1998. Mutated connexin43 proteins inhibit rat glioma cell growth suppression mediated by wild-type connexin43 in a dominant-negative manner. *Int J Cancer* 78: 446-453.
- Oviedo, N.J., Levin, M., 2007. smedinx-11 is a planarian stem cell gap junction gene required for regeneration and homeostasis. *Development* 134: 3121-3131.
- Oviedo, N.J., Morokuma, J., Walentek, P., Kema, I.P., Gu, M.B., Ahn, J.M., Hwang, J.S., Gojobori, T., Levin, M., 2010. Long-range neural and gap junction protein-mediated cues control polarity during planarian regeneration. *Dev Biol* 339: 188-199.
- Paemeleire, K., Martin, P.E., Coleman, S.L., Fogarty, K.E., Carrington, W.A., Leybaert, L., Tuft, R.A., Evans, W.H., Sanderson, M.J., 2000. Intercellular calcium waves in

- HeLa cells expressing GFP-labeled connexin 43, 32, or 26. *Mol Biol Cell* 11: 1815-1827.
- Pai, V.P., Aw, S., Shomrat, T., Lemire, J.M., Levin, M., 2012. Transmembrane voltage potential controls embryonic eye patterning in *Xenopus laevis*. *Development* 139: 313-323.
- Pai, V.P., Lemire, J.M., Pare, J.F., Lin, G., Chen, Y., Levin, M., 2015a. Endogenous Gradients of Resting Potential Instructively Pattern Embryonic Neural Tissue via Notch Signaling and Regulation of Proliferation. *The Journal of Neuroscience* 35: 4366-4385.
- Pai, V.P., Martyniuk, C.J., Echeverri, K., Sundelacruz, S., Kaplan, D.L., Levin, M., 2015b. Genome-wide analysis reveals conserved transcriptional responses downstream of resting potential change in *Xenopus* embryos, axolotl regeneration, and human mesenchymal cell differentiation. *Regeneration* n/a-n/a.
- Palacios-Prado, N., 2009. Heterotypic gap junction channels as voltage-sensitive valves for intercellular signaling. *Proceedings of the ...*
- Palacios-Prado, N., Huetteroth, W., Pereda, A.E., 2014. Hemichannel composition and electrical synaptic transmission: molecular diversity and its implications for electrical rectification. *Front Cell Neurosci* 8: 324.
- Panakova, D., Werdich, A.A., Macrae, C.A., 2010. Wnt11 patterns a myocardial electrical gradient through regulation of the L-type Ca^{2+} channel. *Nature* 466: 874-878.
- Paraguassu-Braga, F.H., Borojevic, R., Bouzas, L.F., Barcinski, M.A., Bonomo, A., 2003. Bone marrow stroma inhibits proliferation and apoptosis in leukemic cells through gap junction-mediated cell communication. *Cell Death Differ* 10: 1101-1108.
- Parak, W.J., Domke, J., George, M., Kardinal, A., Radmacher, M., Gaub, H.E., de Roos, A.D.G., Theuvenet, A.P.R., Wiegand, G., Sackmann, E., Behrends, J.C., 1999. Electrically Excitable Normal Rat Kidney Fibroblasts: A New Model System for Cell-Semiconductor Hybrids. *Biophysical Journal* 76: 1659-1667.
- Paul, D., Yu, K., Bruzzone, R., Gimlich, R., Goodenough, D., 1995. Expression of a dominant negative inhibitor of intercellular communication in the early *Xenopus* embryo causes delamination and extrusion of cells. *Development* 121: 371-381.
- Pearson, R.A., Dale, N., Llaudet, E., Mobbs, P., 2005. ATP released via gap junction hemichannels from the pigment epithelium regulates neural retinal progenitor proliferation. *Neuron* 46: 731-744.
- Peracchia, C., 2004. Chemical gating of gap junction channels; roles of calcium, pH and calmodulin. *Biochim Biophys Acta* 1662: 61-80.
- Peracchia, C., Bernardini, G., Peracchia, L.L., 1983. Is calmodulin involved in the regulation of gap junction permeability? *Pflügers Arch* 399: 152-154.
- Perathoner, S., Daane, J.M., Henrion, U., Seeböhm, G., Higdon, C.W., Johnson, S.L., Nusslein-Volhard, C., Harris, M.P., 2014. Bioelectric signaling regulates size in zebrafish fins. *PLoS genetics* 10: e1004080.
- Pereda, A.E., 2014. Electrical synapses and their functional interactions with chemical synapses. *Nature Reviews Neuroscience* 15: 250-263.

- Pereda, A.E., Curti, S., Hoge, G., Cachope, R., Flores, C.E., Rash, J.E., 2013a. Gap junction-mediated electrical transmission: Regulatory mechanisms and plasticity. *Biochimica et Biophysica Acta (BBA) - Biomembranes* 1828: 134-146.
- Pereda, A.E., Curti, S., Hoge, G., Cachope, R., Flores, C.E., Rash, J.E., 2013b. Gap junction-mediated electrical transmission: regulatory mechanisms and plasticity. *Biochim Biophys Acta* 1828: 134-146.
- Pereira, A., Furlan, F., 2010. Astrocytes and human cognition: Modeling information integration and modulation of neuronal activity. *Progress in Neurobiology* 92: 405-420.
- Pershin, Y.V., Di Ventra, M., 2010. Experimental demonstration of associative memory with memristive neural networks. *Neural Netw* 23: 881-886.
- Pezzulo, G., Levin, M., 2015. Re-membering the body: applications of computational neuroscience to the top-down control of regeneration of limbs and other complex organs. *Integr Biol (Camb)*.
- Pezzulo, G., Rigoli, F., Friston, K., 2015. Active Inference, homeostatic regulation and adaptive behavioural control. *Progress in neurobiology*.
- Pfeiffer, T., Kraushaar, U., Düfer, M., Schönecker, S., Haspel, D., Günther, E., Drews, G., Krippeit-Drews, P., 2011. Rapid functional evaluation of beta-cells by extracellular recording of membrane potential oscillations with microelectrode arrays. *Pflügers Archiv - European Journal of Physiology* 462: 835-840.
- Phelan, P., 2005. Innexins: members of an evolutionarily conserved family of gap-junction proteins. *Biochim Biophys Acta* 1711: 225-245.
- Piccolino, M., Neyton, J., 1984. Decrease of gap junction permeability induced by dopamine and cyclic adenosine 3': 5'-monophosphate in horizontal cells of turtle retina. *The Journal of*
- Pimashkin, A., Gladkov, A., Mukhina, I., Kazantsev, V., 2013. Adaptive enhancement of learning protocol in hippocampal cultured networks grown on multielectrode arrays. *Frontiers in Neural Circuits* 7.
- Pitts, J.D., Finbow, M.E., Kam, E., 1988. Junctional communication and cellular differentiation. *Br J Cancer Suppl* 9: 52-57.
- Pizard, A., Burgon, P.G., Paul, D.L., Bruneau, B.G., Seidman, C.E., Seidman, J.G., 2005. Connexin 40, a target of transcription factor tbx5, patterns wrist, digits, and sternum. *Mol Cell Biol* 25: 5073-5083.
- Pizzuti, A., Flex, E., Mingarelli, R., Salpietro, C., Zelante, L., Dallapiccola, B., 2004. A homozygous GJA1 gene mutation causes a Hallermann-Streiff/ODDD spectrum phenotype. *Hum Mutat* 23: 286.
- Postma, F., Liu, C.H., Dietsche, C., 2011. Electrical synapses formed by connexin36 regulate inhibition-and experience-dependent plasticity. *Proceedings of the*
- Potter, V.R., 1980. Initiation and promotion in cancer formation: the importance of studies on intercellular communication. *Yale J. Biol. Med.* 53: 367-384.
- Ramsdell, A.F., Yost, H.J., 1998. Molecular mechanisms of vertebrate left-right development. *Trends Genet* 14: 459-465.
- Reaume, A.G., De Sousa, P.A., Kilkarni, S., Langille, B.L., Zhu, D., Davies, T.C., Juneja, S.C., Kidder, G.M., Rossant, J., 1995. Cardiac malformation in neonatal mice lacking connexin43. *Science* 267: 1831-1834.

- Reddien, P.W., Sanchez Alvarado, A., 2004. Fundamentals of planarian regeneration. *Annu Rev Cell Dev Biol* 20: 725-757.
- Rela, L., Szczupak, L., 2004. Gap junctions: their importance for the dynamics of neural circuits. *Mol Neurobiol* 30: 341-357.
- Richard, M., Hoch, M., 2015. *Drosophila* eye size is determined by Innexin 2-dependent Decapentaplegic signalling. *Developmental biology*.
- Rimkute, L., Jotautis, V., Marandykina, A., Sveikatiene, R., Antanaviciute, I., Skeberdis, V.A., 2016. The role of neural connexins in HeLa cell mobility and intercellular communication through tunneling tubes. *BMC Cell Biol* 17: 3.
- Robinson, S.R., Hampson, E.C., Munro, M.N., Vaney, D.I., 1993. Unidirectional coupling of gap junctions between neuroglia. *Science* 262: 1072-1074.
- Rogers, K.W., Schier, A.F., 2011. Morphogen gradients: from generation to interpretation. *Ann. Rev. Cell Dev. Biol.* 27: 377-407.
- Rörig, B., Sutor, B., 1996. Nitric oxide-stimulated increase in intracellular cGMP modulates gap junction coupling in rat neocortex. *Neuroreport*.
- Rose, B., Mehta, P.P., Loewenstein, W.R., 1993. Gap-junction protein gene suppresses tumorigenicity. *Carcinogenesis* 14: 1073-1075.
- Rosenbaum, M.B., Blanco, H.H., Elizari, M.V., Lazzari, J.O., Davidenko, J.M., 1982. Electrotonic modulation of the T wave and cardiac memory. *The American journal of cardiology* 50: 213-222.
- Rubin, H., 1985. Cancer as a dynamic developmental disorder. *Cancer Res* 45: 2935-2942.
- Salo, E., Abril, J.F., Adell, T., Cebria, F., Eckelt, K., Fernandez-Taboada, E., Handberg-Thorsager, M., Iglesias, M., Molina, M.D., Rodriguez-Esteban, G., 2009. Planarian regeneration: achievements and future directions after 20 years of research. *Int J Dev Biol* 53: 1317-1327.
- Saraga, F., Ng, L., Skinner, F.K., 2006. Distal Gap Junctions and Active Dendrites Can Tune Network Dynamics. *Journal of Neurophysiology* 95: 1669-1682.
- Sarnat, H.B., and Netsky, M.G., 1985. The brain of the planarian as the ancestor of the human brain. *Can J Neurol Sci*.
- Scarle, S., 2008. Implications of the Turing completeness of reaction-diffusion models. Microsoft.
- Scemes, E., Suadicani, S.O., Dahl, G., Spray, D.C., 2007. Connexin and pannexin mediated cell-cell communication. *Neuron glia biology* 3: 199-208.
- Schiffmann, Y., 1991. An hypothesis: phosphorylation fields as the source of positional information and cell differentiation--(cAMP, ATP) as the universal morphogenetic Turing couple. *Prog Biophys Mol Biol* 56: 79-105.
- Schiffmann, Y., 2005. Induction and the Turing-Child field in development. *Prog Biophys Mol Biol* 89: 36-92.
- Schiffmann, Y., 2008. The Turing-Child energy field as a driver of early mammalian development. *Prog. Biophys. Mol. Biol.* 98: 107-117.
- Schönecker, S., Kraushaar, U., Düfer, M., Sahr, A., Härdtner, C., Guenther, E., Walther, R., Lendeckel, U., Barthlen, W., Krippeit-Drews, P., Drews, G., 2014. Long-term culture and functionality of pancreatic islets monitored using microelectrode arrays. *Integrative Biology* 6: 540-544.

- Schumacher, J.A., Hsieh, Y.W., Chen, S., Pirri, J.K., 2012. Intercellular calcium signaling in a gap junction-coupled cell network establishes asymmetric neuronal fates in *C. elegans*.
- Shahaf, G., Marom, S., 2001. Learning in Networks of Cortical Neurons. *The Journal of Neuroscience* 21: 8782-8788.
- Sherer, N.M., 2013. Long-distance relationships: do membrane nanotubes regulate cell-cell communication and disease progression? *Mol Biol Cell* 24: 1095-1098.
- Sherman, A., Rinzel, J., 1991. Model for synchronization of pancreatic beta-cells by gap junction coupling. *Biophys J* 59: 547-559.
- Shigetomi, E., Kracun, S., Sofroniew, M.V., 2010. A genetically targeted optical sensor to monitor calcium signals in astrocyte processes. *Nature neuroscience*.
- Shomrat, T., Levin, M., 2013. An automated training paradigm reveals long-term memory in planarians and its persistence through head regeneration. *The Journal of experimental biology* 216: 3799-3810.
- Simon, A.M., McWhorter, A.R., Dones, J.A., Jackson, C.L., Chen, H., 2004. Heart and head defects in mice lacking pairs of connexins. *Dev Biol* 265: 369-383.
- Simon, S., 2009. Implications of the Turing completeness of reaction-diffusion models, informed by GPGPU simulations on an Xbox 360: Cardiac arrhythmias, re-entry and the Halting problem. *Computational Biology and Chemistry* 33: 253-260.
- Sims, K., Eble, D.M., Iovine, K.M., 2009a. Connexin43 regulates joint location in zebrafish fins. *Developmental Biology* 327: 410-418.
- Sims, K., Jr., Eble, D.M., Iovine, M.K., 2009b. Connexin43 regulates joint location in zebrafish fins. *Dev Biol* 327: 410-418.
- Sin, W.C., Crespin, S., Mesnil, M., 2012. Opposing roles of connexin43 in glioma progression. *Biochim Biophys Acta* 1818: 2058-2067.
- Slusarski, D.C., Pelegri, F., 2007. Calcium signaling in vertebrate embryonic patterning and morphogenesis. *Dev Biol* 307: 1-13.
- Smendziuk, C.M., Messenberg, A., Vogl, A.W., Tanentzapf, G., 2015. Bi-directional gap junction-mediated soma-germline communication is essential for spermatogenesis. *Development* 142: 2598-2609.
- Sohl, G., Willecke, K., 2004. Gap junctions and the connexin protein family. *Cardiovasc Res* 62: 228-232.
- Solan, J.L., Lampe, P.D., 2014. Specific Cx43 phosphorylation events regulate gap junction turnover in vivo. *FEBS Lett* 588: 1423-1429.
- Soto, A.M., Sonnenschein, C., 2011. The tissue organization field theory of cancer: a testable replacement for the somatic mutation theory. *BioEssays* 33: 332-340.
- Speder, P., Petzoldt, A., Suzanne, M., Noselli, S., 2007. Strategies to establish left/right asymmetry in vertebrates and invertebrates. *Curr Opin Genet Dev* 17: 351-358.
- Spira, M.E., Hai, A., 2013. Multi-electrode array technologies for neuroscience and cardiology. *Nature Nanotechnology* 8: 83-94.
- Stains, J.P., Civitelli, R., 2004. Gap Junctions Regulate Extracellular Signal-regulated Kinase (ERK) Signaling to Affect Gene Transcription. *J Mol. Biol. Cell*.
- Stains, J.P., Civitelli, R., 2005. Gap junctions regulate extracellular signal-regulated kinase signaling to affect gene transcription. *Mol Biol Cell* 16: 64-72.
- Stains, J.P., Lecanda, F., Screen, J., Towler, D.A., Civitelli, R., 2003. Gap junctional communication modulates gene transcription by altering the recruitment of Sp1

- and Sp3 to connexin-response elements in osteoblast promoters. *J Biol Chem* 278: 24377-24387.
- Starich, T.A., Hall, D.H., Greenstein, D., 2014. Two Classes of Gap Junction Channels Mediate Soma-Germline Interactions Essential for Germline Proliferation and Gametogenesis in *Caenorhabditis elegans*. *Genetics*.
- Starich, T.A., Miller, A., Nguyen, R.L., Hall, D.H., Shaw, J.E., 2003. The *caenorhabditis elegans* innexin INX-3 is localized to gap junctions and is essential for embryonic development. *Dev Biol* 256: 403-417.
- Stehberg, J., Moraga-Amaro, R., Salazar, C., Becerra, A., Echeverria, C., Orellana, J.A., Bultynck, G., Ponsaerts, R., Leybaert, L., Simon, F., Saez, J.C., Retamal, M.A., 2012. Release of gliotransmitters through astroglial connexin 43 hemichannels is necessary for fear memory consolidation in the basolateral amygdala. *FASEB J* 26: 3649-3657.
- Stewart, A., Cachat, J., Gaikwad, S., Robinson, K.S.L., Gebhardt, M., Kalueff, A.V., 2013. Perspectives on experimental models of serotonin syndrome in zebrafish. *Neurochemistry international* 62: 893-902.
- Steyn-Ross, M.L., Steyn-Ross, D.A., Wilson, M.T., Sleight, J.W., 2007. Gap junctions mediate large-scale Turing structures in a mean-field cortex driven by subcortical noise. *Physical review. E, Statistical, nonlinear, and soft matter physics* 76: 011916.
- Stoletov, K., Strnadel, J., Zardouzian, E., Momiyama, M., Park, F.D., Kelber, J.A., Pizzo, D.P., Hoffman, R., Vandenberg, S.R., Klemke, R.L., 2013. Role of connexins in metastatic breast cancer and melanoma brain colonization. *J Cell Sci*.
- Stone, J.R., 1997. The spirit of D'arcy Thompson dwells in empirical morphospace. *Math. Biosci.* 142: 13-30.
- Sullivan, R., Huang, G.Y., Meyer, R.A., Wessels, A., Linask, K.K., Lo, C.W., 1998. Heart malformations in transgenic mice exhibiting dominant negative inhibition of gap junctional communication in neural crest cells. *Developmental Biology* 204: 224-234.
- Sundelacruz, S., Levin, M., Kaplan, D.L., 2009. Role of membrane potential in the regulation of cell proliferation and differentiation. *Stem cell reviews and reports* 5: 231-246.
- Sutor, B., Hagerty, T., 2005. Involvement of gap junctions in the development of the neocortex. *Biochim Biophys Acta* 1719: 59-68.
- Takagi, H., Kaneko, K., 2005. Dynamical systems basis of metamorphosis: diversity and plasticity of cellular states in reaction diffusion network. *J Theor Biol* 234: 173-186.
- Tarin, D., 2012. Clinical and Biological Implications of the Tumor Microenvironment. *Cancer Microenviron.*
- Temme, A., Buchmann, A., Gabriel, H.D., Nelles, E., Schwarz, M., Willecke, K., 1997. High incidence of spontaneous and chemically induced liver tumors in mice deficient for connexin32. *Curr Biol* 7: 713-716.
- Ton, Q.V., Iovine, K.M., 2012. Semaphorin3d mediates Cx43-dependent phenotypes during fin regeneration. *Developmental Biology* 366: 195-203.
- Ton, Q.V., Iovine, K.M., 2013. Identification of an *evx1*-Dependent Joint-Formation Pathway during FIN Regeneration. *PLoS ONE* 8.

- Ton, Q.V., Kathryn Iovine, M., 2012. Semaphorin3d mediates Cx43-dependent phenotypes during fin regeneration. *Developmental biology* 366: 195-203.
- Toyofuku, T., Yabuki, M., Otsu, K., Kuzuya, T., Hori, M., Tada, M., 1998. Intercellular calcium signaling via gap junction in connexin-43-transfected cells. *J Biol Chem* 273: 1519-1528.
- Trosko, J.E., 2005. The role of stem cells and gap junctions as targets for cancer chemoprevention and chemotherapy. *Biomed Pharmacother* 59 Suppl 2: S326-331.
- Trosko, J.E., 2007. Gap junctional intercellular communication as a biological "Rosetta stone" in understanding, in a systems biological manner, stem cell behavior, mechanisms of epigenetic toxicology, chemoprevention and chemotherapy. *J Membr Biol* 218: 93-100.
- Tseng, A.S., Beane, W.S., Lemire, J.M., Masi, A., Levin, M., 2010. Induction of vertebrate regeneration by a transient sodium current. *J Neurosci* 30: 13192-13200.
- Tsutsui, H., Jinno, Y., Tomita, A., Okamura, Y., 2014. Rapid evaluation of a protein-based voltage probe using a field-induced membrane potential change. *Biochim Biophys Acta*.
- Turin, L., Warner, A.E., 1980. Intracellular pH in early *Xenopus* embryos: its effect on current flow between blastomeres. *J Physiol* 300: 489-504.
- Turing, A.M., 1953. The chemical basis of morphogenesis. *Philosophical Transactions of the Royal Society B: Biological Sciences* 237: 37-72.
- Turner, C.H., Robling, A.G., Duncan, R.L., Burr, D.B., 2002. Do bone cells behave like a neuronal network? *Calcif Tissue Int* 70: 435-442.
- Umino, Y., Saito, T., 2002. Spatial and temporal patterns of distribution of the gap junctional protein connexin43 during retinal regeneration of adult newt. *J Comp Neurol* 454: 255-262.
- Urschel, S., Höher, T., Schubert, T., Alev, C., Söhl, G., 2006. Protein kinase A-mediated phosphorylation of connexin36 in mouse retina results in decreased gap junctional communication between All amacrine cells. *Journal of biological ...*
- Valiunas, V., Polosina, Y.Y., Miller, H., Potapova, I.A., Valiuniene, L., Doronin, S., Mathias, R.T., Robinson, R.B., Rosen, M.R., Cohen, I.S., Brink, P.R., 2005. Connexin-specific cell-to-cell transfer of short interfering RNA by gap junctions. *J Physiol* 568: 459-468.
- van der Heyden, M.A., Rook, M.B., Hermans, M.M., Rijksen, G., Boonstra, J., Defize, L.H., Destree, O.H., 1998. Identification of connexin43 as a functional target for Wnt signalling. *J Cell Sci* 111: 1741-1749.
- Vandenberg, L.N., Blackiston, D.J., Rea, A.C., Dore, T.M., Levin, M., 2014. Left-right patterning in *Xenopus* conjoined twin embryos requires serotonin signaling and gap junctions. *The International journal of developmental biology* 58: 799-809.
- Vandenberg, L.N., Lemire, J.M., Levin, M., 2012. Serotonin has early, cilia-independent roles in *Xenopus* left-right patterning. *Disease models & mechanisms*.
- Vandenberg, L.N., Levin, M., 2009. Perspectives and open problems in the early phases of left-right patterning. *Semin Cell Dev Biol* 20: 456-463.

- VanHoven, M.K., Huang, S.L.B., Albin, S.D., Bargmann, C.I., 2006. The claudin superfamily protein *nsy-4* biases lateral signaling to generate left-right asymmetry in *C. elegans* olfactory neurons. *Neuron*.
- Verwey, L.J., Edwards, T.M., 2010. Gap junctions and memory: an investigation using a single trial discrimination avoidance task for the neonate chick. *Neurobiol. Learn. Mem.* 93: 189-195.
- Vink, M.J., Suadicani, S.O., Vieira, D.M., Urban-Maldonado, M., Gao, Y., Fishman, G.I., Spray, D.C., 2004. Alterations of intercellular communication in neonatal cardiac myocytes from connexin43 null mice. *Cardiovasc Res* 62: 397-406.
- Vinken, M., Decrock, E., Leybaert, L., Bultynck, G., Himpens, B., Vanhaecke, T., Rogiers, V., 2012. Non-channel functions of connexins in cell growth and cell death. *Biochim Biophys Acta* 1818: 2002-2008.
- Vogel, R., Weingart, R., 1998. Mathematical model of vertebrate gap junctions derived from electrical measurements on homotypic and heterotypic channels. *J Physiol* 510 (Pt 1): 177-189.
- Vogel, R., Weingart, R., 2002. The electrophysiology of gap junctions and gap junction channels and their mathematical modelling. *Biol Cell* 94: 501-510.
- Wade, J.J., McDaid, L.J., Harkin, J., Crunelli, V., Kelso, J.A., 2011. Bidirectional coupling between astrocytes and neurons mediates learning and dynamic coordination in the brain: a multiple modeling approach. *PLoS One* 6.
- Waldo, K.L., Lo, C.W., Kirby, M.L., 1999. Connexin 43 expression reflects neural crest patterns during cardiovascular development. *Developmental Biology* 208: 307-323.
- Waller, B.R., 3rd, McQuinn, T., Phelps, A.L., Markwald, R.R., Lo, C.W., Thompson, R.P., Wessels, A., 2000. Conotruncal anomalies in the trisomy 16 mouse: an immunohistochemical analysis with emphasis on the involvement of the neural crest. *Anat Rec* 260: 279-293.
- Wang, X., Veruki, M.L., Bukoreshtliev, N.V., Hartveit, E., Gerdes, H.H., 2010. Animal cells connected by nanotubes can be electrically coupled through interposed gap-junction channels. *Proc Natl Acad Sci U S A* 107: 17194-17199.
- Wang, Y., Belousov, A.B., 2011. Deletion of neuronal gap junction protein connexin 36 impairs hippocampal LTP. *Neuroscience letters* 502: 30-32.
- Warner, A., 1999. Interactions between growth factors and gap junctional communication in developing systems. *Novartis Found Symp* 219: 60-72.
- Warner, A.E., 1985. The role of gap junctions in amphibian development. *Journal of embryology and experimental morphology* 89 Suppl: 365-380.
- Warner, A.E., Guthrie, S.C., Gilula, N.B., 1984. Antibodies to gap-junctional protein selectively disrupt junctional communication in the early amphibian embryo. *Nature* 311: 127-131.
- Watanabe, M., Iwashita, M., Ishii, M., Kurachi, Y., Kawakami, A., Kondo, S., Okada, N., 2006a. Spot pattern of leopard *Danio* is caused by mutation in the zebrafish connexin41.8 gene. *EMBO Rep* 7: 893-897.
- Watanabe, M., Iwashita, M., Ishii, M., Kurachi, Y., Kawakami, A., Kondo, S., Okada, N., 2006b. Spot pattern of leopard *Danio* is caused by mutation in the zebrafish connexin41.8 gene. *EMBO Rep*.

- Watanabe, M., Kondo, S., 2014. Is pigment patterning in fish skin determined by the Turing mechanism? *Trends in genetics : TIG* 31: 88-96.
- Watanabe, M., Kondo, S., 2015. Is pigment patterning in fish skin determined by the Turing mechanism? *Trends in genetics : TIG* 31: 88-96.
- Webb, R.J., Marshall, F., Swann, K., Carroll, J., 2002. Follicle-stimulating hormone induces a gap junction-dependent dynamic change in [cAMP] and protein kinase a in mammalian oocytes. *Dev Biol* 246: 441-454.
- Weir, M.P., Lo, C.W., 1982. Gap junctional communication compartments in the *Drosophila* wing disk. *Proceedings of the National Academy of Sciences of the United States of America* 79: 3232-3235.
- Weir, M.P., Lo, C.W., 1984. Gap-junctional communication compartments in the *Drosophila* wing imaginal disk. *Developmental Biology* 102: 130-146.
- White, T.W., 2002. Unique and redundant connexin contributions to lens development. *Science* 295: 319-320.
- Wittig, D., Wang, X., Walter, C., Gerdes, H.H., Funk, R.H., Roehlecke, C., 2012. Multi-Level Communication of Human Retinal Pigment Epithelial Cells via Tunneling Nanotubes. *PloS one* 7: e33195.
- Wolszon, L.R., Gao, W.Q., Passani, M.B., Macagno, E.R., 1994. Growth cone "collapse" in vivo: are inhibitory interactions mediated by gap junctions? *J Neurosci* 14: 999-1010.
- Wolvetang, E.J., Pera, M.F., Zuckerman, K.S., 2007. Gap junction mediated transport of shRNA between human embryonic stem cells. *Biochem Biophys Res Commun* 363: 610-615.
- Wong, R.C., Pera, M.F., Pebay, A., 2008. Role of gap junctions in embryonic and somatic stem cells. *Stem cell reviews* 4: 283-292.
- Woodruff, R.I., 2005. Calmodulin transit via gap junctions is reduced in the absence of an electric field. *J Insect Physiol* 51: 843-852.
- Wu, C.L., Shih, M.F., Lai, J.S., Yang, H.T., Turner, G.C., Chen, L., Chiang, A.S., 2011. Heterotypic Gap Junctions between Two Neurons in the *Drosophila* Brain Are Critical for Memory. *Curr Biol*.
- Xin, D., Bloomfield, S., 1997. Tracer coupling pattern of amacrine and ganglion cells in the rabbit retina. *Journal of Comparative Neurology* 383: 512-528.
- Xu, X., Li, W.E., Huang, G.Y., Meyer, R., Chen, T., Luo, Y., Thomas, M.P., Radice, G.L., Lo, C.W., 2001. N-cadherin and Cx43alpha1 gap junctions modulates mouse neural crest cell motility via distinct pathways. *Cell Commun Adhes* 8: 321-324.
- Yamaguchi, M., Yoshimoto, E., Kondo, S., 2007. Pattern regulation in the stripe of zebrafish suggests an underlying dynamic and autonomous mechanism. *Proceedings of the National Academy of Sciences of the United States of America* 104: 4790-4793.
- Yamasaki, H., Enomoto, T., Martel, N., 1984. Intercellular communication, cell differentiation and tumour promotion. *IARC Sci. Publ.* 217-238.
- Yamasaki, H., Mesnil, M., Omori, Y., Mironov, N., Krutovskikh, V., 1995. Intercellular communication and carcinogenesis. *Mutat Res* 333: 181-188.

- Yano, T., Fujimoto, E., Hagiwara, H., Sato, H., Yamasaki, H., Negishi, E., Ueno, K., 2006. Connexin 32 as an anti-invasive and anti-metastatic gene in renal cell carcinoma. *Biol. Pharm. Bull.* 29: 1991-1994.
- Zahler, S., Hoffmann, A., Gloe, T., Pohl, U., 2003. Gap-junctional coupling between neutrophils and endothelial cells: a novel modulator of transendothelial migration. *J. Leukoc. Biol.* 73: 118-126.
- Zhang, W., Green, C., Stott, N.S., 2002. Bone morphogenetic protein-2 modulation of chondrogenic differentiation in vitro involves gap junction-mediated intercellular communication. *J Cell Physiol* 193: 233-243.
- Zhang, Y., Levin, M., 2009. Particle tracking model of electrophoretic morphogen movement reveals stochastic dynamics of embryonic gradient. *Dev Dyn* 238: 1923-1935.
- Zhang, Z.Q., Hu, Y., Wang, B.J., Lin, Z.X., Naus, C.C., Nicholson, B.J., 2003. Effective asymmetry in gap junctional intercellular communication between populations of human normal lung fibroblasts and lung carcinoma cells. *Carcinogenesis*.
- Zoghi, M., 2004a. Cardiac memory: do the heart and the brain remember the same? *J Interv Card Electrophysiol* 11: 177-182.
- Zoghi, M., 2004b. Cardiac memory: do the heart and the brain remember the same? *Journal of interventional cardiac electrophysiology*.
- Zsiros, V., Maccaferri, G., 2008. Noradrenergic modulation of electrical coupling in GABAergic networks of the hippocampus. *The Journal of Neuroscience*.

Figure Legends

Figure 1: Gap junctions form bioelectric circuits in the brain and beyond

(**A**) Neurons are often coupled by gap junctions, which allows electrical activity to propagate and integrate across cells. (**B**) The same scheme, involving ion channels to set V_{mem} levels and gap junctions to communicate bioelectric state to neighboring cells, is present in most somatic tissues. (**C**) Non-neural cells assemble into GJ-coupled networks that have many of the properties of neural networks. Manipulating the function of somatic tissues during pattern formation, by modulating GJ activity, makes use of 2 basic approaches, using pharmacological or genetic techniques to target connectivity (GJ gating, akin to synaptic plasticity) or resting potential (ion channels, akin to intrinsic plasticity). Graphics courtesy of Alexis Pietak and Jeremy Guay.

Figure 2: Information in physiological circuits instructs pattern formation

Gap junctions are key regulators of bioelectric cell state. (**A**) Like gene regulatory networks, which contain numerous feedback loops among gene loci, gap junctions and ion channels both regulate and are regulated by resting potential. This establishes an autonomous layer of physiological dynamics that is coupled to transcriptional cascades, but has its own unique information and functions. Modulating bioelectric dynamics by induced changes of GJC results in large-scale alterations of pattern formation, including hyperinnervation (**B**), left-right organ inversions (**B'**), and multiple head formation in regenerating planaria (**B''**). Graphics courtesy of Alexis Pietak and Jeremy Guay.

Figure 3: Parallelism between brain and body

(A) In the brain, DNA sets the structure of the central nervous system – the hardware. Electrical circuit dynamics process information and store memories, resulting in experience-dependent (and self-organizing) patterns that control muscles resulting in the movement of the animal in 3-dimensional space. Experiences (external sensory stimuli) and reagents targeting GJs, ion channels, and neurotransmitters alter the electric dynamics. Computational pipelines are beginning to be developed and applied to dynamics observed via EEG and MRI imaging, to extract the semantics – the memories represented by these bioelectric states.

(B) In somatic tissues of developing or regenerating organisms, DNA sets the complement of connexins, ion channels, and neurotransmitter machinery in all cells – the hardware. Bioelectric dynamics (slow patterns of resting potential within tissues) process information and establish prepatterns for gene expression and morphogenesis, resulting in chemical signal-dependent (and self-organizing) patterns that control cell functions like migration and differentiation, resulting in anatomical changes that move the body through morphospace. Stimuli (chemical and other signals) and reagents targeting GJs, ion channels, and neurotransmitters alter the electric dynamics. Computational pipelines need to be developed and applied to these dynamics as imaged with voltage-sensitive dyes and GJ tracers, to extract the semantics – the instructive anatomical patterns represented by these bioelectric states.

Figure 4: Stable inheritance of target morphology change after GJ network perturbation

A normal planarian has a head and tail, and regenerates each at the appropriate end of an amputated fragment (**A**). When cut into thirds, and the middle fragment is briefly exposed to octanol, which temporarily blocks long-range bioelectrical signaling between the wound and mature tissues, a 2-headed worm results (**B**). GJC = gap junctional communication. Remarkably, upon further rounds of cutting in plain water (long after the octanol has left the tissues, as confirmed by HPLC), the 2-headed form is recapitulated (**C,D**; images of 2-headed worms provided by Fallon Durant.). This change in the animal's target morphology (the shape to which it regenerates upon damage) appears to be permanent, and persists across the animal's normal reproductive mode (fissioning), despite the fact that the genomic sequence has not been altered. Chromatin modifications alone do not explain this, because the posterior wound cells, which could have been epigenetically reprogrammed to a head fate, are thrown away at each cut: the information encoding a bipolar 2-head animal is present even in the normal gut fragment – it is distributed throughout the body. We propose that this information is a kind of memory, encoded in electrical networks of somatic cells coupled by gap junctions, and is stored at the level of bioelectrical dynamics, not genetics.

Figure 5: Morphogenetic memories visualized as attractors in GJ network state space

GJ dynamics may offer an opportunity to understand information processing, not only molecular biophysics, of pattern formation regulated by physiological cell-cell signaling. One way to visualize planarian regeneration is (**A**) as the function of a large network of electrically coupled cells. Some such networks have been shown (in

computational neuroscience and artificial intelligence research) to have a planaria-like property of holographic memory storage: a trained network can recreate the entire pattern despite deletions of the pattern or of the network components. A well-accepted mathematical paradigm for understanding the global properties of such networks is as an energy landscape, with attractors corresponding to specific stable modes of the network. In our analogy, amputation raises the energy of the system, temporarily pulling it out of the attractor to which the system tends to return. One hypothesis is that these networks are responsible for storing the pattern of a normal planarian, and when damaged, issuing cell-level commands (differentiate, proliferate, and other instructions) that restore the anatomy (in parallel to how recall of complex geometric memories can be triggered by stimuli and induce goal-directed behavior in cognitive science studies of animal behavior). This hypothesis makes a prediction: that coherent changes in patterning will result from experimentally-induced changes of the bioelectric network's topology or dynamics. **(B)** It has been shown (Emmons-Bell et al., 2015) that altering the bioelectric connectivity in *G. dorotocephala* results in regeneration of one of four discrete head types. On this view, amputation of head and tail causes the system to move to an unstable state from its basin of attraction. Partial interruption of gap-junction communication between cells (reduced connectivity and thus altered dynamics of the network) induces head wounds to regenerate (yellow arrows) new heads that resemble closely-related flatworm species that are regions of stability in the regeneration morphospace landscape (left to right: *S. mediterranea*, *D. japonica*, and *P. felina*) as well as heads of the original species in the center basin. The probability of regenerating a certain head shape is proportional to the evolutionary distance from *G. dorotocephala*.

These states are non-permanent (shallow basins of attraction) and over time will remodel into their final morphological state (white dashed arrows) to the deepest and most stable basin of attraction of the original head shape. (C) The same process can be modeled as a neural-like network, with stable modes visualized as stable attractors (a.k.a. memories in neural nets), which lead to specific instructive signals regulating proliferation, migration, and differentiation that induces different but coherent patterning outcomes. Graphics by Alexis Pietak.

Table 1: Channelopathies of GJ genes and the resulting patterning defects

Gap Junction Protein	Morphogenetic role or LOF phenotype	Species	Reference
Innexins	Gonad and germline morphogenesis	<i>C. Elegans</i>	(Starich et al., 2014)
Innexin1,2	Epithelial patterning, foregut development	<i>Drosophila</i>	(Bauer et al., 2002; Bauer et al., 2004)
Innexin 2	Eye size	<i>Drosophila</i>	(Richard and Hoch, 2015)
Innexins	Foregut, cuticle (epithelial) patterning	<i>Drosophila</i>	(Bauer et al., 2004)
Connexin 43	Oculodentodigital dysplasia (ODDD), heart defects (outflow tract and conotruncal), left-right asymmetry randomization, Osteoblast differentiation problems, craniofacial defects, myogenesis	Human, mouse, chick	(Araya et al., 2003; Becker et al., 1999; Britz-Cunningham et al., 1995; Civitelli, 2008; Debeer et al., 2005; Ewart et al., 1997; Lecanda et al., 2000a; Levin and Mercola, 1999; Pizzuti et al., 2004; Reaume et al., 1995)
Connexin 37	Lymphatic system patterning	Mouse	(Kanady et al., 2011; Kanady et al., 2015)
Connexin 45	Cardiac defects (cushion patterning)	Mouse	(Kumai et al., 2000; Nishii et al., 2001)

Connexin 50, Connexin 46	eye patterning (differentiation and proliferation, especially lens),	Mouse	(White, 2002)
Connexin 26	Cochlear development	Mouse	(Chang et al., 2014)
Connexin 41.8	Pigmentation pattern	Zebrafish	(Watanabe et al., 2006a)
Connexin 43	Fin size and pattern regulation Craniofrontonasal syndrome	Zebrafish Mouse	(Davy et al., 2006; Hoptak-Solga et al., 2008; Iovine et al., 2005b; Sims et al., 2009b)
Innexin 4, Innexin 2	Germline differentiation and spermatogenesis	<i>Drosophila</i>	(Smendziuk et al., 2015)
Pannexin 3	Skeletal development	Mouse	(Oh et al., 2015)




Figure 1

(A) Neurons are often coupled by gap junctions, which allows electrical activity to propagate and integrate across cells. (B) The same scheme, involving ion channels to set V_{mem} levels and gap junctions to communicate bioelectric state to neighboring cells, is present in most somatic tissues. (C) Non-neural cells assemble into GJ-coupled networks that have many of the properties of neural networks. Manipulating the function of somatic tissues during pattern formation, by modulating GJ activity, makes use of 2 basic approaches, using pharmacological or genetic techniques to target connectivity (GJ gating, akin to synaptic plasticity) or resting potential (ion channels, akin to intrinsic plasticity). Graphics courtesy of Alexis Pietak and Jeremy Guay.

152x217mm (300 x 300 DPI)




Figure 2

Gap junctions are key regulators of bioelectric cell state. (A) Like gene regulatory networks, which contain numerous feedback loops among gene loci, gap junctions and ion channels both regulate and are regulated by resting potential. This establishes an autonomous layer of physiological dynamics that is coupled to transcriptional cascades, but has its own unique information and functions. Modulating bioelectric dynamics by induced changes of GJC results in large-scale alterations of pattern formation, including hyperinnervation (B), left-right organ inversions (B'), and multiple head formation in regenerating planaria (B''). Graphics courtesy of Alexis Pietak and Jeremy Guay.

152x112mm (300 x 300 DPI)

Acce




Figure 3

(A) In the brain, DNA sets the structure of the central nervous system – the hardware. Electrical circuit dynamics process information and store memories, resulting in experience-dependent (and self-organizing) patterns that control muscles resulting in the movement of the animal in 3-dimensional space. Experiences (external sensory stimuli) and reagents targeting GJs, ion channels, and neurotransmitters alter the electric dynamics. Computational pipelines are beginning to be developed and applied to dynamics observed via EEG and MRI imaging, to extract the semantics – the memories represented by these bioelectric states.

(B) In somatic tissues of developing or regenerating organisms, DNA sets the complement of connexins, ion channels, and neurotransmitter machinery in all cells – the hardware. Bioelectric dynamics (slow patterns of resting potential within tissues) process information and establish prepatterns for gene expression and morphogenesis, resulting in chemical signal-dependent (and self-organizing) patterns that control cell functions like migration and differentiation, resulting in anatomical changes that move the body through morphospace. Stimuli (chemical and other signals) and reagents targeting GJs, ion channels, and neurotransmitters alter the electric dynamics. Computational pipelines need to be developed and applied to

these dynamics as imaged with voltage-sensitive dyes and GJ tracers, to extract the semantics – the instructive anatomical patterns represented by these bioelectric states.

152x193mm (300 x 300 DPI)

Accepted Article




Figure 4

A normal planarian has a head and tail, and regenerates each at the appropriate end of an amputated fragment (A). When cut into thirds, and the middle fragment is briefly exposed to octanol, which temporarily blocks long-range bioelectrical signaling between the wound and mature tissues, a 2-headed worm results (B). GJC = gap junctional communication. Remarkably, upon further rounds of cutting in plain water (long after the octanol has left the tissues, as confirmed by HPLC), the 2-headed form is recapitulated (C,D; images of 2-headed worms provided by Fallon Durant.). This change in the animal's target morphology (the shape to which it regenerates upon damage) appears to be permanent, and persists across the animal's normal reproductive mode (fissioning), despite the fact that the genomic sequence has not been altered. Chromatin modifications alone do not explain this, because the posterior wound cells, which could have been epigenetically reprogrammed to a head fate, are thrown away at each cut: the information encoding a bipolar 2-head animal is present even in the normal gut fragment – it is distributed throughout the body. We propose that this information is a kind of memory, encoded in electrical networks of somatic cells coupled by gap junctions, and is stored at the level of bioelectrical dynamics, not genetics.

152x263mm (300 x 300 DPI)

Accepted Article




Figure 5

GJ dynamics may offer an opportunity to understand information processing, not only molecular biophysics, of pattern formation regulated by physiological cell-cell signaling. One way to visualize planarian regeneration is (A) as the function of a large network of electrically coupled cells. Some such networks have been shown (in computational neuroscience and artificial intelligence research) to have a planaria-like property of holographic memory storage: a trained network can recreate the entire pattern despite deletions of the pattern or of the network components. A well-accepted mathematical paradigm for understanding the global properties of such networks is as an energy landscape, with attractors corresponding to specific stable modes of the network. In our analogy, amputation raises the energy of the system, temporarily pulling it out of the attractor to which the system tends to return. One hypothesis is that these networks are responsible for storing the pattern of a normal planarian, and when damaged, issuing cell-level commands (differentiate, proliferate, and other instructions) that restore the anatomy (in parallel to how recall of complex geometric memories can be triggered by stimuli and induce goal-directed behavior in cognitive science studies of animal behavior). This hypothesis makes a prediction: that coherent changes in patterning will result from experimentally-induced changes of the bioelectric network's topology or dynamics. (B) It has been shown (Emmons-Bell et al., 2015) that altering the bioelectric connectivity in *G. dorotocephala* results in regeneration of one of four discrete head types. On this view, amputation of head and tail causes the system to move to an unstable state from its basin of attraction. Partial interruption of gap-junction communication between cells (reduced connectivity and thus altered dynamics of the network) induces head wounds to regenerate (yellow arrows) new heads that resemble closely-related flatworm species that are regions of stability in the regeneration morphospace landscape (left to right: *S. mediterranea*, *D. japonica*, and *P. felina*) as well as heads of the original species in the center basin. The probability of regenerating a certain head shape is proportional to the evolutionary distance from *G. dorotocephala*. These states are non-permanent (shallow basins of attraction) and over time will remodel into their final morphological state (white dashed arrows) to the deepest and most stable basin of attraction of the original head shape. (C) The same process can be modeled as a neural-like network, with stable modes visualized as stable attractors (a.k.a. memories in neural nets), which lead to specific instructive signals regulating proliferation, migration, and differentiation that induces different but coherent patterning outcomes. Graphics by Alexis Pietak.

152x114mm (300 x 300 DPI)

Accepted Article