



Communicative & Integrative Biology



Taylor & Franci

ISSN: 1942-0889 (Online) Journal homepage: www.tandfonline.com/journals/kcib20

Physiological inputs regulate species-specific anatomy during embryogenesis and regeneration

Kelly G. Sullivan, Maya Emmons-Bell & Michael Levin

To cite this article: Kelly G. Sullivan, Maya Emmons-Bell & Michael Levin (2016) Physiological inputs regulate species-specific anatomy during embryogenesis and regeneration, Communicative & Integrative Biology, 9:4, e1192733, DOI: 10.1080/19420889.2016.1192733

To link to this article: https://doi.org/10.1080/19420889.2016.1192733

9	© 2016 The Author(s). Published with license by Taylor & Francis.© Kelly G. Sullivan, Maya Emmons-Bell, and Michael
+	Levin. View supplementary material 🗹
	Published online: 15 Jul 2016.
	Submit your article to this journal 🗷
hil	Article views: 4162
Q ¹	View related articles 🗹
CrossMark	View Crossmark data ☑
4	Citing articles: 14 View citing articles 🗗

Taylor & Francis Taylor & Francis Group

ARTICLE ADDENDUM 3 OPEN ACCESS

Physiological inputs regulate species-specific anatomy during embryogenesis and regeneration

Kelly G. Sullivan[†], Maya Emmons-Bell[†], and Michael Levin

Allen Discovery Center at Tufts University, Tufts University, Medford, MA, USA

ABSTRACT

A key problem in evolutionary developmental biology is identifying the sources of instructive information that determine species-specific anatomical pattern. Understanding the inputs to large-scale morphology is also crucial for efforts to manipulate pattern formation in regenerative medicine and synthetic bioengineering. Recent studies have revealed a physiological system of communication among cells that regulates pattern during embryogenesis and regeneration in vertebrate and invertebrate models. Somatic tissues form networks using the same ion channels, electrical synapses, and neurotransmitter mechanisms exploited by the brain for information-processing. Experimental manipulation of these circuits was recently shown to override genome default patterning outcomes, resulting in head shapes resembling those of other species in planaria and *Xenopus*. The ability to drastically alter macroscopic anatomy to that of other extant species, despite a wild-type genomic sequence, suggests exciting new approaches to the understanding and control of patterning. Here, we review these results and discuss hypotheses regarding non-genomic systems of instructive information that determine biological growth and form.

ARTICLE HISTORY

Received 4 April 2016 Revised 13 May 2016 Accepted 16 May 2016

KEYWORDS

bioelectricity; epigenetics; regeneration; morphology; morphospace; patterning; planaria; Xenopus

"Treasure your exceptions! Keep them always uncovered and in sight. Exceptions are like the rough brickwork of a growing building which tells that there is more to come and shows where the next construction is to be."

—W. Bateson, The Methods and Scope of Genetics

Introduction: What determines anatomical pattern?

Biological patterning is at the nexus of most of the important problems facing basic biology and biomedicine. Understanding the instructive signals that ensure self-assembly and maintenance of complex 3-dimensional morphology is crucial for basic evolutionary and developmental biology. How do cells, all derived from the same fertilized egg (the original stem cell) and bearing the same DNA, become not only differentiated into distinct cell types, but arranged into stereotypical spatial patterns with no external guidance? This question is at the center of understanding evolutionary

change because development is what links genetics (upon which mutation acts) with form and function (upon which selection operates). Moreover, the identification of and functional control over these processes will lead to transformative strategies for injury repair, reversal of degeneration and aging, and tumor reprogramming.¹ If we understood how the body created its structures in the first place, we could coax it to repeat the process throughout its lifespan as needed. Complex pattern control is thus at the heart of regenerative medicine as well as synthetic bioengineering.^{2,3}

Importantly, pattern control does not solely occur during the feed-forward process by which a fertilized egg's genome progressively unrolls toward an invariant outcome. A view of embryogenesis as terminal shape simply emerging over time from the parallel interactions of molecular components is too limited. Patterning information also has to be stored, processed, and acted upon during *shape homeostasis*: regeneration, remodeling, and maintenance of existing adult structures against

CONTACT Michael Levin michael.levin@tufts.edu Allen Discovery Center at Tufts University, Center for Regenerative and Developmental Biology, Biology Department, Tufts University, 200 Boston Avenue, Suite 4600, Medford, MA 02155-4243, USA.

Addendum to: Emmons-Bell M, Durant F, Hammelman J, Bessonov N, Volpert V, Morokuma J, Pinet K, Adams DS, Pietak A, Lobo D, et al. Gap junctional blockade stochastically induces different species-specific head anatomies in genetically wild-type Girardia dorotocephala flatworms. International Journal of Molecular Sciences 2015; 16:27865–27896; http://dx.doi.org/10.3390/ijms161126065

1 Supplemental data for this article can be accessed on the publisher's website.

© 2016 Kelly G. Sullivan, Maya Emmons-Bell, and Michael Levin. Published with license by Taylor & Francis.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (http://creativecommons.org/licenses/by-nc/3.0/), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. The moral rights of the named author(s) have been assented

[†]These authors equally contributed to this work.

damage, aging, and cancer.4-6 Salamanders readily replace amputated body-parts, growing exactly what's missing and stopping when the correct pattern has been achieved. Grafting the tail of one salamander to the flank of another reveals an even more complex patternhomeostasis capability: over time, the grafted tail remodels to a limb - a structure more appropriate to its new location.⁷ The most important aspect of regeneration and remodeling is that it requires feedback: it cannot simply be an emergent process from hardwired rules because the details of injury are unpredictable for the organism. Somatic plasticity requires an organism to be able to recognize damage and enact repair from a vast variety of starting states, toward the same (correct) target morphology.

How do organs, tissues, and entire bodyplans know what shape they are supposed to build and maintain (Fig. 1A)? It is known that genomes do not specify target morphology directly, but nevertheless it is thought that the origin of a species' specific anatomy, at multiple levels of organization, is ultimately in its genome. The genome is often held to be the entirety of the information medium that determines pattern, and the key factor that differentiates one species' body from another. Thus, the emphasis of modern synthetic biology is almost

exclusively on rewriting the genome.8 This view has to be expanded in several ways toward epigenetic controls of patterning (Fig. 1B). First is chromatin modification, and the genomic regulatory code that drives cell typespecific gene expression and thus cell specification and differentiation. 9,10 Second are biotic signals produced by commensal organisms such as microbes, 11,12 and abiotic aspects of the environment, such as light which determines brain structure and function in some species¹³ and temperature which can determine sex.¹⁴

Feedback between genetic and physical forces is a key determinant of patern formation, enabling gene regulatory circuits to generate specific anatomies. By expressing molecules that couple to specific physical forces (e.g., adhesion proteins), genetic circuits can harness selforganizing physical dynamics such as tensegrity, 15 reaction-diffusion, 16,17 and many others. 18-22 Consistent leftright patterning²³⁻²⁵ is a perfect example of how important physics is to pattern. While a cascade of regulatory molecules (a gene-regulatory network, GRN) is required for maintaining identity of the left and right sides of the body, ²⁶⁻²⁸ it is immediately clear that no genetic circuit can be sufficient to distinguish left from right in space ab *initio*. To establish proper laterality, the cascade of asymmetrically-expressed genes is functionally positioned




Figure 1. Interplay of genetics and physics. (A) The process of developing an organism from a fertilized egg cell involves an interplay of physics and genetics. Transcriptional gene-regulatory networks (GRNs) specify the production of effector proteins that allow coupling to specific physical processes (adhesion, tension, electric propagation, diffusion, etc.). It is the emergent order in those physical processes that ultimately results in a specific 3-dimensional shape of the body and its internal organs. (B) At least 3 codes participate in this process. The genetic code, which maps DNA sequence to protein sequence, controls cell behavior. The Epigenetic code, which maps chromatin state to expression of specific genes, regulates cell type and physiological properties. The bioelectric code maps the distribution of endogenous voltage gradients and electric fields in vivo, and appears to regulate large-scale anatomical patterning. 54,55,75,108

between essential physical processes: the chiral elements that initiate asymmetry and orient it with the other 2 axes, 29-31 and the physical forces that actually execute the bending of asymmetric organs such as the heart and gut. 32,33

More generally, GRNs have to be "painted" onto a pre-existing anatomical structure and integrated with physical effectors of genetic information. Even this picture is likely too simple, as shape is not hardcoded but can be edited by interactions with the environment.³⁴ Trophic memory in deer antlers occurs when damage made to one point in the branched antler structure will cause an ectopic tine in that same location next year, after the antler racks are discarded and re-grown anew.³⁵ We currently lack the conceptual apparatus to propose a plausible model by which genomes and physics allow the growth plate at the scalp to ascertain the 3D location of a wound, remember it for months, and use it to alter the genome-default cell growth decisions during subsequent antler re-growth.

While most of the above-mentioned models have not been addressed with molecular tools, recent work has begun to unravel genetic factors that mediate the differences among species. There are numerous incidences in which simple genetic changes drastically alter the morphology of populations as they diverge from their evolutionary ancestors. Darwin's finches, a group of 14 closely related finch species, are one such example. They display an impressive variety of beak shapes and sizes, and therefore diets, despite having evolved from the same ancestral species. This variation is due, at least in part, to differential expression levels of the genes calmodulin and Bmp4, which define beak length and width, respectively. 36,37 Another compelling example is the evolution of the bat's wing from a mouse-like ancestral arm. Increased expression levels of Bmp2 and Prx1, regulators of bone development, are responsible for the remarkable elongation of the bat forelimb digits and long bones respectively, rendering these animals capable of flight.^{38,39}

Likewise, a number of epigenetic factors have been shown to make significant input into species-specific pattern outcomes. The long-term persistence of epigenetic state has been an area of focus in modern molecular and developmental biology, and work done in this field has added to models of cellular differentiation and stemness. 40 Of especial interest to evolutionary developmental biologists and ecologists are epigenetic modifications that both inform phenotype in an appreciable manner, and are transmitted to offspring through the germ line.^{41,42}

Early botanist Linneaus was the first to describe the puzzling diversity in shape of the toadflax flower (Linaria vulgaris).⁴³ Despite being genetically identical, one morphological variant displays a rounded flower with lobed petals and bilateral symmetry, and the other an acute, starburst flower shape with radial symmetry. Cubas et al. found that the 2 varied only in the heritable methylation status of a single gene (Lcyc), demonstrating that largescale morphological variance, eligible for natural selection, can occur solely as the result of epigenetic state.⁴⁴ Similarly, the alligator weed (Alternatnthera philoxeroides) has a remarkable variety of anatomical configurations, as it is able to dramatically restructure its root system, stem diameter, and internode length given development in a variable environment. 44 Other examples of environmental influence on morphology include species of reptile and fish, where incubation temperatures alter sex ratios of offspring.⁴⁵ Perhaps of even more interest are heritable epigenetic modifications 46-48 in organisms that segregate the germ line early in development. Methylation at a retrotransposon site within the Axin-fused allele produces a distinct kinked-tail phenotype in mice, which can be inherited both maternally and paternally.⁴⁹ For a more detailed account of proposed models and evolutionary significance of epigenetic heritability see refs. 50, 51 and ref. 52.

In addition to genetic sequence, chromatin marking, and environmental stimuli, it is becoming clear that biological pattern is regulated by another key set of processes: physiological signaling among bioelectrical networks.⁵³ In this Perspective, we discuss recent data in the frog and planaria model systems that shed light on the question: how much of the difference among species' morphologies comes from the physiological layer of control?

Bioelectricity: A novel epigenetic layer regulating pattern formation

Bioelectric networks (Fig. 2) consist of 2 basic components.54,55 The first is ion channels (and pumps), which set the resting potentials of cells; these implement bioelectric circuits whose activity regulates the timedependent changes of spatial gradients of resting potential (V_{mem}) across anatomical areas. Downstream of these bioelectric dynamics lie a number of targets, including the movement of neurotransmitters like serotonin, 56-59 and changes in gene expression 60 and chromatin state. 61-63 Endogenous, and experimenter-induced changes in V_{mem} implement intrinsic plasticity; recent functional approaches showed that these dynamics normally implement, and can be used to control, eye induction, organ size, craniofacial patterning, neoplastic transformation, axial polarity, and regenerative capac-

An even more important element of bioelectric networks consists of gap junctions (GJs, electrical




Figure 2. Bioelectric networks and their modulation. Many tissues, not only the nervous system, maintain active electrical communication among cells. Developmental bioelectricity, as in neurons, relies on ion channels and pumps to produce resting potential gradients across their cell membranes, and gap junctional channels to spread those potentials to neighboring cells. Signals within this network are mediated by the transfer of current and small molecules such as neurotransmitters. Because many ion channels and gap junctions are themselves voltage-sensitive, this system can support complex dynamics; the output of these dynamics includes changes in gene expression and cell behavior, mediated by transduction machinery such as neurotransmitter flux. Experimental (and endogenous) modulation of these dynamics can occur via regulation of gap junctional connectivity within the network (targeting gap junctions to alter plasticity of the electrical synapses), changes in ion channel activity (editing of resting potentials as a kind of intrinsic plasticity), or direct effects on the resulting gradients of neurotransmitters. A wide variety of genetic, pharmacological, and optical tools are now available to manipulate physiological networks in developmental or regenerative contexts in vivo. Graphics by Alexis Pietak.

synapses), which allow cells to share their voltage with neighbors, as well as communicate via small molecules such as neurotransmitters. Importantly, gap junctions are themselves voltage-sensitive, allowing them to serve as highly versatile gates for bioelectric circuits that underlie plasticity and computation. 72,73 Gap junction dependent patterns of cell:cell connectivity represent synaptic plasticity - the basis of information storage in the electrical networks of the brain and in non-neural tissues. Disruption of GJ signaling underlies numerous developmental disorders as well as the patterning defect known as cancer.⁷⁴

Brains capitalized on the transistor-like properties of voltage-gated gap junctions (electrical synapses) and ion channels for memory and integrated decision-making, in the control of animal behavior.

However, brains did not invent these tricks *de novo*: they speed-optimized computational properties of slower, "developmental" ionic signaling that cells were using to organize embryogenesis, wound healing, and adaptive physiology long before central nervous systems appeared. 75-80 Evolution capitalized on the unique suitability of bioelectric circuits for guiding complex, flexible, robust outcomes early on; ion channels and neurotransmitter molecules (which move as a result of bioelectric gradients) are present in all somatic cell types, not just excitable nerve and muscle. They are also present in unicellular organisms and predate multicellularity.81-83 The extensive use of rapid bioelectrical signaling in the CNS likely offers a hint of more fundamental somatic functions from which neural dynamics evolved.

Mechanistically, bioelectric circuits in the brain and body use the same, highly conserved set of molecules and pathways. But they use them for very different purposes. The brain harnesses bioelectricity for the control of muscles, moving a body through 3-dimensional space. Developmental bioelectricity controls cell functions such as proliferation and differentiation, 84,85 moving the configuration of a body through morphospace, the abstract mathematical space in which an organism's shape is defined. As the chromatin code helps set cell identity and physiology, and physical forces are exploited by genetic circuits to control morphogenesis, bioelectricity enables the genome to utilize unique computational capabilities and long-range information integration.⁸⁶⁻⁹⁰ Recently we posed the hypothesis that evolution exploits plasticity of non-neural (body-wide) bioelectric networks to process epigenetic information, as does the brain, but whose output is pattern regulation.^{53,91} This hypothesis makes a number of specific predictions which have been borne out by recent results.

Switching species-specific head morphology in planaria

The planarian flatworm is an excellent model system for the study of the generation of form. 92,93 Planarians have extraordinary regenerative abilities, allowing them to reproduce any tissue or organ system given traumatic injury.94 In addition, they are amenable to a variety of molecular biology techniques, and show a diverse array of head shapes across species. 95,96 Alteration of bioelectrical homeostasis in the planarian during regeneration has been shown to lead to dramatic patterning defects, including duplication of anterior polarity, 97 altered scaling of head structures, 98 and most recently, alteration of species-specific head shape.⁹⁹ These data implicate bioelectrical signaling in the maintenance of rational regenerative programs, and indicate an interesting modality with which large-scale morphology may be perturbed experimentally in the laboratory, or on evolutionary timescales.

When the electrical coupling of large populations of cells in the planarian species G. dorotocephala is transiently perturbed using the non-mutagenic long chain alcohol octanol, regenerating fragments grow heads that resemble entirely different species of planarian (Fig. 3A-D, Ai–Di). While the appropriate head morphology of *G*. dorotocephala is triangular with pointed auricles, octanol-treated fragments stochastically regenerate as several discrete shapes belonging to other species. Observed head shapes include those that are rounded, like S. mediterranea, triangular with no auricles like D. japonica, or

flattened with anterior auricles like P. felina, 99 and require multiple coordinated shape changes that are not explained by simple scaling along 1 dimension (Supplement 1). Importantly, this head shape change was not only skin deep - it included coherent, multi-tissue anatomical change, not only editing of the external mor-Immunostaining against synapsin phosphorylated histone 3 (a marker of proliferation) revealed that morphology of the brain and the distribution of the adult stem cell population are also altered to resemble that of the corresponding other species (Fig. 3Aii-iv, Bii-iv, Cii-iv, Dii-iv). The significance of shape alteration was validated using geometric morphometrics, and the shape space defined by canonical variate analysis (CVA) allowed for characterization of morphological transformations (Fig. 3E).

These data indicate that some of the difference among planarian species could be due to the dynamics of signaling among their bioelectric networks, revealing a new epigenetic layer that could have important implications for evolution. Indeed, the frequency of appearance of the various planarian species' heads from random bioelectric network perturbation was proportional to the evolutionary distance among species, suggesting a relationship between morphological speciation and modification of bioelectrical dynamics. The evolutionary implications of bioelectrical controls of patterning have been discussed elsewhere, 53 and offer rich opportunities for future study of the bi-directional interplay between mutations of ion channel / gap junction genes and intrinsic (selforganizing) dynamics of bioelectric circuits. 100-103 One possibility is that the frameworks used to analyze evolutionary dynamics of memory and behavior such as Baldwin-like effects^{46,104} (also mediated by a set of bioelectric circuits, those in the CNS) could be fruitfully applied to pattern regulation in embryogenesis and regeneration, and the variability of developmental outcomes from a specific genome. 105,106

The voltage reporter dye DiBAC4(3) was used to show that bioelectric coupling among cells in worms with inappropriate head morphologies is altered long after octanol is washed out of the body of the worm, revealing long-term changes to the physiological network among cells (plasticity akin to the synaptic plasticity observed in electrical synapses in the CNS^{72,99}). Efforts to build quantitative models of specific head shape outcomes stored in stable bioelectrically-mediated pattern memories in tissue are ongoing in our lab. Importantly, unlike the permanent morphological change (double headedness) induced in D. japonica, 53,107 in the time required for complete cellular turnover in the planarian (roughly




Figure 3. Changes to species-specific head shape in planaria. Physiological determination of species-specific head anatomy in the planarian flatworm. Transient exposure to octanol after amputation in G. dorotocephala results in regeneration of head anatomies resembling other species of planarian. Brain shape and distribution of neoblasts is also altered. Shape change can be quantified using geometric morphometrics, and used to produce a shape space accounting for much of the variation in shape between species. (A) Wild type G. dorotocephala morphology. (Ai) pseudo G. dorotocephala morphotype after octanol treatment. (Aii) brain morphology of pseudo G. dorocephala morphology by anti-synapsin immunostaining. (Aiii) neoblast distribution of pseudo G. dorotocephala morphology by anti-phosphorylated histone 3 immunostaining. (Aiv) wild type G. dorotocephala brain morphology and neoblast distribution. (B) Wild type D. japonica morphology. (Bi) pseudo D. japonica morphotype after octanol treatment. (Bii) brain morphology of pseudo D. japonica morphology by anti-synapsin immunostaining. (Biii) neoblast distribution of pseudo D. japonica morphology by anti-phosphorylated histone 3 immunostaining. (Biv) wild type D. japonica brain morphology and neoblast distribution. (C) Wild type S. mediterranea morphology. (Ci) pseudo S. mediterranea morphotype after octanol treatment. (Cii) brain morphology of pseudo S. mediterranea morphology by anti-synapsin immunostaining. (Ciii) neoblast distribution of pseudo S. mediterranea morphology by anti-phosphorylated histone 3 immunostaining. (Civ) wild type S. mediterranea brain morphology and neoblast distribution. (D) wild type P. felina morphology. (Di) pseudo P. felina morphotype after octanol treatment. (E) CVA analysis of wild type morphologies and pseudo morphologies, with morphometric landmarks shown on a wild type G. dorotocephala head. Figures used with permission from ref. 163.




Figure 4. Morphogenetic state spaces. A conceptual model of shape change driven by physiological network dynamics. Planaria regeneration parallels classical neural network behavior; both can be described in terms of free energy landscapes with multiple attractor states. (A) Behavior of a classical Hopfield neural network trained to reproduce 3 types of patterns, in this case shapes of the letters 'F', 'H', or 'G', which are the 3 stable states of the network's free energy landscape. The state of the Hopfield network's nodes directly relate to the brightness of pixels on a display, generating output. Perturbation of the network from a stable state (red arrow) by deleting (damaging) part of the pattern is akin to moving a ball to an unstable point on the free energy landscape, and leads to regeneration of the closest learned attractor state (blue arrow). In this, such networks' well-known ability to implement memory is analogous to regenerating organisms restoring a specific target morphology upon damage. (B) The parallel concept of planaria regeneration into head shapes of one of 3 different species, which are attractor states of the free energy landscape. Outcome morphology is driven by the dynamic outputs of physiological cellular network. Amputation (red arrow) is akin to moving the system to an unstable point on the free energy landscape. Normal regeneration would return the system to its main attractor, but altering cell network dynamics via gap junction blockade allows for regenerative transitions (blue arrow) to alternative local minima, corresponding to morphospace regions normally occupied by P. felina and S. mediterranea worms. In time, remodeling (green arrow) transfers these morphologies to the global minimum of the wild-type state (G. dorotocephala). (C) Morphospaces are conceptual structures within which distances along specific metrics can represent the differences among species' morphologies. This panel illustrates how variants in the shape of a structure (skull shapes in this case) can be represented in a virtual space describing several orthogonal control parameters. We suggest that the physiologically-induced conversion of an animal with a normal genome into a different species-specific morphology could be modeled by an appropriate bioelectric circuit model whose measured states control relevant parameters forming the axes of a morphospace. Such spaces often include stable attractors corresponding to anatomical configurations that stable to small perturbations of the key parameters. (D) It is possible that the global coordinate axes that facilitate the Thompson transformations are mediated by bioelectric field properties across the organism. Images used with permission as follows: A,B, 163 C, 113 D. 115

30 days), altered head shapes in G. dorotocephala are remodeled back to the species-appropriate morphology, along linear pathways in the shape space defined by CVA.99

The output of bioelectric circuits are spatially-extended instructive signals for cell behaviors such as proliferation, differentiation, and apoptosis. 54,108 Conceptually, we suggest¹⁰⁹ to represent the head shape morphologies as stable

attractors in the state space of the physiological network guiding cell behavior in vivo (Fig. 4A and B). This has the advantage that a significant body of work exists in the computational neuroscience and artificial neural network fields, explaining the dynamics of pattern memories in such networks. 110-112 Each head shape corresponds to a distinct low-energy state of the bioelectric circuit. Amputation temporarily raises the "energy" of the system, which normally relaxes into the standard species-specific shape attractor. However, perturbing the connectivity in the circuit can cause it to instead settle into a different attractor (Fig. 4B). We are currently working on deriving quantitative models of this process that will explain why the above-mentioned shapes are transient (belong to only local minima) while other states (the default head shape, and the 2-head

morphology described below) represent permanent, or very deep, minima.

The conversion among different species-specific shapes is formalized in the notion of morphospace 18,113,114 (Fig. 4C), where specific outcomes of some system (the key canonical variates) can explain a range of observed morphologies. A very early line of thought in this direction was made by D'Arcy Thompson, 115,116 who showed that the bodyplans of distinct species can be converted to those of other species via simple geometric transformations (Fig. 4D). Biochemical gradients have been suggested for the mechanism implementing such spatial axes, 117 and it is also tempting to speculate that these coordinates can also be provided by bioelectrical aspects of large-scale physiological networks. 118,119 Specification and empirical validation of a




Figure 5. Unexplored regions of morphospace reached by editing bioelectric circuits. Briefly altering bioelectric circuit dynamics, despite the presence of a normal genome, can result in drastic alterations of the bodyplan, to regions not currently occupied by extant species. D. japonica worms treated with disruptors of gap junctional connectivity or modulators of ion channel-dependent bioelectric signaling can acquire compound (A), spiky (B), or cup-shaped (C) morphologies instead of the genome-default flat architecture of the planarian. The specimen in panel D has been stained with an antibody to reveal the central nervous system, and cleared to highlight the pocketlike morphology. In some cases, these edits to the normal target morphology can be permanent (E): a flatworm middle fragment that regenerates 2 heads after a brief reduction of bioelectrical coupling among its cells will continue to regenerate as 2-headed in subsequent cuts made in plain water. Even if the "reprogrammed" head tissue is removed, the target morphology information in other fragments of the body have been physiologically altered so that a 2-headed form results. This new worm architecture has distinct behavioral and anatomical structures and is stable across the most common mode of reproduction in this species (fission + regeneration), stretching the definition of speciation. (F) Changes of head shape (red arrowhead) and number (yellow arrow) can occur in the same animal, as can changes in head size (G, red arrow). Editing of large-scale bodyplan can be induced in vertebrate models as well, inducing ectopic eye growth out of gut tissue (red arrow), and ectopic limb growth out of the mouth (I, red arrow). Images used with permission as follows: E,⁵³ G,¹²⁴ H.⁶⁷ Photo in panel I courtesy of Erin Switzer.

quantitative model of a bioelectric circuit and the morphological consequences of its stable modes represents one of the most exciting prospects in this field. A full understanding must also involve explanation of the stochastic nature of this process, perhaps couched in the language of dynamical systems theory, 120-123 to understand the origin of variability. The study of deterministic chaos has long helped analyze many systems where very small initial differences (e.g., physiological noise) can be amplified by feedback loops into distinct large-scale outcomes in a stochastic manner. Such feedback loops are certainly present in bioelectric regulation of organ-level patterning, such as between V_{mem} and the gene Rx1 in eye induction⁶⁷ or V_{mem} and the gene *Notch* in brain patterning.⁶⁸

Interestingly, while perturbing the topology of the physiological network results in head shapes of several extant species (Fig. 3), the same techniques can result in animals that correspond to regions of the morphospace 18,113,114 not exploited by evolution (at least in currently-known variants). For example, planaria can be deviated from their normally flat body architecture to give rise to geometrically very distinct shapes, such as pockets (Fig. 5A-D). Another example is revealed in D. japonica, where a 2-headed form can be produced which is stable – further cycles of amputation (in plain water) of the ectopic head result in 2-headed worms from normal gut fragments, revealing a permanent change of the encoded target morphology. 107 What is to be made of the fact that GJ inhibition induces a change in head number in D. japonica (Fig. 5E) but a change in head shape in G. dorotocephala (Fig. 3)? One possibility is that different species of planaria (which in this case differ by an estimated 100 million years) utilize the outputs of their bioelectric networks for different purposes (determination of head number in *D. japonica*, and of head shape in *G. doro-*




Figure 6. Altering neurotransmitter signaling induces inter-species shape changes in Xenopus tadpoles. (A) Treatment with the β -adrenergic agonist Cimaterol (CIM) stochastically induced 4 distinct head anatomies. Embryos were exposed to CIM from st. 10-45, sacrificed at st. 47 and stained with Alcian blue. (i) Wild-type Xenopus tadpole with major ventral craniofacial cartilages labeled on one side of the face. Meckel's cartilage (m) is outlined in Pink, ceratohyal cartilage (c) is outlined in green, and the branchial arches (ba) are outlined with yellow. (ii) Tadpole with malformed jaw, but otherwise normal craniofacial morphology. (iii) Tadpole with horizontal or "flat" branchial arches and ceratohyal cartiages. (iv) Tadpole whose head anatomy visually resembles that of Rana frogs. (v) Tadpole whose head anatomy resembles that of frogs belonging to the family Microhylidae. (B) Comparison of Pseudo Rana head morphology to that of wild-type Rana species. (i) Graphical output for Canonical Variate analysis of shape data in wild-type and experimentally-derived craniofacial morphologies, showing confidence ellipses for means at an 0.9 probability. Each point represents one individual's face shape data. Data for wildtype Rana frogs were derived from anatomical diagrams of Rana temporaria, 130 Rana dalmatina, 132 and Rana palustris. 131 (ii) ventral view of alcian-blue stained Pseudo rana tadpole with landmarks (black dots) used for morphometric analysis. (iii) Anatomical diagram of ventral craniofacial skeleton of Rana temporaria tadpole. For both ii and iii, protruding nasal cartilages are indicated with red arrows, Meckel's cartilages are outlined in pink, and branchial arches are outlined in yellow. (C) Comparison of Pseudo Microhylidae head morphology to that of wild-type Microhylidae species. (i) Graphical output for Canonical Variate analysis of shape data in wild-type and experimentally-derived craniofacial morphologies, showing confidence ellipses for means at an 0.9 probability. Each point represents one individual's face shape data. Data for wild-type Microhylidae frogs were derived from anatomical diagrams of *Microhyla ornata*, ¹³³ *Gastrophryne* carolinensis. 134 and Dermatonotus muelleri. 135 (ii) ventral view of alcian-blue stained Pseudo Microhylidae tadpole with landmarks (black dots) used for morphometric analysis. (iii) Anatomical diagram of ventral craniofacial skeleton of a Microhyla ornata tadpole. For both ii and iii, ceratohyal cartilages are outlined in green, and branchial arches are outlined in yellow.

tocephala). However, this is unlikely because worms (Fig. 5F) can be created which have both an abnormal number of heads and a different species' head shape. In S. mediterranea, head size (Fig. 5G) is regulated by distribution of voltage gradients, ¹²⁴ echoing the bioelectric control of brain size in Xenopus⁶⁸ and tail size in zebrafish.⁷⁰ In Xenopus, similar modulation of resting potential (intrinsic plasticity, as distinct from the synaptic plasticity edited by gap junctional modulators), can produce aberrant forms possessing ectopic eyes⁶⁷ (Fig. 5H) or limbs (Fig. 5I). One possible implication of these data is that, once predictive, quantitative, biophysical models that link bioelectric circuit modes to anatomical outcomes are developed, it would be possible to approach cell groups as "universal constructors", inducing pattern formation that has never before appeared in biological evolution, for purposes of creating synthetic organisms for specific functions.^{2,125-127}

Frog embryogenesis and neurotransmitter controls

Continuing the theme of plasticity in pattern regulation, it was recently shown that frog development does not merely utilize a hardwired set of deformations that convert a tadpole face into the very different frog face. Rather, it includes a robust and plastic system that can make up for deformed tadpole anatomies (which begin the remodeling process in an abnormal starting state) and correct them to make largely normal froglets. 128 The best-known system that is able to reliably reach specific goal states from diverse starting conditions is the brain, where goal-satisfaction circuits make use of bioelectrical networks (composed of neural cells and non-spiking astrocytes) and the flow of neurotransmitter signals to implement flexible decision-making. We recently began to examine the roles of these neurotransmitters during frog development. In a pharmacological survey screen, we found that treatment of embryos prior to the mature development of the nervous system with drugs modulating catecholamine and glutamatergic signaling induced a wide variety of alterations in craniofacial, eye, muscle, visceral organ, and pigment patterns. 129

Interestingly, we identified several neurotransmitter endpoints that when manipulated pharmacologically, caused the morphology of the head and tail to resemble those of other species. The β -adrenergic agonist Cimaterol altered the shape and placement of the craniofacial cartilages, stochastically inducing 4 distinct head anatomies (Fig. 6A). One such anatomy (which we label as "Pseudo-*Rana*" below) resembled the larval skull of frogs belonging to the genus *Rana* due to proboscis-like nasal cartilages, inset Meckel's cartilages, and compressed branchial arches (Fig. 6B). Another (which we term "Pseudo-*Microhylidae*") resembled the head anatomy

seen in tadpoles belonging to the Microhylid family of frogs due to the dramatic arching of its ceratohyal cartilages and the steep downward slant of its branchial arches (Fig. 6C).

Geometric morphometrics was used to compare experimentally-derived head morphologies to wild-type *Xenopus*, as well as compare "pseudo" head morphologies to the craniofacial anatomy of frogs belonging to the genus or family that they resemble. Principal Component analysis and Canonical Variate analysis (CVA) were run on landmark data taken from alcian-blue images of control and experimental *Xenopus* tadpoles as well as anatomical diagrams of different frog species (Fig. 6Bi, 6Ci). The Rana species analyzed included *Rana temporaria*, ¹³⁰ *Rana palustris*, ¹³¹ and *Rana dalmatina*. ¹³² In the Microhylid family of frogs, we analyzed *Microhyla ornata*, ¹³³ *Gastrophryne carolinensis*, ¹³⁴ and *Dermatonotus Muelleri*. ¹³⁵

In our CVA using nasal, branchial arch, and Meckel's cartilage landmarks, the Pseudo-Rana clustered dramatically closer in morphospace to true rana than wild-type Xenopus on both the CV1 and CV2 axes (Fig. 6B). Movement along the CV1 axis represents anterior movement of the inner branchial arch landmarks and posterior movement of the outer branchial arch landmarks. Movement along the CV2 axis represents movement of the Meckel's cartilages toward the midline, movement of the inner branchial arch landmarks away from the midline, and anterior projection of the nasal cartilage. We can therefore confirm that out Pseudo-Rana tadpoles were indeed more anatomically similar to true Rana tadpoles than to wild-type Xenopus due to the visual features we observed. In our CVA using branchial arch and ceratohyal landmarks, we found that our Pseudo-Microhylidae clustered dramatically closer to true Microhylids than to wild-type Xenopus along CV1 (Fig. 6C). On CV2, pseudo and true Microhylidae diverged equidistantly from wild-type Xenopus. In this analysis, movement along CV2 represents compression of the ceratohyal cartilages toward the midline and movement of the outer branchial arch landmarks away from the midline. Movement along CV1, however, represents anterior movement and bunching of the inner branchial arch landmarks, posterior movement and bunching of the outer branchial arch landmarks, and posterior movement of the outer-most ceratohyal cartilage landmarks. Although pseudo and true Microhylidae diverge on CV2, their locations along CV1 account for the easily visible traits that make them appear Microhylidae-like and different from wild-type, such as notably arched ceratohyal cartilages and steep downward-sloping branchial arches. Therefore, we can still confirm that Pseudo Microhylidae are anatomically closer to true Microhylids than they are to untreated Xenopus.

In addition to the craniofacial alterations caused by Cimaterol, we found another interesting inter-species phenotype during our drug screen. Treatment with the noncompetitive NMDA antagonist Norketamine induced the development of a tail that was morphologically similar to that of a zebrafish (Fig. 6D). As in zebrafish, the portion of the tail containing spinal cord and muscle was truncated and did not continue to the posterior most edge of the tailfin as it does in wild-type Xenopus. In addition, the tailfin was rounded as opposed to the normal pointed shape of a Xenopus tail. These data implicate pre-nervous (and non-neural) neurotransmitter signaling as a key functional player in diverse aspects of vertebrate embryonic development. Echoing the above-mentioned work, neurotransmitters have recently been found to modulate patterning in planaria, 136 the left-right axis in frog and chick, 56,57 cleavage and gastrulation in the sea urchin, ^{137,138} and craniofacial development in frogs and rodents. ^{139,140,141,142}

More broadly however, these data are consistent with the role of neurotransmitters as chemical cues that participate in an active and ongoing interplay with the bioelectric signals in the brain. Neurotransmitters are traditionally known to act as downstream effectors of bioelectricity, yet they also have the power to alter electrical communication between cells by coupling to signaling cascades capable of altering the connectivity of gap junctions. For example, glutamate (via the NMDA receptor 143,144) and serotonin (which stimulates IP3 mediated calcium release¹⁴⁵) raise intracellular calcium concentration, triggering CamKII-mediated phosphorylation of gap junction proteins and altering the permeability of electrical synapses. Other neurotransmitters couple to signaling cascades involving cyclic nucleotides. For example, both nitric oxide (NO) and histamine modulate gap junction coupling via cGMP signaling. 146,147 Similarly, dopamine and noradrenaline make gap junctions less permeable by triggering cAMP-dependent PKA phosphorylation of connexins. 148,149 The bi-directional relationship between neurotransmitters and GJs enable yet another set of feedback loops. Synthesizing the frog and planaria data described above and elsewhere, 89,108,150 it appears that species-specific shapes can be set, and reset, by modulating physiological network topology, specific bioelectric states, and the interconnected neurotransmitter pathways that often mediate between electrical activity and cell functions. 56,59,138,151,152

Conclusion

Recent work has shown that species-specific shape is derived not only from regulatory gene loci and chromatin marking. Large-scale anatomical structure is also a function of physiological signaling among many cell types, and significant (coordinated) changes in morphology can be induced by altering the topology or function of physiological circuits. In some cases, these dynamics can be used to select among morphologies of existing species, although entirely novel forms can also result. Interestingly, manipulation of physiological signals results in a much wider overall range of patterning outcomes (Fig. 5) than has been seen in genetic screens targeting individual gene products. 153-156 This new set of inputs into anatomical structure has several important implications for evolution.

First, these data highlight the importance of truly epigenetic layers of influence over organism anatomy. While ion channel and gap junction proteins are specified in the genome, bioelectric circuits have their own unique and complex dynamics that derive from the fact that ion channels and GJs determine cell voltage but are also themselves regulated by voltage gradients. These feedback loops and the resulting electric circuit state transitions over time are not predictable from the rules governing genetic sequence, transcriptional networks, or chromatin state⁵³ because channels and GJs open and close post-translationally, implementing functional signaling that is invisible to profiling at transcriptional or translational levels. Divergence between genetic state and bioelectric state can be readily seen, as 2 cells with precisely the same complement of ion channels (i.e., identical on any molecular-genetic profile) can be in very different physiological states depending on whether their channels are open or closed. The functional signaling properties of physiological networks are determined by the electrical activity, not the mere presence or absence of specific molecules. This implies a departure from the standard molecular biological paradigm, where cell state is thought to be derivable by proteomic and transcriptomic profiling. Bioelectric information can only be read out in the living state (not in fixed, biochemicallyanalyzed tissue). This is analogous to the content of computer memory, which resides in the flow of energy through the circuit (and is not captured by a full account of the atoms making up the transistor hardware), and to the semantic content of a mature brain, which is neither visible from nor directly determined by the genome. These facts have clear implications for future efforts to fully profile and re-write the signaling that ultimately determines anatomical structure and its modification.

Second, bioelectric networks facilitate robustness of physiological states and the patterning they regulate (via negative feedback loops and long-range state sensing). Morphogenetic functions guided by bioelectric circuits are also robust to mutation in channels and their

transcriptional regulation. Multiple ion channels often contribute to the same V_{mem} state, with extensive compensation by other channel types (which is one factor confounding discovery of these circuits by single gene mutation screens). Moreover, bioelectric circuit modes show significant stability to some perturbations. 103 Thus, individual channel involvement can diverge evolutionarily, as long as the overall signaling dynamic remains.

Third, bioelectric states as control points tend to be powerful "master regulators", allowing the initiation of self-limiting patterning modules as subroutines from a low information content input (trigger); 66,157 this is due to the existence of positive feedback loops, which sustain and amplify bioelectric states once a threshold has been surpassed by a transient bioelectric stimulus (exploited also by action potentials in brain circuits). Such modularity is well known to improve evolvability. 158-160 All of these factors are likely to be exploited by evolution in its exploration of shape spaces. Fourth, these data suggest a complementary view to the usual perspective in which the DNA is the software implemented by a cell's hardware. An alternative view is that the DNA specifies the hardware (the complement of channels, neurotransmitters, and GJ proteins), while the resulting bioelectric activity of these circuits (with spontaneous symmetrybreaking, self-organization, and other complex dynamics) is the software. As in computer science, and as observed in the appearance of distinct animal shapes from the same genome (Figs. 3 and 5), many different types of software can be implemented on the same genetically-specified hardware.

The study of ways in which evolution exploited the computational properties of electric cell networks outside of the central nervous system is an exciting, emerging area of developmental biology. One corollary to this is the possible role of commensal species in determining growth and form, not only physiology. For example, bacteria are known to make compounds that modulate gap junctions and ion channels;¹⁶¹ thus, it is possible that factors produced by other species living within a host organism's body can serve as an additional input to the organism's pattern by editing or altering its endogenous bioelectrical circuits This adds a layer of complexity to the typical "evo-devo" story, but also provides new, tractable, potent control points to be exploited by biomedicine or synthetic bioengineering through the use of novel vectors for bioelectric modulation of host tissue. 162

The list of the molecular signals that propagate through physiological networks is likely to grow rapidly in the following years. However, several players (current, calcium, neurotransmitters) are already implicated. The molecular and algorithmic analogies between how somatic tissues and the brain utilize these same components are a fertile area for novel inquiry, and remain to be tested in specific contexts. It is likely that a mature understanding of the unique properties of bioelectrical signaling, and their integration into the molecular toolkits of developmental biology and synthetic bioengineering, will have transformative effects in a number of areas. In addition to a fuller appreciation of the interplay between genomes and physical forces²⁰ in evolution and development, the increased control over growth and form is sure to have numerous practical applications in biomedicine and beyond.

Glossary of unusual terms

Ectopic – in the case of an organ, located outside the area of the body where it develops during normal embryogenesis.

Target Morphology - the anatomical configuration toward which normal embryogenesis and regeneration coordinate cell behavior; the pattern which, once reached, makes further growth and remodeling stop. Living organisms seek to maintain target morphology against cell turnover, aging, cancer, and injury as a kind of pattern homeostasis.

Plasticity - flexibility that allows past events to alter future behavior of a system. Physiological plasticity enables bioelectric circuits to retain state memory or respond adaptively to future physiological inputs. Anatomical plasticity occurs when animal bodies remodel or regenerate despite diverse types of damage, reaching the same target morphology state from different starting configurations. The ability to respond to unpredictable stimuli with coherent repair programs is contrasted with hardwired responses.

Gap Junctions – Protein channels that enable cell-cell communication by directly connecting the cytoplasm of 2 adjacent cells. This allows ions and small molecules to flow from one cell to another as dictated by electrochemical gradients between the cells. Some gap junction proteins confer gating by electric and chemical signals, and thus both determine and are themselves regulated by bioelectric state. These electrical synapses are thus versatile elements that underlie physiological plasticity in neural and somatic circuits.

Axial polarity - consistent differences along a major body axis (dorsal-ventral, anterior-posterior, or leftright) of the body. Anatomical and histological polarity are preceded by, and established by, bioelectric and chemical gradients which control large-scale pattern.

Neoplastic - a state in which cells begin to ignore the normally tight patterning cues of the body and begin to change toward a "unicellular" phenotype. This developmental disorder is part of the process of carcinogenesis, where cells change shape, overproliferate, and eventually become highly motile and invade distant sites.



Tensegrity - organization of structure consisting of discrete un-stretchable components situated in a continuous web of tension that defines the overall shape of the structure. Tensegrity is one example of essentially physical (non-genetic) regulators of morphogenesis.

Morphospace - an abstract mathematical space in which shape is defined by independent shape metrics along each orthogonal axis. A point in the space represents one possible shape configuration for a biological structure, which may or may not describe an extant animal species. Movement along an axis of the space represents a particular geometric transformation.

Baldwin effect - a mechanism by which learned behavior can affect evolution. Systems which can learn from experience can thereby acquire adaptive states which facilitate evolution to novel fitness peaks. For example, non-genomic (physiological) changes to bioelectric circuits can result in new animal morphologies, which could eventually be canalized into genomic changes by selection and mutation (much as new phenotypes can eventually evolve when animals learn specific behaviors and interact in novel ways with their niche).

Dynamical Systems Theory - a set of mathematical approaches for learning to understand and rationally manipulate the behavior of complex systems. Often such systems feature recursive (feedback) components, resulting in surprising and rich behaviors.

Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Douglas J. Blackiston, Dany Adams, and Justin Guay for assistance with morphometrics, and members of the Levin lab for many useful discussions. We also thank Benjamin Wolfe, Jessica Whited, and Joshua Lapalme for critical revisions of the manuscript.

Funding

M.E.B. was supported by the Arnold and Mabel Beckman Foundation through a Beckman scholar fellowship. M.L. gratefully acknowledges support of the Paul G. Allen Family Foundation, the Templeton World Charity Foundation Inc., the W. M. Keck Foundation, and the G. Harold and Leila Y. Mathers Charitable Foundation.

References

[1] Levin M. The wisdom of the body: future techniques and approaches to morphogenetic fields in regenerative medicine, developmental biology and cancer. Regenerat

- Med 2011; 6:667-73; PMID:22050517; http://dx.doi.org/ 10.2217/rme.11.69
- [2] Doursat R, Sayama H, Michel O. Morphogenetic engineering. Berlin: Springer Berlin Heidelberg; 2012. Morphogenetic engineering: Reconciling self-organization and architecture; p. 1-24.
- [3] Ingber DE, Levin M. What lies at the interface of regenerative medicine and developmental biology? Development 2007; 134:2541-7; PMID:17553905; http://dx.doi. org/10.1242/dev.003707
- [4] Birnbaum KD, Alvarado AS. Slicing across kingdoms: regeneration in plants and animals. Cell 2008; 132:697-710; PMID:18295584; http://dx.doi.org/10.1016/j.cell.2008.01
- [5] Baddour JA, Sousounis K, Tsonis PA. Organ repair and regeneration: an overview. Birth Defects Res C Embryo Today 2012; 96:1-29; PMID:22457174; http://dx.doi. org/10.1002/bdrc.21006
- [6] Tsonis PA. Embryogenesis and carcinogenesis: order and disorder. Anticancer Res 1987; 7:617-23; PMID:3310849
- [7] Farinella-Ferruzza N. The transformation of a tail into a limb after xenoplastic transformation. Experientia 1956; 15:304-5; http://dx.doi.org/10.1007/BF02159624
- [8] Carr PA, Church GM. Genome engineering. Nat Biotechnol 2009; 27:1151-62; PMID:20010598; http://dx. doi.org/10.1038/nbt.1590
- [9] Heinz S, Romanoski CE, Benner C, Glass CK. The selection and function of cell type-specific enhancers. Nat Rev Mol Cell Biol 2015; 16:144-54; PMID:25650801; http://dx.doi.org/10.1038/nrm3949
- [10] Rajagopal J, Stanger BZ. Plasticity in the adult: how should the waddington diagram be applied to regenerating tissues? Dev Cell 2016; 36:133-7; PMID:26812013; http://dx.doi.org/10.1016/j.devcel.2015.12.021
- [11] Coon KL, Vogel KJ, Brown MR, Strand MR. Mosquitoes rely on their gut microbiota for development. Mol Ecol 2014; 23:2727-39; PMID:24766707; http://dx.doi.org/ 10.1111/mec.12771
- [12] Shin SC, Kim SH, You H, Kim B, Kim AC, Lee KA, Yoon JH, Ryu JH, Lee WJ. Drosophila microbiome modulates host developmental and metabolic homeostasis via insulin signaling. Science 2011; 334:670-4; PMID:22053049; http:// dx.doi.org/10.1126/science.1212782
- Sovrano VA, Bertolucci C, Frigato E, Foa A, Rogers LJ. Influence of exposure in ovo to different light wavelengths on the lateralization of social response in zebrafish larvae. Physiol Behav 2016; 157:258-64; PMID:26875515; http:// dx.doi.org/10.1016/j.physbeh.2016.02.016
- [14] Crews D. Temperature-dependent sex determination: the interplay of steroid hormones and temperature. Zoolog Sci 1996; 13:1-13; PMID:8688803; http://dx.doi. org/10.2108/zsj.13.1
- [15] Lobo D, Vico FJ. Evolutionary development of tensegrity structures. Biosystems 2010; 101:167-76; PMID:20619314; http://dx.doi.org/10.1016/j.biosystems.2010.06.005
- [16] Raspopovic J, Marcon L, Russo L, Sharpe J. Modeling digits. Digit patterning is controlled by a Bmp-Sox9-Wnt Turing network modulated by morphogen gradients. Science 2014; 345:566-70; PMID:25082703; http://dx.doi.org/10.1126/science.1252960

- [17] Rogers KW, Schier AF. Morphogen gradients: from generation to interpretation. Ann Rev Cell Dev Biol 2011; 27:377-407; http://dx.doi.org/10.1146/annurevcellbio-092910-154148
- [18] Newman SA, Bhat R. Dynamical patterning modules: a "pattern language" for development and evolution of multicellular form. Int J Dev Biol 2009; 53:693-705; PMID:19378259; http://dx.doi.org/10.1387/ijdb.072481sn
- [19] Newman SA. Physico-genetics of morphogenesis: the hybrid nature of developmental mechanisms. Towards a Theory Dev 2014:95-113; http://dx.doi.org/10.1093/ acprof:oso/9780199671427.003.0006
- [20] Newman SA, Comper WD. 'Generic' physical mechanisms of morphogenesis and pattern formation. Development 1990; 110:1-18; PMID:2081452
- [21] Miller CJ, Davidson LA. The interplay between cell signalling and mechanics in developmental processes. Nat Rev Genet 2013; 14:733-44; PMID:24045690; http://dx. doi.org/10.1038/nrg3513
- [22] von Dassow M, Davidson LA. Physics and the canalization of morphogenesis: a grand challenge in organismal biology. Phys Biol 2011; 8:045002; PMID:21750364; http://dx.doi.org/10.1088/1478-3975/8/4/045002
- [23] Vandenberg LN, Levin M. A unified model for leftright asymmetry? Comparison and synthesis of molecular models of embryonic laterality. Dev Biol 2013; 379:1-15; PMID:23583583; http://dx.doi.org/ 10.1016/j.ydbio.2013.03.021
- [24] Vandenberg LN, Lemire JM, Levin M. It's never too early to get it Right: A conserved role for the cytoskeleton in left-right asymmetry. Commun Integr Biol 2013; 6:e27155; PMID:24505508; http://dx.doi.org/10.4161/ cib.27155
- [25] Speder P, Petzoldt A, Suzanne M, Noselli S. Strategies to establish left/right asymmetry in vertebrates and invertebrates. Curr Opin Genet Dev 2007; 17:351-8; PMID:17643981; http://dx.doi.org/10.1016/ j.gde.2007.05.008
- [26] Levin M, Johnson RL, Stern CD, Kuehn M, Tabin C. A molecular pathway determining left-right asymmetry in chick embryogenesis. Cell 1995; 82:803-14; PMID:7671308; http://dx.doi.org/10.1016/0092-8674 (95)90477-8
- [27] Ramsdell AF, Yost HJ. Molecular mechanisms of vertebrate left-right development. Trends Genet 1998; 14:459-65; PMID:9825674; http://dx.doi.org/10.1016/ S0168-9525(98)01599-6
- [28] Bisgrove BW, Essner JJ, Yost HJ. Multiple pathways in the midline regulate concordant brain, heart and gut left-right asymmetry. Development 2000; 127:3567-79; PMID:10903181
- [29] Basu B, Brueckner M. Cilia: multifunctional organelles at the center of vertebrate left-right asymmetry. Curr Top Dev Biol 2008; 85:151-74; PMID:19147005; http:// dx.doi.org/10.1016/S0070-2153(08)00806-5
- [30] Lobikin M, Wang G, Xu J, Hsieh YW, Chuang CF, Lemire JM, Levin M. Early, nonciliary role for microtubule proteins in left-right patterning is conserved across kingdoms. Proc Natl Acad Sci USA 2012; 109:12586-91; PMID:22802643; http://dx.doi.org/10.1073/pnas.1202659109
- [31] Naganathan SR, Middelkoop TC, Furthauer S, Grill SW. Actomyosin-driven left-right asymmetry:

- molecular torques to chiral self organization. Curr Opin Cell Biol 2016; 38:24-30; PMID:26829488; http://dx.doi. org/10.1016/j.ceb.2016.01.004
- [32] Ramasubramanian A, Chu-Lagraff QB, Buma T, Chico KT, Carnes ME, Burnett KR, Bradner SA, Gordon SS. On the role of intrinsic and extrinsic forces in early cardiac S-looping. Dev Dyn 2013; 242:801-16; PMID:23553909; http://dx.doi.org/ 10.1002/dvdy.23968
- [33] Petzoldt AG, Coutelis JB, Geminard C, Speder P, Suzanne M, Cerezo D, Noselli S. DE-Cadherin regulates unconventional Myosin ID and Myosin IC in Drosophila left-right asymmetry establishment. Development 2012; 139:1874-84; PMID:22491943; http://dx.doi.org/ 10.1242/dev.047589
- [34] Lobo D, Solano M, Bubenik GA, Levin M. A linear-encoding model explains the variability of the target morphology in regeneration. J R Soc Interface 2014; 11:20130918; PMID:24402915; http://dx.doi.org/10.1098/rsif.2013.0918
- [35] Bubenik AB, Pavlansky R. Trophic responses to trauma in growing antlers. J Exp Zool 1965; 159:289-302; PMID:5883952; http://dx.doi.org/10.1002/jez.1401590302
- [36] Abzhanov A, Kuo WP, Hartmann C, Grant BR, Grant PR, Tabin CJ. The calmodulin pathway and evolution of elongated beak morphology in Darwin's finches. Nature 2006; 442:563-7; PMID:16885984; http://dx.doi.org/ 10.1038/nature04843
- [37] Abzhanov A, Protas M, Grant BR, Grant PR, Tabin CJ. Bmp4 and morphological variation of beaks in Darwin's finches. Science 2004; 305:1462-5; PMID:15353802; http://dx.doi.org/10.1126/science.1098095
- [38] Sears KE, Behringer RR, Rasweiler JJt, Niswander LA. Development of bat flight: morphologic and molecular evolution of bat wing digits. Proc Natl Acad Sci U S A 2006; 103:6581-6; PMID:16618938; http://dx.doi.org/ 10.1073/pnas.0509716103
- [39] Cretekos CJ, Wang Y, Green ED, Martin JF, Rasweiler JJt, Behringer RR. Regulatory divergence modifies limb length between mammals. Genes Dev 2008; 22:141-51; PMID:18198333; http://dx.doi.org/ 10.1101/gad.1620408
- [40] Goldberg AD, Allis CD, Bernstein E. Epigenetics: a landscape takes shape. Cell 2007; 128:635-8; PMID:17320500; http://dx.doi.org/10.1016/j.cell.2007.02.006
- [41] Rechavi O, Houri-Ze'evi L, Anava S, Goh WS, Kerk SY, Hannon GJ, Hobert O. Starvation-induced transgenerational inheritance of small RNAs in C. elegans. Cell 2014; 158:277-87; PMID:25018105; http://dx.doi.org/ 10.1016/j.cell.2014.06.020
- [42] Rechavi O, Minevich G, Hobert O. Transgenerational inheritance of an acquired small RNA-based antiviral response in C. elegans. Cell 2011; 147:1248-56; PMID:22119442; http://dx.doi.org/10.1016/j.cell.2011.10.042
- [43] Theissen G. Evolutionary developmental genetics of floral symmetry: the revealing power of Linnaeus' monstrous flower. Bioessays 2000; 22:209-13; PMID:10684579; http://dx.doi.org/10.1002/(SICI)1521-1878(200003)22:3%3c209::AID-BIES1%3e3.0.CO;2-J
- [44] Gao L, Geng Y, Li B, Chen J, Yang J. Genome, Äêwide DNA methylation alterations of Alternanthera philoxeroides in natural and manipulated habitats: implications for epigenetic regulation of rapid responses to

GY

- environmental fluctuation and phenotypic variation. Plant Cell Environ 2010; 33:1820-7; http://dx.doi.org/10.1111/j.1365-3040.2010.02186.x
- [45] Pieau C, Dorizzi M, Richard-Mercier N. Temperature-dependent sex determination and gonadal differentiation in reptiles. Cell Mol Life Sci CMLS 1999; 55:887-900; PMID:10412370; http://dx.doi.org/10.1007/s000180050342
- [46] Jablonka E. Epigenetic inheritance and plasticity: The responsive germline. Prog Biophys Mol Biol 2013; 111:99-107; PMID:22975443; http://dx.doi.org/10.1016/j.pbiomolbio.2012.08.014
- [47] Jablonka E. Epigenetic variations in heredity and evolution. Clin Pharmacol Ther 2012; 92:683-8; PMID:23073209; http://dx.doi.org/10.1038/clpt.2012.158
- [48] Jablonka E, Raz G. Transgenerational epigenetic inheritance: prevalence, mechanisms, and implications for the study of heredity and evolution. Q Rev Biol 2009; 84:131-76; PMID:19606595; http://dx.doi.org/10.1086/598822
- [49] Rakyan VK, Chong S, Champ ME, Cuthbert PC, Morgan HD, Luu KV, Whitelaw E. Transgenerational inheritance of epigenetic states at the murine AxinFu allele occurs after maternal and paternal transmission. Proc Natl Acad Sci 2003; 100:2538-43; http://dx.doi.org/10.1073/pnas.0436776100
- [50] Holliday R. The inheritance of epigenetic defects. Science 1987; 238:163-70; PMID:3310230; http://dx.doi.org/10.1126/science.3310230
- [51] Jablonka E, Lamb MJ. The inheritance of acquired epigenetic variations. J Theor Biol 1989; 139:69-83; PMID:2593687; http://dx.doi.org/10.1016/S0022-5193 (89)80058-X
- [52] Feinberg AP, Irizarry RA. Stochastic epigenetic variation as a driving force of development, evolutionary adaptation, and disease. Proc Natl Acad Sci 2010; 107:1757-64; http:// dx.doi.org/10.1073/pnas.0906183107
- [53] Levin M. Endogenous bioelectrical networks store non-genetic patterning information during development and regeneration. J Physiol 2014; 592:2295-305; PMID:24882814; http://dx.doi.org/10.1113/ jphysiol.2014.271940
- [54] Bates E. Ion Channels in Development and Cancer. Annu Rev Cell Dev Biol 2015; 31:231-47; PMID:26566112; http://dx.doi.org/10.1146/annurev-cellbio-100814-125338
- [55] Funk R. Ion gradients in tissue and organ biology. Biol Sys 2013; 2(1); http://dx.doi.org/10.4172/2329-6577. 1000105
- [56] Fukumoto T, Kema IP, Levin M. Serotonin signaling is a very early step in patterning of the left-right axis in chick and frog embryos. Curr Biol 2005; 15:794-803; PMID:15886096; http://dx.doi.org/10.1016/j.cub.2005 .03.044
- [57] Fukumoto T, Blakely R, Levin M. Serotonin transporter function is an early step in left-right patterning in chick and frog embryos. Dev Neurosci 2005; 27:349-63; PMID:16280633; http://dx.doi.org/10.1159/000088451
- [58] Lobikin M, Lobo D, Blackiston DJ, Martyniuk CJ, Tkachenko E, Levin M. Serotonergic regulation of melanocyte conversion: A bioelectrically regulated network for

- stochastic all-or-none hyperpigmentation. Sci Signal 2015; 8(397):ra99; in press; PMID:26443706
- [59] Blackiston DJ, Anderson GM, Rahman N, Bieck C, Levin M. A novel method for inducing nerve growth via modulation of host resting potential: gap junctionmediated and serotonergic signaling mechanisms. Neurotherapeutics 2015; 12:170-84; PMID:25449797; http:// dx.doi.org/10.1007/s13311-014-0317-7
- [60] Pai VP, Martyniuk CJ, Echeverri K, Sundelacruz S, Kaplan DL, Levin M. Genome-wide analysis reveals conserved transcriptional responses downstream of resting potential change in Xenopus embryos, axolotl regeneration, and human mesenchymal cell differentiation. Regeneration 2015; 3:3–25
- [61] Tseng AS, Carneiro K, Lemire JM, Levin M. HDAC activity is required during Xenopus tail regeneration. PloS One 2011; 6:e26382; PMID:22022609; http://dx. doi.org/10.1371/journal.pone.0026382
- [62] Carneiro K, Donnet C, Rejtar T, Karger BL, Barisone GA, Diaz E, Kortagere S, Lemire JM, Levin M. Histone deacetylase activity is necessary for left-right patterning during vertebrate development. BMC Dev Biol 2011; 11:29; PMID:21599922; http://dx.doi.org/10.1186/1471-213X-11-29
- [63] Chernet BT, Levin M. Transmembrane voltage potential of somatic cells controls oncogene-mediated tumorigenesis at long-range. Oncotarget 2014; 5:3287-306; PMID:24830454; http://dx.doi.org/10.18632/oncotarget.1935
- [64] Levin M, Thorlin T, Robinson KR, Nogi T, Mercola M. Asymmetries in H+/K+-ATPase and cell membrane potentials comprise a very early step in left-right patterning. Cell 2002; 111:77-89; PMID:12372302; http:// dx.doi.org/10.1016/S0092-8674(02)00939-X
- [65] Adams DS, Robinson KR, Fukumoto T, Yuan S, Albertson RC, Yelick P, Kuo L, McSweeney M, Levin M. Early, H+-V-ATPase-dependent proton flux is necessary for consistent left-right patterning of non-mammalian vertebrates. Development 2006; 133:1657-71; PMID:16554361; http://dx.doi.org/10.1242/dev.02341
- [66] Tseng AS, Beane WS, Lemire JM, Masi A, Levin M. Induction of vertebrate regeneration by a transient sodium current. J Neurosci 2010; 30:13192-200; PMID:20881138; http://dx.doi.org/10.1523/JNEUROSCI.3315-10.2010
- [67] Pai VP, Aw S, Shomrat T, Lemire JM, Levin M. Transmembrane voltage potential controls embryonic eye patterning in Xenopus laevis. Development 2012; 139:313-23; PMID:22159581; http://dx.doi.org/10.1242/ dev.073759
- [68] Pai VP, Lemire JM, Pare JF, Lin G, Chen Y, Levin M. Endogenous gradients of resting potential instructively pattern embryonic neural tissue via notch signaling and regulation of proliferation. J Neurosci 2015; 35:4366-85; PMID:25762681; http://dx.doi.org/10.1523/JNEUROS CI.1877-14.2015
- [69] Vandenberg LN, Morrie RD, Adams DS. V-ATPasedependent ectodermal voltage and pH regionalization are required for craniofacial morphogenesis. Dev Dyn 2011; 240:1889-904; PMID:21761475; http://dx.doi.org/ 10.1002/dvdy.22685
- [70] Perathoner S, Daane JM, Henrion U, Seebohm G, Higdon CW, Johnson SL, Nüsslein-Volhard C, Harris MP.

- Bioelectric signaling regulates size in zebrafish fins. PLoS Genet 2014; 10:e1004080; PMID:24453984; http:// dx.doi.org/10.1371/journal.pgen.1004080
- [71] Lobikin M, Chernet B, Lobo D, Levin M. Resting potential, oncogene-induced tumorigenesis, and metastasis: the bioelectric basis of cancer in vivo. Phy Biol 2012; 9:065002; PMID:23196890; http://dx.doi.org/10.1088/ 1478-3975/9/6/065002
- [72] Pereda AE, Curti S, Hoge G, Cachope R, Flores CE, Rash JE. Gap junction-mediated electrical transmission: regulatory mechanisms and plasticity. Biochim Biophys Acta 2013; 1828:134-46; PMID:22659675; http://dx.doi. org/10.1016/j.bbamem.2012.05.026
- [73] Palacios-Prado N, Bukauskas FF. Heterotypic gap junction channels as voltage-sensitive valves for intercellular signaling. Proc Natl Acad Sci U S A 2009; 106:14855-60; PMID:19706392; http://dx.doi. org/10.1073/pnas.0901923106
- [74] Levin M. Gap junctional communication in morphogenesis. Prog Biophys Mol Biol 2007; 94:186-206; PMID:17481700; http://dx.doi.org/10.1016/j.pbiomol bio.2007.03.005
- [75] Mustard J, Levin M. Bioelectrical mechanisms for programming growth and form: taming physiological networks for soft body robotics. Soft Robotics 2014; 1:169-91; http://dx.doi.org/10.1089/soro.2014.0011
- [76] Borgens RB. The role of natural and applied electric fields in neuronal regeneration and development. Prog Clin Biol Res 1986; 210:239-50; PMID:3960913
- [77] Borgens RB. What is the role of naturally produced electric current in vertebrate regeneration and healing. Int Rev Cytol 1982; 76:245-98; PMID:6749746; http://dx. doi.org/10.1016/S0074-7696(08)61793-3
- [78] Mathews AP. Electrical polarity in the hydroids. Am J Physiol 1903; 8:294-9
- [79] Barth LG. Effect of constant electrical current on the regeneration of certain hydroids. Physiol Zool 1934; 7:340-64; http://dx.doi.org/10.1086/physzool.7.3.30152876
- [80] Jaffe LF. The role of ionic currents in establishing developmental pattern. Philos Trans R Soc Lond B Biol Sci 1981; 295:553-66; PMID:6117911; http://dx.doi.org/ 10.1098/rstb.1981.0160
- [81] Buznikov G, Shmukler Y, Lauder J. From oocyte to neuron: do neurotransmitters function in the same way throughout development? Cell Mol Neurobiol 1996; 16:537-59; PMID:8956008; http://dx.doi.org/10.1007/BF02152056
- [82] Keijzer F, van Duijn M, Lyon P. What nervous systems do: early evolution, input-output, and the skin brain thesis. Adapt Behav 2013; 21:67-85; http://dx.doi.org/ 10.1177/1059712312465330
- [83] Liebeskind BJ, Hillis DM, Zakon HH. Evolution of sodium channels predates the origin of nervous systems in animals. Proc Natl Acad Sci U S A 2011; 108:9154-9; PMID:21576472; http://dx.doi.org/10.1073/pnas.1106
- [84] Blackiston DJ, McLaughlin KA, Levin M. Bioelectric controls of cell proliferation: ion channels, membrane voltage and the cell cycle. Cell Cycle 2009; 8:3519-28; http://dx.doi.org/10.4161/cc.8.21.9888
- [85] Sundelacruz S, Levin M, Kaplan DL. Role of membrane potential in the regulation of cell proliferation and differentiation. Stem Cell Rev Rep 2009; 5:231-46;

- PMID:19562527; http://dx.doi.org/10.1007/s12015-009-9080-2
- [86] Levin M. Molecular bioelectricity in developmental biology: new tools and recent discoveries: control of cell behavior and pattern formation by transmembrane potential gradients. Bioessays 2012; 34:205-17; PMID:22237730; http:// dx.doi.org/10.1002/bies.201100136
- [87] Levin M, Stevenson CG. Regulation of cell behavior and tissue patterning by bioelectrical signals: challenges and opportunities for biomedical engineering. Annu Rev Biomed Eng 2012; 14:295-323; PMID:22809139; http:// dx.doi.org/10.1146/annurev-bioeng-071811-150114
- [88] Tseng A, Levin M. Cracking the bioelectric code: Probing endogenous ionic controls of pattern formation. Comm Integr Biol 2013; 6:1-8; http://dx.doi.org/ 10.4161/cib.22595
- [89] Levin M. Reprogramming cells and tissue patterning via bioelectrical pathways: molecular mechanisms and biomedical opportunities. Wiley Interdiscip Rev Sys Biol Med 2013; 5:657-76; PMID:23897652
- [90] Stewart S, Rojas-Munoz A, Izpisua Belmonte JC. Bioelectricity and epimorphic regeneration. Bioessays 2007; 29:1133-7; PMID:17935197; http://dx.doi.org/10.1002/
- [91] Pezzulo G, Levin M. Re-membering the body: applications of computational neuroscience to the top-down control of regeneration of limbs and other complex organs. Integr Biol (Camb) 2015; 7(12):1487-517; PMID:26571046
- [92] Lobo D, Beane WS, Levin M. Modeling planarian regeneration: a primer for reverse-engineering the worm. PLoS Computat Biol 2012; 8:e1002481; PMID:22570595; http://dx.doi.org/10.1371/journal.pcbi.1002481
- [93] Salo E, Abril JF, Adell T, Cebria F, Eckelt K, Fernandez-Taboada E, Handberg-Thorsager M, Iglesias M, Molina MD, Rodríguez-Esteban G. Planarian regeneration: achievements and future directions after 20 years of research. Int J Dev Biol 2009; 53:1317-27; PMID:19247944; http://dx.doi.org/10.1387/ijdb.072414es
- [94] Reddien PW, Alvarado AS. Fundamentals of planarian regeneration. Annu Rev Cell Dev Biol 2004; 20:725-57; PMID:15473858; http://dx.doi.org/10.1146/annurev. cellbio.20.010403.095114
- [95] Alvarado ASn, Newmark PA, Robb SM, Juste Rj. The Schmidtea mediterranea database as a molecular resource for studying platyhelminthes, stem cells and Development 2002; 129:5659-65; regeneration. PMID:12421706; http://dx.doi.org/10.1242/dev.00167
- [96] Sluys R. Global diversity of land planarians (Platyhelminthes, Tricladida, Terricola): a new indicator-taxon in biodiversity and conservation studies. Biodiversity & Conservation 1999; 8:1663-81; http://dx.doi.org/ 10.1023/A:1008994925673
- [97] Oviedo NJ, Morokuma J, Walentek P, Kema IP, Gu MB, Ahn J-M, Hwang JS, Gojobori T, Levin M. Long-range neural and gp junction protein-mediated cues control polarity during planarian regeneration. Dev Biol 2010; 339:188-99; PMID:20026026; http://dx.doi.org/10.1016/ j.ydbio.2009.12.012
- [98] Beane WS, Morokuma J, Lemire JM, Levin M. Bioelectric signaling regulates head and organ size during planarian regeneration. Development 2013;

- 140:313-22; PMID:23250205; http://dx.doi.org/ 10.1242/dev.086900
- [99] Emmons-Bell M, Durant F, Hammelman J, Bessonov N, Volpert V, Morokuma J, Pinet K, Adams DS, Pietak A, Lobo D, et al. Gap junctional blockade stochastically induces different species-specific head anatomies in genetically wild-type Girardia dorotocephala Flatworms. Int J Mol Sci 2015; 16:27865-96; PMID:26610482; http://dx.doi.org/10.3390/ijms161126065
- [100] Cervera J, Alcaraz A, Mafe S. Bioelectrical signals and ion channels in the modeling of multicellular patterns and cancer biophysics. Sci Rep 2016; 6:20403; PMID:26841954; http://dx.doi.org/10.1038/srep20403
- [101] Cervera J, Manzanares JA, Mafe S. Electrical coupling in ensembles of nonexcitable cells: modeling the spatial map of single cell potentials. J Phys Chem B 2015; 119:2968-78; PMID:25622192; http://dx.doi.org/ 10.1021/jp512900x
- [102] Cervera J, Alcaraz A, Mafe S. Membrane potential bistability in nonexcitable cells as described by inward and outward voltage-gated ion channels. J Phys Chem B 2014; 118:12444-50; PMID:25286866; http://dx.doi.org/ 10.1021/jp508304h
- [103] Law R, Levin M. Bioelectric memory: modeling resting potential bistability in amphibian embryos and mammalian cells. Theor Biol Med Model 2015; 12:22; PMID:26472354; http://dx.doi.org/10.1186/s12976-015-
- [104] Bronfman ZZ, Ginsburg S, Jablonka E. Shaping the learning curve: epigenetic dynamics in neural plasticity. Front Integr Neurosci 2014; 8:55; PMID:25071483; http://dx.doi.org/10.3389/fnint.2014.00055
- [105] Vogt G. Stochastic developmental variation, an epigenetic source of phenotypic diversity with far-reaching biological consequences. J Biosci 2015; 40:159-204; PMID:25740150; http://dx.doi.org/10.1007/s12038-015-
- [106] Palmer AR. Developmental plasticity and the origin of novel forms: unveiling cryptic genetic variation via "use and disuse". J Exp Zool B Mol Dev Evol 2012; 318:466-79; PMID:22038780; http://dx.doi.org/10.1002/jez. b.21447
- [107] Oviedo NJ, Morokuma J, Walentek P, Kema IP, Gu MB, Ahn JM, Hwang JS, Gojobori T, Levin M. Long-range neural and gap junction protein-mediated cues control polarity during planarian regeneration. Dev Biol 2010; 339:188-99; PMID:20026026; http://dx.doi.org/10.1016/ j.ydbio.2009.12.012
- [108] Levin M. Molecular bioelectricity: how endogenous voltage potentials control cell behavior and instruct pattern regulation in vivo. Mol Biol Cell 2014; 25:3835-50; PMID:25425556; http://dx.doi.org/10.1091/mbc.E13-
- [109] Friston K, Levin M, Sengupta B, Pezzulo G. Knowing one's place: a free-energy approach to pattern regulation. J R Soc Interface 2015; 12
- [110] Friston K. The free-energy principle: a unified brain theory? Nat Rev Neurosci 2010; 11:127-38; PMID:20068583; http:// dx.doi.org/10.1038/nrn2787
- [111] Friston K. Is the free-energy principle neurocentric? Nat Rev Neurosci 2010; 11:605

- [112] Friston KJ, Stephan KE. Free-energy and the brain. Synthese 2007; 159:417-58; PMID:19325932; http://dx.doi. org/10.1007/s11229-007-9237-y
- [113] Young NM, Hu D, Lainoff AJ, Smith FJ, Diaz R, Tucker AS, Trainor PA, Schneider RA, Hallgrímsson B, Marcucio RS. Embryonic bauplans and the developmental origins of facial diversity and constraint. Development 2014; 141:1059-63; PMID:24550113; http://dx.doi.org/ 10.1242/dev.099994
- [114] Rasskin-Gutman D, Izpisua-Belmonte JC. Theoretical morphology of developmental asymmetries. Bioessays 2004; 26:405-12; PMID:15057938; http://dx.doi.org/ 10.1002/bies.10410
- [115] Thompson DAW, Whyte LL. On growth and form. Cambridge Eng.: The University Press, 1942.
- [116] Stone JR. The spirit of D'arcy Thompson dwells in empirical morphospace. Math Biosci 1997; 142:13-30; PMID:9125858; http://dx.doi.org/10.1016/S0025-5564 (96)00186-1
- [117] Niehrs C. On growth and form: a Cartesian coordinate system of Wnt and BMP signaling specifies bilaterian body axes. Development 2010; 137:845-57; PMID:20179091; http://dx.doi.org/10.1242/dev.039651
- [118] Shi R, Borgens RB. Three-dimensional gradients of voltage during development of the nervous system as invisible coordinates for the establishment of embryonic pattern. Dev Dyn 1995; 202:101-14; PMID:7734729; http://dx.doi.org/10.1002/aja.1002020202
- [119] Burr HS, Northrop FSC. The electro-dynamic theory of life. Quart Rev Biol 1935; 10:322-33; http://dx.doi.org/ 10.1086/394488
- [120] Kaneko K, Tsuda I. Complex systems: chaos and beyond: a constructive approach with applications in life sciences. Berlin; New York: Springer, 2001.
- [121] Furusawa C, Kaneko K. Theory of robustness of irreversible differentiation in a stem cell system: chaos hypothesis. J Theor Biol 2001; 209:395-416; PMID:11319890; http://dx.doi.org/10.1006/jtbi.2001.2264
- [122] Barbi M. Chaos & noise in biology & medicine. Singapore: World Scientific Publishing Company, 1998.
- [123] Elbert T, Ray WJ, Kowalik ZJ, Skinner JE, Graf KE, Birbaumer N. Chaos and physiology: deterministic chaos in excitable cell assemblies. Physiol Rev 1994; 74:1-47; PMID:8295931
- [124] Beane WS, Morokuma J, Lemire JM, Levin M. Bioelectric signaling regulates head and organ size during planarian regeneration. Development 2013; 140:313-22; PMID:23250205; http://dx.doi.org/ 10.1242/dev.086900
- [125] Doursat R, Sayama H, Michel O. A review of morphogenetic engineering. Nat Comput 2013; 12:517-35; http:// dx.doi.org/10.1007/s11047-013-9398-1
- [126] Fernandez JD, Lobo D, Martin GM, Doursat R, Vico FJ. Emergent diversity in an open-ended evolving virtual community. Artif Life 2012; 18:199-222; PMID:22356151; http://dx.doi.org/10.1162/artl_a_00059
- [127] Kamm RD, Bashir R. Creating living cellular machines. Ann Biomed Eng 2014; 42:445-59; PMID:24006130; http://dx.doi.org/10.1007/s10439-013-0902-7
- [128] Vandenberg LN, Adams DS, Levin M. Normalized shape and location of perturbed craniofacial structures

- in the Xenopus tadpole reveal an innate ability to achieve correct morphology. Dev Dyn 2012; 241:863-78; PMID:22411736; http://dx.doi.org/10.1002/dvdy.23770
- [129] Sullivan KG, Levin M. Neurotransmitter signaling pathways required for normal development in Xenopus laevis embryos: a pharmacological survey screen. J Anat 2016; (in press); PMID:27060969
- [130] Parker WK. On the structure and development of the skull of the common Frog (Rana temporaria, L.). Philosoph Tran R Soc Lond 1871; 161:137-211; http://dx.doi. org/10.1098/rstl.1871.0009
- [131] Stone LS. Experiments showing the role of migrating neural crest (mesectoderm) in the formation of head skeleton and loose connective tissue in Rana palustris. Wilhelm Roux' Archiv für Entwicklungsmechanik der Organismen 1929; 118:40-77; http://dx.doi.org/10.1007/ BF02108871
- [132] Pelaz MP, Rougier C. A comparative study of buccal volumes and the branchial skeleton of rana ridibunda and R. dalmatina tadpoles. Copeia 1990:658-665
- [133] Khan MS. The larval hyobranchial skeleton of five anuran species and its ecological correlates. The larval hyobranchial skeleton of five anuran species and its ecological correlates. Herpetozoa 2003; 16:133-140
- [134] Trueb L, Diaz R, Blackburn DC. Osteology and chondrochranial morphology of Gastrophryne carolinensis (Anura: Microhylidae), with a review of the osteological diversity of New World microhylids. Phyllomedusa 2011; 10(2):99-135
- [135] Fabrezi M, Quinzio S, Goldberg J, Sá DRO. The development of dermatonotus muelleri (anura: microhylidae: gastrophryninae). J Herpetol 2012; 46:363-380
- [136] Chan JD, Agbedanu PN, Zamanian M, Gruba SM, Haynes CL, Day TA, Marchant JS. 'Death and axes': unexpected Ca(2+) entry phenologs predict new anti-schistosomal agents. PLoS Pathog 2014; 10:e1003942; PMID:24586156; http://dx.doi.org/10.1371/journal.ppat.1003942
- [137] Buznikov GA, Nikitina LA, Seidler FJ, Slotkin TA, Bezuglov VV, Milosevic I, Lazarević L, Rogac L, Ruzdijić S, Rakić LM. Amyloid precursor protein 96-110 and beta-amyloid 1-42 elicit developmental anomalies in sea urchin embryos and larvae that are alleviated by neurotransmitter analogs for acetylcholine, serotonin and cannabinoids. Neurotoxicol Teratol 2008; 30:503-9; PMID:18565728; http://dx.doi.org/ 10.1016/j.ntt.2008.05.003
- [138] Levin M, Buznikov GA, Lauder JM. Of minds and embryos: left-right asymmetry and the serotonergic controls of pre-neural morphogenesis. Dev Neurosci 2006; 28:171-85; PMID:16679764; http://dx.doi.org/ 10.1159/000091915
- [139] Reisoli E, De Lucchini S, Nardi I, Ori M. Serotonin 2B receptor signaling is required for craniofacial morphogenesis and jaw joint formation in Xenopus. Development 2010; 137:2927-37; PMID:20667918; http://dx.doi. org/10.1242/dev.041079
- [140] Reisoli E, De Lucchini S, Anelli T, Biagioni S, Nardi I, Ori M. Overexpression of 5-HT2B receptor results in retinal dysplasia and defective ocular morphogenesis in Xenopus embryos. Brain Res 2008; 1244:32-9; PMID:18930035; http://dx.doi.org/10.1016/j.brainres.2008.09.067
- [141] Cray JJ Jr, Weinberg SM, Parsons TE, Howie RN, Elsalanty M, Yu JC. Selective serotonin reuptake inhibitor

- exposure alters osteoblast gene expression and craniofacial development in mice. Birth Defects Res A Clin Mol Teratol 2014; 100:912-23; PMID:25308507; http://dx. doi.org/10.1002/bdra.23323
- [142] Moiseiwitsch JR, Lauder JM. Serotonin regulates mouse cranial neural crest migration. Proc Natl Acad Sci U S A 1995; 92:7182-6; PMID:7638165; http://dx.doi.org/ 10.1073/pnas.92.16.7182
- [143] Alev C, Urschel S, Sonntag S, Zoidl G, Fort AG, Hoher T, Matsubara M, Willecke K, Spray DC, Dermietzel R. The neuronal connexin36 interacts with and is phosphorylated by CaMKII in a way similar to CaMKII interaction with glutamate receptors. Proc Natl Acad Sci U S A 2008; 105:20964-9; PMID:19095792; http:// dx.doi.org/10.1073/pnas.0805408105
- [144] Pereda AE, Curti S, Hoge G, Cachope R, Flores CE, Rash JE. Gap junction-mediated electrical transmission: regulatory mechanisms and plasticity. Biochim Biophys Acta 2013; 1828:134-46; PMID:22659675; http://dx.doi. org/10.1016/j.bbamem.2012.05.026
- [145] Rorig B, Sutor B. Serotonin regulates gap junction coupling in the developing rat somatosensory cortex. Eur J Neurosci 1996; 8:1685-95; PMID:8921259; http://dx.doi. org/10.1111/j.1460-9568.1996.tb01312.x
- [146] Rörig B, Sutor B. Nitric oxide-stimulated increase in intracellular cGMP modulates gap junction coupling in rat neocortex. Neuroreport 1996; 7:569-72; PMID:8730831; http:// dx.doi.org/10.1097/00001756-199601310-00046
- [147] Hatton GI, Yang QZ. Synaptically released histamine increases dye coupling among vasopressinergic neurons of the supraoptic nucleus: mediation by H1 receptors and cyclic nucleotides. J Neurosci 1996; 16:123-9; PMID:8613778
- [148] Piccolino M, Neyton J, Gerschenfeld HM. Decrease of gap junction permeability induced by dopamine and cyclic adenosine 3':5'-monophosphate in horizontal cells of turtle retina. J Neurosci 1984; 4:2477-88; PMID:6092564
- [149] Urschel S, Hoher T, Schubert T, Alev C, Sohl G, Worsdorfer P, Asahara T, Dermietzel R, Weiler R, Willecke K. Protein kinase A-mediated phosphorylation of connexin36 in mouse retina results in decreased gap junctional communication between AII amacrine cells. J Biol Chem 2006; 281:33163-71; PMID:16956882; http:// dx.doi.org/10.1074/jbc.M606396200
- Adams DS, Levin M. Endogenous voltage gradients as mediators of cell-cell communication: strategies for investigating bioelectrical signals during pattern formation. Cell Tissue Res 2013; 352:95-122; PMID:22350846; http://dx.doi.org/10.1007/s00441-012-1329-4
- [151] Buznikov GA, Lambert HW, Lauder JM. Serotonin and serotonin-like substances as regulators of early embryogenesis and morphogenesis. Cell Tissue Res 2001; 305:177-86; PMID:11545255; http://dx.doi.org/10.1007/ s004410100408
- [152] Lobikin M, Lobo D, Blackiston DJ, Martyniuk CJ, Tkachenko E, Levin M. Serotonergic regulation of melanocyte conversion: A bioelectrically regulated network for stochastic all-or-none hyperpigmentation. Sci Signal 2015; 8:ra99; PMID:26443706; http://dx.doi.org/ 10.1126/scisignal.aac6609
- [153] Wang Y, Stary JM, Wilhelm JE, Newmark PA. A functional genomic screen in planarians identifies novel regulators of germ cell development. Genes Dev 2010;



- 24:2081-92; PMID:20844018; http://dx.doi.org/10.1101/ gad.1951010
- [154] Reddien PW, Bermange AL, Murfitt KJ, Jennings JR, Sanchez Alvarado A. Identification of genes needed for regeneration, stem cell function, and tissue homeostasis by systematic gene perturbation in planaria. Dev Cell 2005; 8:635-49; PMID:15866156; http://dx.doi.org/ 10.1016/j.devcel.2005.02.014
- [155] Newmark PA. Opening a new can of worms: a largescale RNAi screen in planarians. Dev Cell 2005; 8:623-4; PMID:15866153; http://dx.doi.org/10.1016/j. devcel.2005.04.007
- [156] Gawantka V, Pollet N, Delius H, Vingron M, Pfister R, Nitsch R, Blumenstock C, Niehrs C. Gene expression screening in Xenopus identifies molecular pathways, predicts gene function and provides a global view of embryonic patterning. Mech Dev 1998; 77:95-141; PMID:9831640; http://dx.doi.org/10.1016/S0925-4773(98)00115-4
- [157] Adams DS, Masi A, Levin M. H+ pump-dependent changes in membrane voltage are an early mechanism necessary and sufficient to induce Xenopus tail regeneration. Development 2007; 134:1323-35; PMID:17329365; http:// dx.doi.org/10.1242/dev.02812
- [158] Schlosser G, Wagner GP. Modularity in development and evolution. Chicago: University of Chicago Press, 2004.

- [159] Ravasz E, Somera AL, Mongru DA, Oltvai ZN, Barabasi AL. Hierarchical organization of modularity in metabolic networks. Science 2002; 297:1551-5; PMID:12202830; http://dx.doi.org/10.1126/science.1073374
- [160] Bolker JA. Modularity in development and why it matters to evo-devo. Am Zool 2000; 40:770-6
- [161] Hamilton-Kemp T, Newman M, Collins R, Elgaali H, Yu K, Archbold D. Production of the long-chain alcohols octanol, decanol, and dodecanol by Escherichia coli. Curr Microbiol 2005; 51:82-6; PMID:15991057; http://dx.doi.org/10.1007/s00284-005-4469-x
- [162] Chavez MN, Schenck TL, Hopfner U, Centeno-Cerdas C, Somlai-Schweiger I, Schwarz C, Machens HG, Heikenwalder M, Bono MR, Allende ML, et al. Towards autotrophic tissue engineering: Photosynthetic gene therapy for regeneration. Biomaterials 2016; 75:25-36; PMID:26474040; http://dx.doi.org/ 10.1016/j.biomaterials.2015.10.014
- Emmons-Bell M, Durant F, Hammelman J, Bessonov N, Volpert V, Morokuma J, Pinet K, Adams DS, Pietak A, Lobo D, et al. Gap junctional blockade stochastically induces different species-specific head anatomies in genetically wild-type girardia dorotocephala flatworms. Int J Mol Sci 2015; 16:27865-96; PMID:26610482; http://dx.doi.org/10.3390/ijms161126065