



The stability of memories during brain remodeling: A perspective

Douglas J Blackiston, Tal Shomrat & Michael Levin

To cite this article: Douglas J Blackiston, Tal Shomrat & Michael Levin (2015) The stability of memories during brain remodeling: A perspective, *Communicative & Integrative Biology*, 8:5, e1073424, DOI: [10.1080/19420889.2015.1073424](https://doi.org/10.1080/19420889.2015.1073424)

To link to this article: <https://doi.org/10.1080/19420889.2015.1073424>



© 2015 The Author(s). Published with license by Taylor & Francis Group, LLC© Douglas J Blackiston, Tal Shomrat, and Michael Levin



Published online: 30 Dec 2015.



Submit your article to this journal [↗](#)



Article views: 9287



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 17 View citing articles [↗](#)

The stability of memories during brain remodeling: A perspective

Douglas J Blackiston¹, Tal Shomrat^{2,3}, and Michael Levin^{1,*}

¹Center for Regenerative and Developmental Biology and Department of Biology; Tufts University; Medford, MA USA; ²Department of Neurobiology; Silberman Institute of Life Sciences; The Hebrew University of Jerusalem; Edmond J. Safra Campus; Jerusalem, Israel; ³School of Marine Sciences; Ruppin Academic Center; Michmoret, Israel

One of the most important features of the nervous system is memory: the ability to represent and store experiences, in a manner that alters behavior and cognition at future times when the original stimulus is no longer present. However, the brain is not always an anatomically stable structure: many animal species regenerate all or part of the brain after severe injury, or remodel their CNS toward a new configuration as part of their life cycle. This raises a fascinating question: what are the dynamics of memories during brain regeneration? Can stable memories remain intact when cellular turnover and spatial rearrangement modify the biological hardware within which experiences are stored? What can we learn from model species that exhibit both, regeneration and memory, with respect to robustness and stability requirements for long-term memories encoded in living tissues? In this Perspective, we discuss relevant data in regenerating planaria, metamorphosing insects, and hibernating ground squirrels. While much remains to be done to understand this remarkable process, molecular-level insight will have important implications for cognitive science, regenerative medicine of the brain, and the development of non-traditional computational media in synthetic bioengineering.

Introduction

Biological structures can exhibit considerable morphological plasticity. Many animals can regenerate their original anatomical structure after severe injury,¹ and to reconfigure toward the correct morphology after various grafting operations.²

For example, salamanders can regenerate their limbs, eyes, jaws, hearts, and tails after amputation,^{3,4} and tails grafted to the flank become slowly turned into limbs.⁵ Some remodel their bodies toward a new configuration as a normal part of their life cycle.^{6–8} In some species, this drastic remodeling also involves the nervous system, regenerating or massively rewiring the brain.⁹ For example, planaria regenerate complete new heads after amputation,¹⁰ and salamanders also regenerate their brain after region-specific ablation.^{11–13} Insects tear down their existing CNS and produce one with a different architecture, in the journey from a caterpillar to a butterfly.^{14,15}

However, a central feature of the nervous system is the ability to represent and store memories, allowing animals to adaptively alter behavior and cognition in light of past experience. The presence of memory and recall is revealed when experience alters behavior and cognition at future times when the original stimulus is no longer present. The brain, as the accepted seat of episodic memory, is often thought to be a stable, unchanging structure, which may seem to be a necessary property to ensure long-term stability of encoded information. Thus, animal model species that exhibit both, brain regeneration and learning, confront us with a fascinating set of questions. Could stable memories persist when cellular turnover and spatial rearrangement modify the hardware within which experiences are stored? What are the dynamics of memories when old cells apoptose, and new cells arise from progenitors? What are the mechanistic robustness and stability requirements for encoding of long-term memories in morphologically-dynamic living tissues? Answering these

Keywords: brain, learning, memory, metamorphosis, regeneration, remodeling

© Douglas J Blackiston, Tal Shomrat, and Michael Levin

*Correspondence to: Michael Levin; Email: michael.levin@tufts.edu

Submitted: 06/22/2015

Accepted: 07/13/2015

<http://dx.doi.org/10.1080/19420889.2015.1073424>

This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. The moral rights of the named author(s) have been asserted.

questions (at both, a molecular and computational level) would have numerous profound implications.

First, these issues directly target the fundamental question of the engram^{16,17}: how is information encoded in living tissue and what is the relationship between memory and the spatial location of specific physical modifications in the brain? If specific memories remain stable as cells are replaced and moved, we can learn much about the physico-chemical representations of mental content.¹⁸ Importantly, this is still a major area of debate. Many in the field believe memories to be stored as changes in synaptic connectivity.^{19,20} Nevertheless, new data suggest that other cell properties (such as transcriptional or epigenetic alterations) may be a crucial significant component.^{21,22}

Second, the future of regenerative therapies for brain injury and degeneration^{23,24} depends on our ability to modify brain structure without radically distorting the decades of memory content and personality of the patient. Given recent data implicating non-neuronal cell types in intelligence,^{25,26} a wide range of potential regenerative therapies may be expected to have effects on the rich memory set of adult human brains. It becomes imperative to understand how recall of established memories fares when novel cell types are introduced into the brain, altering network connectivity and perhaps replacing endogenous cell groups.

Third, could we exploit such mechanisms of memory in our efforts to bioengineer artificial hybrid biobots, which would have not only desired structural and physiological properties, but also useful behaviors that are robust to cellular turnover²⁷⁻²⁹? A related issue is that the discovery of such systems would significantly inform efforts to explore non-traditional computational architectures in information science and computer engineering,^{30,31} distinct from the current models of information inextricably dependent on a specific physical location within the digital memory.

Tantalizing findings have already shown that memories can indeed survive extensive remodeling of the brain, indicating transformative opportunities for both fundamental issues in cognitive science

and biomedical engineering. In this Perspective, we discuss relevant data in regenerating planaria, metamorphosing insects, and hibernating ground squirrels (Fig. 1). While much remains to be done, this fascinating intersection of regenerative cell biology and cognitive neuroscience represents a fertile ground for future work of not only deep basic interest but numerous practical applications.

Memories survive CNS rearrangements and remodeling in insects

Holometabolous insects (those which undergo complete metamorphosis) pose an interesting system in which to investigate memory storage and retrieval, as the CNS of these animals undergoes dramatic neurogenesis, pruning, and cell death during the transition from larvae to adulthood. Can memory survive pupation? What benefit would such a robust memory storage system provide the animal? These questions were first formalized nearly one hundred years ago by Andrew Delmar Hopkins, the father of forest entomology, who noted a polyphagous species of pine beetle preferentially infested one host even when multiple hosts were available.³² Hopkins postulated that female scolytid beetles chose to lay their eggs on the pine species they themselves consumed as larvae, a phenomenon which was later to become known as ‘Hopkins’ host-selection principle.’ This hypothesis, as well as that of cross-metamorphic memory in general, has generated large amounts of controversy with numerous reports claiming both support and opposition of the principle.

In support, a variety of studies using lepidoptera have demonstrated a contribution of larval experience on adult behavior,³³⁻³⁶ with adult oviposition preference being shaped by pre-imaginal (i.e. larval) events. It is possible that adults retain some memory of their larval lives in these studies, however, critics have pointed to an alternate mechanism which could explain the results termed the ‘chemical legacy hypothesis’.³⁷ The chemical legacy model proposes that odor contamination on the pupal case or within the hemolymph could sensitize adults to these odors while emerging, affecting the behavior of

the mature insect independent of larval experience. Indeed, numerous studies have directly tested and proven that a chemical legacy can have direct effects on adult behavior. *Drosophila* pupae intentionally contaminated with methanol change their response to methanol as adults, even in the absence of larval training.^{38,39} Similarly, parasitic wasps show an oviposition preference for the aphid host in which they themselves develop.⁴⁰ This behavior was found to be the result of exposure to host cues during eclosion and not from a neutrally-derived larval memory as pupae removed from aphids during pupation fail to show any host preference after emerging as wasps.

However, a number of carefully controlled studies demonstrate memory across metamorphosis while controlling for any chemical contamination. Both weevil and wasp species retain preferences for their larval environments, and contrary to other chemical legacy reports, do not show changes in adult behavior if their pupal cases are intentionally contaminated with odors.^{41,42} Perhaps most convincingly, aversive associative memory also appears to survive from larvae to adult; in studies pairing electric shock and ethyl acetate, both *drosophila* and *manduca* larvae learn to avoid the odor in choice assays and maintain this behavior after emerging as adults.^{43,44} Presentation of odor or shock alone during larval stages had no effect on adults of either species, ruling out any possibility of chemical sensitization/habituation.

How might this memory be stored during the transition for larvae to adult? Olfactory memory is known to reside in the mushroom bodies, paired lobes of the larval and adult insect brain which receive input from the antennal lobes.^{45,46} However, during pupation this structure undergoes extensive remodeling, with the γ neurons completely pruning to the main process and the α'/β' neurons demonstrating partial pruning before establishing adult specific projections.^{47,48} While it is possible a true synaptic memory may persist within a subset of mushroom body neurons, only indirect evidence exists to support this hypothesis and more research is necessary to conclusively confirm or reject the idea. Perhaps




Figure 1. Animal models of memories that survive brain remodeling. Holometabolous insects reorganize their brains during pupation in the transition from larva to adult, with many neurons of the central nervous system pruning to the cell body before the generation of adult specific structures. Planarian species are capable of regenerating their entire brain from a tail fragment in the event of fission or amputation, with new tissue arising from a neoblast stem cell population. Arctic ground squirrels demonstrate a drastic reduction in brain volume during hibernation at near freezing temperatures, which is corrected within hours of arousal. In all 3 of these animal groups, learned behaviors have been observed to survive the striking reorganization of the brain. Photos courtesy of Jerry Friedman (Monarch caterpillar), Ianaré Sévi, Linda Mahoney, and Bering Land Bridge National reserve (top, center, and bottom ground squirrel images respectively), used with permission.

the best support to date comes from research using the *Tenebrio* genus of grain beetle, in which larvae trained to navigate mazes show more rapid learning rates as adults compared to untrained siblings.⁴⁹⁻⁵¹ Analysis of total RNA concentration in the brain revealed an enrichment of RNA in the mushroom bodies in the larva which persisted into adulthood.⁵² However this evidence is still indirect and alternative theories for the survival of memory through metamorphosis are myriad. Non-neural memory, chemoreceptor tuning, and gene silencing

have all been suggested but never formally tested. For the few theories that have been tested, little follow-up work exists. The question of trans-metamorphic memory, postulated nearly a century ago, remains unsolved yet ripe for study given the development of molecular tools for model species such as *Tribolium*.^{53,54}

Memories persist during brain regeneration in planaria

Planaria are bilaterian flatworms, possessing complex internal organs, behavioral repertoires, and remarkable powers

of regeneration.⁵⁵⁻⁶² They are a powerful and increasingly popular model for regenerative biology studies, both in terms of understanding stem cell decision-making⁶³⁻⁶⁷ and physiological controls of large-scale pattern formation.⁶⁸⁻⁷¹ Special emphasis has been placed on understanding the mechanisms guiding CNS regeneration in this model system.^{10,72,73}

Neurons in the planarian brain more closely resemble those of vertebrates than those of advanced invertebrates, exhibiting typical vertebrate features of multipolar shape, dendritic spines with synaptic

boutons, a single axon, expression of vertebrate-like neural proteins, and relatively slow spontaneously generated electrical activity.⁷⁴ Therefore, the planarian is not only the first animal to possess a brain, but may be the ancestor of the vertebrate brain.^{61,75} Some of the most interesting and controversial data on the persistence of memories in regeneration thus came from studies of this model.^{76,77}

Influenced by the ideas of D. O. Hebb and J. Eccles, who postulated synaptic plasticity as the mechanism of memory encoding,^{78,79} James McConnell and Robert Thompson explored the implications of the theory in vivo.⁸⁰ While basal, planarians possess a well-differentiated nervous system and CNS with chemical synapses,^{74,81} satisfying both requirements for synaptic plasticity according to Hebb's model.^{80,82} Having first established planaria as a suitable animal model for the studying of associative learning and memory,⁸³ McConnell decided to explore what would happen if conditioned worms were cut in 2 and then tested following regenerating? Which half, if either, would retain the memory? McConnell and his students trained worms using a classical conditioning protocol (paring light with an electric shock) and after the worms demonstrated learning, they cut the worms in half. After fully regenerating some weeks later, the team retrained the regenerated worms using the same protocol. The results showed that head fragment retained the pre-cut training, but also surprisingly, worms that regenerated from the tails (which lost the original brain) required significantly less training trials to learn ("saving" paradigm) compared to untrained animals.^{77,84}

The astonishing results motivated McConnell to examine another theory of memory - the existence of an "engram," as physical/chemical memory trace which could perhaps be transferred into an untrained animal.^{85,86} McConnell and colleagues conducted a series of experiments where they showed that naïve worms that were fed on trained worms revealed evidences of memory retrieval in a "saving" paradigm, where the fed worms learned more quickly than naïve animals.⁸² McConnell and his colleagues fractionated the planarian extracts in an

effort to identify the particular molecule that actively allows memory transfer and based on their findings hypothesized that RNA was the active agent allowing memory transfer.⁸⁶ Further, naïve worms injected with the RNA extract from trained worms demonstrated the "saving" effect, while RNase treatment abolished it.⁸⁰ This finding was extremely surprising since at the time it was not known that RNA has multiple regulatory functions.⁸⁷ These effects were not limited to just one species. A Russian group conditioned another species of planarians' motor responses to shock paired with vibratory stimuli.⁸⁸ When animals were transected, retention could be observed in both halves some 14 days after sectioning. These experiments pointed to the retention of experientially induced modifications in regenerating tissues. The posterior segments of the worms regenerated heads that retained the conditioned responses. Their results supported those of Corning and John's (1961) with RNase, suggesting on the possibility that RNA inheritance could be involved.⁸⁹⁻⁹¹

McConnell and his colleagues performed a number of essential controls for learning specificity. For example, they tested for pseudoconditioning and sensitization effects due to non-training variables such as the dissection and regeneration processes, and demonstrated that the head is the organ which executes the particular learned behavior which was assayed (2 headed worms learn faster and tail fragments without brain cannot be the conditioned^{80,82}). Later, several labs replicated the results (transfer of memory through brain extracts), in rats, mice and fish.⁸⁵ Cowley and Griesel (1966) found that the male grand-offspring of female rats that were prenatally malnourished performed more poorly than controls on the Hebb-Williams maze, despite the fact that their mothers had been on a standard diet from conception through weaning. That is, the effects of a low protein diet lingered across subsequent, well-fed, generations. These findings shifted the field of memory transfer from planaria to rat as the primary model species.⁸⁵ Following a string of studies with rats and mice create, the field became mired

in controversy the entire line of research was abandoned and forgotten.^{85,92}

Given the technical limitations of the time, the mechanisms underlying the results were never fully identified. However, labs are beginning to reinvestigate this fascinating work using modern molecular advances in the understanding of epigenetic processes⁹³⁻⁹⁶ that may allow the transfer of environmentally-imprinted information transgenerationally.^{97,98} In addition, discoveries of RNAi regulation⁸⁷ and accumulating evidence that links inheritance via noncoding RNAs⁹⁹ to memory formation and behavior¹⁰⁰ support the feasibility of the McConnell's findings. Yet it is unclear whether those epigenetic processes have pivotal roles in acquisition of specific memories and behaviors or whether they function as general cofactors for learning and memory. Additional materials on this fascinating controversy are found in.¹⁰¹⁻¹⁰⁹

During the last decade, planaria have re-emerged as a leading model in the field of molecular biology. The genome of a freshwater planarian (*Schmidtea mediterranea*) has been fully sequenced and molecular techniques including gene-specific RNA interference and *in situ* hybridization are fully functional with the species.^{110,111} Today, planaria are perhaps the only molecularly-tractable system in which memory and complete brain regeneration can be studied in the same animal. To re-examine this issue, our lab developed an automated training and testing device designed to overcome some of the limitations of older work.¹¹² Prior efforts performed manually suffered from 1) the time consuming nature of the experiments which only allowed the experimenter to spend a short time each day training any one worm (thus lowering overall N and providing weak memories), 2) the difficulty of reproducing precisely the same protocol across labs (given the worms' sensitivity to even subtle environmental differences), and 3) the challenge of using human observation of behavior to support highly surprising findings (the need to avoid any chance of subjectivity during scoring and to provide a complete behavioral dataset that can be analyzed by others).

Our device provides 24*7 environmental training to each individual animal in parallel, and uses objective criteria for scoring, while recording all the movements for future analysis. Using this platform, our initial study confirmed the ability of worms to recognize a surface etch pattern in a place learning task, and the persistence of this information across head regeneration.¹¹³ While numerous subsequent studies must be done to improve the protocol and ask questions about the mechanism and specificity of the memory, this early result establishes the planarian as a tractable model within which we can next ask questions like: where is the information encoding learned behavior stored? How many and what kind of cells are needed in order to keep the memory? How is it imprinted on the developing new brain? How does encoding and decoding by naïve tissues work? What is the memory capacity of this system? The plethora of molecular tools, cellular-level analytic methods, and automated behavior analyses enables a rich program of investigation using the basic amputation assay to identify the location and main properties of information storage during brain regeneration.

One possible locus for the cellular basis of memory are the planarian neoblasts (stem cells¹¹⁴), which could be modified through epigenetic changes induced by learning.^{22,115,116} When the new brain develops, neoblasts could potentially imprint the CNS through self-organization mechanisms.¹¹⁷⁻¹¹⁹ A second possibility is that non-coding RNAs implement inheritance.^{99,100} Regardless of the molecular mechanisms required for this process, a complete answer to this question will also require an understanding of the mapping of cognitive content to specific molecular states (encoding and decoding of learned information within RNA, protein, cell networks, or some other mechanism). It is clear that modern techniques and recent findings show great potential for the planarian as an animal model in learning and memory research. Investigating this unique animal, which displays complex behavior and can regenerate its entire brain in only a few days, may provide answers to the enigma of acquisition, storage, and retrieval of memories.

Memory and brain repair in mammals

What happens to memories in mammals whose brains incorporate progeny of (perhaps naïve) stem cells? These are crucial issues because transplants of stem cells into brains is a major medical strategy for stroke and degenerative disease.¹²⁰⁻¹²² The answer to this kind of clinical scenario is unknown. Indeed, even the amount of brain tissue needed for specific cognitive functions is not well-understood, given the (rare) cases of hydrocephaly and greatly reduced brain size with normal cognition.^{123,124} One model system that provides an entry point into addressing the biomedical aspects of this question is what occurs during hibernation.¹²⁵

While there are various examples of neural reorganization in vertebrates, perhaps none are as remarkable as that of the European and arctic ground squirrel family. During the winter months these animals go into a state of torpor, where their metabolism and brain activity slows considerably. Direct measurements during this period reveal body temperatures hovering around 0 degrees Celsius and Na^+ K^+ -ATPase muscle activity decreases up to 60 percent.^{126,127} Under these conditions, massive changes occur within the central nervous system of the animals, with extensive pruning of the dendritic trees within the brain,¹²⁸ including areas necessary for long term memory such as the hippocampus.¹²⁹ Neurons of this region demonstrate multiple changes including spine morphology/number, reduced branching as well as shorter dendritic length, and changes in microtubule assembly/disassembly protein abundance.¹²⁸⁻¹³² In addition to the extensive pruning during torpor, perhaps equally impressive is the fact the brain completely reverses this loss during the first few hours of arousal restoring the original branching density present prior to hibernation.^{131,132}

Given this reorganization it has long been questioned whether memory can survive the severe changes in both temperature and dendritic morphology associated with torpor. Early research into this question examined whether hibernating ground squirrels could identify kin and familiar individuals following 9 months of separation. Interestingly, animals

appeared to remember kin but demonstrated no recognition of previously familiar non-kin individuals,¹³³ suggesting that only memories of siblings persist through hibernation (perhaps serving to avoid possible inbreeding the following season). However, further work determined this recognition was not due to true memory, but was in fact a result of self-reference, or similarity of kin to the subject animals own odor.¹³⁴ While these results did not provide evidence of long-term memory surviving torpor, a similar study proved more promising. Groups of ground squirrels were tested in one of 3 tasks; maze navigation, an operant feeding machine, and the ability to discriminate housemates from strangers. While recall performance was weak on both of the associative learning assays, the animals were successfully able to discriminate familiar from unfamiliar individuals.¹³⁵ It is unclear why the animals could demonstrate memory in one assay but not others, though the authors speculate it could be the result of the complexity of the task or the brain region responsible for the memories. Perhaps most convincingly, in more recent work a number of animals were trained in 2 operant conditioning tasks, jumping between a pair of boxes or crossing through a tube, and found that performance in these tasks was unimpaired following 6 months of hibernation.¹³⁶ This result was the clearest to date, and provided strong evidence that some memories could survive the dendritic pruning associated with torpor.

What mechanism may be responsible for memory persistence? While such studies present many challenges (not the least of which is the 6–12 month time periods needed for hibernation and arousal) a number of interesting observations have been reported. Prior to torpor, adipose tissue of the ground squirrels shows a significant increase in antioxidant levels which are predicted to reduce reactive oxygen damage during the severe shift in metabolic activity associated with arousal.²⁶ While these levels were measured in the liver and plasma, it is possible that the same mechanism could protect persistent neurons in the brain from reactive oxygen damage following hibernation. In addition, a number of mitotically active

immature neurons have been identified within the hippocampus of ground squirrels. The purpose of these cells is not currently known, but they may act as a renewable source of new neurons during the rearrangements associated with hibernation (although it is not clear how this cell population may or may not relate to memory).¹³⁷ Perhaps most compelling is recent work which identified changes in the phosphorylation of tau across torpor, a microtubule associated protein known to play a role in neural plasticity. Phosphorylation of tau increased during the onset of torpor and decreased with arousal, suggesting a possible mechanism for stabilizing neural connections required for long term memory.¹³⁸ What makes this finding particularly provocative is the fact that tau protein shows high phosphorylation in patients suffering from Alzheimer's disease, perhaps positioning the ground squirrel as a model to examine the tau signaling pathways with the hope of identifying new targets for therapeutics.

What mechanisms underlie memory?

The question of memory persistence though brain remodeling is an extreme version of a more basic puzzle: how are memories encoded so that they can be reliably decoded within the lifetime of even an un-altered brain? The current paradigm holds that memories are encoded and stored through modification of synaptic connections and the resulting patterns of activation within neuronal networks. This conception is termed "The Synaptic Plasticity Hypothesis."¹³⁹ The hypothesis was supported by discoveries of activity-dependent plasticity mechanisms such as long-term potentiation (LTP),¹⁴⁰ long term depression (LTD),¹⁴¹ and spike-timing-dependent plasticity (STDP),⁷⁸ which are all mechanisms that clearly correlate to learning and memory behaviors.^{139,142,143} However, the synaptic plasticity hypothesis alone has difficulties in explaining the long-life persistency of memories, since the above-mentioned mechanisms are inherently unstable.^{144,145} In addition, the discovery of memory reconsolidation process¹⁴⁶ and the possible requirement for constant activation of protein PKMzeta for memory maintenance¹⁴⁷ argue against

the concept of memory storage as a hard-wired neural network.

Memories can chemically unwire, and are highly dynamic and sensitive to disruption, even long after consolidation. How can memories survive for a lifetime in a dynamic neural-network? It is also hard to explain how memory transfers offline, from the acquisition organ to the storage regions in the brain (in mammals: from the hippocampus to the neocortex), long after the remembered episode occurred, (⁹⁶ see ¹⁴⁸ for a detailed discussion). A similar process of memory transfer in the brain has been found in octopus^{149,150} indicating that this is a fundamental biological mechanism. Finally, recent work showed that memory traces exist even without the complex activation thought to be determined by the specific synaptic weight^{19,151} and even after erasing the synaptic connectivity related to specific memory.²¹

These recent findings are challenging the conservative point of view and require new ideas. It is likely that although memory is encoded through synaptic plasticity and recalled by activation of the neural network, there is a "blueprint" of the memories that may be kept safely in place by a mechanism other than the synaptic connections themselves. Thus, future studies of the models of survival of memory traces through brain regeneration has the potential to reveal the mechanism of long-term memory maintenance.

Memory in aneural systems

It is possible that encoding in the CNS is such that it is robust to massive rearrangements of the underlying substrate. On the other hand, it is possible that additional mechanisms can support robust memory by providing a "backup" storage medium during brain rearrangement. For example, in planaria, it is possible that memory can be stored outside the brain and imprinted on the nascent brain during regeneration. Feedback between behavioral (brain-mediated) and other cellular memory mechanisms has been demonstrated in a number of interesting contexts. For example, injecting frog eggs with various substances (orange or citral) results in animals that prefer to feed on material containing those substances.¹⁵²

Similarly, exposing eggs to a novel odor together with an endogenous alarm chemical results in larvae that produce escape behavior when presented to the odor alone.¹⁵³ In both of those cases, a novel and as yet unknown mechanism must be at play since the original "learning" takes place in very early embryonic stages, long before a CNS exists. For example, from the Hepper and Waldman study, we must conclude that the intracellular milieu of the frog egg has mechanisms for recognizing various molecules and transmitting this information as a stable memory to the nervous system when it is subsequently formed. This kind of functional coupling between intracellular signaling mechanisms and neural network-mediated behavior suggests that much remains to be discovered about the information sources available to the brain during repair and perhaps even in normal cognition. A more detailed treatment of the parallels between information-processing mechanisms in the CNS and morphogenetic signaling is given in.¹⁵⁴

Interestingly, neural-like computation, decision-making, and memory have been reported well beyond the traditional CNS, including sperm,¹⁵⁵ amoebae,¹⁵⁶ yeast,¹⁵⁷ and plants.¹⁵⁸⁻¹⁶⁴ These appear to be mediated by well-conserved, ubiquitous mechanisms that appear to be also involved in neural information processing, such as cytoskeleton¹⁶⁵ and electrical networks.^{166,167} Single somatic cells perform subtraction, addition, low- and band-pass filtering, normalization, gain control, saturation, amplification, multiplication, and thresholding.¹⁶⁸ It is becoming clear that neural networks have no monopoly on such functions, and indeed fascinating examples of memory and neural-like dynamics have been found in the immune system,^{169,170} bone,^{171,172} heart,^{173,174} and physiological disorders such as diabetes.¹⁷⁵

Why the Mind is in the Brain (or is it?)

"Why is the mind in the head? Because there, and only there, are hosts of possible connections to be performed as time and circumstance demand it"¹⁷⁶

If indeed memory and information processing rely on labile connections within a rich network of signaling activity, it becomes immediately clear (counter McCulloch) than the CNS is not the only game in town. Mechanisms that are responsible for cellular computations necessary to rearrange the body plan during remodeling, regeneration, and metamorphosis include the cytoskeleton, metabolic signaling circuits, and the gene regulatory networks. All of these exhibit (physiological) experience-dependent rewiring and rich feedback loops that can store state information. Indeed, all of the major mechanisms by which nerves function – ion channels, neurotransmitters, and electrical synapses not only exist throughout the body but are now known to be functional drivers of many patterning events during regenerative and developmental pattern regulation.¹⁷⁷⁻¹⁸⁰ It is possible that functional linkage between the memory-keeping mechanisms in the CNS and the encoding of target morphologies during pattern regulation is at least in part mediated by the same bioelectric mechanisms.¹⁸¹⁻¹⁸⁴ Because the molecular components of non-neural bioelectric signaling are increasingly well-characterized,¹⁸⁵⁻¹⁸⁸ it is now possible to specifically test the hypothesis that the somatic and cognitive memory systems are coupled during the above-described examples in which memories survive drastic cellular turnover and rearrangement.

Outlook: implications and future work

Convincing data from insects, planaria, and mammals suggest the ability of memories to survive drastic rearrangement and rebuilding of the CNS. Despite the availability of model systems tractable to both behavioral analysis paradigms and molecular genetics of pattern regulation, this area has not received focused attention and remains fertile ground for new investigation. Suggested lines for investigation in such a research program (currently ongoing in our lab) include:

- New theory and quantitative *in silico* analysis in the field of modeling artificial neural networks under topological change, to learn what kinds of

encodings might allow the type of memory robustness observed in some species;

- Wider surveys of different learning paradigms and kinds of memory persisting in brain-regenerative and metamorphosing organisms, to identify novel and perhaps even more impressive examples of memory persisting through brain remodeling;
- Molecular analyses of the mechanisms by which neural networks may exchange information with surrounding tissues (as would be required for the regenerating planarian brain to be imprinted with information by the remaining body fragment). Such studies could test non-neural bioelectrical signaling^{178,189} and cytoskeletal computation^{190,191};
- Testing the idea that CNS-remodeling operations (regenerative pathways) may function in such a way as to specifically preserve encoded information. If true, this would imply a close relationship between the information-processing algorithms that implement pattern regeneration and those that implement memory.¹⁵⁴ One way to approach this hypothesis is to attempt to apply the computational modeling approaches currently used to understand computation in the CNS to the mechanisms regulating pattern formation; existing examples include neural-like models of intracellular signaling pathways¹⁹² and information-centered models of regeneration¹⁹³;
- Establishment of additional assays at the intersection of pattern regulation and cognition, such as for example regional brain transplants,^{12,194-196} and attempts to demonstrate learning in non-neural cellular networks,¹⁷⁵ which would drive the formulation of specific models of how non-brain tissue can support memory during brain regeneration.

This novel interdisciplinary area, at the intersection of behavioral neuroscience and molecular developmental biology, raises unique challenges both in terms of novel theory that needs to be developed and new approaches at the bench. The impact of significant progress in this area

would be huge, in terms of implications for the basic understanding of how mental content is encoded in cellular structures, the design of new regenerative therapies for radical brain repair in medical contexts, and the engineering of biologically-inspired computational media. Thus, several basic and applied areas of science and biomedicine stand to gain from investigations into a crucial and yet still poorly-understood phenomenon: memory, in its behavioral and morphological aspects.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Daniel Lobo, Vaibhav V. Pai, and Jennifer Hammelman for their helpful suggestions on the manuscript. This work was supported by the Templeton World Charity Foundation (TWC0089/AB55) and the G. Harold and Leila Y. Mathers Charitable Foundation.

References

1. Birnbaum KD, Alvarado AS. Slicing across kingdoms: regeneration in plants and animals. *Cell* 2008; 132:697-710; PMID:18295584; <http://dx.doi.org/10.1016/j.cell.2008.01.040>
2. Lobo D, Solano M, Bubenik GA, Levin M. A linear-encoding model explains the variability of the target morphology in regeneration. *J R Soc Interface* 2014; 11:20130918; PMID:24402915; <http://dx.doi.org/10.1098/rsif.2013.0918>
3. McCusker C, Gardiner DM. The axolotl model for regeneration and aging research: a mini-review. *Gerontology* 2011; 57:565-71; PMID:21372551; <http://dx.doi.org/10.1159/000323761>
4. Maden M. Axolotl/newt. *Methods Mol Biol* 2008; 461:467-80; PMID:19030817; http://dx.doi.org/10.1007/978-1-60327-483-8_32
5. Farinella-Ferruzza N. The transformation of a tail into a limb after xenoplastic transformation. *Experientia* 1956; 15:304-5; <http://dx.doi.org/10.1007/BF02159624>
6. Tissot M, Stocker RF. Metamorphosis in drosophila and other insects: the fate of neurons throughout the stages. *Prog Neurobiol* 2000; 62:89-111; PMID:10821983; [http://dx.doi.org/10.1016/S0301-0082\(99\)00069-6](http://dx.doi.org/10.1016/S0301-0082(99)00069-6)
7. Truman JW. Developmental neuroethology of insect metamorphosis. *J Neurobiol* 1992; 23:1404-22; PMID:1487742; <http://dx.doi.org/10.1002/neu.480231005>
8. Denver RJ. The molecular basis of thyroid hormone-dependent central nervous system remodeling during amphibian metamorphosis. *Comp Biochem Physiol C Pharmacol Toxicol Endocrinol* 1998; 119:219-28; PMID:9826995; [http://dx.doi.org/10.1016/S0742-8413\(98\)00011-5](http://dx.doi.org/10.1016/S0742-8413(98)00011-5)

9. Arlotta P, Berninger B. Brains in metamorphosis: reprogramming cell identity within the central nervous system. *Curr Opin Neurobiol* 2014; 27:208-14; PMID:24800935; <http://dx.doi.org/10.1016/j.conb.2014.04.007>
10. Agata K, Umesono Y. Brain regeneration from pluripotent stem cells in planarian. *Philos Trans R Soc Lond B Biol Sci* 2008; 363:2071-8; PMID:18375378; <http://dx.doi.org/10.1098/rstb.2008.2260>
11. Zupanc GK, Zupanc MM. New neurons for the injured brain: mechanisms of neuronal regeneration in adult teleost fish. *Regen Med* 2006; 1:207-16; PMID:17465804; <http://dx.doi.org/10.2217/17460751.1.2.207>
12. Pietsch P, Schneider CW. Brain Transplantation in Salamanders - an Approach to Memory Transfer. *Brain Res* 1969; 14:707-15; PMID:5822440; [http://dx.doi.org/10.1016/0006-8993\(69\)90210-8](http://dx.doi.org/10.1016/0006-8993(69)90210-8)
13. Berg DA, Kirkham M, Beljajeva A, Knapp D, Habermann B, Ryge J, Tanaka EM, Simon A. Efficient regeneration by activation of neurogenesis in homeostatically quiescent regions of the adult vertebrate brain. *Development* 2010; 137:4127-34; PMID:21068061; <http://dx.doi.org/10.1242/dev.055541>
14. Fährbach SE. Structure of the mushroom bodies of the insect brain. *Ann Rev Entomol* 2006; 51:209-32; PMID:16332210; <http://dx.doi.org/10.1146/annurev.ento.51.110104.150954>
15. Truman JW. Metamorphosis of the central nervous system of *Drosophila*. *J Neurobiol* 1990; 21:1072-84; PMID:1979610; <http://dx.doi.org/10.1002/neu.480210711>
16. Lavond DG, Kim JJ, Thompson RF. Mammalian brain substrates of aversive classical conditioning. *Annu Rev Psychol* 1993; 44:317-42; PMID:8434892; <http://dx.doi.org/10.1146/annurev.ps.44.020193.001533>
17. Bruce D. Fifty years since Lashley's In search of the Engram: refutations and conjectures. *J Hist Neurosci* 2001; 10:308-18; PMID:11770197; <http://dx.doi.org/10.1076/jhin.10.3.308.9086>
18. Black IB. Information in the brain : a molecular perspective. Cambridge, Mass.: MIT Press, 1991.
19. Ryan TJ, Roy DS, Pignatelli M, Arons A, Tonegawa S. Memory. Engram cells retain memory under retrograde amnesia. *Science* 2015; 348:1007-13; PMID:26023136; <http://dx.doi.org/10.1126/science.aaa5542>
20. Bailey CH, Kandel ER. Synaptic remodeling, synaptic growth and the storage of long-term memory in *Aplysia*. *Prog Brain Res* 2008; 169:179-98; PMID:18394474; [http://dx.doi.org/10.1016/S0079-6123\(07\)00010-6](http://dx.doi.org/10.1016/S0079-6123(07)00010-6)
21. Chen S, Cai D, Pearce K, Sun PY, Roberts AC, Glanzman DL. Reinstatement of long-term memory following erasure of its behavioral and synaptic expression in *Aplysia*. *Elife* 2014; 3:e03896; PMID:25402831
22. Zovkic IB, Guzman-Karlsson MC, Sweatt JD. Epigenetic regulation of memory formation and maintenance. *Learn Mem* 2013; 20:61-74; PMID:23322554; <http://dx.doi.org/10.1101/lm.026575.112>
23. Chicha L, Smith T, Guzman R. Stem cells for brain repair in neonatal hypoxia-ischemia. *Childs Nerv Syst* 2014; 30:37-46; PMID:24178233; <http://dx.doi.org/10.1007/s00381-013-2304-4>
24. Gage FH, Temple S. Neural stem cells: generating and regenerating the brain. *Neuron* 2013; 80:588-601; PMID:24183012; <http://dx.doi.org/10.1016/j.neuron.2013.10.037>
25. Porto-Pazos AB, Veiguela N, Mesejo P, Navarrete M, Alvarellos A, Ibanez O, Pazos A, Araque A. Artificial astrocytes improve neural network performance. *PloS one* 2011; 6:e19109; PMID:21526157; <http://dx.doi.org/10.1371/journal.pone.0019109>
26. Goldman SA, Nedergaard M, Windrem MS. Modeling cognition and disease using human glial chimeric mice. *Glia* 2015; 63:1483-93; PMID:26010831; <http://dx.doi.org/10.1002/glia.22862>
27. Kamm RD, Bashir R. Creating living cellular machines. *Ann Biomed Eng* 2014; 42:445-59; PMID:24006130; <http://dx.doi.org/10.1007/s10439-013-0902-7>
28. Doursat R, Sanchez C. Growing fine-grained multicellular robots. *Soft Robotics* 2014; 1:110-21; <http://dx.doi.org/10.1089/soro.2014.0014>
29. Doursat R, Sayama H, Michel O. A review of morphogenetic engineering. *Nat Comput* 2013; 12:517-35; <http://dx.doi.org/10.1007/s11047-013-9398-1>
30. Bull L, Budd A, Stone C, Urokov I, de Lacy Costello B, Adamatzky A. Towards unconventional computing through simulated evolution: control of nonlinear media by a learning classifier system. *Artif Life* 2008; 14:203-22; PMID:18331191; <http://dx.doi.org/10.1162/artl.2008.14.2.203>
31. Adamatzky A, Stepney S. Towards theory of unconventional computing. *Int J Unconv Comput* 2008; 4:1-11
32. Hopkins AD. Economic investigations of the scolytid bark and timber beetles of North America. US Department of Agriculture Program of Work for 1917 1916:353
33. Akhtar Y, Isman MB. Larval exposure to oviposition deterrents alters subsequent oviposition behavior in generalist, *Trichoplusia ni* and specialist, *Plutella xylostella* moths. *J Chem Ecol* 2003; 29:1853-70; PMID:12956511; <http://dx.doi.org/10.1023/A:1024802328458>
34. Chow JK, Akhtar Y, Isman MB. The effects of larval experience with a complex plant latex on subsequent feeding and oviposition by the cabbage looper moth: *Trichoplusia ni* (Lepidoptera: Noctuidae). *Chemoecology* 2005; 15:129-33; <http://dx.doi.org/10.1007/s00049-005-0304-x>
35. Olsson POC, Anderbrant C, Lofstedt C. Experience influences oviposition behaviour in two pyralid moths, *Ephesia cautella* and *Plodia interpunctella*. *Animal Behav* 2006; 72:545-51; <http://dx.doi.org/10.1016/j.anbehav.2005.10.023>
36. Rojars JC, Wyatt TD. The role of pre- and post-imaginal experience in the host-finding and oviposition behaviour of the cabbage moth. *Physiol Entomol* 1999; 24:83-9; <http://dx.doi.org/10.1046/j.1365-3032.1999.00117.x>
37. Corbet SA. Insect chemosensory responses - a chemical legacy hypothesis. *Ecolog Entomol* 1985; 10:143-53; <http://dx.doi.org/10.1111/j.1365-2311.1985.tb00543.x>
38. Barron AB. The life and death of Hopkins' host-selection principle. *J Insect Behav* 2001; 14:725-37; <http://dx.doi.org/10.1023/A:101303332535>
39. Barron AB, Corbet SA. Preimaginal conditioning in *Drosophila* revisited. *Animal Behav* 1999; 58:621-8; PMID:10479377; <http://dx.doi.org/10.1006/anbe.1999.1169>
40. van Emden HF, Sponagl B, Baker T, Ganguly S, Dou-loupaka S. Hopkins 'host selection principle', another nail in its coffin. *Physiol Entomol* 1996; 21:325-8; <http://dx.doi.org/10.1111/j.1365-3032.1996.tb00873.x>
41. Gandolfi M, Mattiacci L, Dorn S. Preimaginal learning determines adult response to chemical stimuli in a parasitic wasp. *Proc Biol Sci* 2003; 270:2623-9; <http://dx.doi.org/10.1098/rspb.2003.2541>
42. Rietdorf K, Steidle JLM. Was Hopkins right? Influence of larval and early adult experience on the olfactory response in the granary weevil *Sitophilus granarius* (Coleoptera, Curculionidae). *Physiol Entomol* 2002; 27:223-7; <http://dx.doi.org/10.1046/j.1365-3032.2002.00289.x>
43. Tully T, Cambiazo V, Kruse L. Memory through metamorphosis in normal and mutant *Drosophila*. *J Neurosci* 1994; 14:68-74; PMID:8283252
44. Blackston DJ, Silva Casey E, Weiss MR. Retention of memory through metamorphosis: can a moth remember what it learned as a caterpillar? *PLoS One* 2008; 3:e1736; PMID:18320055; <http://dx.doi.org/10.1371/journal.pone.0001736>
45. Debelles JS, Heisenberg M. Associative odor learning in *Drosophila* abolished by chemical ablation of mushroom bodies. *Science* 1994; 263:692-5; PMID:8303280; <http://dx.doi.org/10.1126/science.8303280>
46. Wolf R, Wittig T, Liu L, Wustmann G, Eyding D, Heisenberg M. *Drosophila* mushroom bodies are dispensable for visual, tactile, and motor learning. *Learning & Memory* 1998; 5:166-78; PMID:10454381
47. Lee T, Lee A, Luo LQ. Development of the *Drosophila* mushroom bodies: sequential generation of three distinct types of neurons from a neuroblast. *Development* 1999; 126:4065-76; PMID:10457015
48. Marin EC, Watts RJ, Tanaka NK, Ito K, Luo LQ. Developmentally programmed remodeling of the *Drosophila* olfactory circuit. *Development* 2005; 132:725-37; PMID:15659487; <http://dx.doi.org/10.1242/dev.01614>
49. Sheiman IM, Balabanova EF, Martinovich VP, Polikarpova VP, Slododchikova LK. Effects of Luliberin and Its Fragments on the Memory of the Beetle *Tenebrio-Molitor*. *J Evol Biochem Phys+* 1984; 20:252-6
50. Sheiman IM. Training of *Tenebrio Molitor* Larvae at Various Stages of Postembryonal Development. *Zhurnal Obshchei Biol* 1973; 34:470-5
51. Alloway T. Retention of learning through metamorphosis in the grain beetle (*Tenebrio molitor*). *Am Zoologist* 1971; 12:471-7
52. Punzo F, Malatesta RJ. Brain-Rna Synthesis and the Retention of Learning through Metamorphosis in *Tenebrio-Obscurus* (Insecta, Coleoptera). *Comp Biochem Phys A* 1988; 91:675-8; [http://dx.doi.org/10.1016/0300-9629\(88\)90947-4](http://dx.doi.org/10.1016/0300-9629(88)90947-4)
53. Bonneton F. ; When *Tribolium* complements the genetics of *Drosophila*. *Med Sci (Paris)* 2010; 26:297-303; PMID:20346280; <http://dx.doi.org/10.1051/medsci/2010263297>
54. Peel AD. The evolution of developmental gene networks: lessons from comparative studies on holometabolous insects. *Philos Trans R Soc Lond B Biol Sci* 2008; 363:1539-47; PMID:18192180; <http://dx.doi.org/10.1098/rstb.2007.2244>
55. Reddien PW, Sanchez Alvarado A. Fundamentals of planarian regeneration. *Annu Rev Cell Dev Biol* 2004; 20:725-57; PMID:15473858; <http://dx.doi.org/10.1146/annurev.cellbio.20.010403.095114>
56. Slack JM. Regeneration research today. *Dev Dyn* 2003; 226:162-6; PMID:12557195; <http://dx.doi.org/10.1002/dvdy.10232>
57. Sanchez Alvarado A. The freshwater planarian *Schmidtea mediterranea*: embryogenesis, stem cells and regeneration. *Curr Opin Genet Dev* 2003; 13:438-44; PMID:12888018; [http://dx.doi.org/10.1016/S0959-437X\(03\)00082-0](http://dx.doi.org/10.1016/S0959-437X(03)00082-0)
58. Salo E, Baguna J. Regeneration in planarians and other worms: New findings, new tools, and new perspectives. *J Exp Zool* 2002; 292:528-39; PMID:12115936; <http://dx.doi.org/10.1002/jez.90001>
59. Lobo D, Beane WS, Levin M. Modeling planarian regeneration: a primer for reverse-engineering the worm. *PLoS Comput Biol* 2012; 8:e1002481; PMID:22570595; <http://dx.doi.org/10.1371/journal.pcbi.1002481>
60. Nicolas C, Abramson C, Levin M. Analysis of behavior in the planarian model. In: Raffa R, Rawls S, eds. *Planaria: A Model for Drug Action and Abuse*. Austin: RG Landes Co, 2008:83-94.
61. Buttarelli FR, Pellicano C, Pontieri FE. Neuropharmacology and behavior in planarians: translations to mammals. *Comp Biochem Physiol C Toxicol Pharmacol* 2008; 147:399-408; PMID:18294919; <http://dx.doi.org/10.1016/j.cbpc.2008.01.009>
62. Brown F, Park Y. Seasonal variations in sign and strength of gamma-taxis in planarians. *Nature* 1964;

- 202:469-71; PMID:14167828; <http://dx.doi.org/10.1038/202469a0>
63. Handberg-Thorsager M, Fernandez E, Salo E. Stem cells and regeneration in planarians. *Front Biosci* 2008; 13:6374-94; PMID:18508666; <http://dx.doi.org/10.2741/3160>
64. Forsthoefel DJ, Park AE, Newmark PA. Stem cell-based growth, regeneration, and remodeling of the planarian intestine. *Dev Biol* 2011; 356:445-59; PMID:21664348; <http://dx.doi.org/10.1016/j.ydbio.2011.05.669>
65. Wagner DE, Wang IE, Reddien PW. Clonogenic neoblasts are pluripotent adult stem cells that underlie planarian regeneration. *Science* 2011; 332:811-6; PMID:21566185; <http://dx.doi.org/10.1126/science.1203983>
66. Reddien PW, Oviedo NJ, Jennings JR, Jenkin JC, Sanchez Alvarado A. SMEDWI-2 is a PIWI-like protein that regulates planarian stem cells. *Science* 2005; 310:1327-30; PMID:16311336; <http://dx.doi.org/10.1126/science.1116110>
67. Gentile L, Cebria F, Bartscherer K. The planarian flatworm: an in vivo model for stem cell biology and nervous system regeneration. *Dis Model Mech* 2011; 4:12-9; PMID:21135057; <http://dx.doi.org/10.1242/dmm.006692>
68. Beane WS, Morokuma J, Adams DS, Levin M. A Chemical genetics approach reveals H,K-ATPase-mediated membrane voltage is required for planarian head regeneration. *Chem Biol* 2011; 18:77-89; PMID:21276941; <http://dx.doi.org/10.1016/j.chembiol.2010.11.012>
69. Oviedo NJ, Morokuma J, Walentek P, Kema IP, Gu MB, Ahn JM, Hwang JS, Gojbori T, Levin M. Long-range neural and gap junction protein-mediated cues control polarity during planarian regeneration. *Dev Biol* 2010; 339:188-99; PMID:20026026; <http://dx.doi.org/10.1016/j.ydbio.2009.12.012>
70. Beane WS, Morokuma J, Lemire JM, Levin M. Bioelectric signaling regulates head and organ size during planarian regeneration. *Development* 2013; 140:313-22; PMID:23250205; <http://dx.doi.org/10.1242/dev.086900>
71. Lobo D, Levin M. Inferring regulatory networks from experimental morphological phenotypes: a computational method reverse-engineers planarian regeneration. *PLoS Comput Biol* 2015; 11:e1004295; PMID:26042810; <http://dx.doi.org/10.1371/journal.pcbi.1004295>
72. Zhang YF, Ye BP, Wang DY. Molecular actions guiding neural regeneration in planarian. *Neurosci Bull* 2008; 24:329-37; PMID:18839027; <http://dx.doi.org/10.1007/s12264-008-0610-8>
73. Umesono Y, Agata K. Evolution and regeneration of the planarian central nervous system. *Dev Growth Differ* 2009; 51:185-95; PMID:19379275; <http://dx.doi.org/10.1111/j.1440-169X.2009.01099.x>
74. Sarnat HB, Netsky MG. The brain of the planarian as the ancestor of the human brain. *Can J Neurolog Sci* 1985; 12(4):296-302; PMID:4084864
75. Pagán OR. The first brain : the neuroscience of planarians. 2014.
76. Brown HM, Beck EC. Does Learning in Planaria Survive Regeneration. *Federation Proc* 1964; 23:254-
77. McConnell JV, Jacobson AL, Kimble DP. The effects of regeneration upon retention of a conditioned response in the planarian. *J Comparat Physiol Psychol* 1959; 52:1-5; <http://dx.doi.org/10.1037/h0048028>
78. Markram H, Gerstner W, Sjöström PJ. A history of spike-timing-dependent plasticity. *Front Synaptic Neurosci* 2011; 3:4; PMID:22007168; <http://dx.doi.org/10.3389/fnsyn.2011.00004>
79. Hebb DO. The organization of behavior; a neuropsychological theory. New York; Wiley, 1949.
80. McConnell JV. A Manual of psychological experimentation on planarians. *Ann Arbor, Mich.*, 1965.
81. Sarnat HB, Netsky MG. When does a ganglion become a brain? Evolutionary origin of the central nervous system. *Semin Pediatr Neurol* 2002; 9:240-53; PMID:12523550; <http://dx.doi.org/10.1053/spen.2002.32502>
82. McConnell JV. Memory transfer through cannibalism in planarium. *J Neuropsychiat* 1962; 3 suppl 1:542-8
83. Thompson R, McConnell JV. Classical conditioning in the planarian, *Dugesia dorotocephala*. *J Comp Physiol Psychol* 1955; 48:65-8; PMID:14354075; <http://dx.doi.org/10.1037/h0041147>
84. Best J. Protropsychology. *Sci Am* 1963; 208:54-62 62-; PMID:13967782
85. Setlow B. Georges Ungar and memory transfer. *Journal of the history of the neurosciences* 1997; 6:181-92; PMID:11619520; <http://dx.doi.org/10.1080/09647049709525701>
86. Morange M. What history tells us VI. The transfer of behaviours by macromolecules. *J Biosci* 2006; 31:323-7; PMID:17006014; <http://dx.doi.org/10.1007/BF02704104>
87. Smalheiser NR, Manev H, Costa E. RNAi and brain function: was McConnell on the right track? *Trends Neurosci* 2001; 24:216-8; PMID:11250005; [http://dx.doi.org/10.1016/S0166-2236\(00\)01739-2](http://dx.doi.org/10.1016/S0166-2236(00)01739-2)
88. Cherkashin AN, Sheiman IM. Conditioning in planarians and RNA content. *J Biol Psychol* 1967; 9:5-11
89. Cherkashin AN, Sheiman IM, Bogorovskaya GI. Uslonivnye reflexi y planarii i opiti s regeneratsii. *Zhurnal Vysshei Nervnoi Deiatelnosti Imeni I P Pavlova* 1966; XVI:1110-5.
90. Cherkashin AN, Sheiman IM. Effect Produced by Ribonucleose on Nervous Activity of Planaria. *Dokl Akad Nauk SSSR* 1966; 171:996-8; PMID:5994947
91. Cherkashin AN, Sheiman IM. O vliianii ribonukleazy na nervnuu deiatel'nost' planarii. *Dokl Akad Nauk SSSR* 1966; 171:996-8; PMID:5994947
92. Rilling M. The mystery of the vanished citations: James McConnell's forgotten 1960s quest for planarian learning, a biochemical engram, and celebrity. *American Psychologist* 1996; 51:589-98; <http://dx.doi.org/10.1037/0003-066X.51.6.589>
93. Ginsburg S, Jablonka E. Epigenetic learning in non-neural organisms. *J Biosci* 2009; 34:633-46; PMID:19920348; <http://dx.doi.org/10.1007/s12038-009-0081-8>
94. Levenson JM, Sweatt JD. Epigenetic mechanisms in memory formation. *Nat Rev Neurosci* 2005; 6:108-18; PMID:15654323; <http://dx.doi.org/10.1038/nrn1604>
95. Day JJ, Sweatt JD. DNA methylation and memory formation. *Nat Neurosci* 2010; 13:1319-23; PMID:20975755; <http://dx.doi.org/10.1038/nn.2666>
96. Arshavsky YI. "The seven sins" of the Hebbian synapse: can the hypothesis of synaptic plasticity explain long-term memory consolidation? *Prog Neurobiol* 2006; 80:99-113; PMID:17074430; <http://dx.doi.org/10.1016/j.pneurobio.2006.09.004>
97. Dias BG, Ressler KJ. Parental olfactory experience influences behavior and neural structure in subsequent generations. *Nat Neurosci* 2014; 17:89-96; PMID:24292232; <http://dx.doi.org/10.1038/nn.3594>
98. Fehér O, Wang H, Saar S, Mitra PP, Tchernichovski O. De novo establishment of wild-type song culture in the zebra finch. *Nature* 2009; 459:564-8; <http://dx.doi.org/10.1038/nature07994>
99. Rechavi O, Minevich G, Hobert O. Transgenerational inheritance of an acquired small RNA-based antiviral response in *C. elegans*. *Cell* 2011; 147:1248-56; PMID:22119442; <http://dx.doi.org/10.1016/j.cell.2011.10.042>
100. Mercer TR, Dinger ME, Mariani J, Kosik KS, Mehler MF, Mattick JS. Noncoding RNAs in Long-Term Memory Formation. *Neuroscientist* 2008; 14:434-45; PMID:18997122; <http://dx.doi.org/10.1177/1073858408319187>
101. Harper LV. Epigenetic inheritance and the intergenerational transfer of experience. *Psychological bulletin* 2005; 131:340-60; PMID:15869332; <http://dx.doi.org/10.1037/0033-2909.131.3.340>
102. Ungar G. Is there a chemical memory trace. *Israel J Chem* 1975; 14:169-76; <http://dx.doi.org/10.1002/ijch.197500057>
103. Ungar G. Molecular coding of memory. *Life Sciences* 1974; 14:595-604; PMID:4595997; [http://dx.doi.org/10.1016/0024-3205\(74\)90394-4](http://dx.doi.org/10.1016/0024-3205(74)90394-4)
104. Ungar G. Peptides and memory. *Biochem Pharmacol* 1974; 23:1553-8; PMID:4603211; [http://dx.doi.org/10.1016/0006-2952\(74\)90366-9](http://dx.doi.org/10.1016/0006-2952(74)90366-9)
105. Ungar G. Molecular code in memory. *Recherche* 1972; 3:19-8
106. Ungar G. Molecular coding of information in nervous-system. *Naturwissenschaften* 1972; 59:85-8; PMID:5019284; <http://dx.doi.org/10.1007/BF00591779>
107. McConnell JV, Shelby JM. Memory transfer experiments in invertebrates. In: Ungar G, ed. *Molecular mechanisms in memory and learning*. New York: Plenum Press, 1970:71-101.
108. Ungar G. Molecular mechanisms in learning. *Perspect Biol Med* 1968; 11:217-8; PMID:4296038; <http://dx.doi.org/10.1353/pbm.1968.0062>
109. Ungar G. Chemical transfer of learning - its stimulus specificity. *Federation Proceedings* 1966; 25:207-8
110. Elliott SA, Sánchez Alvarado A. The history and enduring contributions of planarians to the study of animal regeneration. *Wiley Interdiscip Rev Dev Biol* 2013; 2:301-26.
111. Newmark P, Sánchez Alvarado A. Not your father's planarian: a classic model enters the era of functional genomics. *Nat Rev Genet* 2002; 3:210-9; PMID:11972158; <http://dx.doi.org/10.1038/nrg759>
112. Blackiston D, Shomrat T, Nicolas CL, Granata C, Levin M. A second-generation device for automated training and quantitative behavior analyses of molecularly-tractable model organisms. *PLoS One* 2010; 5:e14370; PMID:21179424; <http://dx.doi.org/10.1371/journal.pone.0014370>
113. Shomrat T, Levin M. An automated training paradigm reveals long-term memory in planarians and its persistence through head regeneration. *J Exp Biol* 2013; 216:3799-810; PMID:23821717; <http://dx.doi.org/10.1242/jeb.087809>
114. Baguña J. The planarian neoblast: the rambling history of its origin and some current black boxes. *Int J Dev Biol* 2012; 56:19-37; <http://dx.doi.org/10.1387/ijdb.113463jb>
115. Day JJ, Sweatt JD. Cognitive neuroepigenetics: a role for epigenetic mechanisms in learning and memory. *Neurobiol Learn Mem* 2011; 96:2-12; PMID:21195202; <http://dx.doi.org/10.1016/j.nlm.2010.12.008>
116. Day JJ, Sweatt JD. Epigenetic mechanisms in cognition. *Neuron* 2011; 70:813-29; PMID:21658577; <http://dx.doi.org/10.1016/j.neuron.2011.05.019>
117. Harper LV. Epigenetic inheritance and the intergenerational transfer of experience. *Psychol Bull* 2005; 131:340-60; PMID:15869332; <http://dx.doi.org/10.1037/0033-2909.131.3.340>
118. Zovkic IB, Guzman-Karlsson MC, Sweatt JD. Epigenetic regulation of memory formation and maintenance. *Learn Mem* 2013; 20:61-74; PMID:23322554; <http://dx.doi.org/10.1101/lm.026575.112>
119. Lancaster MA, Renner M, Martin CA, Wenzel D, Bicknell LS, Hurler ME, Homfray T, Penninger JM, Jackson AP, Knoblich JA. Cerebral organoids model human brain development and microcephaly. *Nature* 2013; 501:373-9; PMID:23995685; <http://dx.doi.org/10.1038/nature12517>
120. Gao J, Prough DS, McAdoo DJ, Grady JJ, Parsley MO, Ma L, Tarenenko YI, Wu P. Transplantation of primed human fetal neural stem cells improves cognitive function in rats after traumatic brain injury. *Experimental neurology* 2006; 201:281-92; PMID:16904107; <http://dx.doi.org/10.1016/j.expneurol.2006.04.039>

121. Hoane MR, Becerra GD, Shank JE, Tatko L, Pak ES, Smith M, Murashov AK. Transplantation of neuronal and glial precursors dramatically improves sensorimotor function but not cognitive function in the traumatically injured brain. *J Neurotrauma* 2004; 21:163-74; PMID:15000757; <http://dx.doi.org/10.1089/089771504322778622>
122. Darsalia V, Allison SJ, Cusulin C, Monni E, Kuzdas D, Kallur T, Lindvall O, Kokaia Z. Cell number and timing of transplantation determine survival of human neural stem cell grafts in stroke-damaged rat brain. *J Cereb Blood Flow Metab* 2011; 31:235-42; PMID:20531461; <http://dx.doi.org/10.1038/jcbfm.2010.81>
123. Lorber J. Is your brain really necessary? *Nurs Mirror* 1981; 152:29-30; PMID:6909910
124. Lorber J. Is your brain really necessary. *Arch Dis Child* 1978; 53:834
125. Heller HC. Hibernation: neural aspects. *Annu Rev Physiol* 1979; 41:305-21; PMID:373593; <http://dx.doi.org/10.1146/annurev.ph.41.030179.001513>
126. Hut RA, Barnes BM, Daan S. Body temperature patterns before, during, and after semi-natural hibernation in the European ground squirrel. *J Comp Physiol B* 2002; 172:47-58; PMID:11824403; <http://dx.doi.org/10.1007/s003600100226>
127. MacDonald JA, Storey KB. Regulation of ground squirrel Na⁺K⁺-ATPase activity by reversible phosphorylation during hibernation. *Biochem Biophys Res Commun* 1999; 254:424-9; PMID:9918854; <http://dx.doi.org/10.1006/bbrc.1998.9960>
128. Hindle AG, Martin SL. Cytoskeletal regulation dominates temperature-sensitive proteomic changes of hibernation in forebrain of 13-lined ground squirrels. *PLoS One* 2013; 8:e71627; PMID:23951209; <http://dx.doi.org/10.1371/journal.pone.0071627>
129. von der Ohe CG, Darian-Smith C, Garner CC, Heller HC. Ubiquitous and temperature-dependent neural plasticity in hibernators. *J Neurosci* 2006; 26:10590-8; PMID:17035545; <http://dx.doi.org/10.1523/JNEUROSCI.2874-06.2006>
130. Magarinos AM, McEwen BS, Saboureaux M, Pevet P. Rapid and reversible changes in intrahippocampal connectivity during the course of hibernation in European hamsters. *Proc Natl Acad Sci U S A* 2006; 103:18775-80; PMID:17121986; <http://dx.doi.org/10.1073/pnas.0608785103>
131. Popov VI, Bocharova LS, Bragin AG. Repeated changes of dendritic morphology in the hippocampus of ground squirrels in the course of hibernation. *Neuroscience* 1992; 48:45-51; PMID:1584424; [http://dx.doi.org/10.1016/0306-4522\(92\)90336-Z](http://dx.doi.org/10.1016/0306-4522(92)90336-Z)
132. Popov VI, Medvedev NI, Patrusev IV, Ignat'ev DA, Morenkov ED, Stewart MG. Reversible reduction in dendritic spines in CA1 of rat and ground squirrel subjected to hypothermia-normothermia in vivo: A three-dimensional electron microscope study. *Neuroscience* 2007; 149:549-60; PMID:17919827; <http://dx.doi.org/10.1016/j.neuroscience.2007.07.059>
133. Mateo JM, Johnston RE. Retention of social recognition after hibernation in Belding's ground squirrels. *Anim Behav* 2000; 59:491-9; PMID:10715170; <http://dx.doi.org/10.1006/anbe.1999.1363>
134. Mateo JM. Self-referent phenotype matching and long-term maintenance of kin recognition. *Animal Behav* 2010; 80:929-35; <http://dx.doi.org/10.1016/j.anbehav.2010.08.019>
135. Millesi E, Prossinger H, Dittami JP, Fiedler M. Hibernation effects on memory in European ground squirrels (*Spermophilus citellus*). *J Biol Rhythms* 2001; 16:264-71; PMID:11407786; <http://dx.doi.org/10.1177/074873001129001971>
136. Clemens LE, Heldmaier G, Exner C. Keep cool: Memory is retained during hibernation in Alpine marmots. *Physiol Behav* 2009; 98:78-84; PMID:19393672; <http://dx.doi.org/10.1016/j.physbeh.2009.04.013>
137. Popov VI, Kraev IV, Ignat'ev DA, Stewart MG. Suspension of mitotic activity in dentate gyrus of the hibernating ground squirrel. *Neural Plast* 2011; 2011:867525; PMID:21773054; <http://dx.doi.org/10.1155/2011/867525>
138. Stieler JT, Bullmann T, Kohl F, Toien O, Bruckner MK, Hartig W, Barnes BM, Arendt T. The Physiological Link between Metabolic Rate Depression and Tau Phosphorylation in Mammalian Hibernation. *PLoS One* 2011; 6:e14530
139. Martin SJ, Grimwood PD, Morris RGM. Synaptic plasticity and memory: An evaluation of the hypothesis. *Annual Review Of Neuroscience* 2000; 23:649-711; PMID:10845078; <http://dx.doi.org/10.1146/annurev.neuro.23.1.649>
140. Bliss T, Lomo T. Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J Physiol* 1973; 232:331-56; PMID:4727084; <http://dx.doi.org/10.1113/jphysiol.1973.sp010273>
141. Massey PV, Bashir ZI. Long-term depression: multiple forms and implications for brain function. *Trends Neurosci* 2007; 30:176-84; PMID:17335914; <http://dx.doi.org/10.1016/j.tins.2007.02.005>
142. Liu X, Ramirez S, Pang PT, Puryear CB, Govindarajan A, Deisseroth K, Tonegawa S. Optogenetic stimulation of a hippocampal engram activates fear memory recall. *Nature* 2012; 484:381-5; PMID:22441246; <http://dx.doi.org/10.1038/484410a>
143. Kasai H, Fukuda M, Watanabe S, Hayashi-Takagi A, Noguchi J. Structural dynamics of dendritic spines in memory and cognition. *Trends Neurosci* 2010; 33:121-9; PMID:20138375; <http://dx.doi.org/10.1016/j.tins.2010.01.001>
144. Cohen LD, Zuchman R, Sorokina O, Müller A, Dietrich DC, Armstrong JD, Ziv T, Ziv NE. Metabolic turnover of synaptic proteins: kinetics, interdependencies and implications for synaptic maintenance. *PLoS One* 2013; 8:e63191; PMID:23658807; <http://dx.doi.org/10.1371/journal.pone.0063191>
145. Ziv NE, Ahissar E. Neuroscience: New tricks and old spines. *Nature* 2009; 462:859-61; PMID:20016588; <http://dx.doi.org/10.1038/462859a>
146. Nader K, Einarsson EO. Memory reconsolidation: an update. *Ann N Y Acad Sci* 2010; 1191:27-41; PMID:20392274; <http://dx.doi.org/10.1111/j.1749-6632.2010.05443.x>
147. Kwapis JL, Helmstetter FJ. Does PKM(zeta) maintain memory? *Brain Res Bull* 2014; 105:36-45; PMID:24076105; <http://dx.doi.org/10.1016/j.brainresbull.2013.09.005>
148. Frankland PW, Bontempi B. The organization of recent and remote memories. *Nat Rev Neurosci* 2005; 6:119-30; PMID:15685217; <http://dx.doi.org/10.1038/nrn1607>
149. Muntz WRA, Sutherland NS, Young JZ. Simultaneous Shape Discrimination in Octopus after Removal of the Vertical Lobe. *J Exp Biol* 1962; 39:557-66; PMID:13936679
150. Muntz WRA. Intraocular transfer and the function of the optic lobes in Octopus. *Q J Exp Psychol* 1963; 15:116-24; <http://dx.doi.org/10.1080/17470216308416562>
151. Barbour B, Brunel N, Hakim V, Nadal JP. What can we learn from synaptic weight distributions? *Trends Neurosci* 2007; 30:622-9; PMID:17983670; <http://dx.doi.org/10.1016/j.tins.2007.09.005>
152. Hepper PG, Waldman B. Embryonic olfactory learning in frogs. *Q J Exp Psychol* 1992; 44:179-97; PMID:1598418
153. Mathis A, Ferrari MC, Windel N, Messier F, Chivers DP. Learning by embryos and the ghost of predation future. *Proc Biol Sci* 2008; 275:2603-7; PMID:18682368; <http://dx.doi.org/10.1098/rspb.2008.0754>
154. Levin M, Pezzulo G. Re-Membering the Body: applications of computational neuroscience to the top-down control of regeneration of limbs and other complex organs. *Ann Rev Biomed Engineering* 2015; 17:1(1)
155. Alvarez L, Friedrich BM, Gompfer G, Kaupp UB. The computational sperm cell. *Trends Cell Biol* 2013; 198-207; PMID:24342435
156. Zhu L, Aono M, Kim SJ, Hara M. Amoeba-based computing for traveling salesman problem: Long-term correlations between spatially separated individual cells of *Physarum polycephalum*. *Bio Systems* 2013; 112:1-10; PMID:23438635; <http://dx.doi.org/10.1016/j.biosystems.2013.01.008>
157. Caudron F, Barral Y. A super-assembly of Whi3 encodes memory of deceptive encounters by single cells during yeast courtship. *Cell* 2013; 155:1244-57; PMID:24315096; <http://dx.doi.org/10.1016/j.cell.2013.10.046>
158. Gagliano M, Renton M, Depczynski M, Mancuso S. Experience teaches plants to learn faster and forget slower in environments where it matters. *Oecologia* 2014; 175:63-72; PMID:24390479; <http://dx.doi.org/10.1007/s00442-013-2873-7>
159. Gremiaux A, Yokawa K, Mancuso S, Baluska F. Plant anesthesia supports similarities between animals and plants: Claude Bernard's forgotten studies. *Plant Signal Behav* 2014; 9:e27886; PMID:24476640; <http://dx.doi.org/10.4161/psb.27886>
160. Scialdone A, Mugford ST, Feike D, Skeffington A, Borrill P, Graf A, Smith AM, Howard M. Arabidopsis plants perform arithmetic division to prevent starvation at night. *Elife* 2013; 2:e00669; PMID:23805380; <http://dx.doi.org/10.7554/eLife.00669>
161. Trewavas A. How plants learn. *Proc Natl Acad Sci U S A* 1999; 96:4216-8; PMID:10200239; <http://dx.doi.org/10.1073/pnas.96.8.4216>
162. Trewavas A. Aspects of plant intelligence. *Ann Bot (Lond)* 2003; 92:1-20; <http://dx.doi.org/10.1093/aob/mcg101>
163. Bose I, Karmakar R. Simple models of plant learning and memory. *Physica Scripta* 2003; T106:9-12; <http://dx.doi.org/10.1238/Physica.Topical.106a00009>
164. Masi E, Ciszak M, Stefano G, Renna L, Azzarello E, Pandolfi C, Mugnai S, Baluska F, Arecchi FT, Mancuso S. Spatiotemporal dynamics of the electrical network activity in the root apex. *Proc Natl Acad Sci U S A* 2009; 106:4048-53; PMID:19234119; <http://dx.doi.org/10.1073/pnas.0804640106>
165. Sahu S, Ghosh S, Hirata K, Fujita D, Bandyopadhyay A. Multi-level memory-switching properties of a single brain microtubule. *Appl Phys Lett* 2013; 102; <http://dx.doi.org/10.1063/1.4793995>
166. Volkov AG, Carrell H, Adesina T, Markin VS, Jovanov E. Plant electrical memory. *Plant Signal Behav* 2008; 3:490-2; PMID:19704496; <http://dx.doi.org/10.4161/psb.3.7.5684>
167. Inoue J. A simple Hopfield-like cellular network model of plant intelligence. *Prog Brain Res* 2008; 168:169-74; PMID:18166394; [http://dx.doi.org/10.1016/S0079-6123\(07\)68014-5](http://dx.doi.org/10.1016/S0079-6123(07)68014-5)
168. Koch C, Segev I. The role of single neurons in information processing. *Nat Neurosci* 2000; 3 Suppl:1171-7; PMID:11127834; <http://dx.doi.org/10.1038/81444>
169. Cohen IR. The cognitive principle challenges clonal selection. *Immunol Today* 1992; 13:441-4; PMID:1476598; [http://dx.doi.org/10.1016/0167-5699\(92\)90071-E](http://dx.doi.org/10.1016/0167-5699(92)90071-E)
170. Cohen IR. The cognitive paradigm and the immunological homunculus. *Immunol Today* 1992; 13:490-4; PMID:1463581; [http://dx.doi.org/10.1016/0167-5699\(92\)90024-2](http://dx.doi.org/10.1016/0167-5699(92)90024-2)
171. Turner CH, Robling AG, Duncan RL, Burr DB. Do bone cells behave like a neuronal network? *Calcif Tissue Int* 2002; 70:435-42; PMID:12149636; <http://dx.doi.org/10.1007/s00223-001-1024-z>
172. Spencer GJ, Genever PG. Long-term potentiation in bone—a role for glutamate in strain-induced cellular memory? *BMC Cell Biol* 2003; 4:9;

- PMID:12892570; <http://dx.doi.org/10.1186/1471-2121-4-9>
173. Zoghi M. Cardiac memory: do the heart and the brain remember the same? *J Interv Card Electrophysiol* 2004; 11:177-82; PMID:15548883; <http://dx.doi.org/10.1023/B:JICE.0000048567.18088.a2>
 174. Chakravarthy SV, Ghosh J. On Hebbian-like adaptation in heart muscle: a proposal for 'cardiac memory'. *Biol Cybern* 1997; 76:207-15; PMID:9151418
 175. Goel P, Mehta A. Learning theories reveal loss of pancreatic electrical connectivity in diabetes as an adaptive response. *PLoS One* 2013; 8:e70366; PMID:23936417
 176. McCulloch WS. Why the Mind is in the Head? In: Jeffress LA, ed. *Cerebral Mechanisms in Behavior: The Hixon Symposium*, New York: Jon Wiley Publ; 1951. 42-57 p.
 177. Mustard J, Levin M. Bioelectrical Mechanisms for Programming Growth and Form: Taming Physiological Networks for Soft Body Robotics. *Soft Robotics* 2014; 1:169-91
 178. Levin M. Endogenous bioelectrical networks store non-genetic patterning information during development and regeneration. *J Physiol* 2014; 592:2295-305; PMID:24882814
 179. Tseng A, Levin M. Cracking the bioelectric code: Probing endogenous ionic controls of pattern formation. *Commun Integ Biol* 2013; 6:1-8
 180. Levin M. Reprogramming cells and tissue patterning via bioelectrical pathways: molecular mechanisms and biomedical opportunities. *Wiley Interdisciplinary Reviews: Systems Biology and Medicine* 2013; 5: 657-76.
 181. Levin M. The wisdom of the body: future techniques and approaches to morphogenetic fields in regenerative medicine, developmental biology and cancer. *Regen Med* 2011; 6:667-73; PMID:22050517
 182. Levin M. Morphogenetic fields in embryogenesis, regeneration, and cancer: non-local control of complex patterning. *Bio Systems* 2012; 109:243-61; PMID:22542702
 183. Levin M, Buznikov GA, Lauder JM. Of minds and embryos: left-right asymmetry and the serotonergic controls of pre-neural morphogenesis. *Dev Neurosci* 2006; 28:171-85; PMID:16679764
 184. Buznikov G, Shmukler Y, Lauder J. From oocyte to neuron: do neurotransmitters function in the same way throughout development? *Cell Molec Neurobiol* 1996; 16:537-59; PMID:8956008
 185. Stewart S, Rojas-Munoz A, Izpisua Belmonte JC. Bioelectricity and epimorphic regeneration. *Bioessays* 2007; 29:1133-7; PMID:17935197
 186. McCaig CD, Rajnicek AM, Song B, Zhao M. Controlling cell behavior electrically: current views and future potential. *Physiol Rev* 2005; 85:943-78; PMID:15987799
 187. Levin M. Molecular bioelectricity in developmental biology: new tools and recent discoveries: control of cell behavior and pattern formation by transmembrane potential gradients. *Bioessays* 2012; 34:205-17; PMID:22237730
 188. Levin M, Stevenson CG. Regulation of cell behavior and tissue patterning by bioelectrical signals: challenges and opportunities for biomedical engineering. *Annu Rev Biomed Eng* 2012; 14:295-323; PMID:22809139
 189. Levin M. Molecular bioelectricity: how endogenous voltage potentials control cell behavior and instruct pattern regulation in vivo. *Mol Biol Cell* 2014; 25:3835-50; PMID:25425556
 190. Lahoz-Beltra R, Hameroff SR, Dayhoff JE. Cytoskeletal logic: a model for molecular computation via Boolean operations in microtubules and microtubule-associated proteins. *Biosystems* 1993; 29:1-23; PMID:8318677
 191. Rasmussen S, Karampurwala H, Vaidyanath R, Jensen KS, Hameroff S. Computational Connectionism within Neurons - a Model of Cytoskeletal Automata Sub-serving Neural Networks. *Physica D* 1990; 42:428-49; [http://dx.doi.org/10.1016/0167-2789\(90\)90093-5](http://dx.doi.org/10.1016/0167-2789(90)90093-5)
 192. Ling H, Samarasinghe S, Kulasiri D. Novel recurrent neural network for modelling biological networks: oscillatory p53 interaction dynamics. *Biosystems* 2013; 114:191-205; PMID:24012741; <http://dx.doi.org/10.1016/j.biosystems.2013.08.004>
 193. Friston K, Levin M, Sengupta B, Pezzulo G. Knowing one's place: a free-energy approach to pattern regulation. *J R Soc Interface* 2015; 12:pii: 20141383; PMID:25788538; <http://dx.doi.org/10.1098/rsif.2014.1383>
 194. Ray S. Survival of olfactory memory through metamorphosis in the fly *Musca domestica*. *Neuroscience Letters* 1999; 259:37-40; PMID:10027550; [http://dx.doi.org/10.1016/S0304-3940\(98\)00892-1](http://dx.doi.org/10.1016/S0304-3940(98)00892-1)
 195. Maldonado H, Tablante A. Behavioral transfer in praying mantis by injection of brain homogenate. *Physiol Behav* 1976; 16:617-21; PMID:987593; [http://dx.doi.org/10.1016/0031-9384\(76\)90223-7](http://dx.doi.org/10.1016/0031-9384(76)90223-7)
 196. Martin U, Martin H, Lindauer M. Transplantation of a Time-Signal in Honeybees. *J Comparative Physiol* 1978; 124:193-201; <http://dx.doi.org/10.1007/BF00657051>