

17 NOVEMBER 2022

$$MGF_{X}(M) = 0.5 \int_{-\infty}^{\infty} e^{(s+1)N} dN + 0.5 \int_{-\infty}^{\infty} e^{(s-1)M} dN$$

$$y s < 1$$

$$E[X'] = \frac{d^2}{ds^2} MGF_X(Y) = \frac{d}{ds} \left( \frac{d}{ds} \left( 1 - s^2 \right)^{-2} \right) \bigg|_{s=0}$$

$$\sim E[X^2] = \frac{d}{ds} \left( -(1-s^2)^{-2} - 2s \right)$$

$$rightarrow E[X^2] = 2(1-5)^{-2} + 2s(1-5^2)^{-3} \cdot -2 \cdot -2s$$

Find Possible (Xi, Xi) 
$$\frac{1}{3}(\frac{1}{3})^2 = \frac{2}{3}P(Ni) + \frac{1}{3}I(\frac{1}{3})^2$$

1 (0,1), (1,0)  $(\frac{1}{3})(\frac{2}{3})^2 \cdot \frac{1}{3} + (\frac{1}{3})(\frac{2}{3})^{\frac{1}{3}}$ 

2 (0,2), (1,1), (2,0)  $(\frac{1}{3})(\frac{2}{3})^2 \cdot \frac{1}{3} + (\frac{1}{3})(\frac{2}{3})^{\frac{1}{3}}$ 

+  $(\frac{1}{3})(\frac{2}{3})^2 \cdot \frac{1}{3}$ 

$$\frac{3i}{0} \frac{\cancel{5} p(3i) = \sum p(Ni) p(yi)}{\frac{1}{9} \cdot 1 = \frac{1}{5} \cdot \frac{9}{5}}$$

$$\frac{1}{9} \cdot \frac{5}{3} = \frac{1}{9} \cdot \frac{5}{5}$$

$$\frac{1}{9} \cdot \frac{19}{5}$$

$$p(Z \le 2) = p(X+Y \le 2) = \frac{1}{9} \left\{ \frac{9}{9} + \frac{15}{9} + \frac{19}{9} \right\} = \frac{43}{81}$$

• 
$$P(2 + P(x+y \le 2)) = \frac{43}{81}$$

Am

3. fxx (n,y) Shaded enegio fxx(x14)= ¥ n € 0 (1,2) or fx,x (M17) = 2+3)(2+1).1 x € [1/2] y ∈ (0,n) PTO! faylargida fxxx(my) dn £y(y) = fy1973 (0,2) - /yly1fyly]= 2-23 y e (0,2) tx1 y = y (x, Y=y)= (n,y) 3 (24-42) 

$$f_{\gamma}(y) = \begin{cases} \int_{0}^{\infty} f_{x,\gamma}(y,y) dy & y \in (0,1) \\ y = 1 & y = 1 \\ y = 2 & y = 1 \end{cases}$$

$$\int_{0}^{\infty} f_{x,\gamma}(y,y) dy & y \in (0,1)$$

$$\int_{0}^{\infty} f_{x,\gamma}(y,y) dy & y \in (0,1)$$

$$\frac{2}{3}(2-y) = \frac{2}{3} \qquad y \in (971)$$

$$\frac{2}{3}(2-y) \qquad y \in (12)$$

Ann

f(x17=y) = fxx(414)

f(19)

1 11 20

$$f_{X|Y}(X|Y=y) = \begin{cases} \frac{2}{3} & y \in (0,1) \\ \frac{2}{3} & y \in (0,1) \end{cases}$$

$$\frac{2}{3}(2-y) \qquad y \in (0,1)$$

$$3(x)^{(1)}$$
 $f_{X|Y}(X|Y=1) = \begin{cases} 1 & y \in (0,1) \\ \frac{1}{1-y} & y \in (0,2) \end{cases}$ 

Ansa

30)"

FXIY depends ons value of y in X and y are

NOT solve independent.

But  $\mu_{X|Y=7}$  n=2But  $\mu_{X|Y=7}$   $\int n \cdot f_{X|X}(X|Y=Y) dn$  n=01

$$\mu_{X|Y=y} = \begin{cases} \int_{-\infty}^{\infty} x \cdot 1 \cdot dx & y \in [0,1] \\ \int_{-\infty}^{\infty} \frac{1}{1-y} x \cdot n \cdot dx & y \in [1,2] \end{cases}$$

$$\mu_{X|Y} = \int \frac{x^2}{2} \Big|_{1}^{2} \qquad y \in (0,1)$$

$$\frac{1}{1-y} \cdot \frac{x^2}{2} \Big|_{1}^{2} \qquad y \in (0,1)$$

$$\frac{3}{3}$$

$$\frac{3}{2}$$

$$\frac{3}$$

An

or 
$$f_{\times}(n) = \frac{1}{2\sqrt{2n}} e^{-\frac{1}{2}\left\{\frac{n}{2}\right\}^{2}}$$

or 
$$f_{\gamma}(y) = \frac{1}{3\sqrt{2}} e^{-\frac{1}{2}\left\{\left(\frac{y}{3}\right)^{2}\right\}}$$

$$f_{X,Y}(y,y) = \frac{1}{2\sqrt{2n} \cdot 3\sqrt{2n} \int \left(1 - \left(-\frac{1}{2}\right)^{\frac{n}{2}}\right)} \left(\frac{2\sqrt{1 - \left(\frac{1}{2}\right)^{\frac{n}{2}}}}{2\sqrt{2n} \cdot 3\sqrt{2n} \int \left(1 - \left(-\frac{1}{2}\right)^{\frac{n}{2}}\right)} - 2\cdot \left(\frac{2\sqrt{2}}{2}\right) \cdot \left(\frac{2\sqrt{2}}{2$$

$$f_{\times/Y}(n/y) = \frac{1}{9n(\sqrt{2})} \left(\frac{2}{3}\right) + \left(\frac{2}{3}\right) + \left(\frac{2}{3}\right) \left(\frac{2}{3}\right)$$

Again.

Jen clarity)  $f_{x,y}(n,y) = \frac{1}{6\sqrt{3}x}e^{-\frac{1}{2}\cdot\frac{4}{3}\xi\left(\frac{N}{2}\right)^{2} + \left(\frac{y}{3}\right)^{2} + \left(\frac{y}{3}\right)^{3}}$ (4)

$$Con(Z, W) = Con(X+Y, bX)$$

or 
$$Con(\overline{z}/W) = E[bX^2 + bXY]$$

on 
$$Con(R_1W) = bMx^2 + bCon(XY)$$

on 
$$Con(Z/W) = b\{ 6x^2 + Mx^3 + b\{ Pxy 6x6x \}$$
 on  $\mu_x \mu_{y=0}$ .

or 
$$Con(Z_1W) = b \left[ 4 + 0^2 + \frac{-1}{2} \cdot 2 \cdot 3 \right]$$

or 
$$Con(Z,W) = b[1]$$

or 
$$(z,w) = b[1]$$
(4(b)ii)
(1)
(2,w) = b
(2,w) > 0 + b > 0

← Con (z,w) > 0 + b > 0

← Con (z,w) > 0 + b > 0

or 
$$Pz_{,W} = \frac{b}{\sqrt{7.2b}}$$

$$f_{z,w}(3,w) \approx \mathcal{N}(3,w, \mu_{z}=0, \mu_{w}=0, \mu_{z}=0, \mu_{$$

4(c) 
$$\hat{y}_{y|X=N} = E[Y|X=N]$$

Let' toug to find fy/x (Y/X=N) first!

$$f_{\gamma|\chi}(\gamma|\chi=n)=\frac{f_{\chi,\gamma}(\gamma,\gamma)}{f_{\chi}(\chi)}$$

$$\frac{1}{2\sqrt{2}\pi} e^{-\frac{1}{2}\left\{\left(\frac{\chi}{2}\right)^{2}\right\}}$$

4.0

• 
$$f_{Y|X}(Y|X=n) = \frac{1}{\frac{3\sqrt{3}}{2}\sqrt{2n}}e^{-\frac{1}{2}\left\{\frac{y}{3\sqrt{3}} + \frac{x}{2\sqrt{3}}\right\}^{2}}$$

$$e^{-\frac{1}{2}} \left\{ \frac{3\sqrt{3}}{2} \sqrt{2x} e^{-\frac{1}{2}} \right\} \left( \frac{3\sqrt{3}}{2} \sqrt{2x} \right)^{\frac{3}{2}}$$

$$\frac{1}{3} \frac{1}{3} \frac{1}$$

- X, \_\_\_\_>

- 
$$f_{X,Y,Z}(M,Y/3)^{-}$$
  $\frac{1}{y} \cdot \frac{1}{x} \cdot \frac{1}{1} \quad \forall \quad g \in (0,Y)$   
 $x \in (0,L)$ 

:. 
$$E[X^2Y^2Z^2] = \int_{y=0}^{y=0} \int_{y=0}^{y=0} u^2 \cdot y^2 \cdot 3^2 \cdot \frac{1}{ny} dy dy dx$$

or 
$$E[X^2Z^2] = \int_{x=0}^{x=1} x^{\frac{y}{2}} dy dx$$

$$\mathbf{E}\left[X^{2}Y^{2}Z^{2}\right] = \int_{N=2}^{\infty} N \cdot \frac{y^{5}}{15} \begin{vmatrix} y^{5}N \\ y^{5} \end{vmatrix} dn$$

$$= E[X^{2}Y^{2}] = \int_{15}^{\infty} \frac{n^{6}}{15} dn$$

$$\alpha \in [x^2 y^2 z^2] = \frac{n^{\frac{3}{2}}}{105} \Big|_{x=0}^{x=1}$$

6. Random Processes are mappings of outcomes of

à quebabilistic enperiment to signals.

 $X(A,t) = \cos(t)$   $X(C,t) = \begin{cases} 7 & \text{Lt} \text{ is even} \\ 0 & \text{Lt} \text{ is odd} \end{cases}$   $X(B,t) = 2e^{-2t} \qquad \text{where Lt} \text{ surpos } t \text{ the greatest indeger lesson}$  than on equal to t.

Mintually any physical perocess or simulated = phenomenon can be superesented as a standom puocess.

Ey. Bus Voltages in a power guid one terre-varying signals, and there were multiple buses (outcomes) in a power guid or we can measure a single bus voltage in multiple ways. (sample functions).

i=1 V1(t)

i=14 V1(t)

i=14 V1(t)

i=14 V2(t)

i=14 V2(t)

i=14 V2(t)

i=14 V2(t)

Study of pardom Perocesses is important as it can be used to analyze, estimate, peredict, furecast signals which can be very beneficial to us.

Ey. 21 tomacionis lood demand for a regional substation could be peredicted accurately, we can optimize the scheduling of power generation, spot market notes, etc., which can same a guid operation millions of Balloss.

- ender the sales in elloward and

> ing on a sign of the second of the sign of gradina per en la composition de la proposition de la composition de la composition de la composition de la co Proposition de la composition de la co

and the second of the second o 

and the state of an armony of the second of

19 A : 14 and of solar than the

The second of th 

teR+ 7.  $\times$ (4) c ~ enp(1) = fc(c) = e c 20 X(t) is a Rendom Perocess as 7(a) X(t) = X(C,t) is a time vacying signals which are all mapped to the outcome C of a prope probabilistic informat, where C ~ enp (1). have a sample function So X(C,+) can to X(C=2,+) on it can be sampled out time to get a resigne set of X(C, t=t\*) & C resp(1). Souple for. 2 X(C) = +1+ enp(1) C/E [0/00) x(co,t) 4 ×(0,4)= ++1

$$(xt) = \frac{1}{t+1+c}$$
  $t \in \mathbb{R}^{+}$   $c \sim enp(1)$   $c \in [0, \infty)$ 

$$a \times (t) \in \left(0, \frac{1}{t+1}\right)$$

Approach: Find CDF then polf:

$$F_{X(+)}(n) = p(X(+) \le n) = p(\frac{1}{1+1+c} \le n)$$