13

81.5/85

17 NOVEMBER 2022

Great!

1.1

+10/10/10

$$P = MGF_{X}(M) = 0.5 \int_{0.5}^{0.5} e^{(S+1)N} dN + 0.5 \int_{0.5}^{0.5} e^{(S-1)N} dN$$

$$y = S = 1$$

$$y = S = 1$$

$$\sim E[X^2] = \frac{d}{ds} \left( -(1-s^2)^{-2} - 2s \right)$$

$$rightarrow E[X^2] = 2(1-5)^{-2} + 2s(1-5^2)^{-3} \cdot -2 \cdot -2s$$

to the term of the

Finalle (Xi, Yi) 
$$\frac{1}{3}(\frac{2}{3})^{\circ} = \frac{2}{5}P(x_1, y_1) = \frac{2}{5}P($$

$$\frac{3i}{0} \frac{1}{9} \cdot \frac{1}{3} = \frac{1}{3} \cdot \frac{9}{3}$$

$$\frac{1}{9} \cdot \frac{5}{3} = \frac{1}{9} \cdot \frac{5}{9}$$


$$\frac{1}{9} \cdot \frac{19}{3} = \frac{1}{9} \cdot \frac{5}{9}$$

$$p(Z \le 2) = p(X+Y \le 2) = \frac{1}{9} \left\{ \frac{9}{9} + \frac{15}{9} + \frac{19}{9} \right\} = \frac{43}{81}$$

$$P(2 + P(x+y \le 2) = \frac{43}{81}$$

$$P(x+y \le 2) \approx 0.5309$$

Am



$$f_{\gamma}(y) = \begin{cases} \int_{0}^{\infty} f_{x,\gamma}(y,y) dy & y \in (0,1) \\ y = 1 & y = 1 \\ y = 2 & y = 1 \end{cases}$$

$$\int_{0}^{\infty} f_{x,\gamma}(y,y) dy & y \in (0,1)$$

$$\int_{0}^{\infty} f_{x,\gamma}(y,y) dy & y \in (0,1)$$

$$\frac{2}{3}(2-y) = \frac{2}{3} \qquad y \in (971)$$

$$\frac{2}{3}(2-y) \qquad y \in (12)$$

Ann

f(x17=y) = fxx(414)

f(19)

1 11 20

$$f_{X|Y}(X|Y=y) = \begin{cases} \frac{2}{3} & y \in (0,1) \\ \frac{2}{3} & y \in (0,1) \end{cases}$$

$$\frac{2}{3} (2-y) \quad y \in (0,1)$$

$$3(x)^{(1)}$$
 $f_{X|Y}(X|Y=g) = \begin{cases} 1 & y \in (0,1) \\ \frac{1}{1-y} & y \in (1/2) \end{cases}$ 

Anso

Firster independent.

But  $\mu_{X|Y=7} = \int n \cdot f_{X|X}(X|Y=Y) dn$  n=01

$$\mu_{X|Y=Y} = \begin{cases} \int_{-1}^{2} x \cdot 1 \cdot dx & y \in (0,1) \\ x \cdot 1 & y \in (1,2) \end{cases}$$
 wrong x range

$$\mu_{X|Y} = \int \frac{x^2}{2} \Big|_{1}^{2} \qquad y \in (0,1)$$

$$\frac{1}{1-y} \cdot \frac{x^2}{2} \Big|_{1}^{2} \qquad y \in (0,1)$$

$$\frac{3}{3}$$

$$\frac{3}{2}$$

$$\frac{3}$$

An

+14.5/15

or 
$$f_{\times}(n) = \frac{1}{2\sqrt{2n}} e^{-\frac{1}{2}\left\{\left(\frac{n}{2}\right)^{2}\right\}}$$

or 
$$f_{\gamma}(y) = \frac{1}{3\sqrt{2}} e^{-\frac{1}{2}\left\{\left(\frac{y}{3}\right)^{2}\right\}}$$

$$f_{X,Y}(y) = \frac{1}{2\sqrt{2n} \cdot 3\sqrt{2n} \cdot \left(1 - \left(-\frac{1}{2}\right)^{\frac{n}{2}}\right)} = \frac{2\sqrt{1 - \left(\frac{1}{2}\right)^{\frac{n}{2}}}}{2\sqrt{2n} \cdot \left(\frac{1}{2}\right) \cdot \left(\frac{1}{2}\right)^{\frac{n}{2}}} = \frac{2\cdot \left(\frac{1}{2}\right) \cdot \left(\frac{1}{2}\right)$$

$$f_{\times/Y}(n/y) = \frac{1}{\sqrt{3}} \left(\frac{x}{2}\right)^{\frac{1}{2}} + \left(\frac{y}{3}\right)^{\frac{1}{2}} + \left(\frac{x}{2}\right)^{\frac{1}{2}}$$

94 43 6 5 3 7

Again.

Again.

$$f_{x,y}(y) = \frac{1}{6\sqrt{3}x}e^{-\frac{1}{2}\cdot\frac{4}{3}\xi\left(\frac{x}{2}\right)^{2} + \left(\frac{y}{3}\right)^{2} + \left(\frac{y}{3}\right)^{3}}$$

All  $f_{x,y}(y) = \frac{1}{6\sqrt{3}x}e^{-\frac{1}{2}\cdot\frac{4}{3}\xi\left(\frac{x}{2}\right)^{2} + \left(\frac{y}{3}\right)^{2} + \left(\frac{y}{3}\right)^{3}}$ 

$$Con(Z, W) = Con(X+Y, bX)$$

or 
$$Con(\overline{z}/W) = E[bX^2 + bXY]$$

on 
$$Con(R_1W) = bMx^2 + bCon(XY)$$

on 
$$Con(Z/W) = b\{ 6x^2 + Mx^3 + b\{ Pxy 6x 6x \} \}$$
 on  $\mu_X \mu_{Y=0}$ .

or 
$$Con(Z_1W) = b \left[ 4 + 0^2 + \frac{-1}{2} \cdot 2 \cdot 3 \right]$$

or 
$$Con(Z,W) = b[1]$$

or 
$$(z,w) = b[1]$$
(4(b)ii)
(1)
(2,w) = b
(2,w) > 0 + b > 0

← Con (z,w) > 0 + b > 0

← Con (z,w) > 0 + b > 0

or 
$$Pz_{,W} = \frac{b}{\sqrt{7.2b}}$$

4(c) 
$$\hat{y}_{y|X=N} = E[Y|X=N]$$

Let' toug to find fy/x(Y/X=N) first!

$$f_{\gamma|\chi}(\gamma|\chi=n)=\frac{f_{\chi,\gamma}(\gamma,\gamma)}{f_{\chi}(\chi)}$$

$$\frac{1}{2\sqrt{2}\pi} e^{-\frac{1}{2}\left\{\left(\frac{\chi}{2}\right)^{2}\right\}}$$

4.0

• 
$$f_{Y|X}(Y|X=n) = \frac{1}{\frac{3\sqrt{3}}{2}\sqrt{2n}}e^{-\frac{1}{2}\left\{\frac{y}{3\sqrt{3}} + \frac{x}{2\sqrt{3}}\right\}^{2}}$$

$$e^{-\frac{1}{2}} \left\{ \frac{3\sqrt{3}}{2} \sqrt{2x} e^{-\frac{1}{2}} \right\} \left( \frac{3\sqrt{3}}{2} \sqrt{2x} \right)^{\frac{3}{2}}$$

$$\frac{1}{3} \frac{1}{3} \frac{1}$$

- X, \_\_\_\_>

+10/10

But fx, y, z (M, y, 3) = Fato \$263

- 
$$f_{x,y,z}(m,y,3) = \frac{1}{y} \cdot \frac{1}{n} \cdot \frac{1}{1} \quad \forall \quad g \in (0,y)$$

$$n \in (0,1)$$

:. 
$$E[X^2Y^2Z^2] = \int_{y=0}^{y=0} \int_{y=0}^{y=0} x^2 \cdot y^2 \cdot 3^2 \cdot \frac{1}{y^2} dy dy$$

or 
$$E[X^2Z^2] = \int_{x=0}^{x=1} x^{\frac{y}{2}} dy dx$$

$$\mathbf{E}\left[X^{2}Y^{2}Z^{2}\right] = \int_{N=3}^{\infty} N \cdot \frac{y^{5}}{15} \begin{vmatrix} y^{5}N \\ y^{5} \end{vmatrix} dn$$

$$= E[X^{2}Y^{2}] = \int_{15}^{\infty} \frac{n^{6}}{15} dn$$

$$\alpha \in [x^2 y^2 z^2] = \frac{n^{\frac{3}{2}}}{105} \Big|_{x=0}^{x=1}$$



6. Random Priscerces are mappings of outcomes of

à pubabilistic enperiment to signals.

Lt] is even Lt] is odd where LtJ sepass to the quentest indeger lesser than or equal to t.

Mutually any physical perocess or simulated phenomenon can be superesented as a siandom puocess.

G. Bus Voltages in a pomer guid are time-varying signals, and there are multiple buses (outcomes) in a poner gend or we can measure a single bus veltage in multiple ways. (sample functions).

i=1-V; (+) 1=14 V14(t) ie 1 1 14 IZ IFEE 14 Bus System.  $i \in N$ 

Study of Random Perocesses is important as it can be used to analyze, estimate, peredict, furecast signals which can be very beneficial to us.

Ey. 21 tomacionis lood demand for a regional substation could be peredicted accurately, we can optimize the scheduling of power generation, spot market notes, etc., which can same a guid operation millions of Balloss.

- ender the sales in elloward and

> ing on a sign of the second of the sign of gradina per en la composition de la proposition de la composition de la composition de la composition de la co Proposition de la composition de la co

and the second of the second o 

and the state of an armony of the second of

19 A : 14 and of solar than the

The second of th 

7. t E R+  $\times$ (4)  $c \sim enp(1)$ = te(e) = = fc(c) = e c 20 X(t) is a Rendom Perocess as 7(a) X(t) = X(C,t) is a time vacuying signals which are all mapped to the outcome C of a persper probabilistic inferiment, where C ~ enp (1). have a sample function So X(C,+) can to X(C=2,+) on it can be sampled out time to get a resigne set of X(C, t=t\*) & C resp(1). Souple for. 2 X(C) = +1+ enp(1) C/E [0/00)
C/ is a particular x(co,t) h ×(0,4)= +1

$$(xt) = \frac{1}{t+1+c}$$
  $t \in \mathbb{R}^{+}$   $c \sim enp(1)$   $c \in [0, \infty)$ 

$$a \times (t) \in \left(0, \frac{1}{t+1}\right)$$

Approach: Find CDF then polf:

$$F_{X(+)}(n) = p(X(+) \le n) = p(\frac{1}{1+1+c} \le n)$$