

Fall 2025 CPTS530 Homework 5  
MATH 448-548  
Numerical Analysis

Aryan Ritwajeet Jha

November 2025

**Contents**

|                   |           |
|-------------------|-----------|
| <b>Problem 1</b>  | <b>2</b>  |
| <b>Problem 2</b>  | <b>4</b>  |
| <b>Problem 3</b>  | <b>5</b>  |
| <b>Problem 4</b>  | <b>6</b>  |
| <b>Problem 5</b>  | <b>8</b>  |
| <b>Problem 6</b>  | <b>11</b> |
| <b>Problem 7</b>  | <b>12</b> |
| <b>Problem 8</b>  | <b>14</b> |
| <b>Problem 9</b>  | <b>16</b> |
| <b>Appendix</b>   | <b>17</b> |
| <b>References</b> | <b>19</b> |

# Problem 1

## Problem Statement (Textbook 4.4.1):

Show that the norms  $\|\mathbf{x}\|_\infty$ ,  $\|\mathbf{x}\|_2$ , and  $\|\mathbf{x}\|_1$  satisfy the postulates (1), (2), and (3) for norms.

### Norm Postulates [1, 2]:

On a vector space  $V$ , a **norm** is a function  $\|\cdot\|$  from  $V$  to the set of nonnegative reals that obeys these three postulates:

$$\|x\| > 0 \text{ if } x \neq 0, x \in V \quad (1)$$

$$\|\lambda x\| = |\lambda| \|x\| \text{ if } \lambda \in \mathbb{R}, x \in V \quad (2)$$

$$\|x + y\| \leq \|x\| + \|y\| \text{ if } x, y \in V \quad (\text{triangle inequality}) \quad (3)$$

### Solution:

We verify postulates (1), (2), and (3) for  $\|x\|_\infty = \max_i |x_i|$ ,  $\|x\|_2 = \sqrt{\sum_i x_i^2}$ , and  $\|x\|_1 = \sum_i |x_i|$ .

**(a)  $\infty$ -norm:**  $\|x\|_\infty = \max_{1 \leq i \leq n} |x_i|$

(1) If  $x \neq 0$ , then  $\exists i$  with  $x_i \neq 0$ , so  $|x_i| > 0$ . Thus  $\|x\|_\infty = \max_i |x_i| > 0$ .

(2)  $\|\lambda x\|_\infty = \max_i |\lambda x_i| = \max_i |\lambda| |x_i| = |\lambda| \max_i |x_i| = |\lambda| \|x\|_\infty$ .

(3)  $\|x + y\|_\infty = \max_i |x_i + y_i| \leq \max_i (|x_i| + |y_i|) \leq \max_i |x_i| + \max_i |y_i| = \|x\|_\infty + \|y\|_\infty$ .

**(b) 2-norm (Euclidean):**  $\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$

(1) If  $x \neq 0$ , then  $\exists i$  with  $x_i \neq 0$ , so  $\sum_i x_i^2 > 0$ . Thus  $\|x\|_2 = \sqrt{\sum_i x_i^2} > 0$ .

(2)  $\|\lambda x\|_2 = \sqrt{\sum_i (\lambda x_i)^2} = \sqrt{\lambda^2 \sum_i x_i^2} = |\lambda| \sqrt{\sum_i x_i^2} = |\lambda| \|x\|_2$ .

(3) Expanding  $\|x + y\|_2^2$ :

$$\begin{aligned} \|x + y\|_2^2 &= \sum_i (x_i + y_i)^2 = \sum_i x_i^2 + 2 \sum_i x_i y_i + \sum_i y_i^2 \\ &= \|x\|_2^2 + 2 \sum_i x_i y_i + \|y\|_2^2 \end{aligned}$$

Since  $\sum_i x_i y_i \leq \|x\|_2 \|y\|_2$  (dot product bounded by product of norms):

$$\|x + y\|_2^2 \leq \|x\|_2^2 + 2\|x\|_2 \|y\|_2 + \|y\|_2^2 = (\|x\|_2 + \|y\|_2)^2$$

Taking square roots:  $\|x + y\|_2 \leq \|x\|_2 + \|y\|_2$ .

**(c) 1-norm:**  $\|x\|_1 = \sum_{i=1}^n |x_i|$

**(1)** If  $x \neq 0$ , then  $\exists i$  with  $x_i \neq 0$ , so  $|x_i| > 0$ . Thus  $\|x\|_1 = \sum_i |x_i| > 0$ .

**(2)**  $\|\lambda x\|_1 = \sum_i |\lambda x_i| = \sum_i |\lambda| |x_i| = |\lambda| \sum_i |x_i| = |\lambda| \|x\|_1$ .

**(3)** For each term, the triangle inequality for absolute values gives  $|x_i + y_i| \leq |x_i| + |y_i|$ .

Summing over all  $i$ :

$$\|x + y\|_1 = \sum_i |x_i + y_i| \leq \sum_i (|x_i| + |y_i|) = \sum_i |x_i| + \sum_i |y_i| = \|x\|_1 + \|y\|_1$$

All three norms satisfy postulates (1), (2), and (3).

## Problem 2

**Problem Statement (Textbook 4.4.2):**

Show that  $\|\mathbf{x}\|_\infty \leq \|\mathbf{x}\|_2 \leq \|\mathbf{x}\|_1$  for all  $\mathbf{x} \in \mathbb{R}^n$ , and that equalities can occur, even for nonzero vectors.

---

**Solution:****Part 1: Prove**  $\|x\|_\infty \leq \|x\|_2$ Let  $\|x\|_\infty = \max_i |x_i| = |x_k|$  for some index  $k$ . Then:

$$\|x\|_2^2 = \sum_{i=1}^n x_i^2 \geq x_k^2 = \|x\|_\infty^2$$

Taking square roots:  $\boxed{\|x\|_\infty \leq \|x\|_2}$ .**Equality case:** Equality holds when  $x$  has only one nonzero component. For example,  $x = (5, 0, 0)^T$  gives  $\|x\|_\infty = 5$  and  $\|x\|_2 = 5$ .**Part 2: Prove**  $\|x\|_2 \leq \|x\|_1$ First, observe that each component satisfies  $|x_i| \leq \|x\|_1$  (since  $\|x\|_1 = \sum_{j=1}^n |x_j|$  includes all terms).For each  $i$ , we can write:

$$|x_i|^2 = |x_i| \cdot |x_i| \leq |x_i| \cdot \|x\|_1$$

where the inequality holds because  $|x_i| \leq \|x\|_1$ .

Now sum this inequality over all indices:

$$\begin{aligned} \|x\|_2^2 &= \sum_{i=1}^n |x_i|^2 \leq \sum_{i=1}^n |x_i| \cdot \|x\|_1 \\ &= \|x\|_1 \sum_{i=1}^n |x_i| \quad (\text{factor out } \|x\|_1) \\ &= \|x\|_1 \cdot \|x\|_1 = \|x\|_1^2 \end{aligned}$$

Taking square roots of both sides:  $\boxed{\|x\|_2 \leq \|x\|_1}$ .**Equality case:** Equality holds when  $x$  has only one nonzero component. For example,  $x = (0, 3, 0)^T$  gives  $\|x\|_2 = 3$  and  $\|x\|_1 = 3$ .**Conclusion:** The inequality  $\boxed{\|x\|_\infty \leq \|x\|_2 \leq \|x\|_1}$  holds for all  $x \in \mathbb{R}^n$ , with equality possible for nonzero vectors.

## Problem 3

### Problem Statement (Textbook 4.4.3):

(Continuation) Show that  $\|\mathbf{x}\|_1 \leq n\|\mathbf{x}\|_\infty$  and  $\|\mathbf{x}\|_2 \leq \sqrt{n}\|\mathbf{x}\|_\infty$  for  $\mathbf{x} \in \mathbb{R}^n$ .

#### Solution:

**Part 1: Prove**  $\|x\|_1 \leq n\|x\|_\infty$

Since  $\|x\|_\infty = \max_i |x_i|$ , we have  $|x_i| \leq \|x\|_\infty$  for each  $i = 1, 2, \dots, n$ .

Summing over all components:

$$\|x\|_1 = \sum_{i=1}^n |x_i| \leq \sum_{i=1}^n \|x\|_\infty = n\|x\|_\infty$$

Therefore:  $\boxed{\|x\|_1 \leq n\|x\|_\infty}$ .

**Equality case:** Equality holds when all components have the same absolute value. For example,  $x = (1, 1, 1)^T$  gives  $\|x\|_1 = 3 = 3 \cdot 1 = 3\|x\|_\infty$ .

**Part 2: Prove**  $\|x\|_2 \leq \sqrt{n}\|x\|_\infty$

Since  $|x_i| \leq \|x\|_\infty$  for each  $i$ , squaring gives  $|x_i|^2 \leq \|x\|_\infty^2$ .

Summing over all components:

$$\|x\|_2^2 = \sum_{i=1}^n |x_i|^2 \leq \sum_{i=1}^n \|x\|_\infty^2 = n\|x\|_\infty^2$$

Taking square roots:  $\boxed{\|x\|_2 \leq \sqrt{n}\|x\|_\infty}$ .

**Equality case:** Equality holds when all components have the same absolute value. For example,  $x = (2, 2, 2)^T$  gives  $\|x\|_2 = \sqrt{12} = 2\sqrt{3} = \sqrt{3} \cdot 2 = \sqrt{3}\|x\|_\infty$ .

## Problem 4

### Problem Statement (Textbook 4.4.8):

Define

$$\|A\| = \sum_{i=1}^n \sum_{j=1}^n |a_{ij}|$$

Show that this is a matrix norm (that is, a norm on the linear space of all  $n \times n$  matrices). Show that it is not subordinate to any vector norm. Does it conform to Equation (9) and Inequality (10)?

$$\|I\| = 1 \quad (\text{Equation 9})$$

$$\|AB\| \leq \|A\|\|B\| \quad (\text{Inequality 10})$$

### Solution:

#### Part 1: Show $\|A\|$ is a matrix norm

A matrix norm must satisfy three postulates:

- (1) **Positivity:** If  $A \neq 0$ , then  $\exists$  some  $a_{ij} \neq 0$ , so  $|a_{ij}| > 0$ . Thus  $\|A\| = \sum_{i,j} |a_{ij}| > 0$ .
- (2) **Homogeneity:**  $\|\lambda A\| = \sum_{i,j} |\lambda a_{ij}| = \sum_{i,j} |\lambda| |a_{ij}| = |\lambda| \sum_{i,j} |a_{ij}| = |\lambda| \|A\|$ .
- (3) **Triangle inequality:**

$$\begin{aligned} \|A + B\| &= \sum_{i,j} |a_{ij} + b_{ij}| \leq \sum_{i,j} (|a_{ij}| + |b_{ij}|) \\ &= \sum_{i,j} |a_{ij}| + \sum_{i,j} |b_{ij}| = \|A\| + \|B\| \end{aligned}$$

Therefore,  $\|A\|$  is a matrix norm.

#### Part 2: Show it is NOT subordinate to any vector norm

A subordinate (induced) matrix norm satisfies  $\|I\| = 1$ . However, for the identity matrix  $I$  where  $(I)_{ij} = \delta_{ij}$  (Kronecker delta:  $\delta_{ij} = 1$  if  $i = j$ , else 0):

$$\|I\| = \sum_{i=1}^n \sum_{j=1}^n |\delta_{ij}| = \sum_{i=1}^n 1 = n$$

Since  $\|I\| = n \neq 1$  for  $n > 1$ , this norm is NOT subordinate to any vector norm.

#### Part 3: Does it conform to Equation (9)?

Equation (9) requires  $\|I\| = 1$ . As shown above,  $\|I\| = n$ .

No, it does NOT conform to Equation (9).

#### Part 4: Does it conform to Inequality (10)?

We need to check if  $\|AB\| \leq \|A\|\|B\|$ .

*Step 1:* We recall that for matrix multiplication, the  $(i, k)$  entry of  $AB$  is:

$$(AB)_{ik} = \sum_{j=1}^n a_{ij}b_{jk}$$

*Step 2:* Using the triangle inequality for absolute values, we have:

$$|(AB)_{ik}| = \left| \sum_{j=1}^n a_{ij}b_{jk} \right| \leq \sum_{j=1}^n |a_{ij}b_{jk}| = \sum_{j=1}^n |a_{ij}||b_{jk}|$$

*Step 3:* We now compute  $\|AB\|$  by summing over all entries:

$$\begin{aligned} \|AB\| &= \sum_{i=1}^n \sum_{k=1}^n |(AB)_{ik}| \\ &\leq \sum_{i=1}^n \sum_{k=1}^n \sum_{j=1}^n |a_{ij}||b_{jk}| \quad (\text{from Step 2}) \end{aligned}$$

*Step 4:* We rearrange the triple sum. Notice that  $|a_{ij}|$  doesn't depend on  $k$ , and  $|b_{jk}|$  doesn't depend on  $i$ :

$$\begin{aligned} \sum_{i=1}^n \sum_{k=1}^n \sum_{j=1}^n |a_{ij}||b_{jk}| &= \sum_{i=1}^n \sum_{j=1}^n |a_{ij}| \left( \sum_{k=1}^n |b_{jk}| \right) \\ &= \sum_{i=1}^n \sum_{j=1}^n |a_{ij}| \cdot \sum_{k=1}^n \sum_{j=1}^n |b_{jk}| \\ &= \left( \sum_{i=1}^n \sum_{j=1}^n |a_{ij}| \right) \left( \sum_{j=1}^n \sum_{k=1}^n |b_{jk}| \right) \\ &= \|A\| \cdot \|B\| \end{aligned}$$

*Step 5:* Combining Steps 3 and 4, we obtain:  $\|AB\| \leq \|A\|\|B\|$ .

Yes, it DOES conform to Inequality (10).

---

## Problem 5

### Problem Statement (Textbook 4.4.11):

Show that for the vector norm  $\|\mathbf{x}\|_1$  defined in Equation (5), the subordinate matrix norm is

$$\|A\|_1 = \max_{1 \leq j \leq n} \sum_{i=1}^n |a_{ij}|$$

where

$$\|\mathbf{x}\|_1 = \sum_{i=1}^n |x_i| \quad (\text{Equation 5})$$


---

### Solution:

We need to show that the subordinate matrix norm induced by  $\|\mathbf{x}\|_1$  is the maximum absolute column sum.

Recall that the subordinate matrix norm is defined as:

$$\|A\|_1 = \max_{\mathbf{x} \neq \mathbf{0}} \frac{\|A\mathbf{x}\|_1}{\|\mathbf{x}\|_1}$$

Equivalently, we can write:

$$\|A\|_1 = \max_{\|\mathbf{x}\|_1=1} \|A\mathbf{x}\|_1$$



### Part 1: Upper Bound ( $\|A\|_1 \leq \max_j \sum_i |a_{ij}|$ )

For any vector  $\mathbf{x}$  with  $\|\mathbf{x}\|_1 = 1$ , we compute:

$$\begin{aligned}
 \|A\mathbf{x}\|_1 &= \sum_{i=1}^n \left| \sum_{j=1}^n a_{ij} x_j \right| \\
 &\leq \sum_{i=1}^n \sum_{j=1}^n |a_{ij}| |x_j| \quad (\text{triangle inequality}) \\
 &= \sum_{j=1}^n |x_j| \sum_{i=1}^n |a_{ij}| \quad (\text{interchange sums}) \\
 &\leq \sum_{j=1}^n |x_j| \cdot \max_{1 \leq k \leq n} \sum_{i=1}^n |a_{ik}| \\
 &= \left( \sum_{j=1}^n |x_j| \right) \cdot \max_{1 \leq k \leq n} \sum_{i=1}^n |a_{ik}| \\
 &= \|\mathbf{x}\|_1 \cdot \max_{1 \leq k \leq n} \sum_{i=1}^n |a_{ik}| \\
 &= \max_j \sum_{i=1}^n |a_{ij}|
 \end{aligned}$$

This holds for all  $\mathbf{x}$  with  $\|\mathbf{x}\|_1 = 1$ , so:

$$\|A\|_1 \leq \max_j \sum_{i=1}^n |a_{ij}|$$

### Part 2: Lower Bound (Equality is Achieved)

Let  $j^*$  be the column index that achieves the maximum:

$$j^* = \arg \max_{1 \leq j \leq n} \sum_{i=1}^n |a_{ij}|$$

Now choose the test vector  $\mathbf{x}^* = \mathbf{e}_{j^*}$  (the  $j^*$ -th standard basis vector), which has  $\|\mathbf{x}^*\|_1 = 1$ .

Then:

$$A\mathbf{x}^* = A\mathbf{e}_{j^*} = \begin{bmatrix} a_{1,j^*} \\ a_{2,j^*} \\ \vdots \\ a_{n,j^*} \end{bmatrix}$$

Thus:

$$\|A\mathbf{x}^*\|_1 = \sum_{i=1}^n |a_{i,j^*}| = \max_j \sum_{i=1}^n |a_{ij}|$$

This shows that the upper bound is achieved, so:

$$\|A\|_1 \geq \max_j \sum_{i=1}^n |a_{ij}|$$

## Conclusion

Combining Parts 1 and 2:

$$\|A\|_1 = \max_{1 \leq j \leq n} \sum_{i=1}^n |a_{ij}|$$

The subordinate matrix norm induced by the 1-norm is the **maximum absolute column sum**.

---

## Problem 6

### Problem Statement (Textbook 4.4.33):

For any  $n \times n$  matrix  $A$ , define

$$\|A\|_F = \left( \sum_{i=1}^n \sum_{j=1}^n a_{ij}^2 \right)^{1/2}$$

(This is called the **Frobenius** norm.) Is this a subordinate matrix norm? Answer the same question for  $\|A\| = \max_{1 \leq i, j \leq n} |a_{ij}|$ . Prove that these equations define norms on the vector space of all  $n \times n$  matrices.

---

**Solution:**

## Problem 7

### Problem Statement (Textbook 4.6.1b):

Program the Gauss-Seidel method and test it on the following system:

$$\begin{aligned}3x + y + z &= 5 \\3x + y - 5z &= -1 \\x + 3y - z &= 3\end{aligned}$$

Analyze what happens when these systems are solved by simple Gaussian elimination without pivoting.

---

**Solution:**

**System Matrix:**

$$A = \begin{bmatrix} 3 & 1 & 1 \\ 3 & 1 & -5 \\ 1 & 3 & -1 \end{bmatrix}, \quad b = \begin{bmatrix} 5 \\ -1 \\ 3 \end{bmatrix}$$

**True Solution** (using direct method):  $x = y = z = 1$

### Part 1: Gauss-Seidel Method

The Gauss-Seidel iteration is:

$$\begin{aligned}x^{(k+1)} &= \frac{1}{3}(5 - y^{(k)} - z^{(k)}) \\y^{(k+1)} &= \frac{1}{1}(-1 - 3x^{(k+1)} + 5z^{(k)}) \\z^{(k+1)} &= \frac{1}{-1}(3 - x^{(k+1)} - 3y^{(k+1)})\end{aligned}$$

**Diagonal Dominance Check:**

- Row 1:  $|a_{11}| = 3 > |1| + |1| = 2$
- Row 2:  $|a_{22}| = 1 \not> |3| + |-5| = 8$
- Row 3:  $|a_{33}| = |-1| = 1 \not> |1| + |3| = 4$

Matrix is NOT strictly diagonally dominant

**Convergence Results** (see hw05\_p07.jl):

- Starting from  $(1, 1, 1)$ : **Converges in 1 iteration** (already at solution!)
- Starting from  $(0.999, 0.999, 0.999)$ : **Diverges** (NaN)
- Starting from  $(1.01, 1.01, 1.01)$ : **Diverges** (NaN)

- Starting from (0.9, 0.9, 0.9) or (1.1, 1.1, 1.1): **Diverges** (NaN)
- Starting from (0, 0, 0), (5, -1, 3), or (10, 10, 10): **Diverges** (NaN)

**Key Observation:** The method only "converges" when starting exactly at the solution. Even tiny perturbations (0.001 away) cause catastrophic divergence to NaN. This demonstrates **complete instability**.

## Part 2: Gaussian Elimination without Pivoting

Forward elimination proceeds:

**Step 1:** Eliminate column 1 using pivot  $a_{11} = 3$ :

$$\begin{bmatrix} 3 & 1 & 1 \\ 0 & 0 & -6 \\ 0 & 8/3 & -4/3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ -6 \\ 4/3 \end{bmatrix}$$

**Step 2:** Attempt to eliminate column 2 using pivot  $a_{22}$ :

$a_{22} = 0 \text{ (ZERO PIVOT!)}$

**Analysis:**

- Gaussian elimination **fails** due to zero pivot at position (2, 2)
- Rows 2 and 3 need to be swapped (partial pivoting required)
- Without pivoting, the method cannot proceed
- The system has determinant  $\det(A) = 48 \neq 0$ , so the matrix is invertible
- Failure is due to *method limitation*, not matrix singularity

**Summary:** Both methods encounter difficulties with this system:

- **Gauss-Seidel:** Diverges for most starting points (not diagonally dominant)
- **Gaussian Elimination:** Fails without pivoting (zero pivot encountered)

**Implementation:** See Appendix A for complete Julia code listings:

- Gauss-Seidel method: Appendix
- Gaussian elimination: Appendix

## Problem 8

### Problem Statement (Textbook 6.1.1d):

Find the polynomial of least degree that interpolates these sets of data:

$$\begin{array}{cccc} x & 3 & 7 & 12 \\ y & 12 & 146 & 2 \end{array}$$

### Solution:

We find the polynomial of least degree interpolating the data points  $(3, 12)$ ,  $(7, 146)$ , and  $(12, 2)$ .

Since we have 3 data points, the interpolating polynomial has degree at most  $n - 1 = 2$  (quadratic).

### Method: Vandermonde Matrix

We seek a polynomial  $P(x) = a_0 + a_1x + a_2x^2$  such that  $P(x_i) = y_i$  for  $i = 1, 2, 3$ .

This gives the linear system (see Textbook p. 313, Equation 12) [1]:

$$\begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

Substituting our data:

$$\begin{bmatrix} 1 & 3 & 9 \\ 1 & 7 & 49 \\ 1 & 12 & 144 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 12 \\ 146 \\ 2 \end{bmatrix}$$

Solving this system, we obtain:

$$\begin{aligned} a_0 &= -\frac{4211}{18} \approx -233.8667 \\ a_1 &= \frac{1849}{18} \approx 102.7222 \\ a_2 &= -\frac{125}{18} \approx -6.9222 \end{aligned}$$

**Implementation:** See Appendix B (??) for Julia code implementation.

**Final Answer:**

$$P(x) = -\frac{125}{18}x^2 + \frac{1849}{18}x - \frac{4211}{18}$$

Or in decimal form:

$$P(x) = -6.9222x^2 + 102.7222x - 233.8667$$

### Verification:

- $P(3) = 12.000$

- $P(7) = 146.000$
- $P(12) = 2.000$

Maximum interpolation error:  $5.68 \times 10^{-14}$  (numerical precision).

---

## Problem 9

### Problem Statement (Textbook 6.1.8):

Discuss the problem of determining a polynomial of degree at most 2 for which  $p(0)$ ,  $p(1)$ , and  $p(\xi)$  are prescribed,  $\xi$  being any preassigned point.

### Solution:

We seek a polynomial  $p(x) = a_0 + a_1x + a_2x^2$  (degree at most 2) such that:

$$p(0) = y_0, \quad p(1) = y_1, \quad p(\xi) = y_\xi$$

where  $y_0, y_1, y_\xi$  are prescribed values and  $\xi$  is a given point.

### Main Case: Three Distinct Points Required

For a polynomial of degree at most 2, we need three distinct interpolation points. Thus, we require:

$$\xi \neq 0 \text{ and } \xi \neq 1$$

The Vandermonde system is:

$$\begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ 1 & \xi & \xi^2 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} y_0 \\ y_1 \\ y_\xi \end{bmatrix}$$

The determinant is:

$$\det(V) = \xi(\xi - 1)$$

Since  $\xi \neq 0$  and  $\xi \neq 1$ , we have  $\det(V) \neq 0$ , so the system has a **unique solution**.

**Conclusion:** When  $\xi \notin \{0, 1\}$ , there exists a **unique polynomial of degree at most 2** satisfying all three conditions.

### Degenerate Cases

**Note:** If  $\xi = 0$  or  $\xi = 1$ , we only have two distinct interpolation points, which contradicts the requirement for a degree-2 polynomial. In these cases:

- The problem reduces to linear interpolation (degree at most 1)
- The prescribed values must be consistent (e.g., if  $\xi = 0$ , then  $y_0 = y_\xi$ )
- The resulting polynomial is the line through the two distinct points

These cases defeat the original question's intent of determining a degree-2 polynomial.

**Key Insight:** The problem is well-posed if and only if we have three distinct interpolation points, i.e.,  $\xi \notin \{0, 1\}$ .



## Appendix

### A. Problem 7 Code Listings

#### A.1 Gauss-Seidel Method Implementation

```

73  function gauss_seidel(A, b, x0; max_iter=100, tol=1e-10)
74      n = length(b)
75      x = copy(x0)
76      history = [copy(x)]
77
78      for iter in 1:max_iter
79          x_old = copy(x)
80
81          # Gauss-Seidel iteration
82          for i in 1:n
83              sum_val = b[i]
84              for j in 1:n
85                  if j ≠ i
86                      sum_val -= A[i, j] * x[j]
87                  end
88              end
89              x[i] = sum_val / A[i, i]
90          end
91
92          push!(history, copy(x))
93
94          # Check convergence
95          error = norm(x - x_old, Inf)
96          if error < tol
97              return (x, true, iter, history)
98          end
99      end
100
101      return (x, false, max_iter, history)
102  end

```

Figure 1: Julia code implementing the Gauss-Seidel iterative method for Problem 7.

## A.2 Gaussian Elimination without Pivoting

```

166 ✓ function gaussian_elimination_no_pivot(A, b)
167     n = length(b)
168     A_work = copy(A)
169     b_work = copy(b)
170
171     # Forward elimination
172 ✓   for k in 1:n-1
173       println(@sprintf("\nStep %d: Eliminate column %d", k, k))
174
175 ✓       if abs(A_work[k, k]) < 1e-14
176           println(" ⚠ WARNING: Pivot element is near zero!")
177           println(@sprintf("    a_%d%d = %.2e", k, k, A_work[k, k]))
178           return (zeros(n), false, A_work, b_work)
179       end
180
181       println(@sprintf("    Pivot: a_%d%d = %.6f", k, k, A_work[k, k]))
182
183 ✓       for i in k+1:n
184 ✓           if abs(A_work[i, k]) > 1e-14
185               multiplier = A_work[i, k] / A_work[k, k]
186               println(@sprintf("    Row %d: subtract %.6f × Row %d", i,
187                               multiplier, k))
187
188               A_work[i, k:n] -= multiplier * A_work[k, k:n]
189               b_work[i] -= multiplier * b_work[k]
190           end
191       end
192
193       println("\n Matrix after step $k:")
194       display(A_work)
195       println("\n RHS after step $k:")
196       display(b_work)
197   end

```

Figure 2: Julia code implementing Gaussian elimination without pivoting (Part 1).

```

199     # Back substitution
200     x = zeros(n)
201 ✓   for i in n:-1:1
202       x[i] = (b_work[i] - dot(A_work[i, i+1:n], x[i+1:n])) / A_work[i, i]
203   end
204
205   return (x, true, A_work, b_work)
206 end

```

Figure 3: Julia code implementing Gaussian elimination without pivoting (Part 2).

## References

- [1] David Kincaid and Ward Cheney. *Numerical Analysis: Mathematics of Scientific Computing*. American Mathematical Society, Providence, RI, 3rd edition, 2009. Available from ProQuest Ebook Central. <https://ebookcentral.proquest.com/lib/wsu/detail.action?docID=4717637>.
- [2] Alexander Panchenko. Math 448 lecture notes. Washington State University, Fall 2025.