

Fall 2025 CPTS530 Homework 5
MATH 448-548
Numerical Analysis

Aryan Ritwajeet Jha

November 2025

Contents

Problem 1	2
Problem 2	4
Problem 3	5
Problem 4	6
Problem 5	7
Problem 6	8
Problem 7	9
Problem 8	10
Problem 9	11
Appendix	12
References	13

Problem 1

Problem Statement (Textbook 4.4.1):

Show that the norms $\|\mathbf{x}\|_\infty$, $\|\mathbf{x}\|_2$, and $\|\mathbf{x}\|_1$ satisfy the postulates (1), (2), and (3) for norms.

Norm Postulates [1, 2]:

On a vector space V , a **norm** is a function $\|\cdot\|$ from V to the set of nonnegative reals that obeys these three postulates:

$$\|x\| > 0 \text{ if } x \neq 0, x \in V \quad (1)$$

$$\|\lambda x\| = |\lambda| \|x\| \text{ if } \lambda \in \mathbb{R}, x \in V \quad (2)$$

$$\|x + y\| \leq \|x\| + \|y\| \text{ if } x, y \in V \quad (\text{triangle inequality}) \quad (3)$$

Solution:

We verify postulates (1), (2), and (3) for $\|x\|_\infty = \max_i |x_i|$, $\|x\|_2 = \sqrt{\sum_i x_i^2}$, and $\|x\|_1 = \sum_i |x_i|$.

(a) ∞ -norm: $\|x\|_\infty = \max_{1 \leq i \leq n} |x_i|$

(1) If $x \neq 0$, then $\exists i$ with $x_i \neq 0$, so $|x_i| > 0$. Thus $\|x\|_\infty = \max_i |x_i| > 0$.

(2) $\|\lambda x\|_\infty = \max_i |\lambda x_i| = \max_i |\lambda| |x_i| = |\lambda| \max_i |x_i| = |\lambda| \|x\|_\infty$.

(3) $\|x + y\|_\infty = \max_i |x_i + y_i| \leq \max_i (|x_i| + |y_i|) \leq \max_i |x_i| + \max_i |y_i| = \|x\|_\infty + \|y\|_\infty$.

(b) 2-norm (Euclidean): $\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$

(1) If $x \neq 0$, then $\exists i$ with $x_i \neq 0$, so $\sum_i x_i^2 > 0$. Thus $\|x\|_2 = \sqrt{\sum_i x_i^2} > 0$.

(2) $\|\lambda x\|_2 = \sqrt{\sum_i (\lambda x_i)^2} = \sqrt{\lambda^2 \sum_i x_i^2} = |\lambda| \sqrt{\sum_i x_i^2} = |\lambda| \|x\|_2$.

(3) Expanding $\|x + y\|_2^2$:

$$\begin{aligned} \|x + y\|_2^2 &= \sum_i (x_i + y_i)^2 = \sum_i x_i^2 + 2 \sum_i x_i y_i + \sum_i y_i^2 \\ &= \|x\|_2^2 + 2 \sum_i x_i y_i + \|y\|_2^2 \end{aligned}$$

Since $\sum_i x_i y_i \leq \|x\|_2 \|y\|_2$ (dot product bounded by product of norms):

$$\|x + y\|_2^2 \leq \|x\|_2^2 + 2\|x\|_2 \|y\|_2 + \|y\|_2^2 = (\|x\|_2 + \|y\|_2)^2$$

Taking square roots: $\|x + y\|_2 \leq \|x\|_2 + \|y\|_2$.

(c) 1-norm: $\|x\|_1 = \sum_{i=1}^n |x_i|$

(1) If $x \neq 0$, then $\exists i$ with $x_i \neq 0$, so $|x_i| > 0$. Thus $\|x\|_1 = \sum_i |x_i| > 0$.

(2) $\|\lambda x\|_1 = \sum_i |\lambda x_i| = \sum_i |\lambda| |x_i| = |\lambda| \sum_i |x_i| = |\lambda| \|x\|_1$.

(3) For each term, the triangle inequality for absolute values gives $|x_i + y_i| \leq |x_i| + |y_i|$.

Summing over all i :

$$\|x + y\|_1 = \sum_i |x_i + y_i| \leq \sum_i (|x_i| + |y_i|) = \sum_i |x_i| + \sum_i |y_i| = \|x\|_1 + \|y\|_1$$

All three norms satisfy postulates (1), (2), and (3).

Problem 2

Problem Statement (Textbook 4.4.2):

Show that $\|\mathbf{x}\|_\infty \leq \|\mathbf{x}\|_2 \leq \|\mathbf{x}\|_1$ for all $\mathbf{x} \in \mathbb{R}^n$, and that equalities can occur, even for nonzero vectors.

Solution:

Problem 3

Problem Statement (Textbook 4.4.3):

(Continuation) Show that $\|\mathbf{x}\|_1 \leq n\|\mathbf{x}\|_\infty$ and $\|\mathbf{x}\|_2 \leq \sqrt{n}\|\mathbf{x}\|_\infty$ for $\mathbf{x} \in \mathbb{R}^n$.

Solution:

Problem 4

Problem Statement (Textbook 4.4.8):

Define

$$\|A\| = \sum_{i=1}^n \sum_{j=1}^n |a_{ij}|$$

Show that this is a matrix norm (that is, a norm on the linear space of all $n \times n$ matrices). Show that it is not subordinate to any vector norm. Does it conform to Equation (9) and Inequality (10)?

$$\|I\| = 1 \quad (\text{Equation 9})$$

$$\|AB\| \leq \|A\|\|B\| \quad (\text{Inequality 10})$$

Solution:

Problem 5

Problem Statement (Textbook 4.4.11):

Show that for the vector norm $\|\mathbf{x}\|_1$ defined in Equation (5), the subordinate matrix norm is

$$\|A\|_1 = \max_{1 \leq j \leq n} \sum_{i=1}^n |a_{ij}|$$

where

$$\|\mathbf{x}\|_1 = \sum_{i=1}^n |x_i| \quad (\text{Equation 5})$$

Solution:

Problem 6

Problem Statement (Textbook 4.4.33):

For any $n \times n$ matrix A , define

$$\|A\|_F = \left(\sum_{i=1}^n \sum_{j=1}^n a_{ij}^2 \right)^{1/2}$$

(This is called the **Frobenius** norm.) Is this a subordinate matrix norm? Answer the same question for $\|A\| = \max_{1 \leq i, j \leq n} |a_{ij}|$. Prove that these equations define norms on the vector space of all $n \times n$ matrices.

Solution:

Problem 7

Problem Statement (Textbook 4.6.1b):

Program the Gauss-Seidel method and test it on the following system:

$$3x + y + z = 5$$

$$3x + y - 5z = -1$$

$$x + 3y - z = 3$$

Analyze what happens when these systems are solved by simple Gaussian elimination without pivoting.

Solution:

Problem 8

Problem Statement (Textbook 6.1.1d):

Find the polynomial of least degree that interpolates these sets of data:

x	3	7	12
y	12	146	2

Solution:

Problem 9

Problem Statement (Textbook 6.1.8):

Discuss the problem of determining a polynomial of degree at most 2 for which $p(0)$, $p(1)$, and $p(\xi)$ are prescribed, ξ being any preassigned point.

Solution:

Appendix

References

- [1] David Kincaid and Ward Cheney. *Numerical Analysis: Mathematics of Scientific Computing*. American Mathematical Society, Providence, RI, 3rd edition, 2009. Available from ProQuest Ebook Central. <https://ebookcentral.proquest.com/lib/wsu/detail.action?docID=4717637>.
- [2] Alexander Panchenko. Math 448 lecture notes. Washington State University, Fall 2025.