

Fall 2025 CPTS530 Final Project
MATH 448-548
Numerical Analysis

Aryan Ritwajeet Jha

11 December 2025

Contents

Problem 1: Orthogonal Matching Pursuit	2
Problem 2: Fourth-Order Runge-Kutta Method	4
Problem 3: Adams-Bashforth-Moulton Method	5
Appendix	6

Problem 1: Orthogonal Matching Pursuit (OMP)

Problem Statement:

Part 0: Preliminary Information and Data

In part 2, use the matrix

$$A = \begin{bmatrix} 1 & -1/2 & -1/2 \\ 0 & \sqrt{3}/2 & -\sqrt{3}/2 \\ 1 & 3 & 3 \end{bmatrix}$$

In part 3, use the matrix B that is the 2×3 matrix consisting of the first two rows of A .

Singular Value Decomposition (SVD) and Pseudo-Inverse

Given a matrix $M_{m \times n}$ with $\text{rank}(M) = r$, write

$$M = U \Sigma V^T$$

where $U_{m \times m}$ and $V_{n \times n}$ are orthogonal matrices, and $\Sigma_{m \times n}$ is a diagonal matrix with nonzero diagonal entries $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r > 0$ called singular values of M .

Definition. Given M as above, the pseudo-inverse M^+ is defined by

$$M^+ = V \Sigma^+ U^T$$

where $\Sigma_{n \times m}^+$ is the diagonal matrix whose non-zero entries are $\frac{1}{\sigma_1}, \frac{1}{\sigma_2}, \dots, \frac{1}{\sigma_r}$.

Thresholding Operator

The **hard thresholding operator** H_s is defined as follows:

$$\mathbf{x} \rightarrow H_s(\mathbf{x}) \in \mathbb{R}^n$$

$H_s(\mathbf{x})$ sets all entries of $\mathbf{x}$ to zero except for s largest (in absolute value) entries.

The Matrix $A_{\mathcal{S}}$

Given a matrix A and an index set $\mathcal{S}$, denote by $A_{\mathcal{S}}$ the matrix consisting of columns of A with index in $\mathcal{S}$.

Assignments and Questions

1. Implement the pseudocode for the Orthogonal Matching Pursuit (OMP). The pseudocode is contained in a separate handwritten note attached to the Final project assignment. You may use the standard Matlab (or other language) commands to compute SVD and pseudo-inverse.

2. Run the code for $\mathbf{b} = A\mathbf{e}_1$ with $\mathbf{e}_1 = (1, 0, 0)^T$. See how well the obtained vector $\mathbf{x}$ matches $\mathbf{e}_1$.
3. Run the code again, this time using the matrix B instead of A .
4. Compare the results obtained in parts 2, 3. Explain the connection with compressed sensing and sparse recovery.

Solution:

Problem 2: Fourth-Order Runge-Kutta Method

Problem Statement:

Solve Computer problem 1 in Section 8.3. Write your own program from scratch. Then compare with the exact analytic solution that can be obtained by the method of integrating factors studied in Math 315. The exact solution is

$$x(t) = 12 \frac{e^t}{(e^t + 1)^2}$$

Computer Problem 8.3.1

Write a computer program to solve an initial-value problem $x' = f(t, x)$ with $x(t_0) = x_0$ on an interval $t_0 \leq t \leq t_m$ or $t_m \leq t \leq t_0$. Use the fourth-order Runge-Kutta method. Test it on this example:

$$\begin{aligned}(e^t + 1)x' + xe^t - x &= 0 \\ x(0) &= 3\end{aligned}$$

Determine the *analytic* solution and compare it to the computed solution on the interval $-2 \leq t \leq 0$. Use $h = -0.01$.

Solution:

Problem 3: Adams-Bashforth-Moulton Method

Problem Statement:

Solve Computer problem 3 in Section 8.4.

Computer Problem 8.4.3

Compute the solution of

$$\begin{aligned}y' &= -2xy^2 \\ y(0) &= 1\end{aligned}$$

at $x = 1.0$ using $h = 0.25$ and the fourth-order Adams-Bashforth-Moulton method (Problems 8.4.4–5, p. 555) together with the fourth-order Runge-Kutta method. Give the computed solution to five significant digits at 0.25, 0.5, 0.75, and 1.0. Compare your results to the exact solution $y = 1/(1 + x^2)$.

Related Problems from Section 8.4

Problem 8.4.1: Derive the first-order (one-step) Adams-Moulton formula and verify that it is equivalent to the trapezoid rule. (See Section 7.2, p. 481.)

Problem 8.4.2: Verify the correctness of the system of equations in (6).

Problem 8.4.3: Use the method of undetermined coefficients to derive Equation (8).

Problem 8.4.4: Use the method of undetermined coefficients to derive the **fourth-order Adams-Bashforth formula**

$$x_{n+1} = x_n + \frac{h}{24}[55f_n - 59f_{n-1} + 37f_{n-2} - 9f_{n-3}]$$

Problem 8.4.5: Derive the **fourth-order Adams-Moulton formula**

$$x_{n+1} = x_n + \frac{h}{24}[9f_{n+1} + 19f_n - 5f_{n-1} + f_{n-2}]$$

Solution:

Appendix

This appendix contains supplementary material, proofs, and code implementations for the final project problems.