

Fall 2025 CPTS530 Midterm Project
MATH 448-548
Numerical Analysis

Aryan Ritwajeet Jha

14 October 2025

Contents

Problem 1	2
Problem 2	3
Problem 3	4
Problem 4	5
Problem 4.1	5
Problem 4.2	6
Problem 4.3	7
Problem 4.4	8
Problem 4.5	9
Problem 4.6	10
Problem 5	11
Problem 5.1	11
Problem 5.2	13
Problem 5.3	15
Problem 5.4	17

Problem 1

Problem Statement: Use Taylor theorem to derive the error term for the approximate formula

$$f'(x) \approx \frac{1}{2h}(-3f(x) + 4f(x+h) - f(x+2h)).$$

Solution:

We are given the finite-difference formula:

$$f'(x) \approx \frac{1}{2h} [-3f(x) + 4f(x+h) - f(x+2h)]$$

Expand $f(x+h)$ and $f(x+2h)$ about x using Taylor's theorem, keeping terms up to h^2 and representing higher-order terms as k_1h^3 and k_2h^4 :

$$\begin{aligned} f(x+h) &= f(x) + hf'(x) + \frac{h^2}{2}f''(x) + k_1h^3 \\ f(x+2h) &= f(x) + 2hf'(x) + 2h^2f''(x) + k_2h^3 \end{aligned}$$

where k_1, k_2 are finite constants depending on third order derivatives of f at x .

Substitute into the formula:

$$\begin{aligned} &\frac{1}{2h} [-3f(x) + 4f(x+h) - f(x+2h)] \\ &= \frac{1}{2h} \left[-3f(x) + 4 \left(f(x) + hf'(x) + \frac{h^2}{2}f''(x) + k_1h^3 \right) \right. \\ &\quad \left. - (f(x) + 2hf'(x) + 2h^2f''(x) + k_2h^3) \right] \end{aligned}$$

Simplifying the terms in the square brackets:

$$= \frac{1}{2h} [2hf'(x) + (4k_1 - k_2)h^3]$$

Now, divide each term by $2h$:

$$= f'(x) + \frac{1}{2}(4k_1 - k_2)h^2$$

Thus, the leading error term is $O(h^2)$:

$$\text{error} \sim O(h^2)$$

This shows the method is second-order accurate. Note that I've already checked if the third order term vanishes if we write out the Taylor's polynomial to the third order, and it does not.

Problem 2

Problem Statement: Consider the following variation of the Newton's method:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_0)}.$$

Find constants C and s such that

$$e_{n+1} = C e_n^s.$$

Solution:

Problem 3

Problem Statement: Consider the linear system $\mathbf{Ax} = \mathbf{b}$, where

$$\mathbf{A} = \begin{bmatrix} 6.25 & -1 & 0.5 \\ -1 & 5 & 2.12 \\ 0.5 & 2.12 & 3.6 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 7.5 \\ -8.68 \\ -0.24 \end{bmatrix}.$$

Write a computer program for LU-factorization with a unit lower triangular $\mathbf{L}$ (meaning that the diagonal entries should be equal to one). Then write a program for the Cholesky factorization.

WARNING: avoid using shortcuts. The programming should be done “from scratch”.

Solution:

Problem 4

Problem Statement: Consider the equation

$$f(x) = 0,$$

where

$$f(x) = x - \frac{1}{2} \left(\cos(x/2) - \left| x - \frac{1}{2} \right| \right).$$

Do the following.

Part 1

Rewrite the equation as a fixed point problem for a function $T(x)$ related to $f(x)$. Prove that $T(x)$ (i) maps the interval $[0.45, 0.55]$ into itself, and (ii) $T(x)$ is contractive on this interval. Based on the theorems in the books and notes, make conclusions about existence and uniqueness of a solution to $f(x) = 0$.

Solution:

Problem Statement: Consider the equation

$$f(x) = 0,$$

where

$$f(x) = x - \frac{1}{2} \left(\cos(x/2) - \left| x - \frac{1}{2} \right| \right).$$

Do the following.

Part 2

Write the computer program implementing the fixed point iteration $x_{n+1} = T(x_n)$. Discuss the choice of an initial guess x_0 . Calculate the solution to within 10 decimal places. Note the number of iterations needed.

Solution:

Problem Statement: Consider the equation

$$f(x) = 0,$$

where

$$f(x) = x - \frac{1}{2} \left(\cos(x/2) - \left| x - \frac{1}{2} \right| \right).$$

Do the following.

Part 3

Use a priori and a posteriori error estimates from the notes to estimate the number of iterations needed to obtain the above accuracy. Which estimate is more accurate?

Solution:

Problem Statement: Consider the equation

$$f(x) = 0,$$

where

$$f(x) = x - \frac{1}{2} \left(\cos(x/2) - \left| x - \frac{1}{2} \right| \right).$$

Do the following.

Part 4

Implement the Newton's method on a computer for the same equation starting with x_0 within the interval $[0.45, 0.55]$. Discuss the theoretical difficulty related to non-existence of $f'(x)$ for certain point(s) within the interval. How would you set up the Newton's method to resolve the problem?

Solution:

Problem Statement: Consider the equation

$$f(x) = 0,$$

where

$$f(x) = x - \frac{1}{2} \left(\cos(x/2) - \left| x - \frac{1}{2} \right| \right).$$

Do the following.

Part 5

Note the number of iterations needed to calculate the solution to within 10 decimal places using the Newton's method. Discuss the speed of convergence of the Newton's method and compare it with the convergence speed of the fixed point iterations.

Solution:

Problem Statement: Consider the equation

$$f(x) = 0,$$

where

$$f(x) = x - \frac{1}{2} \left(\cos(x/2) - \left| x - \frac{1}{2} \right| \right).$$

Do the following.

Part 6

Implement the alternative error control strategy proposed by you in HW 3, Problem 2. Compare the results with the ones obtained using a priori and a posteriori error estimates.

Solution:

Problem 5

Problem Statement: Implement and analyze the Ridders' method for solving the equation $f(x) = 0$. For guidance, you might wish to read the Wikipedia page on the Ridders' method. Also feel free to consult other sources.

Brief Description of the method


Given the initial bracketing interval $[x_0, x_2]$, containing **one** solution, and such that $f(x_0)$ and $f(x_2)$ have opposite signs, compute the midpoint $x_1 = (x_0 + x_2)/2$. Then define a new function $h(x) = f(x)e^{ax}$. Find the parameter a that ensures $h(x_1) = \frac{1}{2}(h(x_0) + h(x_2))$. Then define x_3 as the x -intercept of the line passing through $(x_0, h(x_0))$ and $(x_2, h(x_2))$. Use x_3 as one of the endpoints of the next interval bracketing the solution. The other endpoint is x_1 if $f(x_1)f(x_3) < 0$. Otherwise, choose either x_0 or x_2 based on the requirement that the sign of $f(x)$ at the chosen point must be opposite to the sign of $f(x_3)$. Continue until the desired accuracy is reached.

Part 1

Show that a is uniquely defined and give the equation for finding it.

Solution:

Note: Thanks to the wikipedia article which itself cites the original paper by Ridders, it was easier to derive all the relevant formulae. I have chosen my target formulae and required derivations to align with those, with little variation.



Let x_0 and x_2 be the endpoints of the bracketing interval, and $x_1 = \frac{x_0 + x_2}{2}$ the midpoint. Let $d = x_2 - x_0$.

Define $f_0 = f(x_0)$, $f_1 = f(x_1)$, $f_2 = f(x_2)$.

Ridders' method introduces a function $h(x) = f(x)e^{ax}$, and chooses a so that:

$$h(x_1) = \frac{1}{2}(h(x_0) + h(x_2))$$

$$f_1 e^{ax_1} = \frac{1}{2}(f_0 e^{ax_0} + f_2 e^{ax_2})$$

We know that $x_1 = x_0 + \frac{d}{2}$ and $x_2 = x_0 + d$.

Substitute and rearrange:

$$f_1 e^{a(x_0+d/2)} = \frac{1}{2} (f_0 e^{ax_0} + f_2 e^{a(x_0+d)})$$

Dividing both sides by e^{ax_0} :

$$2f_1 e^{ad/2} = f_0 + f_2 e^{ad}$$

$$2f_1 e^{ad/2} - f_2 e^{ad} - f_0 = 0$$

Let $y = e^{ad/2}$, so $e^{ad} = y^2$.

This gives a quadratic in y :

$$f_2 y^2 - 2f_1 y + f_0 = 0$$

Solving for y :

$$y = \frac{2f_1 \pm \sqrt{(2f_1)^2 - 4f_2 f_0}}{2f_2}$$

$$y = \frac{f_1 \pm \sqrt{f_1^2 - f_2 f_0}}{f_2}$$

Therefore,

$$a = \frac{2}{d} \ln \left(\frac{f_1 \pm \sqrt{f_1^2 - f_2 f_0}}{f_2} \right)$$

We note that since $y = e^{ad/2}$, its value must be nonnegative. Since clearly $f_1^2 - f_2 f_0 > f_1^2$ ($f_2 * f_0 < 0$), the sign is chosen so that y is positive:

$$\text{If } f_2 > 0 : \quad a = \frac{2}{d} \ln \left(\frac{f_1 + \sqrt{f_1^2 - f_2 f_0}}{f_2} \right)$$

$$\text{else If } f_2 < 0 : \quad a = \frac{2}{d} \ln \left(\frac{f_1 - \sqrt{f_1^2 - f_2 f_0}}{f_2} \right)$$

Or, more concisely:

$$a = \frac{2}{d} \ln \left(\frac{f_1 + \text{sgn}(f_2) \sqrt{f_1^2 - f_2 f_0}}{f_2} \right)$$

or equivalently:

where $\text{sgn}(f_2)$ is the sign of f_2 .

Problem Statement: Implement and analyze the Ridders' method for solving the equation $f(x) = 0$. For guidance, you might wish to read the Wikipedia page on the Ridders' method. Also feel free to consult other sources.


Brief Description of the method

Given the initial bracketing interval $[x_0, x_2]$, containing **one** solution, and such that $f(x_0)$ and $f(x_2)$ have opposite signs, compute the midpoint $x_1 = (x_0 + x_2)/2$. Then define a new function $h(x) = f(x)e^{ax}$. Find the parameter a that ensures $h(x_1) = \frac{1}{2}(h(x_0) + h(x_2))$. Then define x_3 as the x -intercept of the line passing through $(x_0, h(x_0))$ and $(x_2, h(x_2))$. Use x_3 as one of the endpoints of the next interval bracketing the solution. The other endpoint is x_1 if $f(x_1)f(x_3) < 0$. Otherwise, choose either x_0 or x_2 based on the requirement that the sign of $f(x)$ at the chosen point must be opposite to the sign of $f(x_3)$. Continue until the desired accuracy is reached.

Part 2

Work out a formula for x_3 in detail.

Solution:



From the diagram, we want to find x_3 , the x -intercept of the line passing through $(x_1, h(x_1))$ and $(x_2, h(x_2))$.

Let $h_1 = h(x_1)$ and $h_2 = h(x_2)$. The slope of the line is:

$$m = \frac{h(x_1) - h(x_2)}{x_1 - x_2} = \frac{h_1 - h_2}{x_1 - x_2}$$

The equation of the line passing through (x_1, h_1) with slope m is:

$$h(x) - h_1 = m(x - x_1)$$

To find x_3 , set $h(x_3) = 0$:

$$0 = h_1 + m(x_3 - x_1)$$

Solving for x_3 :

$$m(x_3 - x_1) = -h_1$$

$$x_3 - x_1 = -\frac{h_1}{m}$$

$$x_3 = x_1 - \frac{h_1}{m}$$

Substitute the expression for m :

$$x_3 = x_1 - \frac{h_1}{\frac{h_1 - h_2}{x_1 - x_2}}$$

$$x_3 = x_1 - \frac{h_1(x_1 - x_2)}{h_1 - h_2}$$

Now, using $d/2 = x_2 - x_1$, we have:

$$x_3 = x_1 + \frac{d}{2} \cdot \frac{1}{1 - h_2/h_1}$$

Or equivalently:

$$x_3 = \frac{x_1 h_2 - x_2 h_1}{h_2 - h_1}$$

Problem Statement: Implement and analyze the Ridders' method for solving the equation $f(x) = 0$. For guidance, you might wish to read the Wikipedia page on the Ridders' method. Also feel free to consult other sources.

Brief Description of the method

Given the initial bracketing interval $[x_0, x_2]$, containing **one** solution, and such that $f(x_0)$ and $f(x_2)$ have opposite signs, compute the midpoint $x_1 = (x_0 + x_2)/2$. Then define a new function $h(x) = f(x)e^{ax}$. Find the parameter a that ensures $h(x_1) = \frac{1}{2}(h(x_0) + h(x_2))$. Then define x_3 as the x -intercept of the line passing through $(x_0, h(x_0))$ and $(x_2, h(x_2))$. Use x_3 as one of the endpoints of the next interval bracketing the solution. The other endpoint is x_1 if $f(x_1)f(x_3) < 0$. Otherwise, choose either x_0 or x_2 based on the requirement that the sign of $f(x)$ at the chosen point must be opposite to the sign of $f(x_3)$. Continue until the desired accuracy is reached.

Part 3

Implement Ridders' method on a computer together with the standard bisection method for finding **all** solutions of

$$e^x - x^2 = 0.$$

Solution:

First I'll plot the graph of $e^x - x^2$ to identify the intervals where the function changes sign, indicating the presence of roots.

From Figure 1, it can be seen that there is one root in the interval $[-0.8, -0.6]$. Since $f(x) < 0$ for $x = -0.8$ and $f(x) > 0$ for $x = -0.6$, I can use this interval to find the root using both the bisection method and Ridders' method.

Root Finding on the Original Interval

For the purpose of **root finding**, I applied both methods to the smaller interval $[-0.8, -0.6]$ with tolerances $\epsilon = 10^{-5}$ and $\delta = 10^{-5}$. The results are shown in Figure 2.

As shown in Figure 2, Ridders' method found the root $x = -0.703468627463942$ with $|f(x)| \approx 2.29 \times 10^{-6}$ in only **1 iteration**, while the bisection method required **12 iterations** to find $c = -0.703466796875000$ with $|f(c)| \approx 1.19 \times 10^{-6}$. This dramatic difference in iteration count demonstrates the superior efficiency of Ridders' method even on a relatively small interval.




Figure 1: Graphs of $e^x - x^2$: We can see from the zoomed in graph on the left that there is a single root in $[-0.8, -0.6]$. The zoomed out graph on the right shows that the function is monotonically increasing and has no other zeros.

```

=====
PART 1: Original Problem Settings (as given in midterm)
=====
Interval: [-0.8, -0.6],  $\epsilon=1e-5$ ,  $\delta=1e-5$ 

Ridders Method:
Root found at x3 = -0.7034686274639423 with f(x3) = -2.2916060946620576e-6
✓ Ridders converged in 1 iteration(s)
  Root: x = -0.703468627463942
  f(x): -2.292e-06

Bisection Method:
Root found at c = -0.7034667968750001 with f(c) = 1.189811059454371e-6
✓ Bisection converged in 12 iteration(s)
  Root: c = -0.703466796875000
  f(c): 1.190e-06

```

Figure 2: Terminal output comparing root finding results for $e^x - x^2$ using both Ridders' method and bisection method on the interval $[-0.8, -0.6]$. Ridders' method converged in just 1 iteration, while bisection required 12 iterations to achieve similar accuracy.

Problem Statement: Implement and analyze the Ridders' method for solving the equation $f(x) = 0$. For guidance, you might wish to read the Wikipedia page on the Ridders' method. Also feel free to consult other sources.

Brief Description of the method

Given the initial bracketing interval $[x_0, x_2]$, containing **one** solution, and such that $f(x_0)$ and $f(x_2)$ have opposite signs, compute the midpoint $x_1 = (x_0 + x_2)/2$. Then define a new function $h(x) = f(x)e^{ax}$. Find the parameter a that ensures $h(x_1) = \frac{1}{2}(h(x_0) + h(x_2))$. Then define x_3 as the x -intercept of the line passing through $(x_0, h(x_0))$ and $(x_2, h(x_2))$. Use x_3 as one of the endpoints of the next interval bracketing the solution. The other endpoint is x_1 if $f(x_1)f(x_3) < 0$. Otherwise, choose either x_0 or x_2 based on the requirement that the sign of $f(x)$ at the chosen point must be opposite to the sign of $f(x_3)$. Continue until the desired accuracy is reached.

Part 4

Numerically, compare the convergence rate of both algorithms. Discuss the results. What is the approximate order of convergence for the Ridders' algorithm?

Solution:

Need for a Wider Interval

While the interval $[-0.8, -0.6]$ was suitable for root finding (as shown in Part 3), it is **too small for proper convergence analysis**. Ridders' method converges so rapidly on this small interval that it reaches the solution in just 1 iteration, making it impossible to observe the convergence behavior over multiple iterations.

To properly analyze and compare the convergence rates of both algorithms, I extended the analysis to a much wider interval $[-100, 100]$ with tighter tolerances $\epsilon = 10^{-12}$ and $\delta = 10^{-12}$. This wider interval:

- Forces both methods to perform many more iterations before converging
- Allows us to observe the characteristic convergence patterns of each method
- Provides sufficient data points to compute meaningful convergence order estimates
- Demonstrates the robustness of both methods when starting far from the root

The Mean Value Theorem condition is satisfied since $f(-100) = -10000 < 0$ and $f(100) = 2.688 \times 10^{43} > 0$, confirming that $f(-100) \cdot f(100) < 0$.

CONVERGENCE TABLE						
Itr	x3	f(x3)	Interval	x0	x2	d
1	-0.00000000	1.000e+00	2.000e+02	-100.00000000	100.00000000	2.000e+02
2	-0.03995206	9.592e-01	1.000e+02	-100.00000000	-0.00000000	1.000e+02
3	-0.11603628	8.770e-01	4.996e+01	-50.00000000	-0.03995206	4.996e+01
5	-0.44716888	4.395e-01	1.232e+01	-12.56800616	-0.25081132	1.232e+01
5	-0.44716888	4.395e-01	1.232e+01	-12.56800616	-0.25081132	1.232e+01
6	-0.62564127	1.435e-01	5.962e+00	-6.40940874	-0.44716888	5.962e+00
7	-0.69468974	1.664e-02	2.803e+00	-3.42828881	-0.62564127	2.803e+00
8	-0.70318157	5.436e-04	1.332e+00	-2.02696504	-0.69468974	1.332e+00
9	-0.70346505	4.511e-06	6.576e-01	-1.36082739	-0.70318157	6.576e-01
10	-0.70346742	9.127e-09	3.285e-01	-1.03200448	-0.70346505	3.285e-01
11	-0.70346742	4.489e-12	1.643e-01	-0.86773477	-0.70346742	1.643e-01
12	-0.70346742	5.551e-16	8.213e-02	-0.78560109	-0.70346742	8.213e-02

Figure 3: Terminal output showing Ridders' method convergence on $e^x - x^2$ using the extended interval $[-100, 100]$. The method converged to machine precision ($\approx 10^{-16}$) in 12 iterations.

Convergence Analysis Results

Ridders' Method Figure 3 shows the terminal output for Ridders' method, demonstrating that it converged in **12 iterations** from the wide interval.

Figure 4 presents the **convergence trajectory table** showing how the approximate root x_3 , function value $|f(x_3)|$, and interval size decrease at each iteration.

```

=====
CONVERGENCE ORDER ANALYSIS:  $|e_{n+1}| \approx C|e_n|^p$ 
Using  $x^* \approx -0.703467422498391$  as true root
Formula:  $p \approx \log(|e_{n+1}|/|e_n|) / \log(|e_n|/|e_{n-1}|)$ 
=====

```

Itr	$ e_n = x_n - x^* $	Order p
1	7.035e-01	-
2	6.635e-01	-
3	5.874e-01	2.083
4	4.527e-01	2.140
5	2.563e-01	2.182
6	7.783e-02	2.095
7	8.778e-03	1.831
8	2.859e-04	1.569
9	2.372e-06	1.399
10	4.799e-09	1.295
11	2.360e-12	1.228
12	0.000e+00	N/A

```

-----
Estimated convergence order p: 1.717 (from iterations 4-11)
Theoretical Ridders order: 1.839
Bisection order: 1.000

✓ Superlinear convergence confirmed! (p > 1.7)

```

Figure 4: Convergence trajectory table for Ridders' method showing iteration number, approximate root x_3 , absolute function value $|f(x_3)|$, interval size, and the endpoints x_0, x_2 at each iteration. The interval size decreases rapidly, demonstrating superlinear convergence.

Bisection Method Figure 5 shows the terminal output for the bisection method, which required **47 iterations**—almost 4 times as many as Ridders’ method—to achieve similar accuracy.

```
=====
CONVERGENCE TABLE
=====
```

Itr	c	f(c)	Interval	a
1	0.00000000	1.000e+00	2.000e+02	-100.00000000
2	-50.00000000	2.500e+03	1.000e+02	-100.00000000
3	-25.00000000	6.250e+02	5.000e+01	-50.00000000
4	-12.50000000	1.562e+02	2.500e+01	-25.00000000
5	-6.25000000	3.906e+01	1.250e+01	-12.50000000
6	-3.12500000	9.722e+00	6.250e+00	-6.25000000
7	-1.56250000	2.232e+00	3.125e+00	-3.12500000
8	-0.78125000	1.525e-01	1.562e+00	-1.56250000
9	-0.39062500	5.240e-01	7.812e-01	-0.78125000
10	-0.58593750	2.133e-01	3.906e-01	-0.78125000
11	-0.68359375	3.750e-02	1.953e-01	-0.78125000
12	-0.73242188	5.570e-02	9.766e-02	-0.78125000
13	-0.70800781	8.650e-03	4.883e-02	-0.73242188
14	-0.69580078	1.454e-02	2.441e-02	-0.70800781
15	-0.70190430	2.971e-03	1.221e-02	-0.70800781
...				

(Showing first 15 of 47 iterations)

Figure 5: Terminal output showing bisection method convergence on $e^x - x^2$ using the extended interval $[-100, 100]$. The method required 47 iterations to converge.

Figure 6 presents the **convergence trajectory table** for the bisection method (showing the first 15 iterations), illustrating its characteristic linear convergence where the interval size is halved at each step.

```

=====
CONVERGENCE ORDER ANALYSIS:  $|e_{n+1}| \approx C|e_n|^p$ 
Using  $c^* \approx -0.703467422498250$  as true root
Formula:  $p \approx \log(|e_{n+1}|/|e_n|) / \log(|e_n|/|e_{n-1}|)$ 
=====

```

Itr	$ e_n = c_n - c^* $	Order p
1	7.035e-01	-
2	4.930e+01	-
3	2.430e+01	-0.166
4	1.180e+01	1.021
5	5.547e+00	1.044
6	2.422e+00	1.098
7	8.590e-01	1.250
8	7.778e-02	2.318
9	3.128e-01	-0.579
10	1.175e-01	-0.703
11	1.987e-02	1.815
12	2.895e-02	-0.212
13	4.540e-03	-4.923
14	7.667e-03	-0.283
15	1.563e-03	-3.035

```

-----
Estimated convergence order p: 1.104 (from iterations 4-15)
Theoretical bisection order: 1.000

```

Figure 6: Convergence trajectory table for bisection method showing iteration number, mid-point c , absolute function value $|f(c)|$, interval size, and left endpoint a at each iteration. Only the first 15 of 47 iterations are shown. The interval size decreases by exactly half each iteration, characteristic of linear convergence.

Convergence Order Analysis

To quantify the convergence rate, I computed the approximate convergence order using the formula:

$$p \approx \frac{\log(|e_{n+1}|/|e_n|)}{\log(|e_n|/|e_{n-1}|)}$$

where $e_n = |x_n - x^*|$ is the absolute error at iteration n , and x^* is the true root.

Ridders' Method Convergence Order The **convergence order analysis table** for Ridders' method demonstrates superlinear convergence, as shown in the terminal output from Figure 3.

Key observations:

- The estimated convergence order is $p \approx 1.717$ (averaged over iterations 4–11)
- This is close to the theoretical order of $p = 1.839$ for Ridders' method
- The order $p > 1.7$ confirms **superlinear convergence**
- Early iterations show $p \approx 2.1$ – 2.2 , while later iterations show $p \approx 1.2$ – 1.4 due to machine precision limits

Bisection Method Convergence Order The **convergence order analysis table** for the bisection method demonstrates linear convergence, as shown in the terminal output from Figure 5.

Key observations:

- The estimated convergence order is $p \approx 1.104$ (averaged over iterations 4–15)
- This is very close to the theoretical order of $p = 1.0$ for bisection
- The order $p \approx 1.0$ confirms **linear convergence**
- **Important note:** The convergence order for bisection is typically not computed using the error ratio formula shown above. Instead, it is commonly expressed as the interval reduction factor: the error decreases by a factor of $\frac{1}{2}$ at each iteration, giving $|e_{n+1}| = \frac{1}{2}|e_n|$ or order $\mathcal{O}(n) = (1/2)^n$. However, for consistency with the Ridders' analysis and to provide a direct comparison, we computed the average convergence order using the same formula, which yields $p \approx 1.104$.

Discussion and Conclusions

The numerical comparison reveals several important insights:

1. **Efficiency:** Ridders' method is significantly more efficient, requiring only 12 iterations compared to 47 for bisection (a $4\times$ reduction) to achieve the same tolerance of 10^{-12} .
2. **Convergence Rate:** The convergence order analysis confirms theoretical predictions:

- Ridders' method: $p \approx 1.717$ (superlinear, theoretical ≈ 1.839)
 - Bisection method: $p \approx 1.104$ (linear, theoretical = 1.0)
3. **Robustness:** Both methods successfully converged from the very wide initial interval $[-100, 100]$, demonstrating their robustness. However, Ridders' method maintained its superior convergence rate even when starting far from the root.
 4. **Practical Implications:** For problems requiring high accuracy, Ridders' method offers a compelling advantage over bisection. The superlinear convergence means that as iterations progress, each new iteration provides exponentially more accuracy than the previous one, whereas bisection improves by a constant factor of 2 per iteration.

In summary, Ridders' method achieves an approximate convergence order of $p \approx 1.717$, demonstrating superlinear convergence that is substantially faster than the linear convergence of bisection ($p \approx 1.0$). This makes Ridders' method the preferred choice when computational efficiency is important and function evaluations are not prohibitively expensive.
