

Fall 2025 CPTS530 Homework 03 Report

Numerical Analysis

Aryan Ritwajeet Jha

Due: Tuesday, September 30, 2025

Contents

Question 1	2
Question 2	3
Question 3	4
Question 4	5

Question 1

Problem Statement: Suppose that $x > 0$, $y < 0$. Using definition of absolute value and splitting into cases, show that $|x + y| \leq |x - y|$.

Solution:

Given: $x > 0$, $y < 0$

To show: $|x + y| \leq |x - y|$

Visual Representation:




Figure 1: Number line showing the relationship between x , y and $x - y$

Since $x > 0$ and $y < 0$, we can assume that $x - y = d$ where $d > 0$.

Therefore, $\text{RHS} = |x - y| = d$ is always true.

Now we need to consider cases for $|x + y|$ based on whether $x + y$ is positive or negative.

Case I: $x + y \geq 0$

From $x - y = d$, we get $x = y + d$.

Therefore: $x + y = (y + d) + y = 2y + d$

Since $x + y \geq 0$, we have $|x + y| = x + y = 2y + d$.

So $\text{LHS} = 2y + d$ and $\text{RHS} = d$.

Since $y < 0$, we have $2y < 0$, which means $\text{LHS} = 2y + d < d = \text{RHS}$.

Case II: $x + y < 0$

From $x - y = d$, we get $y = x - d$.

Therefore: $x + y = x + (x - d) = 2x - d < 0$

Since $x + y < 0$, we have $|x + y| = -(x + y) = -(2x - d) = d - 2x$.

So $\text{LHS} = d - 2x$ and $\text{RHS} = d$.

Since $x > 0$, we have $2x > 0$, which means $\text{LHS} = d - 2x < d = \text{RHS}$.

Conclusion:

Cases I and II exhaustively cover all possible scenarios. In both cases, we have shown that $\text{LHS} < \text{RHS}$, which means $|x + y| < |x - y|$.

Therefore, we have proven the stronger result: $|x + y| \leq |x - y|$ for $x > 0, y < 0$

Hence Proved. $\square$

Question 2

Problem Statement: Suppose you have an iterative algorithm for finding a fixed point of a function $F(x)$ on a certain interval. Write a short text (2-4 paragraphs) explaining the following:

1. How do you decide when to stop iterating?
2. What is your strategy for making sure that the computed solution is close to the exact solution?

DO NOT use the standard methodology and estimates from the textbook. Try to find other ways of accomplishing the goals above. Discuss briefly the advantages and disadvantages of your error control approach.

Solution:

Question 3

Problem Statement: Do Problem 23 c) from Section 3.4.

Problem 3.4.23 c): Find the order of convergence of the sequence:

$$x_n = \left(1 + \frac{1}{n}\right)^{\frac{1}{2}}$$

Solution:

Question 4

Problem Statement: Do Problem 32 from Section 3.4.

Problem 3.4.32: Let $\frac{1}{2} \leq q \leq 1$, and define $F(x) = 2x - qx^2$. On what interval can it be guaranteed that the method of iteration using F will converge to a fixed point? (This problem is related to Problem 3.2.5, p. 90.)

Related Problem 3.2.5: What is the purpose of the following iteration formula?

$$x_{n+1} = 2x_n - x_n^2 y$$

Identify it as the Newton iteration for a certain function.

Solution: