

Fall 2025 CPTS530 Homework 5
MATH 448-548
Numerical Analysis

Aryan Ritwajeet Jha

November 2025

Contents

Problem 1	2
Problem 2	4
Problem 3	5
Problem 4	6
Problem 5	8
Problem 6	9
Problem 7	10
Problem 8	11
Problem 9	13
Appendix	14
References	15

Problem 1

Problem Statement (Textbook 4.4.1):

Show that the norms $\|\mathbf{x}\|_\infty$, $\|\mathbf{x}\|_2$, and $\|\mathbf{x}\|_1$ satisfy the postulates (1), (2), and (3) for norms.

Norm Postulates [1, 2]:

On a vector space V , a **norm** is a function $\|\cdot\|$ from V to the set of nonnegative reals that obeys these three postulates:

$$\|x\| > 0 \text{ if } x \neq 0, x \in V \quad (1)$$

$$\|\lambda x\| = |\lambda| \|x\| \text{ if } \lambda \in \mathbb{R}, x \in V \quad (2)$$

$$\|x + y\| \leq \|x\| + \|y\| \text{ if } x, y \in V \quad (\text{triangle inequality}) \quad (3)$$

Solution:

We verify postulates (1), (2), and (3) for $\|x\|_\infty = \max_i |x_i|$, $\|x\|_2 = \sqrt{\sum_i x_i^2}$, and $\|x\|_1 = \sum_i |x_i|$.

(a) ∞ -norm: $\|x\|_\infty = \max_{1 \leq i \leq n} |x_i|$

(1) If $x \neq 0$, then $\exists i$ with $x_i \neq 0$, so $|x_i| > 0$. Thus $\|x\|_\infty = \max_i |x_i| > 0$.

(2) $\|\lambda x\|_\infty = \max_i |\lambda x_i| = \max_i |\lambda| |x_i| = |\lambda| \max_i |x_i| = |\lambda| \|x\|_\infty$.

(3) $\|x + y\|_\infty = \max_i |x_i + y_i| \leq \max_i (|x_i| + |y_i|) \leq \max_i |x_i| + \max_i |y_i| = \|x\|_\infty + \|y\|_\infty$.

(b) 2-norm (Euclidean): $\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$

(1) If $x \neq 0$, then $\exists i$ with $x_i \neq 0$, so $\sum_i x_i^2 > 0$. Thus $\|x\|_2 = \sqrt{\sum_i x_i^2} > 0$.

(2) $\|\lambda x\|_2 = \sqrt{\sum_i (\lambda x_i)^2} = \sqrt{\lambda^2 \sum_i x_i^2} = |\lambda| \sqrt{\sum_i x_i^2} = |\lambda| \|x\|_2$.

(3) Expanding $\|x + y\|_2^2$:

$$\begin{aligned} \|x + y\|_2^2 &= \sum_i (x_i + y_i)^2 = \sum_i x_i^2 + 2 \sum_i x_i y_i + \sum_i y_i^2 \\ &= \|x\|_2^2 + 2 \sum_i x_i y_i + \|y\|_2^2 \end{aligned}$$

Since $\sum_i x_i y_i \leq \|x\|_2 \|y\|_2$ (dot product bounded by product of norms):

$$\|x + y\|_2^2 \leq \|x\|_2^2 + 2\|x\|_2 \|y\|_2 + \|y\|_2^2 = (\|x\|_2 + \|y\|_2)^2$$

Taking square roots: $\|x + y\|_2 \leq \|x\|_2 + \|y\|_2$.

(c) 1-norm: $\|x\|_1 = \sum_{i=1}^n |x_i|$

(1) If $x \neq 0$, then $\exists i$ with $x_i \neq 0$, so $|x_i| > 0$. Thus $\|x\|_1 = \sum_i |x_i| > 0$.

(2) $\|\lambda x\|_1 = \sum_i |\lambda x_i| = \sum_i |\lambda| |x_i| = |\lambda| \sum_i |x_i| = |\lambda| \|x\|_1$.

(3) For each term, the triangle inequality for absolute values gives $|x_i + y_i| \leq |x_i| + |y_i|$.

Summing over all i :

$$\|x + y\|_1 = \sum_i |x_i + y_i| \leq \sum_i (|x_i| + |y_i|) = \sum_i |x_i| + \sum_i |y_i| = \|x\|_1 + \|y\|_1$$

All three norms satisfy postulates (1), (2), and (3).

Problem 2

Problem Statement (Textbook 4.4.2):

Show that $\|\mathbf{x}\|_\infty \leq \|\mathbf{x}\|_2 \leq \|\mathbf{x}\|_1$ for all $\mathbf{x} \in \mathbb{R}^n$, and that equalities can occur, even for nonzero vectors.

Solution:

Part 1: Prove $\|x\|_\infty \leq \|x\|_2$

Let $\|x\|_\infty = \max_i |x_i| = |x_k|$ for some index k . Then:

$$\|x\|_2^2 = \sum_{i=1}^n x_i^2 \geq x_k^2 = \|x\|_\infty^2$$

Taking square roots: $\|x\|_\infty \leq \|x\|_2$.

Equality case: Equality holds when x has only one nonzero component. For example, $x = (5, 0, 0)^T$ gives $\|x\|_\infty = 5$ and $\|x\|_2 = 5$.

Part 2: Prove $\|x\|_2 \leq \|x\|_1$

First, observe that each component satisfies $|x_i| \leq \|x\|_1$ (since $\|x\|_1 = \sum_{j=1}^n |x_j|$ includes all terms).

For each i , we can write:

$$|x_i|^2 = |x_i| \cdot |x_i| \leq |x_i| \cdot \|x\|_1$$

where the inequality holds because $|x_i| \leq \|x\|_1$.

Now sum this inequality over all indices:

$$\begin{aligned} \|x\|_2^2 &= \sum_{i=1}^n |x_i|^2 \leq \sum_{i=1}^n |x_i| \cdot \|x\|_1 \\ &= \|x\|_1 \sum_{i=1}^n |x_i| \quad (\text{factor out } \|x\|_1) \\ &= \|x\|_1 \cdot \|x\|_1 = \|x\|_1^2 \end{aligned}$$

Taking square roots of both sides: $\|x\|_2 \leq \|x\|_1$.

Equality case: Equality holds when x has only one nonzero component. For example, $x = (0, 3, 0)^T$ gives $\|x\|_2 = 3$ and $\|x\|_1 = 3$.

Conclusion: The inequality $\|x\|_\infty \leq \|x\|_2 \leq \|x\|_1$ holds for all $x \in \mathbb{R}^n$, with equality possible for nonzero vectors.

Problem 3

Problem Statement (Textbook 4.4.3):

(Continuation) Show that $\|\mathbf{x}\|_1 \leq n\|\mathbf{x}\|_\infty$ and $\|\mathbf{x}\|_2 \leq \sqrt{n}\|\mathbf{x}\|_\infty$ for $\mathbf{x} \in \mathbb{R}^n$.

Solution:

Part 1: Prove $\|x\|_1 \leq n\|x\|_\infty$

Since $\|x\|_\infty = \max_i |x_i|$, we have $|x_i| \leq \|x\|_\infty$ for each $i = 1, 2, \dots, n$.

Summing over all components:

$$\|x\|_1 = \sum_{i=1}^n |x_i| \leq \sum_{i=1}^n \|x\|_\infty = n\|x\|_\infty$$

Therefore: $\boxed{\|x\|_1 \leq n\|x\|_\infty}$.

Equality case: Equality holds when all components have the same absolute value. For example, $x = (1, 1, 1)^T$ gives $\|x\|_1 = 3 = 3 \cdot 1 = 3\|x\|_\infty$.

Part 2: Prove $\|x\|_2 \leq \sqrt{n}\|x\|_\infty$

Since $|x_i| \leq \|x\|_\infty$ for each i , squaring gives $|x_i|^2 \leq \|x\|_\infty^2$.

Summing over all components:

$$\|x\|_2^2 = \sum_{i=1}^n |x_i|^2 \leq \sum_{i=1}^n \|x\|_\infty^2 = n\|x\|_\infty^2$$

Taking square roots: $\boxed{\|x\|_2 \leq \sqrt{n}\|x\|_\infty}$.

Equality case: Equality holds when all components have the same absolute value. For example, $x = (2, 2, 2)^T$ gives $\|x\|_2 = \sqrt{12} = 2\sqrt{3} = \sqrt{3} \cdot 2 = \sqrt{3}\|x\|_\infty$.

Problem 4

Problem Statement (Textbook 4.4.8):

Define

$$\|A\| = \sum_{i=1}^n \sum_{j=1}^n |a_{ij}|$$

Show that this is a matrix norm (that is, a norm on the linear space of all $n \times n$ matrices). Show that it is not subordinate to any vector norm. Does it conform to Equation (9) and Inequality (10)?

$$\|I\| = 1 \quad (\text{Equation 9})$$

$$\|AB\| \leq \|A\|\|B\| \quad (\text{Inequality 10})$$

Solution:

Part 1: Show $\|A\|$ is a matrix norm

A matrix norm must satisfy three postulates:

- (1) **Positivity:** If $A \neq 0$, then $\exists$ some $a_{ij} \neq 0$, so $|a_{ij}| > 0$. Thus $\|A\| = \sum_{i,j} |a_{ij}| > 0$.
- (2) **Homogeneity:** $\|\lambda A\| = \sum_{i,j} |\lambda a_{ij}| = \sum_{i,j} |\lambda| |a_{ij}| = |\lambda| \sum_{i,j} |a_{ij}| = |\lambda| \|A\|$.
- (3) **Triangle inequality:**

$$\begin{aligned} \|A + B\| &= \sum_{i,j} |a_{ij} + b_{ij}| \leq \sum_{i,j} (|a_{ij}| + |b_{ij}|) \\ &= \sum_{i,j} |a_{ij}| + \sum_{i,j} |b_{ij}| = \|A\| + \|B\| \end{aligned}$$

Therefore, $\|A\|$ is a matrix norm.

Part 2: Show it is NOT subordinate to any vector norm

A subordinate (induced) matrix norm satisfies $\|I\| = 1$. However, for the identity matrix I where $(I)_{ij} = \delta_{ij}$ (Kronecker delta: $\delta_{ij} = 1$ if $i = j$, else 0):

$$\|I\| = \sum_{i=1}^n \sum_{j=1}^n |\delta_{ij}| = \sum_{i=1}^n 1 = n$$

Since $\|I\| = n \neq 1$ for $n > 1$, this norm is NOT subordinate to any vector norm.

Part 3: Does it conform to Equation (9)?

Equation (9) requires $\|I\| = 1$. As shown above, $\|I\| = n$.

No, it does NOT conform to Equation (9).

Part 4: Does it conform to Inequality (10)?

We need to check if $\|AB\| \leq \|A\|\|B\|$.

Step 1: We recall that for matrix multiplication, the (i, k) entry of AB is:

$$(AB)_{ik} = \sum_{j=1}^n a_{ij}b_{jk}$$

Step 2: Using the triangle inequality for absolute values, we have:

$$|(AB)_{ik}| = \left| \sum_{j=1}^n a_{ij}b_{jk} \right| \leq \sum_{j=1}^n |a_{ij}b_{jk}| = \sum_{j=1}^n |a_{ij}||b_{jk}|$$

Step 3: We now compute $\|AB\|$ by summing over all entries:

$$\begin{aligned} \|AB\| &= \sum_{i=1}^n \sum_{k=1}^n |(AB)_{ik}| \\ &\leq \sum_{i=1}^n \sum_{k=1}^n \sum_{j=1}^n |a_{ij}||b_{jk}| \quad (\text{from Step 2}) \end{aligned}$$

Step 4: We rearrange the triple sum. Notice that $|a_{ij}|$ doesn't depend on k , and $|b_{jk}|$ doesn't depend on i :

$$\begin{aligned} \sum_{i=1}^n \sum_{k=1}^n \sum_{j=1}^n |a_{ij}||b_{jk}| &= \sum_{i=1}^n \sum_{j=1}^n |a_{ij}| \left(\sum_{k=1}^n |b_{jk}| \right) \\ &= \sum_{i=1}^n \sum_{j=1}^n |a_{ij}| \cdot \sum_{k=1}^n \sum_{j=1}^n |b_{jk}| \\ &= \left(\sum_{i=1}^n \sum_{j=1}^n |a_{ij}| \right) \left(\sum_{j=1}^n \sum_{k=1}^n |b_{jk}| \right) \\ &= \|A\| \cdot \|B\| \end{aligned}$$

Step 5: Combining Steps 3 and 4, we obtain: $\|AB\| \leq \|A\|\|B\|$.

Yes, it DOES conform to Inequality (10).

Problem 5

Problem Statement (Textbook 4.4.11):

Show that for the vector norm $\|\mathbf{x}\|_1$ defined in Equation (5), the subordinate matrix norm is

$$\|A\|_1 = \max_{1 \leq j \leq n} \sum_{i=1}^n |a_{ij}|$$

where

$$\|\mathbf{x}\|_1 = \sum_{i=1}^n |x_i| \quad (\text{Equation 5})$$

Solution:

Problem 6

Problem Statement (Textbook 4.4.33):

For any $n \times n$ matrix A , define

$$\|A\|_F = \left(\sum_{i=1}^n \sum_{j=1}^n a_{ij}^2 \right)^{1/2}$$

(This is called the **Frobenius** norm.) Is this a subordinate matrix norm? Answer the same question for $\|A\| = \max_{1 \leq i, j \leq n} |a_{ij}|$. Prove that these equations define norms on the vector space of all $n \times n$ matrices.

Solution:

Problem 7

Problem Statement (Textbook 4.6.1b):

Program the Gauss-Seidel method and test it on the following system:

$$3x + y + z = 5$$

$$3x + y - 5z = -1$$

$$x + 3y - z = 3$$

Analyze what happens when these systems are solved by simple Gaussian elimination without pivoting.

Solution:

Problem 8

Problem Statement (Textbook 6.1.1d):

Find the polynomial of least degree that interpolates these sets of data:

$$\begin{array}{cccc} x & 3 & 7 & 12 \\ y & 12 & 146 & 2 \end{array}$$

Solution:

We find the polynomial of least degree interpolating the data points $(3, 12)$, $(7, 146)$, and $(12, 2)$.

Since we have 3 data points, the interpolating polynomial has degree at most $n - 1 = 2$ (quadratic).

Method: Vandermonde Matrix

We seek a polynomial $P(x) = a_0 + a_1x + a_2x^2$ such that $P(x_i) = y_i$ for $i = 1, 2, 3$.

This gives the linear system:

$$\begin{bmatrix} 1 & x_1 & x_1^2 \\ 1 & x_2 & x_2^2 \\ 1 & x_3 & x_3^2 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

Substituting our data:

$$\begin{bmatrix} 1 & 3 & 9 \\ 1 & 7 & 49 \\ 1 & 12 & 144 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} 12 \\ 146 \\ 2 \end{bmatrix}$$

Solving this system (using Julia code in `hw05_p08.jl`), we obtain:

$$\begin{aligned} a_0 &= -\frac{4211}{18} \approx -233.8667 \\ a_1 &= \frac{1849}{18} \approx 102.7222 \\ a_2 &= -\frac{125}{18} \approx -6.9222 \end{aligned}$$

Final Answer:

$$P(x) = -\frac{125}{18}x^2 + \frac{1849}{18}x - \frac{4211}{18}$$

Or in decimal form:

$$P(x) = -6.9222x^2 + 102.7222x - 233.8667$$

Verification:

- $P(3) = 12.000$

- $P(7) = 146.000$
- $P(12) = 2.000$

Maximum interpolation error: 5.68×10^{-14} (numerical precision).

Problem 9

Problem Statement (Textbook 6.1.8):

Discuss the problem of determining a polynomial of degree at most 2 for which $p(0)$, $p(1)$, and $p(\xi)$ are prescribed, ξ being any preassigned point.

Solution:

Appendix

References

- [1] David Kincaid and Ward Cheney. *Numerical Analysis: Mathematics of Scientific Computing*. American Mathematical Society, Providence, RI, 3rd edition, 2009. Available from ProQuest Ebook Central. <https://ebookcentral.proquest.com/lib/wsu/detail.action?docID=4717637>.
- [2] Alexander Panchenko. Math 448 lecture notes. Washington State University, Fall 2025.