



Tansley review

50 years of Arabidopsis research: highlights and future directions

Author for correspondence:
Nicholas J. Provart
Tel: +1 416 978 7141
Email: nicholas.provart@utoronto.ca

Received: 26 June 2015
Accepted: 24 August 2015

Nicholas J. Provart¹, Jose Alonso², Sarah M. Assmann³, Dominique Bergmann⁴, Siobhan M. Brady⁵, Jelena Brkljacic⁶, John Browse⁷, Clint Chapple⁸, Vincent Colot⁹, Sean Cutler¹⁰, Jeff Dangl¹¹, David Ehrhardt¹², Joanna D. Friesner¹³, Wolf B. Frommer¹², Erich Grotewold¹⁴, Elliot Meyerowitz¹⁵, Jennifer Nemhauser¹⁶, Magnus Nordborg¹⁷, Craig Pikaard¹⁸, John Shanklin¹⁹, Chris Somerville²⁰, Mark Stitt²¹, Keiko U. Torii¹⁶, Jamie Waese¹, Doris Wagner²² and Peter McCourt¹

¹Department of Cell & Systems Biology/CAGEF, University of Toronto, Toronto, ON M5S 3B2, Canada; ²Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC 27695, USA; ³Department of Biology, Pennsylvania State University, University Park, PA 16802, USA; ⁴Department of Biology, Stanford University, Stanford, CA 94305, USA; ⁵Department of Plant Biology, University of California, Davis, CA 95616, USA; ⁶Arabidopsis Biological Resource Center, The Ohio State University, Columbus, OH 43210, USA; ⁷Institute of Biological Chemistry, Washington State University, Pullman, WA 99164, USA; ⁸Department of Biochemistry, Purdue University, West Lafayette, IN 47907, USA; ⁹Departement de Biologie École Normale Supérieure, Biologie Moléculaire des Organismes Photosynthétiques, F-75230, Paris, France; ¹⁰Department of Botany and Plant Sciences, University of California, Riverside, CA 92507, USA; ¹¹Department of Biology and Howard Hughes Medical Institute, Department of Microbiology and Immunology, University of North Carolina, Chapel Hill, NC 27599, USA; ¹²Department of Plant Biology, Carnegie Institution for Science, Stanford, CA 94305, USA; ¹³Department of Plant Biology, Agricultural Sustainability Institute, University of California, Davis, CA 95616, USA; ¹⁴Center for Applied Plant Science, The Ohio State University, Columbus, OH 43210, USA; ¹⁵Division of Biology and Biological Engineering and Howard Hughes Medical Institute, California Institute of Technology, Pasadena, CA 91125, USA; ¹⁶Department of Biology, University of Washington, Seattle, WA 98195, USA; ¹⁷Gregor Mendel Institute of Molecular Plant Biology, A-1030, Vienna, Austria; ¹⁸Department of Biology, Indiana University, Bloomington, IN 47405, USA; ¹⁹Biology Department, Brookhaven National Laboratory, Upton, NY 11973, USA; ²⁰Energy Biosciences Institute, University of California, Berkeley, CA 94704, USA; ²¹Metabolic Networks Department, Max Planck Institute for Molecular Plant Physiology, D-14476, Potsdam, Germany; ²²Department of Biology, University of Pennsylvania, Philadelphia, PA 19104, USA

Contents

Summary	922	X. Pumps, channels, transporters and the like	931
I. Introduction and a brief survey of 54 033 Arabidopsis publications	922	XI. Cell walls, starch and lipids	932
II. Arabidopsis reverse genetics: paving the way for gene function studies	922	XII. Epigenetics and epigenomics: from genotype to phenotype	933
III. Arabidopsis stock centers	925	XIII. Natural variation and genome-wide association studies	934
IV. Databases and online tools	925	XIV. Gene regulatory networks	934
V. Cell biology	926	XV. Modeling, bioinformatics, systems biology	935
VI. Development	927	XVI. Synthetic biology	936
VII. Hormones	928	XVII. Conclusions and outlook	937
VIII. The plant immune system and Arabidopsis research	929	Acknowledgements	937
IX. Signaling in response to abiotic stress	930	References	937

New Phytologist (2016) **209**: 921–944
doi: 10.1111/nph.13687

Key words: Arabidopsis, development, epigenetics, genome, model system, reverse genetics, signaling, synthetic biology.

Summary

The year 2014 marked the 25th International Conference on Arabidopsis Research. In the 50 yr since the first International Conference on Arabidopsis Research, held in 1965 in Göttingen, Germany, > 54 000 papers that mention *Arabidopsis thaliana* in the title, abstract or keywords have been published. We present herein a citational network analysis of these papers, and touch on some of the important discoveries in plant biology that have been made in this powerful model system, and highlight how these discoveries have then had an impact in crop species. We also look to the future, highlighting some outstanding questions that can be readily addressed in Arabidopsis. Topics that are discussed include Arabidopsis reverse genetic resources, stock centers, databases and online tools, cell biology, development, hormones, plant immunity, signaling in response to abiotic stress, transporters, biosynthesis of cell walls and macromolecules such as starch and lipids, epigenetics and epigenomics, genome-wide association studies and natural variation, gene regulatory networks, modeling and systems biology, and synthetic biology.

I. Introduction and a brief survey of 54 033 Arabidopsis publications

The year 2015 marks the 50th anniversary of the first conference on Arabidopsis research, held in Göttingen, Germany, in 1965. Although Friedrich Laibach (1885–1967) proposed using Arabidopsis as a genetic model organism almost 75 yr ago, it was not until the 1980s after intense discussions among early Arabidopsis proponents that it was widely adopted as such (Laibach, 1943; Meyerowitz, 2001, which lists these Arabidopsis pioneers). Factors such as small genome size, short generation time, ease of crossing, fecundity, and the ability to do mutational screens to saturation in the laboratory have all led to a huge increase in the volume of Arabidopsis research. In the past 50 yr, 54 033 Arabidopsis papers – defined as having Arabidopsis in the title, abstract or keywords in Thomson Reuter's BIOSIS database – covering 406 different biological fields have been published, most in the second half of this time frame (see Fig. 1 for the distribution of some of the more general research areas). The Arabidopsis publications and their citations may be explored interactively at <http://bar.utoronto.ca/50YearsOfArabidopsis/>.

The average Arabidopsis paper from the past 50 yr has been cited 33.8 times, and the number of citations follows a power law distribution, with just a few papers having been cited many times. The most frequently cited Arabidopsis paper describes transforming Arabidopsis by the floral dip method (Clough & Bent, 1998). According to BIOSIS, this paper has been cited 7195 times. Importantly, this collection of Arabidopsis papers has been widely cited outside of the Arabidopsis community (see Fig. 2). In 37 of the past 50 yr > 50% of the cited Arabidopsis papers published each year have been referenced by papers where Arabidopsis was not the focus of the research, as determined by the absence of this species in the taxonomic data available for each paper in the BIOSIS database. A maximum of 88% of cited Arabidopsis papers in 1989 have been referenced by non-Arabidopsis papers. Of 41 682 Arabidopsis papers published in the past 50 yr that have been cited one or more times, 15 388 of these have been cited by a non-Arabidopsis-focused paper.

We now review what we have learned about aspects of plant biology from this small plant that Laibach suggested using as a model in 1943. Many 'firsts' were discovered in Arabidopsis (of course, we cannot cover everything, owing to space limitations – our apologies to the exciting discoveries that we have not included here). But even when initial breakthroughs were made in another species, Arabidopsis research often helped to illuminate their fundamental workings. Fig. 3 highlights some of the discoveries made in Arabidopsis. The first four sections of this review cover resources and methods that have enabled much of this research. The final four sections cover newer research directions that will move plant biology forward in the coming years. The decision to grow and focus a research community around Arabidopsis as a model revolutionized our understanding of plants and, indeed, of all biology (see, e.g. Jones *et al.*, 2008 for a review of how Arabidopsis has had an impact on human health).

II. Arabidopsis reverse genetics: paving the way for gene function studies

Before the genome era, classical 'forward' genetics was the preferred strategy to establish causal relationships between a genotype and a phenotype. This consisted of the generation of large randomly mutagenized populations, the identification among them of individuals with the desired phenotype, mapping and fine mapping the mutations causing the phenotype, followed by sequencing of the corresponding genomic regions and subsequent complementation testing to confirm causality (Alonso & Ecker, 2006). The small genome of Arabidopsis, the inbred nature of the different laboratory strains, and the relative ease of developing genetic and molecular markers made the routine use of these powerful, albeit somewhat laborious, genetic strategies possible in this plant system. With the progression of the Arabidopsis genome project during the late 1990s, thousands of 'interesting' genes were discovered at an unprecedented pace, fueling the desire by plant researchers to test their favorite hypothesis about these genes' potential functions. To be able to do that, however, plants harboring disruptive




Fig. 1 Breakdown of 54 033 *Arabidopsis* publications by research area, from 1965 to 2015 (partial data for 2015). BIOSIS concept codes were used to flag papers in the 54 033 set. If a paper had been assigned multiple codes, the least abundant code was used for this display, out of 46 that were chosen to provide a balance between not being too general (75% of the papers were tagged with '03504': Genetics – Plant) and not being too specific (just one paper was tagged with '64500: Paleobiology'). Concept code terms ranged in prevalence from 11.6% for '51512: Plant physiology – Reproduction' to 1.04% for '13014: Metabolism – Nucleic acids, purines and pyrimidines'. Selected categories that have waxed and waned as research areas over the years are highlighted. All categories and publications may be explored online at <http://bar.utoronto.ca/50YearsOfArabidopsis/>.





Fig. 2 Number of *Arabidopsis* publications per year, categorized by whether or not they have been cited by a paper that was focused on a species other than *Arabidopsis*, as determined by the taxonomic data provided in the BIOSIS database. Eighty-eight per cent of cited *Arabidopsis* papers in 1989 have been referenced by non-*Arabidopsis* papers.

mutations in these genes were urgently needed. Once again, the vibrant *Arabidopsis* research community accepted the new challenge and, soon after, the first large collection of T-DNA mutants was created (Feldmann & Marks, 1987), although initially the idea was that these collections of mutants would simplify the sometimes tedious cloning of the mutated gene in 'forward' genetic screens. The nature of the mutagenic agent in these collections, an *Agrobacterium* plasmid T-DNA of known sequence, also enabled the rapid screening of hundreds of mutant lines not based on their phenotype, but on their sequence

(Alonso & Ecker, 2006). This initial success of the so-called 'reverse genetics', together with the development of extremely efficient methods to generate insertional mutants (T-DNA transformation via vacuum infiltration and later via 'floral dip': Bechtold *et al.*, 1993; Clough & Bent, 1998) and transposon-based mutants (Altmann *et al.*, 1992; Bancroft *et al.*, 1992), resulted in the generation of larger and larger T-DNA and transposon collections that made it possible to find mutations in practically any gene. Mutant identification, however, still required the tedious process of testing large pools of mutants

1st



- Plant genome sequence
- Genome-wide T-DNA collections; TILLING
- Live cell imaging of cellulose synthase
- Optical detection of hormones and metabolites
- Cell-type-specific transcriptomes of an entire plant organ
- First genetic and molecular tests of the floral ABC model
- Elucidation of hormone pathways
- Molecular glue model: auxin, JA, GA, SL, SA, and ABA receptors
- Isolation of plant immune receptor gene
- Idea of 'basal defense'
- Identification of cryptochrome, phototropin, UVB and other receptors
- Amino acid, iron, SWEET and other transporters
- Rewriting of lignin biosynthesis pathway
- Developmental responses to environment
- Regulatory networks for floral homeotic genes, and other processes
- Plant probabilistic functional gene network
- Synthetic biology: hormone pathway engineering
- ...

Fig. 3 Arabidopsis 'firsts' (highlights), ordered according to their mention in this review. ABA, abscisic acid; JA, jasmonic acid; SL, strigolactones; SA, salicylic acid.

one gene at a time (Krysan *et al.*, 1996). This laborious stage in Arabidopsis reverse genetics ended a few years later when new high-throughput ways of sequencing the insertional sites in these collections were established (Sessions *et al.*, 2002), and searchable databases and seeds stocks from individual mutants lines were generated and made publicly available (Alonso *et al.*, 2003). Together, these advances allowed for access to a mutant T-DNA line in the gene of interest by simply ordering the corresponding seeds from a stock center. The ease with which one could obtain the desired mutants had a dramatic impact on the way gene function was approached not only in Arabidopsis, but also in other plant species where the analysis of an Arabidopsis mutant has become a routine and quick method for shedding light on the function of the orthologous gene of interest in less experimentally tractable plant species. Nevertheless, because of the random way in which T-DNA mutants were generated and the insertion sites sequenced, this approach became less and less effective as the number of sequenced lines approached genome saturation. The recent advent of ultra-efficient next-generation sequencing-based approaches is allowing the rapid identification of tens of thousands of new insertional or other kinds of alleles in pre-existing mutant collections (J. Ecker, pers. comm.). As with all the approaches initially developed and tested in Arabidopsis, insertional mutagenesis-based reverse genetics strategies were soon implemented in other plant species and became a fundamental research tool in important crop plants such as rice and maize (reviewed in Jung *et al.*, 2008; Nannas & Dawe, 2015).

Although gene-indexed insertional collections are important pillars of reverse genetics approaches, they have some obvious limitations. For example, a majority of the lines in these collections represent null or hypomorphic alleles and require the stable integration of foreign DNA in the genome of the target plant. Thus, other means of altering the activity of desired genes were developed and, as before, used in Arabidopsis for quick testing and protocol optimization. Large collections of chemically induced mutants were generated and

screened using a technology pioneered in Arabidopsis, TILLING (Targeting Induced Local Lesions in Genomes; Till *et al.*, 2003), and then exported to other plant and animal systems (Till *et al.*, 2006). Since TILLING mutants usually contain a single nucleotide change, a wider array of allelic series could be obtained. Furthermore, because this technology does not rely on plant transformation, it was possible to rapidly translate the protocols developed for Arabidopsis to other plant and animal species that are not easily transformable. As in the case of insertional mutagenesis, the advent of next-generation sequencing has also had an important impact on the throughput of TILLING screening, allowing for the rapid testing of thousands of lines in order to find more and potentially rare mutant alleles (Till *et al.*, 2003). One of the limitations of both insertional mutagenesis and TILLING is the difficulty in combining linked mutations, which represents a significant problem in Arabidopsis and other plant species in which recently duplicated (and often functionally redundant) genes are found in adjacent locations in the genome. This and other caveats, such as the limited diversity of genetic backgrounds available and the inability to manipulate the spatial and temporal effects of the targeted gene's activity, provided a fertile ground for the development of RNA silencing-based approaches such as RNAi and artificial microRNAs (Hilson *et al.*, 2004; Schwab *et al.*, 2006). Once more, the Arabidopsis community played a key role in the implementation of these approaches in other plant systems by setting up the pipelines, generating the expertise and defining the basic rules affecting the efficiency of these reverse genetics approaches.

Finally, new genome editing technologies based on a variety of engineered endonucleases, such as homing endonucleases (Antunes *et al.*, 2012), zinc-finger nucleases (Wright *et al.*, 2005), TALENs (Bedell *et al.*, 2012) and, in recent years, CRISPR-Cas9 (Feng *et al.*, 2014; Li *et al.*, 2014a), have been added to the plant reverse genetics toolbox. With these new technologies, Arabidopsis research continues to play a critical role in the implementation and optimization of the experimental protocols.

III. *Arabidopsis* stock centers

1. Pre-stock center era

The need for a well organized *Arabidopsis* stock collection was recognized long before the *Arabidopsis* Biological Resource Center (ABRC) and the Nottingham *Arabidopsis* Stock Centre (NASC) were established in the early 1990s. The *Arabidopsis* Information Service (AIS) newsletter, initiated in 1964, represented the first version of a curated *Arabidopsis* stock collection (Meyerowitz, 2001). The AIS newsletter volume 24, issued in 1987, and the 'Green Book' (Meyerowitz & Pruitt, 1984) compiled the data for the existing set of stocks for the first time. The collection consisted of *c.* 1000 stocks, mostly natural accessions and individual mutant lines generated by a handful of *Arabidopsis* researchers, including Friedrich Laibach, Gerhard Röbbelen, Maarten Koornneef, and Albert Kranz. An additional 350 stocks (mostly mutants with a few natural accessions) were contributed by George Rédei. According to the Green Book, *c.* 1600 stocks were exchanged among researchers between 1974 and 1987, illustrating that the pioneers of *Arabidopsis* research were well aware of the crucial role of sharing materials and data, which helped set the stage for all future community efforts.

2. Stock center history and data

The vision of establishing two stock centers came from the Multinational *Arabidopsis* Steering Committee (MASC) and was driven both by a number of new resources generated as a result of the development of efficient transformation procedures and by a push to develop a new nonhuman model organism for functional studies (Somerville & Koornneef, 2002; Koornneef & Meinke, 2010). Since their inception in 1991 and 1992, respectively, both the NASC and the ABRC have been supported by their respective national funding agencies, as well as by stock contributions from the *Arabidopsis* community. The policy of sharing donated resources between the two stock centers remains one of the most important strengths of *Arabidopsis* research and one of the main factors contributing to its success. The initial stock center collection comprised seed stocks donated by Albert Kranz (>1000), George Rédei (<1000), Chris Somerville (200) and Maarten Koornneef (150). The donation of 4900 *Arabidopsis* T-DNA insertion lines by Kenneth Feldmann in 1992, and of *c.* 18 000 expressed sequence tags (ESTs) by Chris Somerville and Thomas Newman in 1994–1995, represented the first large seed and DNA donations, respectively (Meinke & Scholl, 2003). Many other large donations followed, contributing to the current ABRC holdings of 959 000 stocks. This 'sharing community' has encompassed >27 000 people in 11 000 laboratories to date. Additional regional centers (INRA, France; GABI, Germany; and RIKEN BRC, Japan) also participate in the distribution of *Arabidopsis* resources, some of which are shared and some of which are unique.

The overall ABRC stock distribution has increased from *c.* 8000 stocks sent in 1992 to >180 000 in 2014, including large sets (NASC shipped 109 311 stocks in 2014; S. May, pers. comm.).

With *c.* 80% of the total distribution, T-DNA pools and individual lines remain the most distributed stock category (Brkljacic *et al.*, 2011), illustrating a high demand for stocks that enable functional analyses. Natural accessions and mapping populations derived from accession crosses represent the second most ordered stock category, with *c.* 10% of the total distribution. The Salk uni-mutant collection of homozygous T-DNA insertion lines donated by Joseph Ecker (O'Malley & Ecker, 2010) permits genome-wide genetic screens to be carried out and represents a highly popular stock that lends itself to high-throughput experiments.

3. Order trends and projections

Stock centers have always closely followed new research trends by soliciting and accepting donations for new types of stocks. An explosion of publications using high-throughput technologies has been followed by the donation of resulting resources and their subsequent release. New additions have included new T-DNA lines and targeted mutagenesis lines as described in the previous section. A set of 1135 natural accessions sequenced by the 1001 Genomes project (Cao *et al.*, 2011; Long *et al.*, 2013; Schmitz *et al.*, 2013) has been available since the summer of 2015, enabling further analyses of the genetic and epigenetic variation of *Arabidopsis*. Based on the statistics showing that the number of orders is tightly linked with the number of papers published on *Arabidopsis* (Fig. 4), it is predicted that the overall number of orders will remain stable, as long as the newly published research resources become available shortly after publication (and researchers continue to use *Arabidopsis*!) As *Arabidopsis* stock collections are expected to grow, an expandable and robust Stock Center Database is envisioned that will be fully integrated within the larger *Arabidopsis* Information Portal framework at Araport.org (Krishnakumar *et al.*, 2015) to help support stock searching and ordering.

IV. Databases and online tools

The *Arabidopsis* community has a long history of organizing genomic and other data in publicly accessible portals. AtDB (Flanders *et al.*, 1998) and its successor, TAIR (Huala *et al.*, 2001), along with other, more specialized portals, initially served to organize sequence, bacterial artificial chromosome, physical maps, and other kinds of data necessary for and generated by the *Arabidopsis* Genome Initiative (*Arabidopsis* Genome Initiative, 2000). Over the following 15 yr, these portals and a collection of other online tools and databases have considerably altered the way that researchers design and interpret their experiments. Notably, the AtGenExpress project, a multinational initiative to document the transcriptome of *Arabidopsis*, generated thousands of gene expression data sets (Schmid *et al.*, 2005; Kilian *et al.*, 2007). These and many other expression data sets from individual researchers have not only helped to understand *Arabidopsis*'s response during development and to different environmental conditions, but have also enabled 'electronic northern' to be performed with tools such as the Bio-Analytic Resource (Toufighi *et al.*, 2005) or Genevestigator (Zimmermann *et al.*, 2004). Coexpression analysis –




Fig. 4 Number of seed stock orders from the Arabidopsis Biological Resource Center (ABRC) from 2001 to 2015 from the five countries that ordered the highest number of seed stocks. The number of orders per year is tabulated by the ABRC from 1 April of the indicated year to 31 March of the following year. The declining number of orders from the USA is being offset by increasing orders from China. The number of seed stocks ordered correlates strongly with the number of Arabidopsis publications per year, from the BIOSIS data described earlier in this review (inset). Since the graph shows only orders from the ABRC, countries ordering exclusively from the Nottingham Arabidopsis Stock Centre (e.g. those in Europe) or from other stock centers are not represented.

identifying genes with similar patterns of expression in a compendium of gene expression data sets – has emerged as a new kind of *in silico* screen to identify genes associated with a particular biological pathway (Usadel *et al.*, 2009). The ability to explore predicted (Geisler-Lee *et al.*, 2007) and documented (Dreze *et al.*, 2011; Mukhtar *et al.*, 2011 and many other smaller studies) protein–protein interactions is also another method for hypothesis generation. Further online and standalone computer tools can be used to help make sense of the large amount of data coming from -omics experiments, such as those for Gene Ontology (GO; Ashburner *et al.*, 2000) enrichment analysis. Tools for doing these and additional kinds of analyses are described by Brady & Provart (2009) and de De Lucas *et al.* (2014). Other useful software tools permit the easy mapping of causal mutations based on next-generation sequencing of pooled segregants (Schneeberger *et al.*, 2009; Austin *et al.*, 2011). Recently, after considerable community discussion and brain-storming mediated by the North American Arabidopsis Steering Committee (NAASC) and its international counterpart MASC, the Arabidopsis Information Portal araport.org (Krishnakumar *et al.*, 2015; International Arabidopsis Informatics Consortium, 2010, 2012) was launched, whose goal it is to integrate the large number of data sets generated in the past decade to further facilitate hypothesis generation at the click of a mouse.

V. Cell biology

For many of the same reasons that Arabidopsis has been a powerful model system to accelerate discovery in other fields of plant biology, Arabidopsis has also transformed plant cell biology. Cell biology can encompass a wide range of studies. Since many subjects that might be considered a part of cell biology, such as signal transduction, transport and the biology of cell walls, are covered elsewhere in this overview, here we focus on how Arabidopsis has contributed to studies of plant cell organization, featuring a handful of highlights.

A central tool in cell biology is microscopy, the technology that literally opened our eyes to the fine structure of life. Plants have a special place in this history, with the term ‘cell’ being credited to Robert Hooke when he applied his lens to cork, the box-like walls reminding him of the rooms occupied by monks, known as cells (Hooke, 1665). Nearly 200 yr later, the botanist Matthias Schleiden concluded that all plant life was in fact composed of cells (Schleiden, 1839), and together with Theodor Schwann articulated the foundational idea that cells were fundamental units of life (Schwann & Schleiden, 1847).

The confluence of Arabidopsis as a molecular genetic model, the discovery of intrinsically fluorescent proteins in jellyfish and other invertebrates, and advances in live cell microscopy addressed the limitations of then state-of-the art immunocytochemistry and enabled a new era of cellular discovery in plant science based on real-time, *in planta* observations.

The power of genetic analysis in Arabidopsis alone was revolutionary for plant cell biology. Prominent examples include screens for cell development and pattern using root hairs, trichomes, pollen, guard cells, roots and hypocotyl cells as models. These screens uncovered or helped to reveal important new insights into the function of essential players in cellular growth and morphology, such as cell wall biosynthetic enzymes and regulators (Arioli *et al.*, 1998; Nicol *et al.*, 1998; Favery *et al.*, 2001), the arp2/3 complex (Li *et al.*, 1999; Le *et al.*, 2003; Mathur *et al.*, 2003), plant-specific microtubule-associated proteins (Shoji *et al.*, 2004), and insights into the molecular basis of plant cell polarity, including novel cell polarity factors (Friml *et al.*, 2004; Dong *et al.*, 2009).

As powerful as genetic analysis can be, what can be learned is defined and limited by the phenotypes that can be observed and measured. Here, advances in microscopy have combined with genetics to push cell biology forward. An early example in plant biology was the immunocytochemical visualization of the inter-phase microtubule cytoskeleton as a phenotype in Arabidopsis, a screen that yielded the essential cytoskeletal protein MOR1 (Whittington *et al.*, 2001). This was a heroic screen that was not repeated owing to its difficulty. Genetic tagging with fluorescent proteins greatly simplified the marking of specific proteins and cellular structures. Protein localization alone could now be used as a phenotype, allowing for identification of peptide sequences that allow for subcellular targeting and the creation of useful subcellular markers (Cutler *et al.*, 2000). The distribution of such subcellular markers has since been used in direct microscopy screens to identify mutations or small molecules that perturb subcellular morphology and function (Hicks & Raikhel, 2012; Renna *et al.*, 2013). We can

anticipate more such screens in the future, especially as platforms for automated imaging and image analysis are developed further.

Fresh insights are being made into the age-old mystery of cytoskeletal organization, with the interphase microtubule cytoskeleton emerging as a self-organizing system (Deinum & Mulder, 2013) that may be directed by a variety of mechanisms including localized regulation of polymer dynamics (Ambrose *et al.*, 2011) and regulation of essential activities such as microtubule severing by states of polymer interaction. New plant-specific mechanisms for basic cellular processes are being discovered, including discovery of novel pathways for cellular polarity (Dong *et al.*, 2009), endocytosis (Gadeyne *et al.*, 2014), and vacuolar trafficking (Sanmartín *et al.*, 2007).

Arabidopsis is now coming into its own as a model for cell biological processes that are shared across a wide range of biological diversity. This is in part because cells can be easily observed *in situ*, permitting investigation of cellular behavior in a native developmental context. The architecture of plant cells is also helpful. The large central vacuole pushes cytosol and organelles into a thin shell at the cell periphery, producing a kind of natural optical section that increases signal to background, aiding detection of labeled single molecular complexes. Finally, plant cells display variations on basic cellular processes that allow for new experimental opportunities. An example is the architecture of cytokinesis, where the large phragmoplast is much more easily studied than in the more obscure midbody, which is thought to play a similar role as the phragmoplast in completing cytokinesis in animal cells.

In looking to the future, new experimental opportunities will be opened by the invention of methods to manipulate cells and their molecular components in time and space, as exemplified by the recent development of tools to regulate protein localization and activity by light in neuronal systems (Fenno *et al.*, 2011), and the continued innovation of imaging methods and tools to observe dynamic biological processes across greater ranges of time and space and with an expanding range of molecular and material measurements, including optical detection of metabolites, hormones and protein activity (Okumoto *et al.*, 2012; Michele *et al.*, 2013).

VI. Development

The study of plant development goes back centuries (Wolff, 1759), but a plausible case could be made that most of what we know now about the mechanisms of cellular differentiation, pattern formation, control of cell division, and the relationships between environment and plant development has been learned from experiments, largely with *Arabidopsis thaliana*, that have been done since the early 1990s.

The current systems biological approach to development has leveraged the power of Arabidopsis as a genetic model system (Rédei, 1975; Meyerowitz & Pruitt, 1985) through the addition of numerous tools built and shared among members of the Arabidopsis research community and our discovery and appreciation of other, more hidden attributes of this plant; some of these discoveries arose from the completion of the annotated genome sequence in 2000 (Arabidopsis Genome Initiative, 2000). For example, the complementarity between small RNAs and their

target genes enabled a revolution in understanding how miRNAs regulate spatial and temporal aspects of development. The particular history of genome duplications and losses simplified some regulatory systems (e.g. RETINOBLASTOMA-RELATED proteins and heterotrimeric G proteins, notwithstanding three recently discovered XLGs: Chakravorty *et al.*, 2015) while expanding others (like Polycomb complexes), allowing the roles of broadly conserved genes and processes to be approached in ways that are impossible in animals.

Our concept of development now is both more finely dissected and more integrated than it was 25 yr ago. Today, instead of monitoring gene expression of a few key regulators, we characterize transcriptional responses of entire genomes, and can do so within individual cells (Brady *et al.*, 2007; Yadav *et al.*, 2009; Adrian *et al.*, 2015; Efroni *et al.*, 2015). We are now modeling gene regulatory networks within cells and tissues in real time with fluorescent reporters and new methods of microscopy (Roeder *et al.*, 2010; Robinson *et al.*, 2011), and integrating multiple signaling modes (particularly hormone signaling) into these transcriptional networks.

Signaling, especially cell-to-cell communication, is a topic of long-term interest that has undergone conceptual revisions, often as a result of technical advances. We now appreciate that such communication is mediated not only by classical plant hormones, but also by newly discovered ones such as brassinosteroids and strigolactones, and that the classical hormone responses have considerable degrees of crosstalk (see the Hormones section). The diversity of signaling molecules, their modes of transport, and the degree to which they control each other are far greater than we imagined 25 yr ago. For example, investigating cell fate and patterning in the Arabidopsis root led to the identification of small regulatory RNAs and mobile transcription factors, whose movement and interaction dictate radial tissue organization (Nakajima *et al.*, 2001; Carlsbecker *et al.*, 2010). Peptides of a growing number of families (Katsir *et al.*, 2011) interact with numerous cell-surface receptors, and the resulting landscape demands new ways of understanding crosstalk and specificity. Beyond these chemical signals, we are also reappreciating the role of mechanical forces as a critical feedback leading to morphogenesis in developing plants, but now with the physical tools to manipulate forces and computational tools to understand their contributions (Kierzkowski *et al.*, 2012; Sampathkumar *et al.*, 2014).

An example of how technical advance and community effort have led to conceptual change in plant development is in the understanding of flower development. In 1980 there was no model of regulatory gene interaction to specify floral organs, but with the conceptual elegance of the ABC model and the cloning of the first ABC genes (in 1990, in *Antirrhinum* as well as *Arabidopsis*; Sommer *et al.*, 1990; Yanofsky *et al.*, 1990), it became clear that organ specification is a transcriptionally regulated process, and that combinatorial action of transcription factors leads to specific cellular differentiation (Bowman *et al.*, 1991; Coen & Meyerowitz, 1991). Work on auxin transport, cell division, and cell–cell communication then showed that these functions interact to create organs with patterned cells (Lampugnani *et al.*, 2013) in particular organ shapes and sizes. The addition of the integration of different

signaling modalities – hormone interactions, peptide signals, and mechanical signals, all feeding back on each other – has led to new hypotheses and computational models with predictive capabilities (Heisler *et al.*, 2010; Sahlin *et al.*, 2011), and a real understanding of the mechanisms and integrated systems that result in flowers (Smyth & Banks, 2014).

Another area where the understanding of mechanism has created a revolution is in the developmental responses of plants to their environments. As sessile organisms, plants use control of development to adapt, and plant size, branching, growth rate and time of flowering have long been known to depend on the abiotic environment. The past quarter-century has seen explanations for the mechanism of flowering time calculation (Jaeger *et al.*, 2013), including chromatin-based vernalization (described in more detail in the Epigenetics and epigenomics section) in response to temperature (Song *et al.*, 2013), the discovery of the long-known graft-transmissible flowering signal (Wigge *et al.*, 2005; Corbesier *et al.*, 2007; tomato FT (SFT) shown to be mobile signal: Lifschitz *et al.*, 2006; rice FT (Hd3a) shown to be mobile signal at same time as Arabidopsis FT: Tamaki *et al.*, 2007), and of the mechanisms by which day-length is measured to allow for seasonal flowering (Suárez-López *et al.*, 2001), all from work on Arabidopsis that was initially based on studies of single mutants and the genes responsible, and then progressed to studies of networks of interacting genes, cells and regulatory feedback loops, which have since been combined into a predictive set of models.

VII. Hormones

Phytohormones are a collection of structurally diverse small molecules that act at a distance to control plant growth, development and environmental responses. The 20th century was a golden age for phytohormone discovery that exploited sensitive bioassays coupled with purification to identify and elucidate the chemical structures of the major phytohormones. Further, biosynthetic pathways have been fully decoded, often using Arabidopsis (summarized in Davies, 2004; Davies writes in the preface to the 3rd edition that Arabidopsis has gone from a mention in the 1995 2nd edition to centre-stage in every chapter of the 3rd edition). Unsuccessful ‘pray and spray’ approaches to biochemically dissect mechanisms of action have given way to saturating mutational screens in Arabidopsis, which, quite remarkably, have revealed that most phytohormones signal through a limited number of mechanisms (see Fig. 5).

Early ideas on plant hormone signaling were heavily influenced by animal models of plasma membrane receptors that signal through cytoplasmic proteins which then transduce information to nuclear transcription factors. Surprisingly, this paradigm appears to hold only for the hormone brassinosteroid (BR). The major components of the BR pathway were defined in Arabidopsis, capitalizing on the characteristic dwarf phenotype of BR biosynthetic and signaling mutants (Li & Chory, 1997). Loss of function *BRI1* (*Brassinosteroid Insensitive 1*) alleles are BR-insensitive, and




Fig. 5 Abridged mechanisms of hormone signaling as elucidated in Arabidopsis. ER, endoplasmic reticulum; RAF K, RAPIDLY ACCELERATED FIBROSARCOMA; EIN2, ETHYLENE INSENSITIVE 2; GSK, GLYCOGEN SYNTHASE KINASE; LRR, leucine rich repeat; PM, plasma membrane; BR, brassinosteroid; JA, jasmonic acid; GA, gibberellic acid; SL, strigolactone; IAA, indole-3-acetic acid; JA-Ile, jasmonic acid-isoleucine conjugate; F-box receptor, F-box domain-containing receptor; SCF, SKP/CULLIN/F-box-containing complex; PP2C, PROTEIN PHOSPHATASE 2C; ABA, abscisic acid; SnRK, SUCROSE NONFERMENTING-1 (SNF1)-RELATED KINASE.

extensive analyses established that BRI1 is a plasma membrane-anchored LRR-kinase that binds BR along with a coreceptor BAK1 (Chang *et al.*, 1993; Li *et al.*, 2002). Activation of the BR receptor complex triggers a cascade of phosphorylation events that control transcription. Although the BR response pathway is the most 'animal'-like of the plant pathways, curiously animals perceive steroids primarily through soluble nuclear hormone receptors, which lack homologs in plants; Arabidopsis has therefore provided a window into alternate mechanisms of steroid action.

A second mechanism of hormone signaling derives from bacterial signaling modules and probably originated from the chloroplast genomes that were internalized early during plant evolution. Both ethylene (ET) and cytokinins (CKs) are perceived by endoplasmic reticulum (ER) membrane-anchored receptors that are homologous to bacterial two-component histidine kinases (Chang *et al.*, 1993; Inoue *et al.*, 2001). To identify mutations in these and other pathway components, Arabidopsis researchers took advantage of two classic physiological responses: the ET-controlled triple response of etiolated seedlings and the CK-induced greening of callus tissue (Bleecker *et al.*, 1988; Inoue *et al.*, 2001). For CK signaling, the pathway appears to be more like those seen in bacteria, which use a phosphorelay system from receptor to gene expression. In the case of ET, however, the pathway is novel, involving an 'animal'-like RAF kinase (CTR1) and an ER membrane-localized Nrap homolog (EIN2) between the 'bacterial'-like receptor (ETR family) and plant-specific transcription factors (EIN3 family) (Guzmán & Ecker, 1990; Roman *et al.*, 1995). Aside from defining the core components in ET signaling, Arabidopsis enabled rapid construction of double mutant lines and epistasis tests, which unequivocally ordered the steps of the pathway (Kieber *et al.*, 1993). Moreover, Arabidopsis enables quick construction of complex genotypes, which is onerous in many plants; this was essential to the rapid progress, as it allowed investigators to make functional insights despite the high redundancy that is typical of plant genomes. In this context, the ability to screen large numbers facilitated the isolation of rare dominant mutations that reveal phenotypes in spite of redundancy (Bleecker *et al.*, 1988; Inoue *et al.*, 2001).

The third and largest mechanism of plant hormone signaling involves soluble cytoplasmic receptors that modulate protein–protein interactions. Small molecule control of protein–protein interactions had previously been described with immunosuppressant drugs, but its role as a general signaling mechanism emerged from Arabidopsis. This 'plant'-like mechanism is used in the perception of auxin, jasmonates (JA), GAs, strigolactones (SL), salicylic acid (SA) and ABA (Lumba *et al.*, 2010). Auxin-regulated transcription is controlled by the AUX/IAA family of small corepressor proteins, whose stability is controlled by ubiquitin-mediated proteolysis via a canonical SKP/CULLIN/F-box-containing complex (SCF) mechanism, which itself turns out to play a hugely important role in a wide variety of plant processes, such as photomorphogenesis, circadian response and flower development (reviewed in Smalle & Vierstra, 2004). Auxin binds directly to the SCF's F-box, TIR1 or related Auxin-signaling F-Boxes, and creates a modified binding surface that stabilizes AUX/IAA binding, acting as a 'molecular glue' (Dharmasiri *et al.*, 2005;

Kepinski & Leyser, 2005; Tan *et al.*, 2007). This promotes AUX/IAA ubiquitylation by the SCF and subsequent degradation. Although the molecular details differ from hormone to hormone, auxin illustrates a general mechanism: plant hormones directly stabilize protein–protein interactions that are coupled to control of enzyme action. Auxin, JA and SA control E3-ligase action and repressor degradation. GA and, most likely, SL also control degradation, although not by direct binding to an SCF, but by stabilizing complexes that in turn trigger degradation of repressors (Dill *et al.*, 2001; Stirnberg *et al.*, 2002). Although ABA does not signal through ubiquitin-mediated proteolysis, it still regulates phosphatase activity by stabilizing protein complexes between receptors and clade A PP2Cs (Ma *et al.*, 2009; Park *et al.*, 2009). It is noteworthy that all the receptors, with the exception of BR, are encoded by more than a single gene, and redundancy therefore complicated their genetic identification. While classic ABA-insensitive screens defined key signaling components, *bona fide* receptors never emerged from genetic analyses because of redundancy (Koornneef *et al.*, 1984; Merlot *et al.*, 2002). The identification of a receptor-selective agonist illustrated how chemical probes can combat this redundancy when combined with the power of genetic analysis of mechanism (Park *et al.*, 2009). The SL, ABA and SCF-type receptors all belong to large gene families that hint at the existence of undiscovered hormones. If true, clever genetic screens will help define these functions.

VIII. The plant immune system and Arabidopsis research

Plant breeders have, since days that were fast on the heels of the re-discovery of Mendel's laws (Biffen 1905), been remarkably successful in keeping crops one step ahead of pathogen evolution, which follows a classic 'arms race' model. Furthermore, the efforts of H. H. Flor and others provided a genetic concept for how recognition of pathogens required specific molecules encoded by the pathogen acting as single triggers of disease resistance gene products (Flor, 1971). However, the relative speed by which microbes and insects with short generation times can evolve to avoid plant immune system detection, combined with large-scale monoculture agricultural practices, gives pathogens and pests a never-ending advantage in this game of immunological hide and seek.

Arabidopsis research has been the key instrument in the remarkable progress that has been made in our understanding of the molecular details of the plant immune system. Arabidopsis research has also been an important cornerstone in the bid to describe how plants respond to chewing pests. A few milestones are outlined here, focusing mostly on genetics and older breakthroughs. A detailed review of this history was published in 2010 (Nishimura & Dangel, 2010). Currently, Arabidopsis remains the primary organism in which to derive new principles regarding how plants and microbes interact. This was not always the case. In the mid-to-late 1980s, several of the initial grant proposals from those looking to apply Arabidopsis as a model for plant–pathogen interactions were met with skepticism that a noncrop plant with such a short generation time would not have real pathogens.

Pioneering work from Shauna Somerville (Tsuji & Somerville, 1988), Alan Slusarenko (Koch & Slusarenko, 1990) and Eric Holub (summarized in Holub & Beynon, 1996) laid this short-sightedness to rest. By 1991, the first small workshop was organized at the Max-Planck Institute for Plant Breeding Research in Cologne, Germany, to discuss progress to date and to collaborate around shared bacterial and oomycete pathogen strains and standard protocols between labs *well before the submission of papers*. Attendees at that workshop agreed that the very open flow of ideas, reagents, detailed protocols and people among laboratories that had already been established in this subdiscipline was instrumental in driving further rapid progress. By 1993 there was significant momentum in the field and progress was made in establishing useful pathosystems with which to look for Arabidopsis mutants altered in their responses to infection (Dangl, 1993).

Important milestones (by no means an exhaustive list) were quickly achieved: isolation of the first intracellular immune system receptors, including several from Arabidopsis using both mutagenesis (Bent *et al.*, 1994; Mindrinos *et al.*, 1994) and natural variation (Grant *et al.*, 1995), revealed that the ability to respond to and stop infection was mediated by a superfamily of related proteins, now called NLR proteins after their nucleotide-binding, leucine-rich repeat domains. Subsequently, it became apparent that NLRs were often encoded at loci that had been the object of plant breeding for many decades, the so-called disease resistance or R genes.

The notion that a susceptible parent could give rise to mutants that were even more susceptible defined the concept of 'basal defense' and led to the identification of many key immune system cornerstones (Glazebrook *et al.*, 1997). Equally compelling was the isolation of mutants defining host loci required for successful systemic acquired resistance (Cao *et al.*, 1994) and host loci required by pathogens for successful pathogen colonization (Vogel & Somerville, 2000). A critical breakthrough that unified disparate thinking about microbe- or plant-derived small signal molecules that triggered very rapid plant cell biological responses was the definition and isolation of the flagellin receptor (Gómez-Gómez & Boller, 2000). Equally important was the finding that many pathogens deliver virulence effectors into plant cells to disrupt immune system signaling (Gopalan *et al.*, 1996; Leister *et al.*, 1996). A reasonably clear conceptual view of the plant immune system was at hand (Dangl & Jones, 2001; Jones & Dangl, 2006).

Breathtaking progress continues. Plant immunology has expanded beyond binary interactions of one pathogen and the plant to encompass the plant's above- and below-ground microbiota (Vorholt, 2012; Bulgarelli *et al.*, 2013). Similarly, progress in plant immunology has overlapped with the dissection of the molecular switches guiding normal growth and development (Belkadir *et al.*, 2014). There is still no better test bed for the study of the plant immune system than Arabidopsis, once thought a useless little weed with no pathogens worthy of study.

IX. Signaling in response to abiotic stress

Counted among abiotic stresses are nonoptimal light environments, sub- or supraoptimal temperatures, drought, and nonideal soil conditions caused by nutrient limitation, salinity, or heavy

metal content. Other abiotic parameters that affect plant growth are those related to atmospheric constituents, including soil O₂ profiles, ambient humidity and CO₂ concentrations, and atmospheric pollutants such as ozone. Although impacts of abiotic stresses on plant physiological attributes have been studied for many decades, it was not until the development of Arabidopsis as a model system that the mechanistic underpinnings of many abiotic stress sensing and response pathways began to be uncovered. It is impossible to do justice to the countless discoveries made in this area based on the study of this small yet mighty weed. Instead, a small sampling of noteworthy findings that exemplify the essentiality of key Arabidopsis-based tools is presented.

The ease of forward genetic screens in Arabidopsis enabled the identification of the cryptochrome, phototropin and Zeitlupe blue light sensors (Ahmad & Cashmore, 1993; Christie *et al.*, 1998; Kim *et al.*, 2007b; note that Koornneef *et al.*, 1980 also identified *cry1* (which they called *hy4*) in an Arabidopsis genetic screen for light-inhibited hypocotyl elongation mutants); in all three cases, classical genetics approaches were central to definitive photoreceptor identification and to discrimination of which light responses are governed by which photoreceptor (Briggs, 2014). The 'new(est) kid on the block', the UV-B photoreceptor UVR8, essential in plant protective responses to damaging UV stress, was also first identified in Arabidopsis, starting with screens for mutants exhibiting UV-B hypersensitivity (Kliebenstein *et al.*, 2002; Rizzini *et al.*, 2011).

Another major plant photoreceptor, the R/FR sensor phytochrome, was first identified in lettuce (Borthwick *et al.*, 1952) as the sensing agent and cloned from oats (Hershey *et al.*, 1985), followed by work in tobacco identifying it as a photoreceptor that controls flowering in response to day-length. However, whole-genome sequencing in Arabidopsis confirmed the presence of five phytochrome (*PHY*) genes, and a wealth of studies on induced, natural and insertional *phy* mutations has allowed partitioning of the multitude of plant R/FR responses between and among the five *PHY* gene products (Franklin & Quail, 2010). The complete genome sequence of Arabidopsis also ushered in the -omics era. There is a plethora of publications on Arabidopsis transcriptome and proteome responses to abiotic stressors. More recently, impacts of stress on the metabolome (Weckwerth, 2011), and Barbara McClintock's prescient speculation that the epigenome is stress-regulated (McClintock, 1984) have also been experimentally evaluated in Arabidopsis (Chinnusamy & Zhu, 2009; Iwasaki & Paszkowski, 2014).

One example of the power of transcriptome analysis is seminal work demonstrating the importance of the CBF regulon in cold acclimation (Thomashow, 2010). The identification of CBF target genes and gene cascades in Arabidopsis illustrates the value of a model species in which extensive annotation of promoter and other regulatory elements can enhance the comprehensive elucidation of transcriptional networks containing hundreds of individual elements. As a complement to transcriptome measures of RNA abundance, methods have recently been developed, initially in Arabidopsis, to elucidate RNA structures *in vivo* on a genome-wide basis (Ding *et al.*, 2014). As many abiotic stresses (e.g. temperature, heavy metals) are known to affect folding of RNA *in vitro*, it will be

of interest to determine *in vivo* stress-regulated RNA 'structure-omes'. Yet another -ome related to RNA is the translome, defined as the set of mRNAs undergoing active translation. Low oxygen titer, as can occur in the soil during flooding stress, has wholesale impacts on cellular physiology as a result of its detrimental effect on cellular energy charge, and this has been illustrated by analysis of low O₂ effects on the translome (Branco-Price *et al.*, 2008).

Also of particular note are targeted approaches first developed in Arabidopsis that allow -omes to be analyzed in single cell types (Dinnyen *et al.*, 2008; Misra *et al.*, 2014). With regard to ABA signaling and drought stress, for example, the single most analyzed cell type is the guard cell. To date, -omics analyses of guard cells have been largely conducted in Arabidopsis and, to a lesser extent, in its brassicaceous relative, *Brassica napus*. And yet another -omics approach, chemical genomics, facilitated identification of the RCAR/PYR1/PYL soluble receptors for ABA (Cutler *et al.*, 2010; Raghavendra *et al.*, 2010). Indeed, identification of receptors for multiple plant hormones is one of the crowning achievements of Arabidopsis research (see the Hormones section).

While low soil water potentials initiate ABA and drought responses, nonoptimal soil mineral profiles also impact plant growth. Elegant biochemical genetic experiments in Arabidopsis have revealed that some inorganic nutrients, such as NO₃⁻, are sensed by the membrane protein that also transports them (Ho *et al.*, 2009). On the other hand, Na⁺ and heavy metals are toxic ions for which dedicated receptors have not been identified. Nevertheless, the proteins that bind and either extrude or sequester these toxic ions are essential to the survival of sessile plants subjected to adverse edaphic conditions. Our understanding of mechanisms underlying tolerance of such unfavorable conditions has particularly benefited from one of the more recent tools in the Arabidopsis arsenal: genomes from the Arabidopsis 1001 Genomes project, accompanied by extensive information on environmental conditions at the collection sites of these Arabidopsis accessions (Hancock *et al.*, 2011). This information, combined with development of statistical methods for genome-wide association study (GWAS) analysis (Segura *et al.*, 2012), is providing an unsurpassed resource for elucidation of the allelic diversity that can result from selective pressure exerted by environmental conditions (Assmann, 2013). One compelling illustration of the power of this approach comes from analysis of allelic diversity at the *AtHKT1* locus. The *AtHKT1* transporter reduces Na⁺ toxicity in the shoot via recovery of Na⁺ from the xylem sap. GWAS and functional analyses have shown that a particular allele of *AtHKT1* confers improved salinity tolerance and is prevalent in accessions found in saline environments (Rus *et al.*, 2006; Baxter *et al.*, 2010).

Finally, it is crucial to note that mechanisms of stress tolerance first identified in Arabidopsis have been introduced into crop species with resulting agronomic improvements. For example, overexpression of either native or Arabidopsis CBF transcription factors in a number of crops has been shown to improve cold tolerance, and improved salinity tolerance is conferred in crops upon expression of Arabidopsis or native transporters that promote Na⁺ extrusion or seclusion (Zhang *et al.*, 2004; Reguera *et al.*, 2012).

X. Pumps, channels, transporters and the like

The molecular nature of transporters was largely a mystery in plants before Arabidopsis started to be used as a model system. Most of what was known in the late 1980s was based on radiotracer studies and electrophysiology. Classical biochemistry led to the identification of a few transporters, the prime example of which is the plastidic triose phosphate translocator originally identified from spinach (Flügge *et al.*, 1989). But Arabidopsis rapidly took over as the key species in which transporter genes have been first identified. The proton ATPase protein was identified from oat and subsequently cloned simultaneously from *Nicotiana plumbaginifolia* and Arabidopsis (Schaller & Sussman, 1988; Boutry *et al.*, 1989; Harper *et al.*, 1989; Pardo & Serrano, 1989). At the same time, Sauer & Tanner (1989) identified the first sugar transporters from *Chlorella*. Just 1 yr later they had the homolog from Arabidopsis (Sauer *et al.*, 1990). A genetic trick rapidly changed the landscape: suppression cloning in yeast mutants. In 1992, two groups identified KAT1 and AKT1 potassium channels from Arabidopsis using yeast mutants (Anderson *et al.*, 1992; Sentenac *et al.*, 1992). In the same year, the first sucrose transporter was identified from spinach, followed by one from potato in 1993, and then in 1994 the Arabidopsis genes (Riesmeier *et al.*, 1992, 1993; Sauer & Stolz, 1994). Amino acid, iron, auxin, and many other transporters were then all identified first from Arabidopsis (Frommer *et al.*, 1993; Hsu *et al.*, 1993; Bennett *et al.*, 1996; Eide *et al.*, 1996; Gälweiler *et al.*, 1998). A unique exception was the identification of the sulfate transporter from a tropical grass *Stylosanthes* (Smith *et al.*, 1995). Arabidopsis became the most efficient system to clone transporter genes, and to study the localization, regulation and physiology by efficient access to mutants. Some of these important proteins were also studied in crops, yet discovery of new transporters is still occurring in Arabidopsis. And the value of a transporter identified from Arabidopsis is immediate – it explains quantitative trait loci (QTLs), and physiological roles discovered in Arabidopsis can be used to bring crops with improved properties to the field.

The use of Arabidopsis as a discovery tool continues to be productive: recently, a whole suite of novel sugar transporters, the SWEETs, has been identified in Arabidopsis. These have been shown to play roles across dicots in nectar secretion, in phloem loading, pollen nutrition and seed filling (Chen *et al.*, 2010, 2012, 2015; Lin *et al.*, 2014; Eom *et al.*, 2015). The application potential in the context of crop yield here is apparent, but surprisingly SWEETs also serve as pathogen susceptibility loci in rice, opening up the possibility of engineering resistance via TALENs or CRISPRs (Chen *et al.*, 2010; Li *et al.*, 2012). Vacuolar sugar storage is another example where Arabidopsis research lay the basis for discoveries with major biotechnological relevance: the sugar we eat is mainly derived from three sources: vacuole-stored sugars from sugarcane and sugarbeet or from fermented corn starch. The fermentation for winemaking critically depends on vacuolar sugars in grapes. Little was known about the transporters involved in vacuolar storage of sugars. A couple of years ago, Neuhaus's group identified a vacuolar monosaccharide transporter named TMT. TMT served as bait to identify a sucrose-transporting homolog (TST) in sugarbeet that is important for sugar accumulation in tap roots (Jung *et al.*, 2015).

In summary, *Arabidopsis* has been the workhorse that has not only served as an ideal system for identifying transporters, but also allowed for rapid identification of the role of the transporters and channels *in planta*. Importantly, the broad knowledge base derived from *Arabidopsis* has laid the groundwork for biotech applications, as summarized in detail elsewhere (Schroeder *et al.*, 2013).

XI. Cell walls, starch and lipids

The vast majority of what humans obtain from plants consists of cell walls, starch, sugar or lipids. Thus, many of the discoveries from *Arabidopsis* research have concerned synthesis, structure and function of these components.

1. Starch

Starch synthesis in *Arabidopsis* is similar to that in other plants. *Arabidopsis* has been important in identifying and characterizing enzymes involved in the synthesis of amylopectin and amylose, and the trimming and modification of nascent glucans to allow them to crystallize into a granule, and in identifying genes that control the initiation and number of starch granule per plastid (Crumpton-Taylor *et al.*, 2012; Seung *et al.*, 2015). By contrast, the pathway for degradation of starch in *Arabidopsis* is very different from that in germinating cereals, which is based on α -amylases (Smith, 2012). Screens for *starch excess* (*sex*) mutants in combination with reverse genetics revealed that solubilization of glucans at the granule surface is triggered by a unique cycle of phosphorylation and dephosphorylation (catalyzed by *SEX1/GWD* and *PWD*, and *PIPKIS1*, *LSF1* and *LSF2*, respectively), followed by a β -amylolytic attack to release maltose that is exported to the cytosol via a maltose exporter and converted to sucrose.

Source-limited plants regulate the rate of starch degradation to match the length of the night, thereby avoiding deleterious periods of carbon starvation at the end of the night (Stitt & Zeeman, 2012). Wild-type plants adjust the rates of both starch synthesis and degradation to maintain this pattern of starch turnover across a wide range of conditions. Control of synthesis has been attributed to allosteric and thioredoxin-dependent redox regulation of ADP-glucose pyrophosphorylase (Mugford *et al.*, 2014). The rate of starch degradation is regulated by the biological clock, which sets a maximum rate to ensure that starch is not exhausted before dawn (Smith, 2012; Stitt & Zeeman, 2012). These insights are leading to the development of new whole-plant growth models in which assumptions or rigid parameterization are replaced by modules that predict carbon allocation on the basis of the underlying regulatory pathways (Chew *et al.*, 2014).

As noted by Smith (2012), information from *Arabidopsis* cannot always simply be extrapolated to crop plants. However, the depth of understanding of *Arabidopsis* starch metabolism now informs and guides research in many crop plants.

2. Cell wall polysaccharides

Mutants in cell wall polysaccharide and proteoglycan composition have been isolated by screening for alterations in sugar composition

of the polysaccharides, but reverse genetics is the preferred approach because many of the proteins of interest can be identified by sequence similarity to related proteins (Velasquez *et al.*, 2011). The mutants have been essential in deciphering the biochemical mechanisms by which cell walls are synthesized and deposited. An example of the types of perplexing observations from mutant analysis that reflect the incompletely understood complexity of cell walls was the surprising discovery that the major hemicellulose, xyloglucan, is dispensable (Zabotina *et al.*, 2012). However, mutations that alter the structure of xyloglucan are deleterious (Kong *et al.*, 2015).

For many proteins implicated in cell wall synthesis and assembly, the purified proteins cannot be assayed by conventional biochemical assays, but the function can be convincingly tested or observed in mutants or transgenic plants. For example, cellulose synthase cannot be reliably assayed in cell-free extracts, but the development of transgenic plants that express fluorescently labeled protein allows the individual complexes to be observed while synthesizing cellulose in live cells (Paredes *et al.*, 2006). Live-cell imaging revealed that the pattern of cellulose deposition is controlled by association of the cellulose synthase complexes with microtubules and also revealed that this process can be disrupted by several herbicides and by mutations in a number of genes, such as *CSII/POM2*, *KOR* and *COB*, that appear to participate in the overall process. Thus, although many questions remain, such as what controls the amount of cellulose or the properties of the microfibrils, there is steady progress (McFarlane *et al.*, 2014).

Cell wall synthesis and deposition is part of the plant cell cycle, the response to pathogenesis and herbivory, and many aspects of morphogenesis. Several proteins that participate in the regulation of cell wall structure and function have been identified (Wolf *et al.*, 2012), but this aspect of plant biology is in its infancy.

3. Lignin

Lignification mutants of *Arabidopsis* were identified primarily by anatomical/histochemical identification of plants with reduced lignin deposition and screens for plants with reduced concentrations of the easily scored leaf metabolite, sinapoylmalate, the synthesis of which shares many steps in common with lignin (Vanholme *et al.*, 2010). Reverse genetic approaches have also contributed substantially, particularly for those enzymes encoded by multigene families, such as PAL and CAD, and where coexpression analyses (see the Databases and online tools section) suggested a potential role for previously unidentified catalysts, such as CSE. Along with other studies, the analysis of these mutants resulted in an almost total rewriting of a pathway that was thought to be well understood (Vanholme *et al.*, 2010).

Some mutants defective in lignin biosynthesis exhibit dwarfism and a collapsed xylem phenotype indicative of the importance of the polymer to cell wall rigidification and overall plant growth (Bonawitz *et al.*, 2014). Others, such as *cadc*, *cadd*, *fah1* and *comt1*, lead only to minor perturbations in development, even though, in these plants, lignin monomer composition is substantially altered, in some cases including noncanonical lignin subunits. Analysis of these mutants and other experiments have demonstrated the plasticity of lignin

composition that is an outcome of the nontemplate-mediated mode of synthesis of this biopolymer and have provided important insights into how lignin modification can be used to enhance the conversion of biomass in biofuel production scenarios.

Recent studies have begun to describe the complex regulatory machinery that orchestrates lignin deposition and maintains phenylpropanoid homeostasis (Bonawitz *et al.*, 2014; Taylor-Teeple *et al.*, 2015). The recent discovery that inactivation of three laccases leads to loss of lignin (Zhao *et al.*, 2013) has stimulated a reanalysis of the pathway of polymerization.

4. Lipids

Forward genetic approaches based on 'brute force' screens identified mutants with altered fatty acid, glycerolipid and wax compositions. These mutants facilitated studies directed at resolving uncertainties in some aspects of the pathways of fatty acid and lipid modification and facilitated identification and cloning of the corresponding genes. More importantly, mutant lines provided new insights into membrane structure and function (Wallis & Browse, 2010). The identification of mutants that globally affect oil biosynthesis led to the identification of critical transcription factors such as WRINKLED1 and LEC2 (Baud & Lepiniec, 2010). By contrast, discoveries in sphingolipid metabolism and function leveraged reverse genetics. Completion of the Arabidopsis genome sequence allowed comprehensive annotation of putative lipid-metabolism genes (Li-Beisson *et al.*, 2013), including homologs of yeast genes encoding enzymes of sphingolipid synthesis that were tested genetically by identifying T-DNA insertion mutants and biochemically by yeast expression (Markham *et al.*, 2013).

Transgenic approaches have been useful in many areas of lipid biology, particularly in seed lipid metabolism through experiments that engineer the fatty acid composition of vegetable oils. Efforts to increase the nutritional value of food oils and, alternatively, to produce unusual, industrially useful oil components have contributed to basic scientific knowledge (Nguyen *et al.*, 2010; Lu *et al.*, 2011). Recent studies highlighted differences between leaf and seed lipid metabolism that point the way to converting vegetative tissues of crop plants to accumulate oil (Fan *et al.*, 2014).

XII. Epigenetics and epigenomics: from genotype to phenotype

Compared with humans or other model systems, Arabidopsis has an unprecedented repertoire of chromatin regulatory proteins for methylating DNA, chemically modifying histones, or altering nucleosome spacing to control DNA accessibility to DNA-binding proteins (Pikaard & Mittelsten Scheid, 2014). Having multiple paralogous genes with partially overlapping functions presumably allows for fine-tuning of transcriptional programs that bring about physiological or developmental responses in a changeable environment. A consequence of having such a rich integrated network of partially redundant and overlapping functions is that null mutants in Arabidopsis chromatin regulatory protein genes are typically viable, allowing them to be studied. By contrast, corresponding mutations in metazoans are often embryo-lethal.

Genetic screens in Arabidopsis have uncovered biological roles of chromatin regulatory proteins in the control of developmental phase transitions, in cell identity switches and in activation of stress responses. One paradigm that emerged is that opposing types of chromatin regulators – those that inhibit and those that promote access to critical regions in the genomic DNA – together ensure that transcriptional activation of master regulatory proteins occurs at the correct stage and in the correct cell type and environment. Examples include activation of the floral homeotic genes in the developing flower, or silencing of the flowering time repressor *FLC* after prolonged cold (Goodrich *et al.*, 1997; Bastow *et al.*, 2004; Carles *et al.*, 2005; Aichinger *et al.*, 2009; Wu *et al.*, 2012; Hepworth & Dean, 2015, see this review for an overview of 'lessons' from *FLC*). In this manner, epigenetic mechanisms prevent 'unlicensed' gene activation that could be detrimental for growth or reproductive success.

Arabidopsis has been a particularly rich model system for understanding of DNA methylation and its regulation, with three cytosine methyltransferases responsible for maintaining CG, CHG and CHH methylation patterns (where H is a nucleotide other than G), and a fourth enzyme responsible for *de novo* methylation (Law & Jacobsen, 2010). Methylcytosine patterns can be perpetuated through multiple rounds of cell division, providing a basis for inheritance of epigenetic information. A fascinating aspect of *de novo* methylation is that noncoding RNAs specify the sites of DRM2 action via an elaborate process known as the RNA-directed DNA methylation pathway, whose elucidation by numerous laboratories represents a major effort over the past decade. Critical to the pathway are two plant-specific RNA polymerases, discovered in Arabidopsis and abbreviated as Pol IV and Pol V, that generate noncoding RNAs that guide the process. DNA methylation is not permanent (apart from a few cases, such as for *SUPERMAN* and *FWA*, where there does seem to be epigenetic inheritance, as discussed in Henderson & Jacobsen, 2007) and can be passively removed through DNA replication or actively through enzymatic activities. Indeed, discoveries of ROS1 and DEMETER as DNA demethylases were important achievements (Choi *et al.*, 2002; Gong *et al.*, 2002). Maternal expression of *DEMETER* was shown to demethylate and derepress *MEDEA* and other methylated target genes whose maternal-specific expression was hitherto unexplained. There is crosstalk between DNA methylation and histones, and Arabidopsis has played an important role in attaining this understanding (Lindroth *et al.*, 2004; Johnson *et al.*, 2007). An important example of a mechanism for crosstalk between DNA methylation and histone methylation came from studies of the partnership between the H3K9 methyl transferase KYP/SUVH4 and the CHG methyltransferase CMT3.

Epigenomics, the genome-wide study of chromatin and DNA transactions, is a new field of research. It emerged following the advent of genomics in the late 1990s and the parallel development of DNA microarray technologies, replaced a decade later by a flurry of high-throughput DNA sequencing-based approaches. For the first time, it became possible to investigate systematically the transcriptional activity and chromatin state of genes as well as

nongenic sequences along large genomic regions or entire genomes. Indeed, pioneering work in *Arabidopsis* has led to major advances in our understanding of how genomes are functionally organized, notably through the first detailed molecular characterization in any eukaryote of a large region of heterochromatin, the first genome-wide mapping of cytosine methylation at single nucleotide resolution, and the integrative analysis of the genomic distribution of DNA methylation, histone modifications and histone variants (Lippman *et al.*, 2004; Cokus *et al.*, 2008; Lister *et al.*, 2008; Roudier *et al.*, 2011; Yelagandula *et al.*, 2014). Equipped with this new knowledge and with the development of methods to investigate cell type-specific epigenomes as well as the three-dimensional organization of the genome in the nucleus, attention is now turning to the dynamics of the *Arabidopsis* epigenome in response to endogenous or exogenous cues during development and growth.

XIII. Natural variation and GWAS

Arabidopsis is arguably unique among model organisms in that variation among so-called ecotypes was of central interest from the very beginning (Laibach, 1943). Classical linkage mapping approaches have a long history in *Arabidopsis*; and studies using biparental recombinant inbred line populations of *Arabidopsis* ecotypes have identified several genes underlying QTLs (Weinig *et al.*, 2002; Alonso-Blanco *et al.*, 2009; Keurentjes *et al.*, 2011).

Genome-wide association is rapidly becoming the default approach for studying the genetics of natural variation (Vilhjálmsón & Nordborg, 2012). *A. thaliana* was one of first nonhuman organisms for which GWAS became possible (Atwell *et al.*, 2010), and it has played (and continues to play) an important role in developing statistical methodology appropriate for heavily structured populations (Zhao *et al.*, 2007; Korte *et al.*, 2012; Segura *et al.*, 2012; Vilhjálmsón & Nordborg, 2012). There were two main reasons for picking *Arabidopsis*: one good and one bad. The bad reason was that, just as in humans, linkage disequilibrium was extensive enough for it to be conceivable to carry out GWAS cheaply, using sparse single nucleotide polymorphism (SNP) markers, without resequencing entire genomes (Nordborg *et al.*, 2002). And, just as in humans, rapidly decreasing sequencing costs have nullified this advantage, while the major disadvantage of extensive linkage disequilibrium remains: the difficulty of identifying the causal site(s) among many highly associated ones (Atwell *et al.*, 2010).

However, the good reason for picking *Arabidopsis* remains a very good one: the availability of naturally occurring inbred ('pure') lines, which make it possible to capture local (and presumably locally adapted) genotypes, and grow them with replication under a variety of conditions. The generation and characterization of genome-wide polymorphism data was a natural extension of this (Nordborg *et al.*, 2005; Clark *et al.*, 2007). At present, several hundred lines have been sequenced, in the sense of having been aligned and compared to the existing reference genome (*Arabidopsis* Genome Initiative, 2000; Cao *et al.*, 2011; Gan *et al.*, 2011; Long *et al.*, 2013; Schmitz *et al.*, 2013), and a publication describing a set of well over 1000 is in progress

(www.1001genomes.org). When combined with the equally large collection for which dense SNP data are available (Horton *et al.*, 2012), over 2000 densely genotyped lines will be available.

The utilization of these lines is still far from ubiquitous, but is increasing. Studies involving hundreds of natural lines are time-consuming and many papers have yet to appear. Several studies have demonstrated the power of using GWAS as a tool for dissecting natural variation (Todesco *et al.*, 2010; Chao *et al.*, 2012; Karasov *et al.*, 2014), as well as for functional genomics (Todesco *et al.*, 2010; Chao *et al.*, 2012; Meijón *et al.*, 2014). GWAS is also likely to play an important role in interpreting genomic data sets, such as transcriptome or epigenome variation data (Schmitz *et al.*, 2013; Dubin *et al.*, 2015).

Where the *Arabidopsis* GWAS efforts have already had an impact is as a model for other studies, in particular in terms of statistical analysis. Unlike in humans, GWAS in *Arabidopsis* and maize were never carried out in 'case-control' settings (cases are people with a disease, and controls are a carefully selected set of healthy individuals from the same population), and it was thus immediately obvious that population structure would be a problem (Thornsberry *et al.*, 2001). The solution proposed (Yu *et al.*, 2006; Zhao *et al.*, 2007) – an adaptation of the classical 'animal model' from quantitative genetics (Henderson, 1984) – has since become the default approach in human GWAS as well as in other organisms. Indeed, GWAS in crops are now progressing much faster than in *Arabidopsis*, for the obvious reason that quantitative traits are of central interest in breeding (Huang *et al.*, 2011; Tian *et al.*, 2011).

The *Arabidopsis* germplasm collection is, of course, not only of interest for GWAS. The first effort to characterize polymorphism patterns on a large scale revealed a species-wide pattern of isolation by distance, with linkage disequilibrium decaying similarly to that in humans (Nordborg *et al.*, 2005). This work was followed by the generation of the first genome-wide SNP data (Clark *et al.*, 2007), which revealed interesting – and as yet unexplained – patterns of polymorphism. This work led directly to the generation of large, species-wide SNP data (Kim *et al.*, 2007a; Horton *et al.*, 2012), which, in addition to GWAS, have been used to search for patterns of climate adaptation (Hancock *et al.*, 2011; Fournier-Level *et al.*, 2011). The extensive sequence data currently being generated are certain to generate new insights, about *Arabidopsis* and about adaptation more generally.

XIV. Gene regulatory networks

Gene expression is controlled by large protein complexes that include transcription factors and coregulators acting in a combinatorial fashion and organized into gene regulatory grids, consisting of all the possible connections between transcription factors and the corresponding target genes, in what is also known as the protein–DNA interaction (PDI) space. Gene regulatory networks provide a temporal and/or spatial manifestation of the gene regulatory grid (Mejia-Guerra *et al.*, 2012). The past decades have witnessed a complete cataloguing of *Arabidopsis* transcription factors, and the development of tools and approaches to investigate individual and genome-wide transcription factor–target gene

interactions. These include transcription factor-centered chromatin immunoprecipitation (ChIP)-based techniques (e.g. ChIP-chip and ChIP-Seq) and gene-centered approaches such as yeast one-hybrid. When combined with genome-wide expression experiments, for example contrasting plants that harbor a mutant or a wild-type allele for the particular transcription factor gene, these approaches provide powerful tools to probe plant gene regulatory networks (Moreno-Risueno *et al.*, 2010).

Not surprisingly, most studies so far have centered on mutants with interesting developmental phenotypes, resulting in a loosely associated collection of subnetworks that are starting to provide a first glimpse of what the Arabidopsis gene regulatory grid may look like (Heyndrickx *et al.*, 2014). For example, the control of flower development provides one of the earliest examples of how homeotic genes encoding a small set of transcription factors, primarily from the MADS-box family, combinatorially specify different floral organs, resulting in what was initially known as the ABC model (Weigel & Meyerowitz, 1994), which has since evolved into the ABC(E) model (Ó'Maoiléidigh *et al.*, 2014). Direct targets for SEPALLATA3 (SEP3) (Kaufmann *et al.*, 2009) and APETALA1 (AP1) (Kaufmann *et al.*, 2010a) have been identified, providing important insights into how plants control developmental switches (Kaufmann *et al.*, 2010b). Stomatal cell-fate specification in the Arabidopsis leaf is in part controlled by combinatorial interactions between the basic helix–loop–helix (bHLH) factors SCREAM (SCRM), SCRM2, SPEECHLESS (SPCH), MUTE and FAMA (Pillitteri & Torii, 2012). The MYB transcription factor FOUR LIPS (FLP) controls the final symmetric cellular division that results in the formation of two guard cells by directly inhibiting the expression of several cell division genes (Xie *et al.*, 2010). Cell division genes are also directly controlled by the MYB and bHLH transcription factors GL1 and GL3 in the control of another epidermal differentiation, the formation of leaf hairs, or trichomes (Morohashi & Grotewold, 2009). Modeling of root gene regulatory networks involving the SHORT ROOT transcription factor and other components is described in the Modeling, bioinformatics, and systems biology section.

The Arabidopsis gene regulatory networks studied are not limited to developmental processes. For example, a circadian clock-controlled nitrate-responsive gene regulatory network includes the bZIP1 transcription factor (Gutiérrez *et al.*, 2008). In turn, bZIP1-controlled genes include several direct targets that are transiently bound by this transcription factor (hit-and-run transcriptional control), and which could have been easily missed in conventional ChIP-Seq experiments (Para *et al.*, 2014). In contrast to transcription factor-centered approaches, genome-wide yeast one-hybrid experiments have uncovered a comprehensive gene regulatory network for secondary cell wall biosynthesis (Taylor-Teeple *et al.*, 2015), which is likely to inform our understanding of the regulation of similarly essential processes in other plants.

Resources that integrate information on experimentally validated PDIs are valuable because finding the information from the literature and from public DNA-sequence repositories is laborious. One such resource in Arabidopsis is provided by the AGRIS (<http://arabidopsis.med.ohio-state.edu/>) knowledge base. The AtRegNet module in AGRIS currently displays over 16 000 experimentally

identified interactions between 94 transcription factors and the corresponding target genes. Clearly, while significant progress has been made, information on the targets of Arabidopsis transcription factors (estimated to be *c.* 2000) is in its infancy, but will rapidly expand with new data sets from protein-binding microarray studies (Weirauch *et al.*, 2014) and ChIP-Seq experiments with large numbers of Arabidopsis transcription factors (J. Ecker, unpublished).

XV. Modeling, bioinformatics, systems biology

The reference *A. thaliana* genome sequence and the facile nature of high-throughput gene expression profiling via microarray analyses resulted in a whole suite of data sets profiling the expression of individual genes across a range of cell, tissue and organ types. These and tools used to query them are described in the Databases and online tools section. Further functional genomic methodologies resulted in compilations of gene interactions including protein–protein interactions and protein–DNA interactions as described in the Gene regulatory networks section. In the midst of these massive data sets, however, elucidating biological novelty can still be an onerous task. In this section a subset of modeling and computational approaches that have been critical in the biological interpretation of these big data will be highlighted.

1. Gene expression in space and time

The Arabidopsis root has several developmental features that enable observation of cell type development in space and time. First, the root's stem cell niche is located at the root tip. Initial or stem cells divide and give rise to daughter cells. These daughter cells successively displace slightly older cells, resulting in a continuous file of cells along the root's longitudinal axis. Furthermore, most cell types are arranged in a rotationally symmetrical manner, which allows the observation of most cell types along the root's radial axis. In order to capture cell type-specific transcriptional profiles in the Arabidopsis root, fluorescent-activated cell sorting was coupled with transcriptional profiling (Birnbaum *et al.*, 2003; Brady *et al.*, 2007). Additionally, individual sections of the Arabidopsis root were captured and gene expression profiled, with each section representing a developmental time point. The cell type-specific transcription profiles often captured expression in only a subset of developmental stages. Furthermore, each section or developmental time point contained multiple cell types. These data could be analyzed with a series of bilinear equations that describe the proportion of a given cell type in a particular longitudinal section, in addition to the amount of expression of a gene in a given cell type at a specific developmental stage (Cartwright *et al.*, 2009). Using an iterative algorithm for finding approximate roots to these systems of bilinear equations, high-resolution spatiotemporal expression patterns of each gene could be reconstructed. These data may be visualized in Bio-Analytic Resource's EFP BROWSER (Winter *et al.*, 2007) and are a useful resource for plant biologists to determine when and where a given gene of interest is expressed in the Arabidopsis root at high spatiotemporal resolution.

2. Modeling root development and hormone responsiveness

Plant hormones have long been known to regulate plant growth, development and response to the environment. A variety of parameters have been described for passive and active transport of these hormones. The hormone auxin, for instance, is critical for root development but is transported by distinct transporters in distinct cell types and often in distinct directions (towards or away from the root tip). Modeling approaches have facilitated our understanding of how hormones regulate developmental processes by integrating two-dimensional and three-dimensional models of roots in conjunction with particular regulatory genes. In order to determine how a nested feedback circuit involving the transcription factors SHORT-ROOT, (SHR), SCARECROW (SCR), and the cell cycle mediators RETINOBLASTOMA-RELATED (RBR) and cyclin CYCLIND6;1 (CYCD6;1) precisely control the division of just a single stem cell daughter in a longitudinal file, a series of coupled ordinary differential equations (ODEs) was overlaid on a two-dimensional multicellular model of a root (Cruz-Ramirez *et al.*, 2012). These ODEs describe the wiring of this nested feedback circuit in addition to the influence of auxin on expression of *CYCD6;1*. The two-dimensional multicellular model of the root incorporated measurements of auxin transporter distribution and dynamics. Using these, a model was developed that explains how, in the stem cell daughter, the system is switched to the 'on' state to effect an asymmetric cell division (Cruz-Ramirez *et al.*, 2012). Similarly, in order to prevent the LAX3 transporter from being transiently expressed in multiple cell files, and thus not facilitating efficient lateral root emergence, modeling predicted that the auxin transporter PIN3 must be induced first, followed by LAX3 (Peret *et al.*, 2013). Experimental validation demonstrated that the model prediction was indeed correct. Recent reviews of root systems biology include those by Hill *et al.* (2013) and Wachsman *et al.* (2015). Finally, Boolean models have proved to be very useful in terms of predicting essential components of signal transduction networks in guard cell ABA and light response, with model predictions readily evaluated with Arabidopsis mutants (Li *et al.*, 2006).

3. Mapping transcriptional regulatory cascades

Developmental genetic studies have revealed hormone-mediated signaling via transcriptional cascades to be of prime importance in plant growth and development. The plant hormone ET controls fruit ripening, cell elongation and a multitude of other processes in part via the ETHYLENE INSENSITIVE3 (EIN3) transcription factor. Identifying the downstream targets of this transcription factor and their temporal regulation was recently characterized using a combination of ChIP coupled with sequencing and temporal whole-genome expression profiling (Chang *et al.*, 2013). Importantly, using the Dynamic Regulatory Events Miner (DREM; Schulz *et al.*, 2012), a series of four distinct waves of transcriptional regulation were identified. While the first wave of gene expression was quite variable and 'noisy', subsequent waves were less variable (Chang *et al.*, 2013).

Probabilistic functional gene networks utilize and integrate multiple, different data set types. The rationale for integrating these data sets and providing measures of their likelihood of interaction will enhance model accuracy and coverage. The AraNet network was generated using such an approach by integrating mRNA coexpression patterns from microarray data sets, known protein–protein interaction data sets, protein sequence features including protein domains, and similarity of phylogenetic profiles or the genomic context of bacterial or archaeobacterial homologs (Lee *et al.*, 2010). Finally, diverse gene–gene associations were incorporated from yeast, fly, worm and human genes based on orthology (Lee *et al.*, 2010). Importantly, AraNet was able to generate potential gene functions for over 7465 genes lacking functional annotation.

XVI. Synthetic biology

1. How Arabidopsis is helping plant synthetic biology bloom

The newly emerged discipline of synthetic biology aims at engineering genetic, signaling and/or metabolic pathways in a predictable manner in order to create novel functionalities (Purnick & Weiss, 2009).

Synthetic biology in Arabidopsis can be classified into two major approaches: engineering plant networks in heterologous contexts; and engineering novel networks in plants with genetic components from other organisms, as depicted in Fig. 6. These two approaches are complementary and synergistic, and together they promote a systems-level understanding of plant biology. A growing list of new functions engineered in yeast with plant proteins highlight the potential of the first approach, including small molecule-induced protein degradation (Nishimura *et al.*, 2009; Havens *et al.*, 2012) or proximity (Liang *et al.*, 2011), light-induced gene expression (Shimizu-Sato *et al.*, 2002), or protein splicing and sender–receiver communication (Tyszkiewicz & Muir, 2008). Analysis of signaling components in a heterologous setting can generate new hypotheses about pathway function in its original context, particularly where redundancy, shared signaling components and feedback play significant roles. For example, recapitulation of auxin signaling in yeast revealed that the rate of auxin-induced degradation is highly variable across the large Arabidopsis Aux/IAA family (Havens *et al.*, 2012); and that the rate of degradation of Aux/IAs is among the most effective tuning knobs for altering downstream transcriptional dynamics (Pierre-Jerome *et al.*, 2014). This led to the




Fig. 6 The main approaches for synthetic biology in Arabidopsis.

hypothesis that the rate of auxin-induced Aux/IAA turnover was acting as a checkpoint for auxin-regulated developmental events. Indeed, when a set of degradation rate variants in IAA14 was engineered in yeast and expressed in plants, a striking correlation was discovered between the dynamics of IAA14 turnover and the dynamics of lateral root initiation (Guseman *et al.*, 2015).

Recently, there have been several compelling demonstrations of the second approach: synthetic engineering of plant networks (Sun *et al.*, 2014; Park *et al.*, 2015). One remarkable example is the recent report of the re-engineered ABA receptor (Park *et al.*, 2015). A screen of PYR1 variants led to identification of a hexuple mutant receptor with a switch in ligand recognition from ABA to an existing agrochemical compound for blight pathogens, mandipropamid (Park *et al.*, 2015). When transgenic Arabidopsis expressing the PYR1^{MANDI} variant are exposed to mandipropamid, they have an ABA-like response, including transcriptional responses, stomatal closure and drought tolerance.

Synthetic biology relies on a set of tools that are increasingly becoming accessible to any molecular biologist. This could revolutionize both basic plant science and translational research for crops and energy biomass feedstock. The long history of such work in Arabidopsis lays the foundation for rapid gains in the near future.

VII. Conclusions and outlook

Research using the model plant *A. thaliana* has clearly transformed the field of plant biology in the past 50 yr and this journey continues. As mentioned in the Databases and online tools section, new frameworks to house existing and yet-to-be-generated Arabidopsis data like Aport.org are in the works. In the fall of 2014, the NAASC developed a funding proposal to National Science Foundation (NSF) entitled 'Arabidopsis Research and Training for the 21st Century (ART-21)'. This recently funded proposal will engage community members over the next 5 yr in activities to capitalize upon the excellence in Arabidopsis research and facilitate new training directions for the next generation of diverse Arabidopsis research scientists. Collectively, the goal is to identify emerging technologies, such as network modeling (Bastien *et al.*, 2015; Dong *et al.*, 2015; Leal Valentim *et al.*, 2015) or digital image analysis (Montenegro-Johnson *et al.*, 2015; Reuille *et al.*, 2015), needed for 21st-century biology, and skills relevant to various types of organizations, not only to academia.

Criticisms have emerged with respect to the future role of Arabidopsis research in the plant sciences. The first involves the belief that 'Arabidopsis is not a real plant'; instead, we should be putting our efforts into crops. A second criticism is that many important crops will not work like Arabidopsis, hence why bother working on this 'little plant' anymore? The fault of this logic lies in the misconceived view of a model genetic system. Model systems are not created to explain everything but to give researchers a touchstone or reference for comparison. Without Arabidopsis, how would plant scientists ask, 'Is it like Arabidopsis'? For example, differences clearly exist between the way the methylome(s) function in maize vs Arabidopsis (Li *et al.*, 2014b). However, without the

Arabidopsis baseline, how would we know this? But far more often than not, knowledge gained in Arabidopsis serves all of us well as a basis for follow-up experiments in other model plant species (e.g. in tomato: Piquerez *et al.*, 2014; since 2007, MASC has documented many translational examples in its annual reports, available at <http://arabidopsisresearch.org>). Although the NSF-funded Arabidopsis 2010 Project, which aimed to determine a function for all Arabidopsis genes, was wildly successful, 7540 Arabidopsis genes still have no GO biological process term associated with them, and 5944 have no GO molecular function term assigned (T. Berardini, TAIR, pers. comm., 21 May 2015). Determining the function of these will surely lead to a treasure-trove of new insights for plant biology in general. Interestingly, around the time of the publishing of the first draft of Arabidopsis genome, Lee Hartwell and colleagues suggested that genetic interactions would be essential to truly understanding the genotype–phenotype relationship in any organism (Hartman *et al.*, 2001). Hints of this vision are now clear from functional studies in yeast where double mutants can be systematically constructed (Tong *et al.*, 2004). It is difficult to see how parallel approaches in plants will be possible without Arabidopsis. It is ironic that, as governments around the world cut back on basic research (including Arabidopsis research) under the guise of 'cost' savings, such progress on a mechanistic understanding of how a plant or, for that matter, life works is within our reach. Perhaps with respect to Arabidopsis research funding, agencies should remember Winston Churchill's famous quote: 'This is not the end. It is not even the beginning of the end. But it is, perhaps, the end of the beginning'.

Acknowledgements

Citation and bibliographic data for Arabidopsis publications were provided by Thomson Reuters, from their BIOSIS database, with funding for this provided by the North American Arabidopsis Steering Committee. N.J.P. would also like to thank David Guttman, Darrell Desveaux, and Zane Caplansky for helpful discussions regarding the literature analysis section of this review.

References

- Adrian J, Chang J, Ballenger CE, Bargmann BOR, Alassimone J, Davies KA, Lau OS, Matos JL, Hachez C, Lanctot A *et al.* 2015. Transcriptome dynamics of the stomatal lineage: birth, amplification, and termination of a self-renewing population. *Developmental Cell* 33: 107–118.
- Ahmad M, Cashmore AR. 1993. *HY4* gene of *A. thaliana* encodes a protein with characteristics of a blue-light photoreceptor. *Nature* 366: 162–166.
- Aichinger Ernst, Villar CBR, Farrona S, Reyes JC, Hennig L, Köhler C. 2009. CHD3 proteins and polycomb group proteins antagonistically determine cell identity in Arabidopsis. *PLoS Genetics* 5: e1000605.
- Alonso JM, Ecker JR. 2006. Moving forward in reverse: genetic technologies to enable genome-wide phenomic screens in Arabidopsis. *Nature Reviews. Genetics* 7: 524–536.
- Alonso JM, Stepanova AN, Leisse TJ, Kim CJ, Chen H, Shinn P, Stevenson DK, Zimmerman J, Barajas P, Cheuk R *et al.* 2003. Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* 301: 653–657.
- Alonso-Blanco C, Aarts MGM, Bentsink L, Keurentjes JJB, Reymond M, Vreugdenhil D, Koornneef M. 2009. What has natural variation taught us about plant development, physiology, and adaptation? *Plant Cell* 21: 1877–1896.

- Altmann T, Schmidt R, Willmitzer L. 1992. Establishment of a gene tagging system in *Arabidopsis thaliana* based on the maize transposable element Ac. *Theoretical and Applied Genetics* 84: 371–383.
- Ambrose C, Allard JF, Cytrynbaum EN, Wasteneys GO. 2011. A CLASP-modulated cell edge barrier mechanism drives cell-wide cortical microtubule organization in *Arabidopsis*. *Nature Communications* 2: 430.
- Anderson JA, Huprikar SS, Kochian LV, Lucas WJ, Gaber RF. 1992. Functional expression of a probable *Arabidopsis thaliana* potassium channel in *Saccharomyces cerevisiae*. *Proceedings of the National Academy of Sciences, USA* 89: 3736–3740.
- Antunes MS, Jeff Smith J, Jantz D, Medford JI. 2012. Targeted DNA excision in *Arabidopsis* by a re-engineered homing endonuclease. *BMC Biotechnology* 12: 86.
- Arabidopsis Genome Initiative. 2000. Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408: 796–815.
- Arioli T, Peng L, Betzner AS, Burn J, Wittke W, Herth W, Camilleri C, Höfte H, Plazinski J, Birch R *et al.* 1998. Molecular analysis of cellulose biosynthesis in *Arabidopsis*. *Science* 279: 717–720.
- Ashburner M, Ball CA, Blake JA, Botstein D, Heather Butler J, Cherry M, Davis AP, Dolinski K, Dwight SS, Eppig JT *et al.* 2000. Gene ontology: tool for the unification of biology. *Nature Genetics* 25: 25–29.
- Assmann SM. 2013. Natural variation in abiotic stress and climate change responses in *Arabidopsis*: implications for twenty-first-century agriculture. *International Journal of Plant Sciences* 174: 3–26.
- Atwell S, Huang YS, Vilhjálmsson BJ, Willems G, Horton M, Li Y, Meng D, Platt A, Tarone AM, Hu TT *et al.* 2010. Genome-wide association study of 107 phenotypes in *Arabidopsis thaliana* inbred lines. *Nature* 465: 627–631.
- Austin RS, Vidaurre D, Stamatiou G, Breit R, Provart NJ, Bonetta D, Zhang J, Fung P, Gong Y, Wang PW *et al.* 2011. Next-generation mapping of *Arabidopsis* genes. *Plant Journal* 67: 715–725.
- Bancroft I, Bhatt AM, Sjödin C, Scofield S, Jones JD, Dean C. 1992. Development of an efficient two-element transposon tagging system in *Arabidopsis thaliana*. *Molecular & General Genetics* 233: 449–461.
- Bastien R, Douady S, Moulia B. 2015. A unified model of shoot tropism in plants: photo-gravi- and proprioception. *PLoS Computational Biology* 11: e1004037.
- Bastow R, Mylne JS, Lister C, Lippman Z, Martienssen RA, Dean C. 2004. Vernalization requires epigenetic silencing of FLC by histone methylation. *Nature* 427: 164–167.
- Baud S, Lepiniec L. 2010. Physiological and developmental regulation of seed oil production. *Progress in Lipid Research* 49: 235–249.
- Baxter I, Brazelton JN, Danni Y, Huang YS, Lahner B, Yakubova E, Li Y, Bergelson J, Borevitz JO, Nordborg M *et al.* 2010. A coastal cline in sodium accumulation in *Arabidopsis thaliana* is driven by natural variation of the sodium transporter AtHKT1;1. *PLoS Genetics* 6: e1001193.
- Bechtold N, Ellis J, Pelletier G. 1993. In planta agrobacterium mediated gene transfer by infiltration of adult *Arabidopsis thaliana* plants. *Comptes Rendus de l'Académie Des Sciences. Série 3. Sciences de La Vie* 316: 1194–1199.
- Bedell VM, Wang Y, Campbell JM, Poshusta TL, Starker CG, Krug RG, Tan W, Penheiter SG, Ma AC, Leung AYH *et al.* 2012. In vivo genome editing using a high-efficiency TALEN system. *Nature* 491: 114–118.
- Belkhadir Y, Yang L, Hetzel J, Dangel JL, Chory J. 2014. The growth-defense pivot: crisis management in plants mediated by LRR-RK surface receptors. *Trends in Biochemical Sciences* 39: 447–456.
- Bennett MJ, Marchant A, Green HG, May ST, Ward SP, Millner PA, Walker AR, Schulz B, Feldmann KA. 1996. *Arabidopsis AUX1* gene: a permease-like regulator of root gravitropism. *Science* 273: 948–950.
- Bent AF, Kunkel BN, Dahlbeck D, Brown KL, Schmidt R, Giraudat J, Leung J, Staskawicz BJ. 1994. *RPS2* of *Arabidopsis thaliana*: a leucine-rich repeat class of plant disease resistance genes. *Science* 265: 1856–1860.
- Biffen RH. 1905. Mendel's laws of inheritance and wheat breeding. *Journal of Agricultural Science* 1: 4–48.
- Birnbaum K, Shasha DE, Wang JY, Jung JW, Lambert GM, Galbraith DW, Benfey PN. 2003. A gene expression map of the *Arabidopsis* root. *Science* 302: 1956–1960.
- Bleecker AB, Estelle MA, Somerville C, Kende H. 1988. Insensitivity to ethylene conferred by a dominant mutation in *Arabidopsis thaliana*. *Science* 241: 1086–1089.
- Bonawitz ND, Kim JI, Tobimatsu Y, Ciesielski PN, Anderson NA, Ximenes E, Maeda J, Ralph J, Donohoe BS, Ladisch M *et al.* 2014. Disruption of mediator rescues the stunted growth of a lignin-deficient *Arabidopsis* mutant. *Nature* 509: 376–380.
- Borthwick HA, Hendricks SB, Parker MW, Toole EH, Toole VK. 1952. A reversible photoreaction controlling seed germination. *Proceedings of the National Academy of Sciences, USA* 38: 662–666.
- Boutry M, Michelet B, Goffeau A. 1989. Molecular cloning of a family of plant genes encoding a protein homologous to plasma membrane H⁺-translocating ATPases. *Biochemical and Biophysical Research Communications* 162: 567–574.
- Bowman JL, Smyth DR, Meyerowitz EM. 1991. Genetic interactions among floral homeotic genes of *Arabidopsis*. *Development* 112: 1–20.
- Brady SM, Orlando DA, Lee JY, Wang JY, Koch J, Dinneny JR, Mace D, Ohler U, Benfey PN. 2007. A high-resolution root spatiotemporal map reveals dominant expression patterns. *Science* 318: 801–806.
- Brady SM, Provart NJ. 2009. Web-queryable large-scale data sets for hypothesis generation in plant biology. *Plant Cell* 21: 1034–1051.
- Branco-Price C, Kaiser KA, Jang CJH, Larive CK, Bailey-Serres J. 2008. Selective mRNA translation coordinates energetic and metabolic adjustments to cellular oxygen deprivation and reoxygenation in *Arabidopsis thaliana*. *Plant Journal* 56: 743–755.
- Briggs Winslow R. 2014. Phototropism: some history, some puzzles, and a look ahead. *Plant Physiology* 164: 13–23.
- Brkljacic J, Grotewold E, Scholl R, Mockler T, Garvin DF, Vain P, Brutnell T, Sibout R, Bevan M, Budak H *et al.* 2011. Brachypodium as a model for the grasses: today and the future. *Plant Physiology* 157: 3–13.
- Bulgarelli D, Schlaeppli K, Spaepen S, Ver Loren van Themaat E, Schulze-Lefert P. 2013. Structure and functions of the bacterial microbiota of plants. *Annual Review of Plant Biology* 64: 807–838.
- Cao H, Bowling SA, Gordon S, Dong X. 1994. Characterization of an *Arabidopsis* mutant that is non-responsive to inducers of systemic acquired resistance. *Plant Cell* 6: 1583–1592.
- Cao J, Schneeberger K, Ossowski S, Gunther T, Bender S, Fitz J, Koenig D, Lanz C, Stegle O, Lippert C *et al.* 2011. Whole-genome sequencing of multiple *Arabidopsis thaliana* populations. *Nature Genetics* 43: 956–963.
- Carles CC, Choffnes-Inada D, Reville K, Lertpiriyapong K, Fletcher JC. 2005. *ULTRAPETALAI* encodes a SAND domain putative transcriptional regulator that controls shoot and floral meristem activity in *Arabidopsis*. *Development* 132: 897–911.
- Carlsbecker A, Lee J-Y, Roberts CJ, Dettmer J, Lehesranta S, Zhou J, Lindgren O, Moreno-Risueno MA, Vatén A, Thitamadee S *et al.* 2010. Cell signalling by microRNA165/6 directs gene dose-dependent root cell fate. *Nature* 465: 316–321.
- Cartwright DA, Brady SM, Orlando DA, Sturmels B, Benfey PN. 2009. Reconstructing spatiotemporal gene expression data from partial observations. *Bioinformatics* 25: 2581–2587.
- Chakravorty D, Gookin TE, Milner M, Yunqing Y, Assmann SM. 2015. Extra-large G proteins (XLGs) expand the repertoire of subunits in *Arabidopsis* heterotrimeric G protein signaling. *Plant Physiology* 169: 512–529.
- Chang C, Kwok SF, Bleecker AB, Meyerowitz EM. 1993. *Arabidopsis* ethylene-response gene ETR1: similarity of product to two-component regulators. *Science* 262: 539–544.
- Chang KN, Zhong S, Weirauch MT, Hon G, Pelizzola M, Li H, Huang S-SC, Schmitz RJ, Urich MA, Kuo D *et al.* 2013. Temporal transcriptional response to ethylene gas drives growth hormone cross-regulation in *Arabidopsis*. *eLife* 2: e00675.
- Chao D-Y, Silva A, Baxter I, Huang YS, Nordborg M, Danku J, Lahner B, Yakubova E, Salt DE. 2012. Genome-wide association studies identify heavy metal ATPase3 as the primary determinant of natural variation in leaf cadmium in *Arabidopsis thaliana*. *PLoS Genetics* 8: e1002923.
- Chen LQ, Lin IW, Qu XQ, Sosso D, McFarlane HE, Londoño A, Samuels AL, Frommer WB. 2015. A cascade of sequentially expressed sucrose transporters in the seed coat and endosperm provides nutrition for the *Arabidopsis* embryo. *Plant Cell* 27: 607–619.
- Chen LQ, Qu XQ, Hou BH, Sosso D, Osorio S, Fernie AR, Frommer WB. 2012. Sucrose efflux mediated by SWEET proteins as a key step for phloem transport. *Science* 335: 207–211.

- Chen L-Q, Hou B-H, Lalonde S, Takanaga H, Hartung ML, Qu X-Q, Guo W-J, Kim J-G, Underwood W, Chaudhuri B *et al.* 2010. Sugar transporters for intercellular exchange and nutrition of pathogens. *Nature* 468: 527–532.
- Chew YH, Wenden B, Flis A, Mengin V, Taylor J, Davey CL, Tindal C, Thomas H, Ougham H, de Reffye P *et al.* 2014. Multiscale digital Arabidopsis predicts individual organ and whole-organism growth. *Proceedings of the National Academy of Sciences, USA* 111: E4127–E4136.
- Chinnusamy V, Zhu J-K. 2009. Epigenetic regulation of stress responses in plants. *Current Opinion in Plant Biology* 12: 133–139.
- Choi Y, Gehring M, Johnson L, Hannon M, Harada JJ, Goldberg RB, Jacobsen SE, Fischer RL. 2002. DEMETER, a DNA glycosylase domain protein, is required for endosperm gene imprinting and seed viability in Arabidopsis. *Cell* 110: 33–42.
- Christie JM, Reymond P, Powell GK, Bernasconi P, Raibekas AA, Liscum E, Briggs WR. 1998. Arabidopsis NPH1: a flavoprotein with the properties of a photoreceptor for phototropism. *Science* 282: 1698–1701.
- Clark RM, Schweikert G, Toomajian C, Ossowski S, Zeller G, Shinn P, Warthmann N, Hu TT, Fu G, Hinds DA *et al.* 2007. Common sequence polymorphisms shaping genetic diversity in *Arabidopsis thaliana*. *Science* 317: 338–342.
- Clough SJ, Bent AF. 1998. Floral dip: a simplified method for agrobacterium-mediated transformation of *Arabidopsis thaliana*. *Plant Journal* 16: 735–743.
- Coen ES, Meyerowitz EM. 1991. The war of the whorls: genetic interactions controlling flower development. *Nature* 353: 31–37.
- Cokus SJ, Feng S, Zhang X, Chen Z, Merriman B, Haudenschild CD, Pradhan S, Nelson SF, Pellegrini M, Jacobsen SE. 2008. Shotgun bisulphite sequencing of the Arabidopsis genome reveals DNA methylation patterning. *Nature* 452: 215–219.
- Corbesier L, Vincent C, Jang S, Fornara F, Fan Q, Searle I, Giakountis A, Farrona S, Gissot L, Turnbull C *et al.* 2007. FT protein movement contributes to long-distance signaling in floral induction of Arabidopsis. *Science* 316: 1030–1033.
- Crumpton-Taylor M, Grandison S, Png KMY, Bushby AJ, Smith AM. 2012. Control of starch granule numbers in Arabidopsis chloroplasts. *Plant Physiology* 158: 905–916.
- Cruz-Ramirez A, Diaz-Trivino S, Blilou I, Grieneisen VA, Sozzani R, Zamioudis C, Miskolczi P, Nieuwland J, Benjamins R, Dhonukshe P *et al.* 2012. A bistable circuit involving SCARECROW-RETINOBLASTOMA integrates cues to inform asymmetric stem cell division. *Cell* 150: 1002–1015.
- Cutler SR, Ehrhardt DW, Griffiths JS, Somerville CR. 2000. Random GFP:cDNA fusions enable visualization of subcellular structures in cells of Arabidopsis at a high frequency. *Proceedings of the National Academy of Sciences, USA* 97: 3718–3723.
- Cutler SR, Rodriguez PL, Finkelstein RR, Abrams SR. 2010. Absciscic acid: emergence of a core signaling network. *Annual Review of Plant Biology* 61: 651–679.
- Dangl JL. 1993. The emergence of *Arabidopsis thaliana* as a model for plant-pathogen interactions. *Advances in Plant Pathology* 10: 127–155.
- Dangl JL, Jones JD. 2001. Plant pathogens and integrated defence responses to infection. *Nature* 411: 826–833.
- Davies PJ. 2004. *Plant hormones: biosynthesis, signal transduction, action!*. Dordrecht, the Netherlands: Kluwer Academic Publishers.
- De Lucas M, Provart NJ, Brady SM. 2014. Bioinformatic tools in Arabidopsis research. *Methods in Molecular Biology* 1062: 97–136.
- Deinum EE, Mulder BM. 2013. Modelling the role of microtubules in plant cell morphology. *Current Opinion in Plant Biology* 16: 688–692.
- Dharmasiri N, Dharmasiri S, Estelle M. 2005. The F-box protein TIR1 is an auxin receptor. *Nature* 435: 441–445.
- Dill A, Jung HS, Sun TP. 2001. The DELLA motif is essential for gibberellin-induced degradation of RGA. *Proceedings of the National Academy of Sciences, USA* 98: 14162–14167.
- Ding Y, Tang Y, Kwok CK, Zhang Y, Bevilacqua PC, Assmann SM. 2014. *In vivo* genome-wide profiling of RNA secondary structure reveals novel regulatory features. *Nature* 505: 696–700.
- Dinneny JR, Long TA, Wang JY, Jung JW, Mace D, Pointer S, Barron C, Brady SM, Schiefelbein J, Benfey PN. 2008. Cell identity mediates the response of Arabidopsis roots to abiotic stress. *Science* 320: 942–945.
- Dong X, Jiang Z, Peng Y-L, Zhang Z. 2015. Revealing shared and distinct gene network organization in Arabidopsis immune responses by integrative analysis. *Plant Physiology* 167: 1186–1203.
- Dong J, MacAlister CA, Bergmann DC. 2009. BASL controls asymmetric cell division in Arabidopsis. *Cell* 137: 1320–1330.
- Dreze M, Carvunis AR, Charlotiaux B, Galli M, Pevzner SJ, Tasan M, Ahn Y-Y, Braun P, Ecker J, Vidal M *et al.* 2011. Evidence for network evolution in an Arabidopsis interactome map. *Science* 333: 601–607.
- Dubin MJ, Zhang P, Meng D, Remigereau M-S, Osborne EJ, Casale FP, Drewe P, Kahles A, Vilhjálmsson B, Jagoda J *et al.* 2015. DNA methylation variation in Arabidopsis has a genetic basis and shows evidence of local adaptation. *eLife* 4: e05255.
- Efroni I, Ip P-L, Nawy T, Mello A, Birnbaum KD. 2015. Quantification of cell identity from single-cell gene expression profiles. *Genome Biology* 16: 9.
- Eide D, Broderius M, Fett J, Guerinot ML. 1996. A novel iron-regulated metal transporter from plants identified by functional expression in yeast. *Proceedings of the National Academy of Sciences, USA* 93: 5624–5628.
- Eom J-S, Chen L-Q, Sosso D, Julius BT, Lin IW, Xiao-Qing Q, Braun DM, Frommer WB. 2015. SWEETs, transporters for intracellular and intercellular sugar translocation. *Current Opinion in Plant Biology* 25: 53–62.
- Fan J, Yan C, Roston R, Shanklin J, Changcheng X. 2014. Arabidopsis lipins, PDAT1 acyltransferase, and sdp1 triacylglycerol lipase synergistically direct fatty acids toward B-oxidation, thereby maintaining membrane lipid homeostasis. *Plant Cell* 26: 4119–4134.
- Favery B, Ryan E, Foreman J, Linstead P, Boudonck K, Steer M, Shaw P, Dolan L. 2001. KOJAK encodes a cellulose synthase-like protein required for root hair cell morphogenesis in Arabidopsis. *Genes & Development* 15: 79–89.
- Feldmann KA, Marks MD. 1987. Agrobacterium-mediated transformation of germinating seeds of *Arabidopsis thaliana*: a non-tissue culture approach. *Molecular and General Genetics* 208: 1–9.
- Feng Z, Mao Y, Xu N, Zhang B, Wei P, Yang DL, Wang Z, Zhang Z, Zheng R, Yang L *et al.* 2014. Multigeneration analysis reveals the inheritance, specificity, and patterns of CRISPR/Cas-induced gene modifications in Arabidopsis. *Proceedings of the National Academy of Sciences, USA* 111: 4632–4637.
- Fenno L, Yizhar O, Deisseroth K. 2011. The development and application of optogenetics. *Annual Review of Neuroscience* 34: 389–412.
- Flanders DJ, Weng S, Petel FX, Cherry JM. 1998. AtDB, the *Arabidopsis thaliana* database, and graphical-web-display of progress by the Arabidopsis Genome Initiative. *Nucleic Acids Research* 26: 80–84.
- Flor HH. 1971. Current status of the gene-for-gene concept. *Annual Review of Phytopathology* 9: 275–296.
- Flügge UI, Fischer K, Gross A, Sebald W, Lottspeich F, Eckerskorn C. 1989. The triose phosphate-3-phosphoglycerate-phosphate translocator from spinach chloroplasts: nucleotide sequence of a full-length cDNA clone and import of the in vitro synthesized precursor protein into chloroplasts. *EMBO Journal* 8: 39–46.
- Fournier-Level A, Korte A, Cooper MD, Nordborg M, Schmitt J, Wilczek AM. 2011. A map of local adaptation in *Arabidopsis thaliana*. *Science* 334: 86–89.
- Franklin KA, Quail PH. 2010. Phytochrome functions in Arabidopsis development. *Journal of Experimental Botany* 61: 11–24.
- Friml J, Yang X, Michniewicz M, Weijers D, Quint A, Tietz O, Benjamins R, Ouwerkerk PBF, Ljung K, Sandberg G *et al.* 2004. A PINOID-dependent binary switch in apical-basal PIN polar targeting directs auxin efflux. *Science* 306: 862–865.
- Frommer WB, Hummel S, Riesmeier JW. 1993. Expression cloning in yeast of a cDNA encoding a broad specificity amino acid permease from *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* 90: 5944–5948.
- Gadeyne A, Sánchez-Rodríguez C, Vanneste S, Di Rubbo S, Zaubner H, Vanneste K, Van Leene J, De Winne N, Eeckhout D, Persiau G *et al.* 2014. The TPLATE adaptor complex drives clathrin-mediated endocytosis in plants. *Cell* 156: 691–704.
- Gälweiler L, Guan C, Müller A, Wisman E, Mendgen K, Yephremov A, Palme K. 1998. Regulation of polar auxin transport by AtPIN1 in Arabidopsis vascular tissue. *Science* 282: 2226–2230.
- Gan X, Stegle O, Behr J, Steffen JG, Drewe P, Hildebrand KL, Lyngsoe R, Schultheiss SJ, Osborne EJ, Sreedharan VT *et al.* 2011. Multiple reference genomes and transcriptomes for *Arabidopsis thaliana*. *Nature* 477: 419–423.

- Geisler-Lee J, O'Toole N, Ammar R, Provart NJ, Millar AH, Geisler M. 2007. A predicted interactome for Arabidopsis. *Plant Physiology* 145: 317–329.
- Glazebrook J, Rogers EE, Ausubel FM. 1997. Use of Arabidopsis for genetic dissection of plant defense responses. *Annual Review of Genetics* 31: 547–569.
- Gómez-Gómez L, Boller T. 2000. FLS2: an LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in Arabidopsis. *Molecular Cell* 5: 1003–1011.
- Gong Z, Morales-Ruiz T, Ariza RR, Roldán-Arjona T, David L, Zhu JK. 2002. ROS1, a repressor of transcriptional gene silencing in Arabidopsis, encodes a DNA glycosylase/lyase. *Cell* 111: 803–814.
- Goodrich J, Puangsomlee P, Martin M, Long D, Meyerowitz EM, Coupland G. 1997. A polycomb-group gene regulates homeotic gene expression in Arabidopsis. *Nature* 386: 44–51.
- Gopalan S, Bauer DW, Alfano JR, Loniello AO, He SY, Collmer A. 1996. Expression of the *Pseudomonas syringae* avirulence protein AvrB in plant cells alleviates its dependence on the hypersensitive response and pathogenicity (Hrp) secretion system in eliciting genotype-specific hypersensitive cell death. *Plant Cell* 8: 1095–1105.
- Grant MR, Godiard L, Straube E, Ashfield T, Lewald J, Sattler A, Innes RW, Dangel JL. 1995. Structure of the Arabidopsis RPM1 gene enabling dual specificity disease resistance. *Science* 269: 843–846.
- Guseman JM, Hellmuth A, Lancot A, Feldman TP, Moss BL, Klavins E, Calderon Villalobos LI, Nemhauser JL. 2015. Auxin-induced degradation dynamics set the pace for lateral root development. *Development* 142: 905–909.
- Gutiérrez RA, Stokes TL, Thum K, Xu X, Obertello X, Katari MS, Tanurdzic M, Dean A, Nero DC, McClung CR *et al.* 2008. Systems approach identifies an organic nitrogen-responsive gene network that is regulated by the master clock control gene *CCA1*. *Proceedings of the National Academy of Sciences, USA* 105: 4939–4944.
- Guzmán P, Ecker JR. 1990. Exploiting the triple response of Arabidopsis to identify ethylene-related mutants. *Plant Cell* 2: 513–523.
- Hancock AM, Brachi B, Faure N, Horton MW, Jarymowycz LB, Sperone FG, Toomajian C, Roux F, Bergelson J. 2011. Adaptation to climate across the *Arabidopsis thaliana* genome. *Science* 334: 83–86.
- Harper JF, Surowy TK, Sussman MR. 1989. Molecular cloning and sequence of cDNA encoding the plasma membrane proton pump (H⁺-ATPase) of *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* 86: 1234–1238.
- Hartman JL, Garvik B, Hartwell L. 2001. Principles for the buffering of genetic variation. *Science* 291: 1001–1004.
- Havens KA, Guseman JM, Jang SS, Pierre-Jerome E, Bolten N, Klavins E, Nemhauser JL. 2012. A synthetic approach reveals extensive tunability of auxin signaling. *Plant Physiology* 160: 135–142.
- Heisler MG, Hamant O, Krupinski P, Uyttewaald M, Ohno C, Jönsson H, Traas J, Meyerowitz EM. 2010. Alignment between PIN1 polarity and microtubule orientation in the shoot apical meristem reveals a tight coupling between morphogenesis and auxin transport. *PLoS Biology* 8: e1000516.
- Henderson CR. 1984. *Applications of linear models in animal breeding*. Guelph, Canada: University of Guelph Press.
- Henderson IR, Jacobsen SE. 2007. Epigenetic inheritance in plants. *Nature* 447: 418–424.
- Hepworth J, Dean C. 2015. Flowering Locus C's Lessons: conserved Chromatin Switches Underpinning Developmental Timing and Adaptation. *Plant Physiology* 168: 1237–1245.
- Hershey HP, Barker RF, Idler KB, Lissimore JL, Quail PH. 1985. Analysis of cloned cDNA and genomic sequences for phytochrome: complete amino acid sequences for two gene products expressed in etiolated avena. *Nucleic Acids Research* 13: 8543–8559.
- Heyndrickx KS, Van de Velde J, Wang C, Weigel D, Vandepoele K. 2014. A functional and evolutionary perspective on transcription factor binding in *Arabidopsis thaliana*. *Plant Cell* 26: 3894–3910.
- Hicks GR, Raikhel NV. 2012. Small molecules present large opportunities in plant biology. *Annual Review of Plant Biology* 63: 261–282.
- Hill K, Porco S, Lobet G, Zappala S, Mooney S, Draye X, Bennett MJ. 2013. Root systems biology: integrative modeling across scales, from gene regulatory networks to the rhizosphere. *Plant Physiology* 163: 1487–1503.
- Hilson P, Allemeersch J, Altmann T, Aubourg S, Avon A, Beynon J, Bhalerao RP, Bitton F, Caboche M, Cannoot B *et al.* 2004. Versatile gene-specific sequence tags for Arabidopsis functional genomics: transcript profiling and reverse genetics applications. *Genome Research* 14: 2176–2189.
- Ho C-H, Lin S-H, Heng-Cheng H, Tsay Y-F. 2009. CHL1 functions as a nitrate sensor in plants. *Cell* 138: 1184–1194.
- Holub EB, Beynon JL. 1996. Symbiology of mouse ear cress (*Arabidopsis thaliana*) and oomycetes. *Advances in Botanical Research* 24: 228–273.
- Hooke R. 1665. *Micrographia: or some physiological descriptions of minute bodies made by magnifying glasses?: with observations and inquiries thereupon*. London, UK: The Royal Society.
- Horton MW, Hancock AM, Huang YS, Toomajian C, Atwell S, Adam Auton N, Muliyil W *et al.* 2012. Genome-wide patterns of genetic variation in worldwide *Arabidopsis thaliana* accessions from the RegMap panel. *Nature Genetics* 44: 212–216.
- Hsu L-C, Chiou T-J, Chen L, Bush DR. 1993. Cloning a plant amino acid transporter by functional complementation of a yeast amino acid transport mutant. *Proceedings of the National Academy of Sciences, USA* 90: 7441–7445.
- Huala E, Dickerman AW, Garcia-Hernandez M, Weems D, Reiser L, LaFond F, Hanley D, Kiphart D, Zhuang M, Huang W *et al.* 2001. The Arabidopsis Information Resource (TAIR): a comprehensive database and web-based information retrieval, analysis, and visualization system for a model plant. *Nucleic Acids Research* 29: 102–105.
- Huang X, Zhao Y, Wei X, Li C, Wang A, Zhao Q, Li W, Guo Y, Deng L, Zhu C *et al.* 2011. Genome-wide association study of flowering time and grain yield traits in a worldwide collection of rice germplasm. *Nature Genetics* 44: 32–39.
- Inoue T, Higuchi M, Hashimoto Y, Seki M, Kobayashi M, Kato T, Tabata S, Shinozaki K, Kakimoto T. 2001. Identification of CRE1 as a cytokinin receptor from Arabidopsis. *Nature* 409: 1060–1063.
- International Arabidopsis Informatics Consortium. 2010. An International bioinformatics infrastructure to underpin the Arabidopsis Community. *Plant Cell* 22: 2530–2536.
- International Arabidopsis Informatics Consortium. 2012. Taking the next step: building an Arabidopsis information portal. *Plant Cell* 24: 2248–2256.
- Iwasaki M, Paszkowski J. 2014. Epigenetic memory in plants. *EMBO Journal* 33: 1987–1998.
- Jaeger KE, Pullen N, Lamzin S, Morris RJ, Wigge PA. 2013. Interlocking feedback loops govern the dynamic behavior of the floral transition in Arabidopsis. *Plant Cell* 25: 820–833.
- Johnson LM, Bostick M, Zhang X, Kraft E, Henderson I, Callis J, Jacobsen SE. 2007. The SRA methyl-cytosine-binding domain links DNA and histone methylation. *Current Biology* 17: 379–384.
- Jones AM, Chory J, Dangel JL, Estelle M, Jacobsen SE, Meyerowitz EM, Nordborg M, Weigel D. 2008. The impact of arabidopsis on human health: diversifying our portfolio. *Cell* 133: 939–943.
- Jones JDG, Dangel JL. 2006. The plant immune system. *Nature* 444: 323–329.
- Jung K-H, An G, Ronald PC. 2008. Towards a better bowl of rice: assigning function to tens of thousands of rice genes. *Nature Reviews Genetics* 9: 91–101.
- Jung B, Ludewig F, Schulz A, Meißner G, Wöstefeld N, Flügge U-I, Pommerrenig B, Wirsching P, Sauer N, Koch W *et al.* 2015. Identification of the transporter responsible for sucrose accumulation in sugar beet taproots. *Nature Plants* 1: 14001.
- Karasov TL, Kniskern JM, Gao L, DeYoung BJ, Ding J, Dubiella U, Lastra RO, Nallu S, Roux F, Innes RW *et al.* 2014. The long-term maintenance of a resistance polymorphism through diffuse interactions. *Nature* 512: 436–440.
- Katsir L, Davies KA, Bergmann DC, Laux T. 2011. Peptide signaling in plant development. *Current Biology* 21: R356–R364.
- Kaufmann K, Muñoz JM, Jauregui R, Airolidi CA, Smaczniak C, Krajewski P, Angenent GC. 2009. Target genes of the MADS transcription factor SEPALLATA3: integration of developmental and hormonal pathways in the Arabidopsis flower. *PLoS Biology* 7: e1000090.
- Kaufmann K, Pajaro A, Angenent GC. 2010b. Regulation of transcription in plants: mechanisms controlling developmental switches. *Nature Reviews. Genetics* 11: 830–842.

- Kaufmann K, Wellmer F, Muino JM, Ferrier T, Wuest SE, Kumar V, Serrano-Mislata A, Madueño F, Krajewski P, Meyerowitz EM *et al.* 2010a. Orchestration of floral initiation by APETALA1. *Science* 328: 85–89.
- Kepinski S, Leyser O. 2005. The Arabidopsis F-Box protein TIR1 is an auxin receptor. *Nature* 435: 446–451.
- Keurentjes JJB, Willems G, van Eeuwijk F, Nordborg M, Koornneef M. 2011. A comparison of population types used for QTL mapping in *Arabidopsis thaliana*. *Plant Genetic Resources* 9: 185–188.
- Kieber JJ, Rothenberg M, Roman G, Feldmann KA, Ecker JR. 1993. CTR1, a negative regulator of the ethylene response pathway in Arabidopsis, encodes a member of the Raf family of protein kinases. *Cell* 72: 427–441.
- Kierzkowski D, Nakayama N, Routier-Kierzkowska A-L, Weber A, Bayer E, Schorderet M, Reinhardt D, Kuhlemeier C, Smith RS. 2012. Elastic domains regulate growth and organogenesis in the plant shoot apical meristem. *Science* 335: 1096–1099.
- Kilian J, Whitehead D, Horak J, Wanke D, Weinl S, Batistic O, D'Angelo C, Bornberg-Bauer E, Kudla J, Harter K. 2007. The AtGenExpress global stress expression data set: protocols, evaluation and model data analysis of UV-B light, drought and cold stress responses. *Plant Journal* 50: 347–363.
- Kim W-Y, Fujiwara S, Suh S-S, Kim J, Kim Y, Han L, David K, Putterill J, Nam HG, Somers DE. 2007b. ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA in blue light. *Nature* 449: 356–360.
- Kim S, Plagnol V, Hu TT, Toomajian C, Clark RM, Ossowski S, Ecker JR, Weigel D, Nordborg M. 2007a. Recombination and linkage disequilibrium in *Arabidopsis thaliana*. *Nature Genetics* 39: 1151–1155.
- Kliebenstein DJ, Lim JE, Landry LG, Last RL. 2002. Arabidopsis UVR8 regulates ultraviolet-B signal transduction and tolerance and contains sequence similarity to human regulator of chromatin condensation 1. *Plant Physiology* 130: 234–243.
- Koch E, Slusarenko A. 1990. Arabidopsis is susceptible to infection by a downy mildew fungus. *Plant Cell* 2: 437–445.
- Kong Y, Peña MJ, Renna L, Avci U, Pattathil S, Tuomivaara ST, Xuemei Li RW-D, Brandizzi F, Hahn MG *et al.* 2015. Galactose-depleted xyloglucan is dysfunctional and leads to dwarfism in Arabidopsis. *Plant Physiology* 167: 1296–1306.
- Koornneef M, Meinke D. 2010. The development of Arabidopsis as a model plant. *Plant Journal* 61: 909–921.
- Koornneef M, Reuling G, Karssen CM. 1984. The isolation and characterization of abscisic acid-insensitive mutants of *Arabidopsis thaliana*. *Physiologia Plantarum* 61: 377–383.
- Koornneef M, Rolff E, Spruit CJP. 1980. Genetic control of light-inhibited hypocotyl elongation in *Arabidopsis thaliana* (L.) Heynh. *Zeitschrift Für Pflanzenphysiologie* 100: 147–160.
- Korte A, Vilhjálmsson BJ, Segura V, Platt A, Long Q, Nordborg M. 2012. A mixed-model approach for genome-wide association studies of correlated traits in structured populations. *Nature Genetics* 44: 1066–1071.
- Krishnakumar V, Hanlon MR, Contrino S, Ferlanti ES, Karamycheva S, Kim M, Rosen BD, Cheng C-Y, Moreira W, Mock SA *et al.* 2015. Araport: the Arabidopsis information portal. *Nucleic Acids Research* 43: D1003–D1009.
- Krysan PJ, Young JC, Tax F, Sussman MR. 1996. Identification of transferred DNA insertions within Arabidopsis genes involved in signal transduction and ion transport. *Proceedings of the National Academy of Sciences, USA* 93: 8145–8150.
- Laibach F. 1943. *Arabidopsis thaliana* (L.) Heynh. Als Objekt Für Genetische Und Entwicklungsphysiologische Untersuchungen. *Botanisches Archiv* 44: 439–455.
- Lampugnani ER, Kilinc A, Smyth DR. 2013. Auxin controls petal initiation in Arabidopsis. *Development* 140: 185–194.
- Law JA, Jacobsen SE. 2010. Establishing, maintaining and modifying DNA methylation patterns in plants and animals. *Nature Reviews Genetics* 11: 204–220.
- Le J, El-Assal SE-D, Basu D, Saad ME, Szymanski DB. 2003. Requirements for Arabidopsis ATARP2 and ATARP3 during epidermal development. *Current Biology* 13: 1341–1347.
- Leal Valentim F, van Mourik S, Posé D, Kim MC, Schmid M, van Ham RCHJ, Busscher M, Sanchez-Perez GF, Molenaar J, Angenent GC *et al.* 2015. A quantitative and dynamic model of the arabidopsis flowering time gene regulatory network. *PLoS ONE* 10: e0116973.
- Lee I, Ambaru B, Thakkar P, Marcotte EM, Rhee SY. 2010. Rational association of genes with traits using a genome-scale gene network for *Arabidopsis thaliana*. *Nature Biotechnology* 28: 149–156.
- Leister RT, Ausubel FM, Katagiri F. 1996. Molecular recognition of pathogen attack occurs inside of plant cells in plant disease resistance specified by the Arabidopsis genes RPS2 and RPM1. *Proceedings of the National Academy of Sciences, USA* 93: 15497–15502.
- Li S, Assmann SM, Albert R. 2006. Predicting essential components of signal transduction networks: a dynamic model of guard cell abscisic acid signaling. *PLoS Biology* 4: e312.
- Li J, Chory J. 1997. A putative leucine-rich repeat receptor kinase involved in brassinosteroid signal transduction. *Cell* 90: 929–938.
- Li Q, Eichten SR, Hermanson PJ, Zaunbrecher VM, Song J, Wendt J, Rosenbaum H, Madzima TF, Sloan AE, Huang J *et al.* 2014b. Genetic perturbation of the maize methylome. *Plant Cell* 26: 4602–4616.
- Li H, Lin Y, Heath RM, Zhu MX, Yang Z. 1999. Control of pollen tube tip growth by a rop GTPase-dependent pathway that leads to tip-localized calcium influx. *Plant Cell* 11: 1731–1742.
- Li T, Liu B, Spalding MH, Weeks DP, Yang B. 2012. High-efficiency TALEN-based gene editing produces disease-resistant rice. *Nature Biotechnology* 30: 390–392.
- Li J, Wen J, Lease KA, Doke JT, Tax FE, Walker JC. 2002. BAK1, an Arabidopsis LRR receptor-like protein kinase, interacts with BRI1 and modulates brassinosteroid signaling. *Cell* 110: 213–222.
- Li J-F, Zhang D, Sheen J. 2014a. Cas9-based genome editing in Arabidopsis and tobacco. *Methods in Enzymology* 546: 459–472.
- Liang FS, Ho WQ, Crabtree GR. 2011. Engineering the ABA Plant Stress Pathway for Regulation of Induced Proximity. *Science Signalling* 4: rs2.
- Li-Beisson Y, Shorrosh B, Beisson F, Andersson MX, Arondel V, Bates PD, Baud S, Bird D, DeBono A, Durrett TP *et al.* 2013. Acyl-lipid metabolism. *Arabidopsis Book* 11: e0161.
- Lifschitz E, Eviatar T, Rozman A, Shalit A, Goldshmidt A, Amsellem Z, Alvarez JP, Eshed Y. 2006. The tomato FT ortholog triggers systemic signals that regulate growth and flowering and substitute for diverse environmental stimuli. *Proceedings of the National Academy of Sciences, USA* 103: 6398–6403.
- Lin IW, Sosso D, Chen L-Q, Gase K, Kim S-G, Kessler D, Klinkenberg PM, Gorder MK, Houe B-H, Qu X-Q *et al.* 2014. Nectar secretion requires sucrose phosphate synthases and the sugar transporter SWEET9. *Nature* 508: 546–549.
- Lindroth AM, Shultz D, Jasencakova Z, Fuchs J, Johnson L, Schubert D, Patnaik D, Pradhan S, Goodrich J, Schubert I *et al.* 2004. Dual histone H3 methylation marks at lysines 9 and 27 required for interaction with CHROMOMETHYLASE3. *EMBO Journal* 23: 4286–4296.
- Lippman Z, Gendrel A-V, Black M, Vaughn MW, Dedhia N, McCombie WR, Lavine K, Mittal V, May B, Kasschau KD *et al.* 2004. Role of transposable elements in heterochromatin and epigenetic control. *Nature* 430: 471–476.
- Lister R, O'Malley RC, Tonti-Filippini J, Gregory BD, Berry CC, Millar AH, Ecker JR. 2008. Highly integrated single-base resolution maps of the epigenome in Arabidopsis. *Cell* 133: 523–536.
- Long Q, Rabanal FA, Meng D, Huber CD, Farlow A, Platzer A, Zhang Q, Vilhjálmsson BJ, Korte A, Nizhynska V *et al.* 2013. Massive genomic variation and strong selection in *Arabidopsis thaliana* lines from Sweden. *Nature Genetics* 45: 884–890.
- Lu C, Napier JA, Clemente TE, Cahoon EB. 2011. New frontiers in oilseed biotechnology: meeting the global demand for vegetable oils for food, feed, biofuel, and industrial applications. *Current Opinion in Biotechnology* 22: 252–259.
- Lumba S, Cutler S, McCourt P. 2010. Plant nuclear hormone receptors: a role for small molecules in protein-protein interactions. *Annual Review of Cell and Developmental Biology* 26: 445–469.
- Ma Y, Zostkiewicz I, Korte A, Moes D, Yang Y, Christmann A, Grill E. 2009. Regulators of PP2C phosphatase activity function as abscisic acid sensors. *Science* 324: 1064–1068.
- Markham JE, Lynch DV, Napier JA, Dunn TM, Cahoon EB. 2013. Plant sphingolipids: function follows form. *Current Opinion in Plant Biology* 16: 350–357.

- Mathur J, Mathur N, Kernebeck B, Hülskamp M. 2003. Mutations in actin-related proteins 2 and 3 affect cell shape development in *Arabidopsis*. *Plant Cell* 15: 1632–1645.
- McClintock B. 1984. The significance of responses of the genome to challenge. *Science* 226: 792–801.
- McFarlane HE, Döring A, Persson S. 2014. The cell biology of cellulose synthesis. *Annual Review of Plant Biology* 65: 69–94.
- Meijón M, Satbhai SB, Tsuchimatsu T, Busch W. 2014. Genome-wide association study using cellular traits identifies a new regulator of root development in *Arabidopsis*. *Nature Genetics* 46: 77–81.
- Meinke D, Scholl R. 2003. The preservation of plant genetic resources. Experiences with *Arabidopsis*. *Plant Physiology* 133: 1046–1050.
- Mejia-Guerra MK, Pomeranz M, Morohashi K, Grotewold E. 2012. From plant gene regulatory grids to network dynamics. *Biochimica et Biophysica Acta* 1819: 454–465.
- Merlot S, Mustilli A-C, Genty B, North H, Lefebvre V, Sotta B, Vavasseur A, Giraudat J. 2002. Use of infrared thermal imaging to isolate *Arabidopsis* mutants defective in stomatal regulation. *Plant Journal* 30: 601–609.
- Meyerowitz EM. 2001. Prehistory and history of *Arabidopsis* research. *Plant Physiology* 125: 15–19.
- Meyerowitz EM, Pruitt RE. 1984. *Genetic Variations of Arabidopsis thaliana*. Pasadena, CA, USA: California Institute of Technology.
- Meyerowitz EM, Pruitt RE. 1985. *Arabidopsis thaliana* and plant molecular genetics. *Science* 229: 1214–1218.
- Michele RD, Ast C, Loqué D, Ho CH, Andrade SLA, Lanquar V, Grossmann G, Gehne S, Kumke MU, Frommer WB. 2013. Fluorescent sensors reporting the activity of ammonium transporters in live cells. *eLife* 2: e00800.
- Mindrinis M, Katagiri F, Yu G-L, Ausubel FM. 1994. The *A. thaliana* disease resistance gene RPS2 encodes a protein containing a nucleotide-binding site and leucine-rich repeats. *Cell* 78: 1089–1099.
- Misra BB, Assmann SM, Chen S. 2014. Plant single-cell and single-cell-type metabolomics. *Trends in Plant Science* 19: 637–646.
- Montenegro-Johnson TD, Stamm P, Strauss S, Topham AT, Tsagris M, Wood ATA, Smith RS, Bassel GW. 2015. Digital single-cell analysis of plant organ development using 3DCellAtlas. *Plant Cell* 27: 1018–1033.
- Moreno-Risueno MA, Busch W, Benfey PN. 2010. Omics meet networks - using systems approaches to infer regulatory networks in plants. *Current Opinion in Plant Biology* 13: 126–131.
- Morohashi K, Grotewold E. 2009. A systems approach reveals regulatory circuitry for *Arabidopsis* trichome initiation by the GL3 and GL1 selectors. *PLoS Genetics* 5: e1000396.
- Mugford ST, Fernandez O, Brinton J, Flis A, Krohn N, Encke B, Feil R, Sulpice R, Lunn JE, Stitt M *et al.* 2014. Regulatory properties of ADP glucose pyrophosphorylase are required for adjustment of leaf starch synthesis in different photoperiods. *Plant Physiology* 166: 1733–1747.
- Mukhtar MS, Carvunis A-R, Dreze M, Eppe P, Steinbrenner J, Moore J, Tasan M, Galli M, Hao T, Nishimura MT *et al.* 2011. Independently evolved virulence effectors converge onto hubs in a plant immune system network. *Science* 333: 596–601.
- Nakajima K, Sena G, Nawy T, Benfey PN. 2001. Intercellular movement of the putative transcription factor SHR in root patterning. *Nature* 413: 307–311.
- Nannas NJ, Dawe RK. 2015. Genetic and genomic toolbox of *Zea mays*. *Genetics* 199: 655–669.
- Nguyen HT, Mishra G, Whittle E, Pidkowiak MS, Bevan SA, Merlo AO, Walsh TA, Shanklin J. 2010. Metabolic engineering of seeds can achieve levels of omega-7 fatty acids comparable with the highest levels found in natural plant sources. *Plant Physiology* 154: 1897–1904.
- Nicol F, His I, Jauneau A, Vernhettes S, Canut H, Höfte H. 1998. A plasma membrane-bound putative endo-1,4-beta-D-glucanase is required for normal wall assembly and cell elongation in *Arabidopsis*. *EMBO Journal* 17: 5563–5576.
- Nishimura MT, Dangl JL. 2010. *Arabidopsis* and the plant immune system. *Plant Journal* 61: 1053–1066.
- Nishimura K, Fukagawa T, Takisawa H, Kakimoto T, Kanemaki M. 2009. An auxin-based degron system for the rapid depletion of proteins in non-plant cells. *Nature Methods* 6: 917–922.
- Nordborg M, Borevitz JO, Bergelson J, Berry CC, Chory J, Hagenblad J, Kreitman M, Maloof JN, Noyes T, Oefner PJ *et al.* 2002. The extent of linkage disequilibrium in *Arabidopsis thaliana*. *Nature Genetics* 30: 190–193.
- Nordborg M, Hu TT, Ishino Y, Jhaveri J, Toomajian C, Zheng H, Bakker E, Calabrese P, Gladstone J, Goyal R *et al.* 2005. The pattern of polymorphism in *Arabidopsis thaliana*. *PLoS Biology* 3: e196.
- Okumoto S, Jones A, Frommer WB. 2012. Quantitative imaging with fluorescent biosensors. *Annual Review of Plant Biology* 63: 663–706.
- O'Malley RC, Ecker JR. 2010. Linking genotype to phenotype using the *Arabidopsis* unimutant collection. *Plant Journal* 61: 928–940.
- Ó'Maoléidigh DS, Graciet E, Wellmer F. 2014. Gene networks controlling *Arabidopsis thaliana* flower development. *New Phytologist* 201: 16–30.
- Para A, Li Y, Marshall-Colón A, Varala K, Francoeur NJ, Moran TM, Edwards MB, Hackley C, Bargmann BOR, Birnbaum KD *et al.* 2014. Hit-and-run transcriptional control by bZIP1 mediates rapid nutrient signaling in *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* 111: 10371–10376.
- Pardo JM, Serrano R. 1989. Structure of a plasma membrane H⁺-ATPase gene from the plant *Arabidopsis thaliana*. *Journal of Biological Chemistry* 264: 8557–8562.
- Paredes AR, Somerville CR, Ehrhardt DW. 2006. Visualization of cellulose synthase demonstrates functional association with microtubules. *Science* 312: 1491–1495.
- Park SY, Peterson FC, Mosquana A, Yao J, Volkman BF, Cutler SR. 2015. Agrochemical control of plant water use using engineered abscisic acid receptors. *Nature* 520: 545–548.
- Park S-Y, Fung P, Nishimura N, Jensen DR, Fujii H, Zhao Y, Lumba S, Santiago J, Rodrigues A, Chow T-F *et al.* 2009. Abscissic acid inhibits type 2C protein phosphatases via the PYR/PYL family of START proteins. *Science* 324: 1068–1071.
- Peret B, Middleton AM, French AP, Larrieu A, Bishopp A, Njo M, Wells DM, Porco S, Mellor N, Band LR *et al.* 2013. Sequential induction of auxin efflux and influx carriers regulates lateral root emergence. *Molecular Systems Biology* 9: 699.
- Pierre-Jerome E, Jang SS, Havens KA, Nemhauser JL, Klavins E. 2014. Recapitulation of the forward nuclear auxin response pathway in yeast. *Proceedings of the National Academy of Sciences, USA* 111: 9407–9412.
- Pikaard CS, Mittelsten Scheid O. 2014. Epigenetic regulation in plants. *Cold Spring Harbor Perspectives in Biology* 6: a019315.
- Pillitteri LJ, Torii KU. 2012. Mechanisms of stomatal development. *Annual Review of Plant Biology* 63: 591–614.
- Piquerez SJM, Harvey SE, Beynon JL, Ntoukakis V. 2014. Improving crop disease resistance: lessons from research on *Arabidopsis* and tomato. *Frontiers in Plant Science* 5: 671.
- Purnick PEM, Weiss R. 2009. The second wave of synthetic biology: from modules to systems. *Nature Reviews Molecular Cell Biology* 10: 410–422.
- Raghavendra AS, Gonugunta VK, Christmann A, Grill E. 2010. ABA perception and signalling. *Trends in Plant Science* 15: 395–401.
- Rédei GP. 1975. *Arabidopsis* as a genetic tool. *Annual Review of Genetics* 9: 111–127.
- Reguera M, Peleg Z, Blumwald E. 2012. Targeting metabolic pathways for genetic engineering abiotic stress-tolerance in crops. *Biochimica et Biophysica Acta* 1819: 186–194.
- Renna L, Stefano G, Majeran W, Micalella C, Meinel T, Giglione C, Brandizzi F. 2013. Golgi traffic and integrity depend on N-myristoyl transferase-1 in *Arabidopsis*. *Plant Cell* 25: 1756–1773.
- Reuille PB, Routier-Kierzkowska AL, Kierzkowski D, Bassel GW, Schüpbach T, Tauriello G, Bajpai N, Strauss S, Weber A, Kiss A *et al.* 2015. MorphoGraphX: a platform for quantifying morphogenesis in 4D. *eLife* 4: e05864.
- Riesmeier JW, Hirner B, Frommer WB. 1993. Potato sucrose transporter expression in minor veins indicates a role in phloem loading. *Plant Cell* 5: 1591–1598.
- Riesmeier JW, Willmitzer L, Frommer WB. 1992. Isolation and characterization of a sucrose carrier cDNA from spinach by functional expression in yeast. *EMBO Journal* 11: 4705–4713.

- Rizzini L, Favory J-J, Cloix C, Faggionato D, O'Hara A, Kaiserli E, Baumeister R, Schäfer E, Nagy F, Jenkins GI *et al.* 2011. Perception of UV-B by the Arabidopsis UVR8 protein. *Science* 332: 103–106.
- Robinson S, Barbier de Reuille P, Chan J, Bergmann D, Prusinkiewicz P, Coen E. 2011. Generation of spatial patterns through cell polarity switching. *Science* 333: 1436–1440.
- Roeder AHK, Chickarmane V, Cunha A, Obara B, Manjunath BS, Meyerowitz EM. 2010. Variability in the control of cell division underlies sepal epidermal patterning in *Arabidopsis thaliana*. *PLoS Biology* 8: e1000367.
- Roman G, Lubarsky B, Kieber JJ, Rothenberg M, Ecker JR. 1995. Genetic analysis of ethylene signal transduction in *Arabidopsis thaliana*: five novel mutant loci integrated into a stress response pathway. *Genetics* 139: 1393–1409.
- Roudier F, Ahmed I, Bérard C, Sarazin A, Mary-Huard T, Cortijo S, Bouyer D, Caillieux E, Duvernios-Berther E, Al-Shikhley L *et al.* 2011. Integrative epigenomic mapping defines four main chromatin states in Arabidopsis. *EMBO Journal* 30: 1928–1938.
- Rus A, Baxter I, Muthukumar B, Gustin J, Lahner B, Yakubova E, Salt DE. 2006. Natural variants of AtHKT1 enhance Na⁺ accumulation in two wild populations of Arabidopsis. *PLoS Genetics* 2: e210.
- Sahlin P, Melke P, Jönsson H. 2011. Models of sequestration and receptor cross-talk for explaining multiple mutants in plant stem cell regulation. *BMC Systems Biology* 5: 2.
- Sampathkumar A, Yan A, Krupinski P, Meyerowitz EM. 2014. Physical forces regulate plant development and morphogenesis. *Current Biology* 24: R475–R483.
- Sanmartín M, Ordóñez A, Sohn EJ, Robert S, Sánchez-Serrano JJ, Surpin MA, Raikhel NV, Rojo E. 2007. Divergent functions of VTI12 and VTI11 in trafficking to storage and lytic vacuoles in Arabidopsis. *Proceedings of the National Academy of Sciences, USA* 104: 3645–3650.
- Sauer N, Friedländer K, Gräml-Wicke U. 1990. Primary structure, genomic organization and heterologous expression of a glucose transporter from *Arabidopsis thaliana*. *EMBO Journal* 9: 3045–3050.
- Sauer N, Stolz J. 1994. SUC1 and SUC2: two sucrose transporters from *Arabidopsis thaliana*; expression and characterization in Baker's yeast and identification of the histidine-tagged protein. *Plant Journal* 6: 67–77.
- Sauer N, Tanner W. 1989. The hexose carrier from *Chlorella*. cDNA cloning of a eukaryotic H⁺ cotransporter. *FEBS Letters* 259: 43–46.
- Schaller GE, Sussman MR. 1988. Isolation and sequence of tryptic peptides from the proton-pumping ATPase of the oat plasma membrane. *Plant Physiology* 86: 512–516.
- Schleiden MJ. 1839. Contributions to phytogenesis. *Archiv für Anatomie, Physiologie und wissenschaftliche Medizin* 1838: 137–176.
- Schmid M, Davison TS, Henz SR, Pape UJ, Demar M, Vingron M, Schölkopf B, Weigel D, Lohmann JU. 2005. A gene expression map of *Arabidopsis thaliana* development. *Nature Genetics* 37: 501–506.
- Schmitz RJ, Schultz MD, Urich MA, Nery JR, Pelizzola M, Libiger O, Alix A, McCosh RB, Chen H, Schork NJ *et al.* 2013. Patterns of population epigenomic diversity. *Nature* 495: 193–198.
- Schneeberger K, Ossowski S, Lanz C, Juul T, Petersen AH, Nielsen KL, Jørgensen J-E, Weigel D, Andersen SU. 2009. SHOREmap: simultaneous mapping and mutation identification by deep sequencing. *Nature Methods* 6: 550–551.
- Schroeder JI, Delhaize E, Frommer WB, Gueriot ML, Harrison MJ, Herrera-Estrella L, Horie T, Kochain LV, Munns R, Nishizawa NK *et al.* 2013. Using membrane transporters to improve crops for sustainable food production. *Nature* 497: 60–66.
- Schulz MH, Devanny WE, Gitter A, Zhong S, Ernst J, Bar-Joseph Z. 2012. DREM 2.0: improved reconstruction of dynamic regulatory networks from time-series expression data. *BMC Systems Biology* 6: 104.
- Schwab R, Ossowski S, Riester M, Warthmann N, Weigel D. 2006. Highly specific gene silencing by artificial microRNAs in Arabidopsis. *Plant Cell* 18: 1121–1133.
- Schwann T, Schleiden MJ. 1847. *Microscopical researches into the accordance in the structure and growth of animals and plants*. London, UK: The Sydenham Society.
- Segura V, Vilhjálmsson BJ, Platt A, Korte A, Seren Ü, Long Q, Nordborg M. 2012. An efficient multi-locus mixed-model approach for genome-wide association studies in structured populations. *Nature Genetics* 44: 825–830.
- Sentenac H, Bonneaud N, Minet M, Lacroute F, Salmon JM, Gaymard F, Grignon C. 1992. Cloning and expression in yeast of a plant potassium ion transport system. *Science* 256: 663–665.
- Sessions A, Burke E, Presting G, Aux G, McElver J, Patton D, Dietrich B, Ho P, Bacwaden J, Ko C *et al.* 2002. A high-throughput arabidopsis reverse genetics system. *Plant Cell* 14: 2985–2994.
- Seung D, Soyk S, Coiro M, Maier BA, Eicke S, Zeeman SC. 2015. PROTEIN TARGETING TO STARCH is required for localising GRANULE-BOUND STARCH SYNTHASE to starch granules and for normal amylose synthesis in Arabidopsis. *PLoS Biology* 13: e1002080.
- Shimizu-Sato S, Huq E, Tepperman JM, Quail PH. 2002. A light-switchable gene promoter system. *Nature Biotechnology* 20: 1041–1044.
- Shoji T, Narita NN, Hayashi K, Hayashi K, Asada J, Hamada T, Sonobe S, Nakajima K, Hashimoto T. 2004. Plant-specific microtubule-associated protein SPIRAL2 is required for anisotropic growth in Arabidopsis. *Plant Physiology* 136: 3933–3944.
- Smalle J, Vierstra RD. 2004. The ubiquitin 26S proteasome proteolytic pathway. *Annual Review of Plant Biology* 55: 555–590.
- Smith AM. 2012. Starch in the Arabidopsis plant. *Starch - Stärke* 64: 421–434.
- Smith FW, Ealing PM, Hawkesford MJ, Clarkson DT. 1995. Plant members of a family of sulfate transporters reveal functional subtypes. *Proceedings of the National Academy of Sciences, USA* 92: 9373–9377.
- Smyth DR, Banks JA. 2014. Editorial overview: plant morphogenesis-new understanding of its organization and evolution. *Current Opinion in Plant Biology* 17: iv–viii.
- Somerville C, Koornneef M. 2002. A fortunate choice: the history of arabidopsis as a model plant. *Nature Reviews Genetics* 3: 883–889.
- Sommer H, Beltrán JP, Huijser P, Pape H, Lönning WE, Saedler H, Schwarz-Sommer Z. 1990. Deficiens, a homeotic gene involved in the control of flower morphogenesis in *Antirrhinum majus*: the protein shows homology to transcription factors. *EMBO Journal* 9: 605–613.
- Song J, Irwin J, Dean C. 2013. Remembering the prolonged cold of winter. *Current Biology* 23: R807–R811.
- Stirnberg P, van De Sande K, Ottoline Leyser HM. 2002. MAX1 and MAX2 control shoot lateral branching in Arabidopsis. *Development* 129: 1131–1141.
- Stitt M, Zeeman SC. 2012. Starch turnover: pathways, regulation and role in growth. *Current Opinion in Plant Biology* 15: 282–292.
- Suárez-López P, Wheatley K, Robson F, Onouchi H, Valverde F, Coupland G. 2001. CONSTANS mediates between the circadian clock and the control of flowering in Arabidopsis. *Nature* 410: 1116–1120.
- Sun B, Looi LS, Guo S, He Z, Gan ES, Huang J, Xu Y, Wee WY, Ito T. 2014. Timing mechanism dependent on cell division is invoked by polycomb eviction in plant stem cells. *Science* 343: 1248559.
- Tamaki S, Matsuo S, Wong HL, Yokoi S, Shimamoto K. 2007. Hd3a protein is a mobile flowering signal in rice. *Science* 316: 1033–1036.
- Tan X, Calderon-Villalobos LIA, Sharon M, Zheng C, Robinson CV, Estelle M, Zheng N. 2007. Mechanism of auxin perception by the TIR1 ubiquitin ligase. *Nature* 446: 640–645.
- Taylor-Teeple M, Lin L, de Lucas M, Turco G, Toal TW, Gaudinier A, Young NF, Trabucco GM, Veling MT, Lamothe R *et al.* 2015. An Arabidopsis gene regulatory network for secondary cell wall synthesis. *Nature* 517: 571–575.
- Thomashow MF. 2010. Molecular basis of plant cold acclimation: insights gained from studying the CBF cold response pathway. *Plant Physiology* 154: 571–577.
- Thornberry JM, Goodman MM, Doebley J, Kresovich S, Nielsen D, Buckler ES. 2001. Dwarf8 polymorphisms associate with variation in flowering time. *Nature Genetics* 28: 286–289.
- Tian F, Bradbury PJ, Brown PJ, Hung H, Sun Q, Flint-Garcia S, Rocheford TR, McMullen MD, Holland JB, Buckler ES. 2011. Genome-wide association study of leaf architecture in the maize nested association mapping population. *Nature Genetics* 43: 159–162.
- Till BJ, Reynolds SH, Greene EA, Codoño CA, Enns LC, Johnson JE, Burtner C, Odden AR, Young K, Taylor NE *et al.* 2003. Large-scale discovery of induced point mutations with high-throughput TILLING. *Genome Research* 13: 524–530.
- Till BJ, Zerr T, Comai L, Henikoff S. 2006. A protocol for TILLING and ecotilling in plants and animals. *Nature Protocols* 1: 2465–2477.

- Todesco M, Balasubramanian S, Hu TT, Traw MB, Horton M, Eppe P, Kuhns C, Sureshkumar S, Schwartz C, Lanz C *et al.* 2010. Natural allelic variation underlying a major fitness trade-off in *Arabidopsis thaliana*. *Nature* **465**: 632–636.
- Tong AHY, Lesage G, Bader GD, Ding H, Xu H, Xin X, Young J, Berriz GF, Brost RL, Chang M *et al.* 2004. Global mapping of the yeast genetic interaction network. *Science* **303**: 808–813.
- Toufighi K, Brady SM, Austin R, Ly E, Provart NJ. 2005. The Botany Array Resource: e-Northern, Expression Angling, and promoter analyses. *Plant Journal* **43**: 153–163.
- Tsuji J, Somerville SC. 1988. *Xanthomonas campestris* P_v. campestris-induced chlorosis in *Arabidopsis thaliana*. *Arabidopsis Information Service* **26**: 1–8.
- Tyszkiewicz AB, Muir TW. 2008. Activation of protein splicing with light in yeast. *Nature Methods* **5**: 303–305.
- Usadel B, Obayashi T, Mutwil M, Giorgi FM, Bassel GW, Tanimoto M, Chow A, Steinhauser D, Persson S, Provart NJ. 2009. Co-expression tools for plant biology: opportunities for hypothesis generation and caveats. *Plant, Cell & Environment* **32**: 1633–1651.
- Vanholme R, Demedts B, Morreel K, Ralph J, Boerjan W. 2010. Lignin biosynthesis and structure. *Plant Physiology* **153**: 895–905.
- Velasquez SM, Ricardi MM, Dorosz JG, Fernandez PV, Nadra AD, Pol-Fachin L, Egelund J, Gille S, Harholt J, Ciana M *et al.* 2011. O-glycosylated cell wall proteins are essential in root hair growth. *Science* **332**: 1401–1403.
- Vilhjálmsdóttir BJ, Nordborg M. 2012. The nature of confounding in genome-wide association studies. *Nature Reviews Genetics* **14**: 1–2.
- Vogel J, Somerville S. 2000. Isolation and characterization of powdery mildew-resistant *Arabidopsis* mutants. *Proceeding of the National Academy of Sciences, USA* **97**: 1897–1902.
- Vorholt JA. 2012. Microbial life in the phyllosphere. *Nature Reviews Microbiology* **10**: 828–840.
- Wachsman G, Sparks EE, Benfey PN. 2015. Genes and networks regulating root anatomy and architecture. *New Phytologist* **208**: 26–38.
- Wallis JG, Browse J. 2010. Lipid biochemists salute the genome. *Plant Journal* **61**: 1092–1106.
- Weckwerth W. 2011. Green systems biology – from single genomes, proteomes and metabolomes to ecosystems research and biotechnology. *Journal of Proteomics* **75**: 284–305.
- Weigel D, Meyerowitz EM. 1994. The ABCs of floral homeotic genes. *Cell* **78**: 203–209.
- Weinig C, Ungerer MC, Dorn LA, Kane NC, Toyonaga Y, Halldorsdóttir SS, Mackay TFC, Purugganan MD, Schmitt J. 2002. Novel loci control variation in reproductive timing in *Arabidopsis thaliana* in natural environments. *Genetics* **162**: 1875–1884.
- Weirauch MT, Yang A, Albu M, Cote AG, Montenegro-Montero A, Drewe P, Najafabadi HS, Lambert SA, Sihninder M, Cook K *et al.* 2014. Determination and inference of eukaryotic transcription factor sequence specificity. *Cell* **158**: 1431–1443.
- Whittington AT, Vugrek O, Wei KJ, Hasenbein NG, Sugimoto K, Rashbrooke MC, Wasteney GO. 2001. MOR1 is essential for organizing cortical microtubules in plants. *Nature* **411**: 610–613.
- Wigge PA, Kim MC, Jaeger KE, Busch W, Schmid M, Lohmann JU, Weigel D. 2005. Integration of spatial and temporal information during floral induction in *Arabidopsis*. *Science* **309**: 1056–1059.
- Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ. 2007. An 'Electronic Fluorescent Pictograph' browser for exploring and analyzing large-scale biological data sets. *PLoS ONE* **2**: e718.
- Wolf S, Hématy K, Höfte H. 2012. Growth control and cell wall signaling in plants. *Annual Review of Plant Biology* **63**: 381–407.
- Wolff CF. 1759. *Theoria Generationis*. MD thesis, University of Halle, Halle, Germany.
- Wright DA, Townsend JA, Winfrey RJ, Irwin PA, Rajagopal J, Lonosky PM, Hall BD, Jondle MD, Voytas DF. 2005. High-frequency homologous recombination in plants mediated by zinc-finger nucleases. *Plant Journal* **44**: 693–705.
- Wu M-F, Sang Y, Bezhan S, Yamaguchi N, Han S-K, Li Z, Yanhui S, Slewinski TL, Wagner D. 2012. SWI2/SNF2 chromatin remodeling atpases overcome polycomb repression and control floral organ identity with the LEAFY and SEPALLATA3 transcription factors. *Proceedings of the National Academy of Sciences, USA* **109**: 3576–3581.
- Xie Z, Lee E, Lucas JR, Morohashi K, Li D, Murray JAH, Sack FD, Grotewold E. 2010. Regulation of cell proliferation in the stomatal lineage by the *Arabidopsis* MYB FOUR LIPS via direct targeting of core cell cycle genes. *Plant Cell* **22**: 2306–2321.
- Yadav RK, Girke T, Pasala S, Xie M, Reddy GV. 2009. Gene expression map of the *Arabidopsis* shoot apical meristem stem cell niche. *Proceedings of the National Academy of Sciences, USA* **106**: 4941–4946.
- Yanofsky MF, Ma H, Bowman JL, Drews GN, Feldmann KA, Meyerowitz EM. 1990. The protein encoded by the *Arabidopsis* homeotic gene *Agamous* resembles transcription factors. *Nature* **346**: 35–39.
- Yelagandula R, Stroud H, Holec S, Zhou K, Feng S, Zhong X, Muthurajan UM, Nie X, Kawashima T, Groth M *et al.* 2014. The histone variant H2A.W defines heterochromatin and promotes chromatin condensation in *Arabidopsis*. *Cell* **158**: 98–109.
- Yu J, Pressoir G, Briggs WH, Vroh Bi I, Yamasaki M, Doebley JF, McMullen MD, Gaut BS, Nielsen DM, Holland JB *et al.* 2006. A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. *Nature Genetics* **38**: 203–208.
- Zabotina OA, Avci U, Cavalier D, Pattathil S, Chou Y-H, Eberhard S, Danhof L, Keegstra K, Hahn MG. 2012. Mutations in multiple XXT genes of *Arabidopsis* reveal the complexity of xyloglucan biosynthesis. *Plant Physiology* **159**: 1367–1384.
- Zhang JZ, Creelman RA, Zhu J-K. 2004. From laboratory to field. using information from *Arabidopsis* to engineer salt, cold, and drought tolerance in crops. *Plant Physiology* **135**: 615–621.
- Zhao K, Aranzana MJ, Kim S, Lister C, Shindo C, Tang C, Toomajian C, Zheng H, Dean C, Marjoram P *et al.* 2007. An *Arabidopsis* example of association mapping in structured samples. *PLoS Genetics* **3**: e4.
- Zhao Q, Nakashima J, Chen F, Yin Y, Chunxiang F, Yun J, Shao H, Wang X, Wang Z-Y, Dixon RA. 2013. Laccase is necessary and nonredundant with peroxidase for lignin polymerization during vascular development in *Arabidopsis*. *Plant Cell* **25**: 3976–3987.
- Zimmermann P, Hirsch-Hoffmann M, Hennig L, Gruissem W. 2004. GENEVESTIGATOR. *Arabidopsis* microarray database and analysis toolbox. *Plant Physiology* **136**: 2621–2632.