

# Impact of vaccination on nasopharyngeal carriage pneumococcal genotypic diversity among children in The Gambia: a longitudinal study

Khan D<sup>1</sup> , Worwui A<sup>1</sup> , Senghore M<sup>1</sup> , Ebruke C<sup>1</sup> , Okoi C<sup>1</sup> , Foster Nyarko E<sup>1</sup> , L o SW<sup>2</sup> , Gladstone RA<sup>2</sup> , Klugman KP<sup>3</sup> , McGee L<sup>4</sup> , Breiman RF<sup>3</sup> , Bentley SD<sup>2</sup> , Adegbola R<sup>5</sup> , Antonio M<sup>1</sup> , Kwambana-Adams BA<sup>1</sup>

1. Medical Research Council Unit The Gambia at the London School of Hygiene and Tropical Medicine, Atlantic Road, Fajara, The Gambia; 2. The Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge CB10 1SA, UK; 3. Global Health Institute, Emory University, Atlanta GA, USA; 4. Respiratory Diseases Branch, Center for Disease Control and Prevention, Atlanta, GA, USA; 5. GlaxoSmithKline Vaccines, Wavre, Belgium

## Background

- The nasopharynx serves as a reservoir for pneumococcal acquisition, carriage and transmission
- Pneumococcal conjugate vaccines (PCVs) decrease the carriage of vaccine type (VT) pneumococcal serotypes and increase the carriage of non vaccine type (NVT) serotypes
- However, the impact that vaccination has on the genotypic diversity and stability of *Streptococcus pneumoniae* (*S. pneumoniae*) in the nasopharynx remains unclear

## Aim

To investigate the impact of vaccination on the genotypic diversity and stability of *S. pneumoniae* in the nasopharynx

## Method

- The villages were split into three groups of nine villages based on vaccination status, and nasopharyngeal swabs were collected over time (figure 1)
- Whole genome sequencing was performed on *S. pneumoniae* and in silico sequence type (ST) and in silico serotype derived




Figure 1: Flowchart of infant recruitment and sample collection

## Results

- We obtained serotype and ST data for 323, 284 and 391 *S. pneumoniae* from group one, two and three respectively (Table 1)
- There was an increase in certain STs such as ST847 post vaccination (figures 1&2)
- S. pneumoniae* from consecutive sampling points that differed in serotype also differed in genotype except in 2 cases – 15B/15C→17F: ST910 and 9L→19A: ST1735 (figure 3); indicative of serotype switching

Table 1: Serotype distribution and diversity of pneumococcal isolates in the three groups

Group 1			Group 2			Group 3		
serotype	No of specimens	No of STs	serotype	No. of specimens	No. of STs	serotype	No. of specimens	No. of STs
n (%)			n (%)			n (%)		
4	6 (1.9)	1	4	1 (0.4)	1	4	0 (0.0)	0
6B	33 (10.2)	5	6B	16 (5.6)	3	6B	12 (3.1)	5
9V	9 (2.8)	1	9V	3 (1.1)	2	9V	1 (0.3)	1
14	22 (6.8)	2	14	12 (4.2)	1	14	1 (0.3)	1
18C	4 (1.2)	1	18C	0 (0.0)	0	18C	2 (0.5)	1
19F	22 (6.8)	6	19F	3 (1.1)	3	19F	8 (2.0)	4
23F	15 (4.6)	4	23F	6 (2.1)	1	23F	1 (0.3)	1
NVT	212 (65.6)	59	NVT	243 (85.6)	58	NVT	364 (93.1)	114
Total	323 (100)	79	Total	284 (100)	69	Undetermined	2 (0.5)	2
Total							391 (100)	129




Figure 1: Box plot showing ST count plotted against vaccination period.  
\*pre vac: weeks 0-9, vac: weeks 10-17, post vac: weeks 18-54

## Discussion

- Vaccine pressure may have lead to the rise in ST847 of serotype 19A
- Apart from two cases, a change in serotype was always associated with a change in ST indicative of the acquisition of a new strain.




Figure 2: ST distribution of Serotype 19A in groups 2 (A) and 3 (B)




Figure 3: Serotype and ST data for participants C12202 and P12403.

## Acknowledgements

We would like to thank the Medical Research Council Unit The Gambia for funding the research. This study would not have been possible without the invaluable contributions of the study participants, their families and the communities of the Foni Districts. We thank them for their contribution.