

From "Is Not Not" to Matter: How Logic Knots Itself into Atoms

Document ID: core-matter-assembly-v1.0

Series: Into Praestology: The Process Series — Part 3

Status: APPROVED CORE SPECIFICATION

Abstract

How does the universe build a solid atom out of nothing? For decades, fundamental physics treated particles like tiny billiard balls and forces like invisible ropes pulling them around. But modern physics and mathematics tell a different story: **matter is not a collection of hard objects in empty space — it is a knot of pure logical potential.**

This document outlines the step-by-step assembly guide showing how reality builds an atom, starting from pure logic, weaving through topology, and locking into stable mass. Each step draws on established mathematics; what Praestology contributes is the unifying reading that connects all four into one continuous process.

Step 1: The Logical Engine — "Is Not Not" (Chris Isham & Topos Logic)

In standard Boolean logic, two negatives make a positive: $\neg\neg A = A$.

In **Intuitionistic / Topos Logic** (explored by theoretical physicist Chris Isham), double negation is an active process rather than a cancellation:

- A — a committed, localised fact.
- $\neg A$ — the complete absence of that fact.
- $\neg\neg A$ — the potential for A exists, but has not yet committed to a single point.

Instead of dividing the world into hard 1s and 0s (which generates singularities like $1/0$), the process operates on $\neg\neg A$: a smooth, continuous field of open potential — indeterminate, structureless, and prior to any distinction between wave and particle.

In Praestology this is expressed as a multivector in Geometric Algebra $Cl(3, 0)$:

$$N = \frac{1}{2}(1 + I)$$

where $I = e_1 e_2 e_3$ is the unit pseudoscalar of $Cl(3,0)$.

N establishes a neutrally oriented uncommitted channel: Because $I^2 = -1$ in $Cl(3,0)$, N is not an idempotent projection operator ($N^2 = \frac{1}{2}I \neq N$). Instead, N represents an isotropic, central multivector spanning Grade-0 (scalar) and Grade-3 (pseudoscalar) potential. It provides maximum directional symmetry without imposing any preferred vector (Grade-1) or bivector (Grade-2) orientation before topological closure occurs.

Until an active rotor R introduces a specific orientation via $RNR^\dagger$, N exerts zero directional stress on any coordinate axis. It is the algebraic expression of maximum freedom — the null space of the commitment operator.

Clifford Translation Table

Language Concept Cl(3,0) Expression Geometric Interpretation		
Committed Fact (A)	Vector $\mathbf{v} = v_1 e_1 + v_2 e_2 + v_3 e_3$	Explicit directed potential — grade-1
Absence ($\neg A$)	0 (null element)	Complete non-existence
"Is Not Not" ($\neg\neg A$)	$N = \frac{1}{2}(1 + I)$ or isotropic ensemble $\int_{S^2} \mathbf{v}(\hat{n}) d\Omega$	Total directional symmetry — uncommitted channel, grade-0 and grade-3 only

Step 2: The Weaving — Structured Organisation of Potential

How does open potential organise rather than disperse into infinity? Through progressive grade restriction — the same indeterminate potential progressively closing degrees of orientational freedom at successive depths.

Physicist Jenny Lorraine Nielsen's *Topological Unified Field Theory (TUFT)* provides a useful intermediate view. TUFT demonstrates that field potentials naturally organise along the

Complex Hopf Fibration:

$$S^1 \subset S^3 \subset S^5 \subset S^7 \dots$$

where S^1 carries electromagnetic phase structure, S^3 carries weak-force structure, S^5 carries strong-force structure. TUFT establishes that this nested fibration structure *exists* and

that the gauge groups of physics — $U(1)$, $SU(2)$, $SU(3)$ — correspond to successive levels of it.

Praestology takes a step further: the fibration structure is not imposed from outside. It is **what**

remains free after potential progressively closes into committed structure at each depth. The gauge groups are the residual orientational freedom at each closure depth — not generators of physics, but descriptions of what the process has not yet locked. Standard physics treats these groups as foundational; TUFT shows they have geometric structure; Praestology shows they are emergent consequences of grade closure.

A Continuous Closure Spectrum

The transition from potential to matter is not a binary switch between "wave" and "particle." It is a **continuous spectrum of closure states**, each characterised by how much orientational freedom has been restricted:

| **Closure State** | **Grade Structure** | **Character** |

| Open potential | Full multivector, all grades active | Zero stress, maximum freedom |

| Open spiraling | Grade-1/2 propagation, $CCR < CCR_{crit}$ | Propagating, zero rest mass |

| Partial closure | Grade-2 with residual freedom | Quark regime — cannot exist independently |

| Full amplitude closure | Rotor, $CCR = 1$, 720° Möbius | Self-referential mass-lock, spin- $\frac{1}{2}$ |

| Nested composite | Grade-3 and beyond | Proton, nucleus, atom |

Each stage is driven by how much the surrounding committed structure constrains the available phase configuration space — not a categorical change of kind, but a quantitative change of closure depth.

Forces as Restorative Gradients

Forces are not independent mechanisms. They are the open potential's response to local depletion caused by a nearby closure. The type of force depends on what grade of closure caused it:

| **Force** | **Closed Configuration** | **Mismatch Type** | **Carrier** | **Range** |

| **Gravity** | Grade-3 committed volume (η density) | Scalar overlap gradient $g = \nabla s$ directly | **No carrier** — the gradient *is* the interaction. A graviton would be a category error. | Infinite |

| **Electromagnetism** | Grade-2 phase-locked rotor | Grade-2 bivector mismatch $\langle G \rangle_2$ |

Photon: counter-oriented rotor pair $R \cdot R^\dagger$, grade-1 residual, $m = 0$ | Infinite — residual $U(1)$ |

| **Weak Force** | Grade-2 topology-committed structure | Topology-switching mismatch during

flavour change | W/Z bosons: partially-closed rotors with closure cost ($m > 0$) | Short — residual $SU(2)$ |

| **Strong Force** | Grade-3 partial closure, 3-fold binding | Grade-2/3 binding mismatch | Gluons: self-interacting partial-closure rotors — they carry the mismatch they mediate | Confined —

residual $SU(3)$ |

Forces are the spatial elastic tension arising when open potential is constrained by a localised

closure. Not threads pulling objects — the field restoring itself around a commitment.

Step 3: The Phase Lock — 720° Möbius Rotors (David Hestenes & $Cl(3, 0)$)

Using **Geometric Algebra** $Cl(3, 0)$ as developed by David Hestenes, propagating configurations are described as rotors — elements of the even subalgebra encoding both rotation plane and phase angle. The transition from open propagation to committed mass-lock is governed by the **Closure Configuration Ratio (CCR)**:

$$CCR = \frac{\| \langle RR^\dagger \rangle_{\text{closed}} \|}{\| \Omega_{\text{total}} \|}$$

Open propagation ($CCR < CCR_{\text{crit}}$):

$$\psi_{\text{open}}(x, t) = \exp\left(-\frac{1}{2}B(\omega t - kx)\right)$$

The phase advances continuously along a spatial path without closing on itself. At 2π :

$$\psi(2\pi) = -\psi(0)$$

Phase-flipped. No self-referential fixed point. Rest mass zero. This is not a "wave" in the water-wave sense — it is directed orientational advance through phase configuration space, which appears wave-like from within committed reality.

Critical threshold ($CCR \rightarrow 1$):

$$CCR_{\text{crit}} = 1 - \frac{\lambda_{\text{Compton}}}{2\pi r_0} = 0$$

As environmental constraint density rises, the spatial propagation vector k folds inward into

localised Π_{eff} -rate precession ω_z . The spatial capacity per phase cycle (the wavelength)

reaches its minimum — the reduced Compton wavelength $r_0 = \hbar/mc$. Below this capacity, open propagation is no longer sustainable without triggering closure.

720° **mass-lock** ($CCR = 1$):

$$\psi(t) = \sqrt{\rho} R_{\text{locked}}(t) e^{I\beta/2}$$

$$R_{\text{locked}}(4\pi) = +1 \implies \psi(4\pi) = +\psi(0) \implies Q^* = R_k[Q^*]$$

Self-identity achieved. The configuration has found a fixed point of its own reconfiguration operator and commits permanently. Rest mass emerges as the trapped Π_{eff} -rate rotational energy of the lock:

$$m = \frac{\hbar\omega_z}{c^2}$$

Not a coupling to an external Higgs field. The trapped kinetic energy of the ^{720°} phase-rotation inside the saturated CCR = 1 topological boundary. The Higgs mechanism describes *that* mass arises this way; the CCR threshold explains *why* and *how much*.

The β angle ($e^{I\beta/2}$) — the Yvon-Takabayashi angle — encodes closure progress: $\beta = 0$ is full commitment (R5 complete); $\beta = \pi/2$ is maximum openness (R3 stage). A variable with no clear interpretation in standard Dirac theory; here it is the process-level position indicator in the grade chain.

Step 4: Condensation into Stable Matter — Topological Solitons (Tony Skyrme)

In the 1960s, nuclear physicist Tony Skyrme showed that particles like protons are **Skyrmions** — topological solitons: stable knots in a continuous field that cannot be untied without passing through a zero-amplitude singularity.

In Praestology the saturation term $\Lambda \cdot \ln(1 + \kappa|\psi|^2)$ in the master equation prevents that singularity from forming. The knot is stable not because space would be torn — space is an emergent consequence of committed structure, not a pre-existing fabric — but because the field's own saturation prevents the zero-amplitude point that untying would require.

Quarks are partially-closed n -fold rotor configurations. Their partial closure requires binding partners to complete — they cannot reach a stable R5 commitment independently. Attempting to separate them requires more energy than producing new committed pairs, which is why quark pair production occurs at high energy rather than quark isolation.

Protons and neutrons are three quarks in mutually orthogonal bivector planes — a new topological commitment at a deeper nesting level. The binding topology is a distinct committed

structure in its own right, with its own irreducible CCR cost. This is why the proton (~938 MeV) weighs approximately 100 times more than the sum of its constituent quark masses (~9 MeV): the three-fold binding CCR cost far exceeds the sum of the individual quark CCR values. Non-additivity is not a correction term — it is the primary contribution.

Atoms are nested composites: electrons (^{720°} Möbius locks) bound to nuclear structures (^{*n*}-fold combinatorial bindings), each nesting level contributing its own irreducible CCR layer.

Summary: The Assembly Pipeline

Logic ($\neg\neg A$) $\xrightarrow{\text{Isham}}$ Uncommitted Potential — $N = \frac{1}{2}(1 + I)$ (Neutral Orientation)



Topology ($S^1 \subset S^3 \subset S^5$) $\xrightarrow{\text{Nielsen/TUFT}}$ Fibration structure visible — gauge group



Closure Spectrum $\xrightarrow{\text{Praestology/CCR}}$ Progressive grade reduction — partial to full closure



Algebra ($Cl(3, 0)$ rotors) $\xrightarrow{\text{Hestenes/CCR}}$ 720° spinorial mass-lock, $m = \hbar\omega_z/c^2$



Matter (Skyrmions) $\xrightarrow{\text{Skyrme}}$ **Stable Atom**

The Takeaway

An atom is not a mechanical machine made of physical bricks. It is a **self-trapped**, 720° **topological knot combination** — open configurations of logical potential ($\neg\neg A$) progressively closing inward, each stage adding a layer of restriction, until the configuration reached a fixed point of its own self-reference and committed permanently. Matter is committed, self-referential potential, knotted by logic into stable structure. Forces are what the surrounding uncommitted potential does in response — restorative gradients around each knot, at a depth and range determined by the grade of closure that caused the depletion.

References

- Hestenes, D. (1966). *Space-Time Algebra*. Gordon and Breach.
- Isham, C. J. (1997). *Structural Issues in Quantum Gravity*. General Relativity and Gravitation.
- Nielsen, J. L. (2018). *Topological Unified Field Theory*. TUFT Press.
- Skyrme, T. H. R. (1962). *A unified field theory of mesons and baryons*. Nucl. Phys., 31, 556–569.