

Chapter 12 worked solutions – Further calculus

Solutions to Exercise 12A

1a

x	-1	-0.7	-0.5	-0.2	0	0.3	0.6	0.8	1
$\frac{dy}{dx}$	undefined	1.40	1.16	1.02	1	1.05	1.25	1.67	undefined

1b

x	-1	-0.7	-0.5	-0.2	0	0.3	0.6	0.8	1
$\frac{dy}{dx}$	undefined	1.40	1.16	1.02	1	1.05	1.25	1.67	undefined

2a

$$\frac{d}{dx}(\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$$

2b

$$\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$$

2c

$$\begin{aligned} \frac{d}{dx}(\sin^{-1} 2x) \\ = \frac{1}{\sqrt{1-(2x)^2}} \times \frac{d}{dx}(2x) \\ = \frac{2}{\sqrt{1-4x^2}} \end{aligned}$$

2d

$$\begin{aligned} \frac{d}{dx}(\tan^{-1} 3x) \\ = \frac{1}{1+(3x)^2} \times \frac{d}{dx}(3x) \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{3}{1 + 9x^2}$$

2e

$$\begin{aligned} & \frac{d}{dx}(\cos^{-1} 5x) \\ &= \frac{-1}{\sqrt{1 - (5x)^2}} \times \frac{d}{dx}(5x) \\ &= -\frac{5}{\sqrt{1 - 25x^2}} \end{aligned}$$

2f

$$\begin{aligned} & \frac{d}{dx}(\sin^{-1} -x) \\ &= \frac{1}{\sqrt{1 - (-x)^2}} \times \frac{d}{dx}(-x) \\ &= \frac{-1}{\sqrt{1 - x^2}} \end{aligned}$$

2g

$$\begin{aligned} & \frac{d}{dx}(\sin^{-1} x^2) \\ &= \frac{1}{\sqrt{1 - (x^2)^2}} \times \frac{d}{dx}(x^2) \\ &= \frac{2x}{\sqrt{1 - x^4}} \end{aligned}$$

2h

$$\begin{aligned} & \frac{d}{dx}(\tan^{-1} x^3) \\ &= \frac{1}{1 + (x^3)^2} \times \frac{d}{dx}(x^3) \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{3x^2}{1 + x^6}$$

2i

$$\begin{aligned} & \frac{d}{dx}(\tan^{-1}(x+2)) \\ &= \frac{1}{1+(x+2)^2} \times \frac{d}{dx}(x+2) \\ &= \frac{1}{x^2+4x+5} \end{aligned}$$

2j

$$\begin{aligned} & \frac{d}{dx}(\cos^{-1}(1-x)) \\ &= \frac{-1}{\sqrt{1-(1-x)^2}} \times \frac{d}{dx}(1-x) \\ &= \frac{-1}{\sqrt{1-(1-2x+x^2)}} \times -1 \\ &= \frac{1}{\sqrt{2x-x^2}} \end{aligned}$$

2k

$$\begin{aligned} & \frac{d}{dx}(x \sin^{-1} x) \\ &= \sin^{-1} x \times \frac{d}{dx}(x) + x \times \frac{d}{dx}(\sin^{-1} x) \\ &= \sin^{-1} x + \frac{x}{\sqrt{1-x^2}} \end{aligned}$$

2l

$$\begin{aligned} & \frac{d}{dx}((1+x^2)\tan^{-1} x) \\ &= \tan^{-1} x \times \frac{d}{dx}(1+x^2) + (1+x^2) \times \frac{d}{dx}(\tan^{-1} x) \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \tan^{-1} x \times 2x + (1 + x^2) \times \frac{1}{1 + x^2} \\
 &= 2x \tan^{-1} x + 1
 \end{aligned}$$

2m

$$\begin{aligned}
 &\frac{d}{dx} \left(\sin^{-1} \frac{1}{5} x \right) \\
 &= \frac{1}{\sqrt{1 - \left(\frac{1}{5} x\right)^2}} \times \frac{d}{dx} \left(\frac{1}{5} x \right) \\
 &= \frac{1}{\sqrt{1 - \frac{x^2}{25}}} \times \frac{1}{5} \\
 &= \frac{1}{\sqrt{25 - x^2}}
 \end{aligned}$$

2n

$$\begin{aligned}
 &\frac{d}{dx} \left(\tan^{-1} \frac{1}{4} x \right) \\
 &= \frac{1}{1 + \left(\frac{1}{4} x\right)^2} \times \frac{d}{dx} \left(\frac{1}{4} x \right) \\
 &= \frac{1}{1 + \frac{x^2}{16}} \times \frac{1}{4} \\
 &= \frac{4}{16 + x^2}
 \end{aligned}$$

2o

$$\begin{aligned}
 &\frac{d}{dx} (\cos^{-1} \sqrt{x}) \\
 &= \frac{-1}{\sqrt{1 - \sqrt{x}^2}} \times \frac{d}{dx} (\sqrt{x}) \\
 &= \frac{-1}{\sqrt{1 - \sqrt{x}^2}} \times \frac{d}{dx} \left(x^{\frac{1}{2}} \right)
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \frac{-1}{\sqrt{1 - \sqrt{x^2}}} \times \frac{1}{2}x^{-\frac{1}{2}} \\
 &= \frac{-1}{2\sqrt{x}\sqrt{1-x}} \\
 &= -\frac{1}{2\sqrt{x-x^2}}
 \end{aligned}$$

2p

$$\begin{aligned}
 &\frac{d}{dx}(\tan^{-1}\sqrt{x}) \\
 &= \frac{1}{1+(\sqrt{x})^2} \times \frac{d}{dx}(\sqrt{x}) \\
 &= \frac{1}{1+(\sqrt{x})^2} \times \frac{d}{dx}(x^{\frac{1}{2}}) \\
 &= \frac{1}{1+(\sqrt{x})^2} \times -\frac{1}{2}x^{-\frac{1}{2}} \\
 &= \frac{1}{2\sqrt{x}(1+x)}
 \end{aligned}$$

2q

$$\begin{aligned}
 &\frac{d}{dx}\left(\tan^{-1}\frac{1}{x}\right) \\
 &= \frac{1}{1+\left(\frac{1}{x}\right)^2} \times \frac{d}{dx}\left(\frac{1}{x}\right) \\
 &= \frac{1}{1+\frac{1}{x^2}} \times -\frac{1}{x^2} \\
 &= -\frac{1}{x^2+1}
 \end{aligned}$$

- 3 For this question you only need to provide the gradient. We also provide the equation of the tangent for your benefit.

Chapter 12 worked solutions – Further calculus

3a

$$\frac{dy}{dx} = \frac{2}{1+x^2}$$

At $x = 0$, $y = 2 \tan^{-1} 0 = 0$ and $\frac{dy}{dx} = \frac{2}{1+0^2} = 2$. Hence, the tangent will be given by

$$y - y_1 = m(x - x_1)$$

$$y - 0 = 2(x - 0)$$

$$y = 2x$$

3b

$$\frac{dy}{dx} = \frac{\sqrt{3}}{\sqrt{1-x^2}}$$

At $x = \frac{1}{2}$, $y = \sqrt{3} \sin^{-1} \frac{1}{2} = \frac{\sqrt{3}}{6}$ and $\frac{dy}{dx} = \frac{\sqrt{3}}{\sqrt{1-\frac{1}{2}^2}} = \frac{\sqrt{3}}{\sqrt{\frac{3}{4}}} = 2$. Hence, the tangent will be

given by

$$y - y_1 = m(x - x_1)$$

$$y - \frac{\sqrt{3}}{6} = 2(x - 0)$$

$$y = 2x + \frac{\sqrt{3}}{6}$$

3c

$$\frac{dy}{dx} = \frac{2}{1+4x^2}$$

At $x = -\frac{1}{2}$, $y = \tan^{-1} \left(2 \left(-\frac{1}{2} \right) \right) = -\frac{\pi}{4}$ and $\frac{dy}{dx} = \frac{2}{1+4\left(-\frac{1}{2}\right)^2} = \frac{2}{2} = 1$. Hence, the tangent will be given by

$$y - y_1 = m(x - x_1)$$

$$y - \left(-\frac{\pi}{4} \right) = 1(x - 0)$$

$$y = x - \frac{\pi}{4}$$

Chapter 12 worked solutions – Further calculus

3d

$$\frac{dy}{dx} = \frac{-\frac{1}{2}}{\sqrt{1 - \left(\frac{x}{2}\right)^2}} = -\frac{1}{\sqrt{4 - x^2}}$$

At $x = \sqrt{3}$, $y = \cos^{-1} \frac{\sqrt{3}}{2} = \frac{\pi}{6}$ and $\frac{dy}{dx} = -\frac{1}{\sqrt{4 - \sqrt{3}^2}} = -1$. Hence, the tangent will be

given by

$$y - y_1 = m(x - x_1)$$

$$y - \frac{\pi}{6} = -1(x - 0)$$

$$y = -x + \frac{\pi}{6}$$

4a

$$\frac{dy}{dx} = -\frac{1}{\sqrt{1 - (3x)^2}} \times 2 \times 3 = -\frac{6}{\sqrt{1 - 9x^2}}$$

When $x = 0$

$$y = 2 \cos^{-1} 3(0) = 2 \left(\frac{\pi}{2}\right) = \pi,$$

$$\frac{dy}{dx} = -\frac{6}{\sqrt{1-9(0)^2}} = -6 \text{ and } m_{norm} = -\frac{1}{\left(\frac{dy}{dx}\right)} = \frac{1}{6}$$

For the tangent:

$$y - y_1 = m(x - x_1)$$

$$y - \pi = -6(x - 0)$$

$$y = -6x + \pi$$

For the normal:

$$y - y_1 = m(x - x_1)$$

$$y - \pi = \frac{1}{6}(x - 0)$$

$$y = \frac{1}{6}x + \pi$$

Chapter 12 worked solutions – Further calculus

4b

$$\frac{dy}{dx} = \frac{1}{\sqrt{1 - \left(\frac{x}{2}\right)^2}} \times \frac{1}{2} = \frac{1}{\sqrt{4 - x^2}}$$

When $x = \sqrt{2}$

$$y = \sin^{-1} \frac{\sqrt{2}}{2} = \frac{\pi}{4},$$

$$\frac{dy}{dx} = \frac{1}{\sqrt{4 - (\sqrt{2})^2}} = \frac{1}{\sqrt{2}} \text{ and } m_{norm} = -\frac{1}{\left(\frac{dy}{dx}\right)} = -\sqrt{2}$$

For the tangent:

$$y - y_1 = m(x - x_1)$$

$$y - \frac{\pi}{4} = \frac{1}{\sqrt{2}}(x - \sqrt{2})$$

$$y = \frac{x}{\sqrt{2}} - 1 + \frac{\pi}{4}$$

For the normal:

$$y - y_1 = m(x - x_1)$$

$$y - \frac{\pi}{4} = -\sqrt{2}(x - \sqrt{2})$$

$$y = -x\sqrt{2} + 2 + \frac{\pi}{4}$$

5a

$$\begin{aligned} & \frac{d}{dx}(\sin^{-1} x + \cos^{-1} x) \\ &= \frac{d}{dx}(\sin^{-1} x) + \frac{d}{dx}(\cos^{-1} x) \\ &= \frac{1}{\sqrt{1-x^2}} - \frac{1}{\sqrt{1-x^2}} \\ &= 0 \end{aligned}$$

- 5b As the gradient of the function is zero, it must be flat along its entire domain and hence constant. Letting $x = 0$ gives $\sin^{-1} 0 + \cos^{-1} 0 = 0 + \frac{\pi}{2} = \frac{\pi}{2}$, hence the function has a constant value of $\frac{\pi}{2}$.

Chapter 12 worked solutions – Further calculus

6a

$$\begin{aligned}
 & \frac{d}{dx}(\cos^{-1} x + \cos^{-1}(-x)) \\
 &= -\frac{1}{\sqrt{1-x^2}} - \frac{d}{dx}(-x) \\
 &= -\frac{1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1-x^2}} \\
 &= 0
 \end{aligned}$$

As the gradient of the function is zero, it must be flat along its entire domain and hence constant. Letting $x = 0$ gives $\cos^{-1} 0 + \cos^{-1} 0 = \frac{\pi}{2} + \frac{\pi}{2} = \pi$, hence the function has a constant value of π .

6b

$$\begin{aligned}
 & \frac{d}{dx}(2 \sin^{-1} \sqrt{x} - \sin^{-1}(2x-1)) \\
 &= \frac{1}{\sqrt{1-\sqrt{x}^2}} \times 2 \times \frac{1}{2} x^{-\frac{1}{2}} + \frac{1}{\sqrt{1-(2x-1)^2}} \times (-2) \\
 &= \frac{1}{\sqrt{x}\sqrt{1-x}} - \frac{2}{\sqrt{1-(4x^2-4x+1)}} \\
 &= \frac{1}{\sqrt{x-x^2}} - \frac{2}{\sqrt{4x-4x^2}} \\
 &= \frac{1}{\sqrt{x-x^2}} - \frac{2}{2\sqrt{x-x^2}} \\
 &= \frac{1}{\sqrt{x-x^2}} - \frac{1}{\sqrt{x-x^2}} \\
 &= 0
 \end{aligned}$$

As the gradient of the function is zero, it must be flat along its entire domain and hence constant. Letting $x = 0$ gives $2 \sin^{-1} \sqrt{0} - \sin^{-1}(2(0)-1) = 2 \sin^{-1} 0 - \sin^{-1}(-1) = 0 - \frac{\pi}{2} = \frac{\pi}{2}$, hence the function has a constant value of $\frac{\pi}{2}$.

7a

$$\begin{aligned}
 f'(x) &= 1 \times \tan^{-1} x + x \times \frac{1}{1+x^2} - \frac{1}{2} \times \frac{2x}{1+x^2} \\
 &= \tan^{-1} x + \frac{x}{1+x^2} - \frac{x}{1+x^2} \\
 &= \tan^{-1} x
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

Hence

$$f''(x) = \frac{1}{1+x^2}$$

- 7b $f''(-1) = \frac{1}{2} > 0$, hence the curve is concave up

- 8 By the quotient rule

$$\begin{aligned}\frac{dy}{dx} &= \frac{x \times \frac{d}{dx}(\sin^{-1} x) - \sin^{-1} x \times \frac{d}{dx}(x)}{x^2} \\ &= \frac{x \times \frac{1}{\sqrt{1-x^2}} - \sin^{-1} x}{x^2}\end{aligned}$$

$$\text{When } x = \frac{1}{2}$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{\frac{1}{2} \times \frac{1}{\sqrt{1-\left(\frac{1}{2}\right)^2}} - \sin^{-1}\left(\frac{1}{2}\right)}{\left(\frac{1}{2}\right)^2}\end{aligned}$$

$$= \frac{\frac{1}{2} \times \frac{1}{\sqrt{3}} - \frac{\pi}{6}}{\frac{1}{4}}$$

$$= \frac{\frac{1}{2} \times \frac{2}{\sqrt{3}} - \frac{\pi}{6}}{\frac{1}{4}}$$

$$= \frac{\frac{1}{\sqrt{3}} - \frac{\pi}{6}}{\frac{1}{4}}$$

$$= 4 \left(\frac{\sqrt{3}}{3} - \frac{\pi}{6} \right)$$

$$= \frac{2}{3} (2\sqrt{3} - \pi)$$

Chapter 12 worked solutions – Further calculus

9a

$$\begin{aligned}
 & \frac{d}{dx} \left(x \cos^{-1} x - \sqrt{1-x^2} \right) \\
 &= x \times -\frac{1}{\sqrt{1-x^2}} + \cos^{-1} x - \frac{1}{2} \times (-2x) \times (1-x^2)^{-\frac{1}{2}} \\
 &= -\frac{x}{\sqrt{1-x^2}} + \cos^{-1} x + \frac{x}{\sqrt{1-x^2}} \\
 &= \cos^{-1} x
 \end{aligned}$$

9b

$$\begin{aligned}
 & \frac{d}{dx} (\sin^{-1} e^{3x}) \\
 &= \frac{1}{\sqrt{1-(e^{3x})^2}} \times \frac{d}{dx} (e^{3x}) \\
 &= \frac{3e^{3x}}{\sqrt{1-e^{6x}}}
 \end{aligned}$$

9c

$$\begin{aligned}
 & \frac{d}{dx} \left(\sin^{-1} \frac{1}{4}(2x-3) \right) \\
 &= \frac{1}{\sqrt{1-\left(\frac{1}{4}(2x-3)\right)^2}} \times \frac{d}{dx} \left(\frac{1}{4}(2x-3) \right) \\
 &= \frac{1}{\sqrt{1-\frac{1}{16}(4x^2-12x+9)}} \times \frac{2}{4} \\
 &= \frac{2}{4} \times \frac{1}{\frac{1}{4}\sqrt{16-(4x^2-12x+9)}} \\
 &= \frac{2}{\sqrt{7-4x^2+12x}}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

9d

$$\begin{aligned}
 & \frac{d}{dx} \left(\tan^{-1} \frac{1}{1-x} \right) \\
 &= \frac{1}{1 + \left(\frac{1}{1-x} \right)^2} \times \frac{d}{dx} \left(\frac{1}{1-x} \right) \\
 &= \frac{(1-x)^2}{(1-x)^2 + 1} \times -1 \times -1 \times (1-x)^{-2} \\
 &= \frac{1}{(1-x)^2 + 1} \\
 &= \frac{1}{1 - 2x + x^2 + 1} \\
 &= \frac{1}{x^2 - 2x + 2}
 \end{aligned}$$

9e

$$\begin{aligned}
 & \frac{d}{dx} (\sin^{-1} e^x) \\
 &= \frac{1}{\sqrt{1 - (e^x)^2}} \times \frac{d}{dx} (e^x) \\
 &= \frac{e^x}{\sqrt{1 - e^{2x}}}
 \end{aligned}$$

9f

$$\begin{aligned}
 & \frac{d}{dx} \left(\log_e \sqrt{\sin^{-1} x} \right) \\
 &= \frac{d}{dx} \left(\sqrt{\sin^{-1} x} \right) \\
 &= \frac{1}{2} \times (\sin^{-1} x)^{-\frac{1}{2}} \times \frac{d}{dx} (\sin^{-1} x) \\
 &= \frac{1}{2 \sin^{-1} x} \times \frac{d}{dx} (\sin^{-1} x) \\
 &= \frac{1}{2\sqrt{1-x^2} \sin^{-1} x}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

9g

$$\begin{aligned}
 & \frac{d}{dx}(\sin^{-1} \sqrt{\log_e x}) \\
 &= \frac{1}{\sqrt{1 - (\log_e x)^{\frac{1}{2}}}} \times \frac{d}{dx}((\log_e x)^{\frac{1}{2}}) \\
 &= \frac{1}{\sqrt{1 - \log_e x}} \times \frac{1}{2} \times (\log_e x)^{-\frac{1}{2}} \times \frac{d}{dx}(\log_e x) \\
 &= \frac{1}{\sqrt{\log_e x (1 - \log_e x)}} \times \frac{1}{2} \left(\frac{1}{x} \right) \\
 &= \frac{1}{2x\sqrt{\log_e x (1 - \log_e x)}}
 \end{aligned}$$

9h

$$\begin{aligned}
 & \frac{d}{dx}(\sqrt{x} \sin^{-1} \sqrt{1-x}) \\
 &= \sin^{-1} \sqrt{1-x} \times \frac{d}{dx}(\sqrt{x}) + \sqrt{x} \times \frac{d}{dx}(\sin^{-1} \sqrt{1-x}) \\
 &= \sin^{-1} \sqrt{1-x} \times \frac{1}{2\sqrt{x}} + \sqrt{x} \times \frac{1}{\sqrt{1 - (\sqrt{1-x})^2}} \times \frac{d}{dx}(\sqrt{1-x}) \\
 &= \frac{1}{2\sqrt{x}} \times \sin^{-1} \sqrt{1-x} + \sqrt{x} \times \frac{1}{\sqrt{1 - (1-x)}} \times \frac{1}{2} \times -1 \times \frac{1}{\sqrt{1-x}} \\
 &= \frac{1}{2\sqrt{x}} \times \sin^{-1} \sqrt{1-x} + \sqrt{x} \times \frac{1}{\sqrt{x}} \times \frac{1}{2} \times -1 \times \frac{1}{\sqrt{1-x}} \\
 &= \frac{1}{2\sqrt{x}} \times \sin^{-1} \sqrt{1-x} - \frac{1}{2\sqrt{1-x}}
 \end{aligned}$$

9i

$$\begin{aligned}
 & \frac{d}{dx}\left(\tan^{-1} \frac{x+2}{1-2x}\right) \\
 &= \frac{1}{1 + \left(\frac{x+2}{1-2x}\right)^2} \times \frac{d}{dx}\left(\frac{x+2}{1-2x}\right) \\
 &= \frac{1}{1 + \left(\frac{x+2}{1-2x}\right)^2} \times \frac{(1-2x) - (-2)(x+2)}{(1-2x)^2}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \frac{1}{1 + \left(\frac{x+2}{1-2x}\right)^2} \times \frac{1-2x+2x+4}{(1-2x)^2} \\
 &= \frac{5}{(1-2x)^2 + (x+2)^2} \\
 &= \frac{5}{1 - 4x + 4x^2 + x^2 + 4x + 4} \\
 &= \frac{5}{5x^2 + 5} \\
 &= \frac{1}{1 + x^2}
 \end{aligned}$$

10a i

$$\begin{aligned}
 y' &= 2 \times \frac{1}{\sqrt{1-x^2}} \times \sin^{-1} x = \frac{2 \sin^{-1} x}{\sqrt{1-x^2}} \\
 y'' &= \frac{2 \times \frac{1}{\sqrt{1-x^2}} \times \sqrt{1-x^2} - \frac{1}{2} \times (-2x) \times (1-x^2)^{-\frac{1}{2}} \times 2 \sin^{-1} x}{(\sqrt{1-x^2})^2} \\
 &= \frac{2 + \frac{2x \sin^{-1} x}{\sqrt{1-x^2}}}{1-x^2} \\
 &= \frac{2(1+x \sin^{-1} x)}{(1-x^2)^{\frac{3}{2}}}
 \end{aligned}$$

10a ii

$$\begin{aligned}
 &(1-x^2)y'' - xy' - 2 \\
 &= (1-x^2) \left(\frac{2 + \frac{2x \sin^{-1} x}{\sqrt{1-x^2}}}{1-x^2} \right) - x \left(\frac{2 \sin^{-1} x}{\sqrt{1-x^2}} \right) - 2 \\
 &= 2 + \frac{2x \sin^{-1} x}{\sqrt{1-x^2}} - \frac{2x \sin^{-1} x}{\sqrt{1-x^2}} - 2 \\
 &= 0
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

10b

$$y' = \frac{1}{\sqrt{1-x^2}} e^{\sin^{-1} x}$$

$$\begin{aligned} y'' &= -\frac{1}{2}(-2x)(1-x^2)^{-\frac{3}{2}} e^{\sin^{-1} x} + \frac{1}{\sqrt{1-x^2}} \frac{1}{\sqrt{1-x^2}} e^{\sin^{-1} x} \\ &= \frac{x e^{\sin^{-1} x}}{(1-x^2)^{\frac{3}{2}}} + \frac{e^{\sin^{-1} x}}{1-x^2} \end{aligned}$$

$$\begin{aligned} (1-x^2)y'' - xy' - y &= (1-x^2) \left(\frac{x e^{\sin^{-1} x}}{(1-x^2)^{\frac{3}{2}}} + \frac{e^{\sin^{-1} x}}{1-x^2} \right) - x \left(\frac{1}{\sqrt{1-x^2}} e^{\sin^{-1} x} \right) - e^{\sin^{-1} x} \\ &= \frac{x e^{\sin^{-1} x}}{\sqrt{1-x^2}} + e^{\sin^{-1} x} - \frac{x}{\sqrt{1-x^2}} e^{\sin^{-1} x} - e^{\sin^{-1} x} \\ &= 0 \end{aligned}$$

11a The range is the same as that of $\sin^{-1} x$ which is $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$

11b

$$y = \sin^{-1} 2x$$

$$2x = \sin y$$

$$x = \frac{1}{2} \sin y$$

11c

$\frac{dx}{dy} = \frac{1}{2} \cos y$, as the square root function always returns positive numbers, and as the division of one number by another positive is always positive, this derivative function will always be positive.

Chapter 12 worked solutions – Further calculus

11d

$$\begin{aligned}
 \frac{dy}{dx} &= \frac{1}{\frac{dx}{dy}} \\
 &= \frac{1}{\frac{1}{2} \cos y} \\
 &= \frac{2}{\cos y} \\
 &= \frac{2}{\sqrt{1 - \sin^2 y}} \\
 &= \frac{2}{\sqrt{1 - (2x)^2}} \\
 &= \frac{2}{\sqrt{1 - 4x^2}}
 \end{aligned}$$

Note that taking the derivative normally yields

$$\frac{dy}{dx} = \frac{2}{\sqrt{1 - (2x)^2}} = \frac{2}{\sqrt{1 - 4x^2}}$$

12a

$$\begin{aligned}
 y &= \sin^{-1} \frac{1}{2} x \\
 \frac{x}{2} &= \sin y \\
 x &= 2 \sin y \\
 \frac{dy}{dx} &= \frac{1}{\frac{dx}{dy}} \\
 &= \frac{1}{2 \cos y} \\
 &= \frac{1}{2 \sqrt{1 - \left(\frac{x}{2}\right)^2}} \\
 &= \frac{1}{\sqrt{4 - x^2}}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

12b

$$y = \cos^{-1}(x - 1)$$

$$x - 1 = \cos y$$

$$x = \cos y + 1$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{1}{\frac{dx}{dy}} \\ &= \frac{1}{-\sin y} \\ &= \frac{1}{-\sqrt{1 - \cos^2 y}} \\ &= \frac{1}{-\sqrt{1 - (x - 1)^2}} \\ &= \frac{-1}{\sqrt{1 - (x^2 - 2x + 1)}} \\ &= \frac{-1}{\sqrt{2x - x^2}}\end{aligned}$$

12c

$$y = \tan^{-1} \sqrt{x}$$

$$\sqrt{x} = \tan y$$

$$x = \tan^2 y$$

$$\frac{dx}{dy} = 2 \times \frac{1}{1 + y^2} \times \tan y$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{1}{\frac{dx}{dy}} \\ &= \frac{1}{2 \times \frac{1}{1 + y^2} \times \tan y} \\ &= \frac{1 + y^2}{2 \tan y} \\ &= \frac{1 + (\tan^{-1} \sqrt{x})^2}{2 \tan(\tan^{-1} \sqrt{x})} \\ &= \frac{1 + (\tan^{-1} \sqrt{x})^2}{2\sqrt{x}}\end{aligned}$$

Chapter 12 worked solutions – Further calculus

- 13a From the domain of the inverse cosine function we know that $-1 \leq x^2 \leq 1$. Furthermore, $x^2 \geq 0$ for all real x . Hence $0 \leq x^2 \leq 1$.

Hence the domain is $-1 \leq x \leq 1$

- 13b $f(-x) = \cos^{-1}(-x)^2 = \cos^{-1}x^2 = f(x)$, hence the function is even and symmetric about the line $x = 0$

13c

$$\begin{aligned}f'(x) &= \frac{-1}{\sqrt{1-(x^2)^2}} \times \frac{d}{dx}(x^2) \\&= \frac{-1}{\sqrt{1-(x^2)^2}} \times 2x \\&= -\frac{2x}{\sqrt{1-x^4}}\end{aligned}$$

- 13d $f''(x) = 2\cos^{-1}x^2 + 4x^2\cos^{-1}x^2$

$$f'(0) = 0$$

$$f''(0) = -\pi < 0$$

Hence at $x = 0$ the derivative is zero and the curve is concave down, so it is a maximum turning point.

13e


When $x = -1$, $f'(-1) = -\frac{2(-1)}{\sqrt{1-(-1)^4}} = \frac{2}{\sqrt{0}}$ which is undefined.

When $x = 1$, $f'(1) = -\frac{2(1)}{\sqrt{1-(1)^4}} = \frac{2}{\sqrt{0}}$ which is undefined.

This means that the gradient is vertical at that point.

Chapter 12 worked solutions – Further calculus

13f



14a

$$\tan \angle TEP = \frac{4}{x}$$

$$\angle TEP = \tan^{-1} \frac{4}{x}$$

$$\tan \angle BEP = \frac{3}{x}$$

$$\angle BEP = \tan^{-1} \frac{3}{x}$$

Hence

$$\theta = \angle TEP - \angle BEP = \tan^{-1} \frac{4}{x} - \tan^{-1} \frac{3}{x}$$

14b

$$\begin{aligned} \frac{d\theta}{dx} &= \frac{1}{\sqrt{1 - \left(\frac{4}{x}\right)^2}} \times \frac{d}{dx} \left(\frac{4}{x}\right) - \frac{1}{\sqrt{1 - \left(\frac{3}{x}\right)^2}} \times \frac{d}{dx} \left(\frac{3}{x}\right) \\ &= \frac{1}{\sqrt{1 - \frac{16}{x^2}}} \times \left(-\frac{4}{x^2}\right) - \frac{1}{\sqrt{1 - \frac{9}{x^2}}} \times \left(-\frac{3}{x^2}\right) \\ &= \frac{-4}{x\sqrt{x^2 - 16}} + \frac{3}{x\sqrt{x^2 - 9}} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

When $x = 2\sqrt{3}$

$$\begin{aligned}\frac{d\theta}{dx} &= \frac{-4}{(2\sqrt{3})\sqrt{(2\sqrt{3})^2 - 16}} + \frac{3}{(2\sqrt{3})\sqrt{(2\sqrt{3})^2 - 9}} \\ &= 0\end{aligned}$$

Hence there is a stationary point

When $x = \sqrt{3}$, $\frac{d\theta}{dx} > 0$ and when $x = 3\sqrt{3}$, $\frac{d\theta}{dx} < 0$. Hence this is a maximum turning point.

14c When $x = 2\sqrt{3}$

$$\theta = \tan^{-1} \frac{4}{2\sqrt{3}} - \tan^{-1} \frac{3}{2\sqrt{3}}$$

$\tan \theta$

$$\begin{aligned}&= \tan \left(\tan^{-1} \frac{4}{2\sqrt{3}} - \tan^{-1} \frac{3}{2\sqrt{3}} \right) \\ &= \frac{\tan \left(\tan^{-1} \frac{4}{2\sqrt{3}} \right) - \tan \left(\tan^{-1} \frac{3}{2\sqrt{3}} \right)}{1 + \tan \left(\tan^{-1} \frac{4}{2\sqrt{3}} \right) \tan \left(\tan^{-1} \frac{3}{2\sqrt{3}} \right)} \\ &= \frac{\frac{4}{2\sqrt{3}} - \frac{3}{2\sqrt{3}}}{1 + \frac{4}{2\sqrt{3}} \frac{3}{2\sqrt{3}}} \\ &= \frac{\frac{1}{2\sqrt{3}}}{1 + \frac{1}{2}} \\ &= \frac{1}{4\sqrt{3}} \\ &= \frac{\sqrt{3}}{12}\end{aligned}$$

Hence

$$\theta = \tan^{-1} \frac{\sqrt{3}}{12}$$

Chapter 12 worked solutions – Further calculus

15a

$\theta = \angle OPA$ (equal alternate angles on parallel lines)

$$\tan \theta = \tan \angle OPA = \frac{OA}{AP} = \frac{6}{x}$$

Hence

$$\theta = \tan^{-1} \frac{6}{x}$$

15b

$$\begin{aligned}\frac{d\theta}{dx} &= \frac{1}{1 + \left(\frac{6}{x}\right)^2} \times \frac{d}{dx}\left(\frac{6}{x}\right) \\ &= \frac{1}{1 + \frac{36}{x^2}} \times \left(-\frac{6}{x^2}\right) \\ &= \frac{-6}{x^2 + 36}\end{aligned}$$

$$\text{Now } \frac{dx}{dt} = 600$$

Hence

$$\begin{aligned}\frac{d\theta}{dt} &= \frac{d\theta}{dx} \times \frac{dx}{dt} \\ &= \frac{-6}{x^2 + 36} \times 600 \\ &= \frac{-3600}{x^2 + 36}\end{aligned}$$

15c When $x = 3$,

$$\frac{d\theta}{dt} = \frac{-3600}{3^2 + 36} = -80$$

Hence it is travelling at 80 radians per hour which is $\frac{1}{45}$ rad/s.

- 16a Note that $\tan^{-1} x$ is defined for all values of x , the only value for which the function is undefined is when $x = 0$ as $\frac{1}{x}$ is undefined when $x = 0$. Hence the domain is all x such that $x \neq 0$.

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 f(-x) &= \tan^{-1}(-x) + \tan^{-1}\left(\frac{1}{-x}\right) \\
 &= -\tan^{-1}(x) - \tan^{-1}\left(\frac{1}{x}\right) \\
 &= -\left(\tan^{-1}(x) + \tan^{-1}\left(\frac{1}{x}\right)\right) \\
 &= -f(x)
 \end{aligned}$$

Hence the function is odd so it has rotational symmetry about the origin.

16b

$$\begin{aligned}
 f'(x) &= \frac{1}{1+x^2} + \frac{1}{1+\left(\frac{1}{x}\right)^2} \times \frac{d}{dx}\left(\frac{1}{x}\right) \\
 &= \frac{1}{1+x^2} + \frac{1}{1+\frac{1}{x^2}} \times \left(-\frac{1}{x^2}\right) \\
 &= \frac{1}{1+x^2} - \frac{1}{1+x^2} \\
 &= 0
 \end{aligned}$$

16c

$$\begin{aligned}
 f(x) &= \int f'(x) dx \\
 &= \int 0 dx \\
 &= C
 \end{aligned}$$

Now as this function has a point of discontinuity at $x = 0$, this constant may be different on either side of the discontinuity.

For $x > 0$, substitute in $x = 1$.


$$C = f(1) = \tan^{-1} 1 + \tan^{-1} 1 = \frac{\pi}{4} + \frac{\pi}{4} = \frac{\pi}{2}$$

For $x < 0$, substitute in $x = -1$.

$$c = f(-1) = -f(1) = -\frac{\pi}{2} \text{ (as the function is odd)}$$

$$\text{Hence for } x > 0, f(x) = \frac{\pi}{2} \text{ and for } x < 0, f(x) = -\frac{\pi}{2}$$

Chapter 12 worked solutions – Further calculus



17a

$$\left| \frac{1}{x} \right| \leq 1$$

$$|x| > 1$$


Hence the domain is $x > 1$ or $x < -1$.

17b

$$\begin{aligned}
 f'(x) &= \frac{-1}{\sqrt{1 - \left(\frac{1}{x}\right)^2}} \times \frac{d}{dx}\left(\frac{1}{x}\right) \\
 &= \frac{-1}{\sqrt{1 - \frac{1}{x^2}}} \times -\frac{1}{x^2} \\
 &= \frac{1}{x^2 \sqrt{1 - \frac{1}{x^2}}} \\
 &= \frac{1}{\sqrt{x^4} \sqrt{1 - \frac{1}{x^2}}} \\
 &= \frac{1}{\sqrt{x^2} \sqrt{x^2 - 1}} \\
 &= \frac{1}{|x| \sqrt{x^2 - 1}}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

17c

17d When $x > 1$,

$$f'(x) = \frac{1}{x\sqrt{x^2 - 1}}$$

When $x < 1$,

$$f'(x) = -\frac{1}{x\sqrt{x^2 - 1}}$$

17e For $x > 1$ we use the equation

$$f'(x) = \frac{1}{x\sqrt{x^2 - 1}}$$

Hence $x > 0$ and $\sqrt{x^2 - 1} > 0$ thus $f'(x) > 0$ For $x < -1$ we use the equation

$$f'(x) = -\frac{1}{x\sqrt{x^2 - 1}} = \frac{1}{-x\sqrt{x^2 - 1}}$$


Hence $-x > 0$ and $\sqrt{x^2 - 1} > 0$ thus $f'(x) > 0$

$$17\text{f i } \lim_{x \rightarrow \infty} \cos^{-1} \frac{1}{x} = \lim_{x \rightarrow \infty} \cos^{-1} 0^+ = \frac{\pi^+}{2}$$

$$17\text{f ii } \lim_{x \rightarrow \infty} \cos^{-1} \frac{1}{x} = \lim_{x \rightarrow \infty} \cos^{-1} 0^- = \frac{\pi^-}{2}$$

Chapter 12 worked solutions – Further calculus

17g



$$\begin{aligned}
 18a \quad & \frac{d}{dx} (\tan^{-1} e^{3x}) \\
 &= \frac{1}{1 + (e^{3x})^2} \times \frac{d}{dx} (e^{3x}) \\
 &= \frac{1}{1 + (e^{3x})^2} \times 3e^{3x} \\
 &= \frac{3e^{3x}}{1 + e^{6x}}
 \end{aligned}$$

$$\begin{aligned}
 18b \quad & \frac{d}{dx} (\sin^{-1} x^3) \\
 &= \frac{1}{\sqrt{1 - (x^3)^2}} \times \frac{d}{dx} (x^3) \\
 &= \frac{1}{\sqrt{1 - (x^3)^2}} \times 3x^2 \\
 &= \frac{3x^2}{\sqrt{1 - x^6}}
 \end{aligned}$$

$$\begin{aligned}
 18c \quad & \frac{d}{dx} (\cos^{-1} \log_e x) \\
 &= \frac{-1}{\sqrt{1 - (\log_e x)^2}} \times \frac{d}{dx} (\log_e x) \\
 &= \frac{-1}{\sqrt{1 - (\log_e x)^2}} \times \frac{1}{x} \\
 &= -\frac{1}{x \sqrt{1 - (\log_e x)^2}}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

19a The restrictions on x are

$$-1 \leq x \leq 1$$

and

$$-1 \leq \sqrt{1 - x^2} \leq 1$$

But $\sqrt{1 - x^2} \geq 0$, so

$$0 \leq \sqrt{1 - x^2} \leq 1$$

$$0 \leq 1 - x^2 \leq 1$$

$$-1 \leq -x^2 \leq 0$$

$$0 \leq x^2 \leq 1$$

Hence the domain is $-1 \leq x \leq 1$.

19b

$$\begin{aligned} g'(x) &= \frac{1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1-(\sqrt{1-x^2})^2}} \times \frac{d}{dx}(\sqrt{1-x^2}) \\ &= \frac{1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1-(1-x^2)}} \times \left(\frac{1}{2} \times -2x \times (1-x^2)^{-\frac{1}{2}} \right) \\ &= \frac{1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{x^2}} \times \left(-\frac{x}{\sqrt{1-x^2}} \right) \\ &= \frac{1}{\sqrt{1-x^2}} - \frac{1}{|x|} \times \frac{x}{\sqrt{1-x^2}} \\ &= \frac{1}{\sqrt{1-x^2}} - \frac{x}{|x|\sqrt{1-x^2}} \end{aligned}$$

19c $g(x)$ is constant when $g'(x) = 0$, hence

$$g'(x) = 0$$

$$\frac{1}{\sqrt{1-x^2}} - \frac{x}{|x|\sqrt{1-x^2}} = 0$$

$$\frac{1}{\sqrt{1-x^2}} = \frac{x}{|x|\sqrt{1-x^2}}$$

For $x \neq 1$ and $x \neq -1$

$$|x|\sqrt{1-x^2} = x\sqrt{1-x^2}$$

Chapter 12 worked solutions – Further calculus

$$|x| = x$$

$$|x| - x = 0$$

This is true for $x \geq 0$. But $x \neq 0$ so the function is constant for $x > 0$.

20

$\tan^{-1} \frac{x+2}{1-2x}$ is simply $\tan^{-1} x + \tan^{-1} 2$ for $x < \frac{1}{2}$, and is $\tan^{-1} x + \tan^{-1} 2 - \pi$ for $x > \frac{1}{2}$

(this can be proven by applying tan to both equations and then using double angle formulae to show they have the same result)

- 21a As $\sin x$ is defined for all real x , and because $-1 \leq \sin x \leq 1$, this function will be defined for all real x .

Now since $-1 \leq \sin x \leq 1$

$$\sin^{-1}(-1) \leq \sin^{-1}(\sin x) \leq \sin^{-1}(1)$$

$$-\frac{\pi}{2} \leq \sin^{-1}(\sin x) \leq \frac{\pi}{2}$$

So the range is

$$-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$$

21b

$$\begin{aligned} f'(x) &= \frac{d}{dx} (\sin^{-1}(\sin(x))) \\ &= \frac{1}{\sqrt{1 - (\sin x)^2}} \times \frac{d}{dx} (\sin x) \\ &= \frac{1}{\sqrt{1 - \sin^2 x}} \times \cos x \\ &= \frac{\cos x}{\sqrt{\cos^2 x}} \\ &= \frac{\cos x}{|\cos x|} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

21c When $\cos x = 0$, $f'(x) = \frac{0}{|\cos x|}$ which is undefined. Hence $f'(x)$ is not defined when $\cos x = 0$.

21d Since $|\cos x| = \cos x$ or $|\cos x| = -\cos x$. Hence

$$f'(x) = \frac{\cos x}{\cos x} = 1$$


or

$$f'(x) = \frac{\cos x}{-\cos x} = -1$$


Now $f'(x) = 1$ when $\cos x > 0$ which is when $-\frac{\pi}{2} + 2\pi n < x < -\frac{\pi}{2} + 2\pi n$ for all integers n .

Now $f'(x) = -1$ when $\cos x < 0$ which is when $\frac{\pi}{2} + 2\pi n < x < \frac{3\pi}{2} + 2\pi n$ for all integers n .

21e



22a



Chapter 12 worked solutions – Further calculus

Solutions to Exercise 12B

1 Answers will vary

2a

$$\int -\frac{1}{\sqrt{1-x^2}} dx = \cos^{-1} x + C$$

2b

$$\begin{aligned} & \int \frac{1}{\sqrt{4-x^2}} dx \\ &= \int \frac{1}{2\sqrt{1-\frac{x^2}{4}}} dx \\ &= \int \frac{1}{2} \times \frac{1}{\sqrt{1-\left(\frac{x}{2}\right)^2}} dx \\ &= \sin^{-1}\left(\frac{x}{2}\right) + C \end{aligned}$$

2c

$$\begin{aligned} & \int \frac{1}{9+x^2} dx \\ &= \int \frac{1}{9} \frac{1}{1+\frac{x^2}{9}} dx \\ &= \frac{1}{3} \int \frac{1}{3} \frac{1}{1+\left(\frac{x}{3}\right)^2} dx \\ &= \frac{1}{3} \tan^{-1}\left(\frac{x}{3}\right) + C \end{aligned}$$

Chapter 12 worked solutions – Further calculus

2d

$$\begin{aligned} & \int \frac{1}{\sqrt{\frac{4}{9} - x^2}} dx \\ &= \int \frac{1}{\frac{2}{3} \sqrt{1 - \left(\frac{3}{2}x\right)^2}} dx \\ &= \int \frac{3}{2} \frac{1}{\sqrt{1 - \left(\frac{3}{2}x\right)^2}} dx \\ &= \sin^{-1}\left(\frac{3}{2}x\right) + C \end{aligned}$$

2e

$$\begin{aligned} & \int \frac{1}{2+x^2} dx \\ &= \int \frac{1}{2} \frac{1}{1 + \left(\frac{x}{\sqrt{2}}\right)^2} dx \\ &= \frac{\sqrt{2}}{2} \tan^{-1} \frac{x}{\sqrt{2}} + C \end{aligned}$$

2f

$$\begin{aligned} & \int -\frac{1}{\sqrt{5-x^2}} dx \\ &= \int \frac{1}{\sqrt{5}} \frac{-1}{\sqrt{1 - \left(\frac{x}{\sqrt{5}}\right)^2}} dx \\ &= \cos^{-1}\left(\frac{x}{\sqrt{5}}\right) + C \end{aligned}$$

Chapter 12 worked solutions – Further calculus

3a

$$\begin{aligned}
 & \int_0^3 \frac{1}{\sqrt{9-x^2}} dx \\
 &= \int_0^3 \frac{1}{3\sqrt{1-\left(\frac{x}{3}\right)^2}} dx \\
 &= \left[\sin^{-1} \frac{x}{3} \right]_0^3 \\
 &= \sin^{-1} 1 - \sin^{-1} 0 \\
 &= \frac{\pi}{2}
 \end{aligned}$$

3b

$$\begin{aligned}
 & \int_0^2 \frac{1}{4+x^2} dx \\
 &= \int_0^2 \frac{1}{4} \frac{1}{1+\left(\frac{x}{2}\right)^2} dx \\
 &= \frac{1}{2} \left[\tan^{-1} \left(\frac{x}{2} \right) \right]_0^2 \\
 &= \frac{1}{2} (\tan^{-1} 1 - \tan^{-1} 0) \\
 &= \frac{\pi}{8}
 \end{aligned}$$

3c

$$\begin{aligned}
 & \int_0^1 \frac{1}{\sqrt{2-x^2}} dx \\
 &= \int_0^1 \frac{1}{\sqrt{2}} \frac{1}{\sqrt{1-\left(\frac{x}{\sqrt{2}}\right)^2}} dx \\
 &= \left[\sin^{-1} \left(\frac{x}{\sqrt{2}} \right) \right]_0^1
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \sin^{-1}\left(\frac{1}{\sqrt{2}}\right) - \sin^{-1}\left(\frac{0}{\sqrt{2}}\right)$$

$$= \frac{\pi}{4}$$

3d

$$\begin{aligned} & \int_{\frac{1}{2}}^{\frac{1}{2}\sqrt{3}} \frac{\frac{1}{2}}{\frac{1}{4} + x^2} dx \\ &= \int_{\frac{1}{2}}^{\frac{1}{2}\sqrt{3}} \frac{2}{1 + 4x^2} dx \\ &= \int_{\frac{1}{2}}^{\frac{1}{2}\sqrt{3}} \frac{2}{1 + (2x)^2} dx \\ &= [\tan^{-1}(2x)]_{\frac{1}{2}}^{\frac{1}{2}\sqrt{3}} \\ &= \tan^{-1}(2\sqrt{3}) - \tan^{-1}(1) \\ &= \frac{\pi}{12} \end{aligned}$$

3e

$$\begin{aligned} & \int_{\frac{\sqrt{3}}{6}}^{\frac{1}{6}} \frac{-1}{\sqrt{\frac{1}{9} - x^2}} dx \\ &= \int_{\frac{\sqrt{3}}{6}}^{\frac{1}{6}} \frac{-1}{\frac{1}{3}\sqrt{1 - (3x)^2}} dx \\ &= \int_{\frac{\sqrt{3}}{6}}^{\frac{1}{6}} \frac{-3}{\sqrt{1 - (3x)^2}} dx \\ &= [\cos^{-1}(3x)]_{\frac{\sqrt{3}}{6}}^{\frac{1}{6}} \\ &= \cos^{-1}\frac{1}{2} - \cos^{-1}\frac{\sqrt{3}}{2} \\ &= \frac{\pi}{6} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

3f

$$\begin{aligned}
 & \int_{-\frac{3}{4}\sqrt{2}}^{\frac{3}{4}} \frac{1}{\sqrt{\frac{9}{4} - x^2}} dx \\
 &= \int_{-\frac{3}{4}\sqrt{2}}^{\frac{3}{4}} \frac{1}{\frac{3}{2}\sqrt{1 - \left(\frac{2x}{3}\right)^2}} \\
 &= \int_{-\frac{3}{4}\sqrt{2}}^{\frac{3}{4}} \frac{2}{3\sqrt{1 - \left(\frac{2x}{3}\right)^2}} \\
 &= \left[\sin^{-1}\left(\frac{2x}{3}\right) \right]_{-\frac{3}{4}\sqrt{2}}^{\frac{3}{4}} \\
 &= \sin^{-1}\frac{1}{2} - \sin^{-1}\left(-\frac{\sqrt{2}}{2}\right) \\
 &= \frac{5\pi}{12}
 \end{aligned}$$

4a

$$y' = (1 - x^2)^{-\frac{1}{2}} = \frac{1}{\sqrt{1 - x^2}}$$

$$y = \sin^{-1} x + C$$

Substituting in $(0, \pi)$

$$\pi = \sin^{-1} 0 + C$$

$$\pi = 0 + C$$

$$C = \pi$$

Hence

$$y = \sin^{-1} x + \pi$$

4b

$$y' = 4(16 + x^2)^{-1}$$

$$= \frac{1}{4(16 + x^2)}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{4}{4 \times 16} \times \frac{1}{1 + \left(\frac{x}{4}\right)^2}$$

$$= \frac{1}{4} \times \frac{1}{1 + \left(\frac{x}{4}\right)^2}$$

$$y = \tan^{-1} \frac{x}{4} + C$$

Substituting in $(-4, 0)$

$$0 = \tan^{-1} -1 + C$$

$$0 = -\frac{\pi}{4} + C$$

$$C = \frac{\pi}{4}$$

Hence

$$y = \tan^{-1} \frac{x}{4} + \frac{\pi}{4}$$

5a

$$y' = \frac{1}{\sqrt{36 - x^2}} = \frac{1}{6\sqrt{1 - \left(\frac{x}{6}\right)^2}}$$

$$y = \sin^{-1} \left(\frac{x}{6}\right) + C$$

$$\text{When } x = 3, y = \frac{\pi}{6}$$

$$\frac{\pi}{6} = \sin^{-1} \frac{1}{2} + C$$

$$C = 0$$

$$y = \sin^{-1} \left(\frac{x}{6}\right)$$

$$\text{When } x = 3\sqrt{3}$$

$$y = \sin^{-1} \frac{\sqrt{3}}{2} = \frac{\pi}{3}$$

Chapter 12 worked solutions – Further calculus

5b

$$y' = \frac{2}{4+x^2} = \frac{1}{2} \frac{1}{1+\left(\frac{x}{2}\right)^2}$$

$$y = \tan^{-1} \frac{x}{2} + C$$

$$\text{When } x = 2, y = \frac{\pi}{3}$$

$$\frac{\pi}{3} = \tan^{-1} 1 + C$$

$$\frac{\pi}{3} = \frac{\pi}{4} + C$$

$$C = \frac{\pi}{12}$$

$$y = \tan^{-1} \frac{x}{2} + \frac{\pi}{12}$$

$$\text{When } x = \frac{2}{\sqrt{3}}$$

$$y = \tan^{-1} \frac{1}{\sqrt{3}} + \frac{\pi}{12} = \frac{\pi}{4}$$

6a

$$\begin{aligned} & \int \frac{1}{\sqrt{1-4x^2}} dx \\ &= \int \frac{1}{\sqrt{1-(2x)^2}} dx \\ &= \frac{1}{2} \sin^{-1} 2x + C \end{aligned}$$

6b

$$\begin{aligned} & \int \frac{1}{1+16x^2} dx \\ &= \int \frac{1}{1+(4x)^2} dx \\ &= \frac{1}{4} \tan^{-1} 4x + C \end{aligned}$$

Chapter 12 worked solutions – Further calculus

6c

$$\begin{aligned} & \int -\frac{1}{\sqrt{1-2x^2}} dx \\ &= \int -\frac{1}{\sqrt{1-(\sqrt{2}x)^2}} dx \\ &= \frac{1}{\sqrt{2}} \cos^{-1} \sqrt{2}x + C \end{aligned}$$

6d

$$\begin{aligned} & \int \frac{1}{\sqrt{4-9x^2}} dx \\ &= \int \frac{1}{2\sqrt{1-\left(\frac{3x}{2}\right)^2}} dx \\ &= \frac{1}{3} \sin^{-1} \frac{3x}{2} + C \end{aligned}$$

6e

$$\begin{aligned} & \int \frac{1}{25+9x^2} dx \\ &= \int \frac{1}{25} \frac{1}{1+\left(\frac{3x}{5}\right)^2} dx \\ &= \frac{1}{15} \tan^{-1} \frac{3x}{5} + C \end{aligned}$$

6f

$$\begin{aligned} & \int -\frac{1}{\sqrt{3-4x^2}} dx \\ &= \int -\frac{1}{\sqrt{3}} \frac{1}{\sqrt{1-\left(\frac{2x}{\sqrt{3}}\right)^2}} dx \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{1}{2} \cos^{-1} \frac{2x}{\sqrt{3}} + C$$

7a

$$\begin{aligned} & \int_0^{\frac{1}{6}} \frac{1}{\sqrt{1 - 9x^2}} dx \\ &= \int_0^{\frac{1}{6}} \frac{1}{\sqrt{1 - (3x)^2}} dx \\ &= \frac{1}{3} [\sin^{-1} 3x]_0^{\frac{1}{6}} \\ &= \frac{\pi}{18} \end{aligned}$$

7b

$$\begin{aligned} & \int_{\frac{1}{2}}^{\frac{1}{2}\sqrt{3}} \frac{2}{1 + 4x^2} dx \\ &= \int_{\frac{1}{2}}^{\frac{1}{2}\sqrt{3}} \frac{2}{1 + (2x)^2} dx \\ &= [\tan^{-1}(2x)]_{\frac{1}{2}}^{\frac{1}{2}\sqrt{3}} \\ &= \frac{\pi}{12} \end{aligned}$$

7c

$$\begin{aligned} & \int_{-\frac{1}{2}\sqrt{3}}^{\frac{1}{2}} \frac{1}{\sqrt{1 - 3x^2}} dx \\ &= \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{1}{\sqrt{1 - (\sqrt{3}x)^2}} dx \\ &= \frac{1}{\sqrt{3}} [\sin^{-1} \sqrt{3}x]_{-\frac{1}{2}}^{\frac{1}{2}} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{2\pi}{9}\sqrt{3}$$

7d

$$\begin{aligned} & \int_{-\frac{3}{4}}^{\frac{3}{2\sqrt{2}}} \frac{1}{\sqrt{9 - 4x^2}} dx \\ &= \int_{-\frac{3}{4}}^{\frac{3}{2\sqrt{2}}} \frac{1}{3\sqrt{1 - \left(\frac{2x}{3}\right)^2}} dx \\ &= \frac{1}{2} \left[\sin^{-1} \left(\frac{2x}{3} \right) \right]_{-\frac{3}{4}}^{\frac{3}{2\sqrt{2}}} \\ &= \frac{5\pi}{24} \end{aligned}$$

7e


$$\begin{aligned} & \int_{-\frac{1}{2}}^{\frac{3}{2}} \frac{1}{3 + 4x^2} dx \\ &= \int_{-\frac{1}{2}}^{\frac{3}{2}} \frac{1}{3} \frac{1}{1 + \left(\frac{2x}{\sqrt{3}}\right)^2} dx \\ &= \frac{\sqrt{3}}{6} \left[\tan^{-1} \frac{2x}{\sqrt{3}} \right]_{-\frac{1}{2}}^{\frac{3}{2}} \\ &= \frac{\pi}{12}\sqrt{3} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

7f

$$\begin{aligned}
 & \int_{\frac{1}{2}\sqrt{10}}^{\frac{1}{2}\sqrt{30}} \frac{1}{5 + 2x^2} dx \\
 &= \int_{\frac{1}{2}\sqrt{10}}^{\frac{1}{2}\sqrt{30}} \frac{1}{5} \frac{1}{1 + \left(\frac{\sqrt{2}x}{\sqrt{5}}\right)^2} dx \\
 &= \frac{\sqrt{5}}{5\sqrt{2}} \left[\tan^{-1} \frac{\sqrt{2}x}{\sqrt{5}} \right]_{\frac{1}{2}\sqrt{10}}^{\frac{1}{2}\sqrt{30}} \\
 &= \frac{\pi}{120} \sqrt{10}
 \end{aligned}$$

8a



Chapter 12 worked solutions – Further calculus

8b


$$\begin{aligned} & \frac{d}{dx} \left(x \sin^{-1} x + \sqrt{1 - x^2} \right) \\ &= \frac{d}{dx} (x \sin^{-1} x) + \frac{d}{dx} (\sqrt{1 - x^2}) \\ &= \frac{d}{dx} (x) \sin^{-1} x + x \frac{d}{dx} (\sin^{-1} x) + \frac{d}{dx} ((1 - x^2)^{\frac{1}{2}}) \\ &= \sin^{-1} x + \frac{x}{\sqrt{1 - x^2}} + \frac{1}{2} \times -2x \times \frac{1}{\sqrt{1 - x^2}} \\ &= \sin^{-1} x + \frac{x}{\sqrt{1 - x^2}} - \frac{x}{\sqrt{1 - x^2}} \\ &= \sin^{-1} x \end{aligned}$$

8c

$$\begin{aligned} & \int_0^{\frac{1}{2}} \sin^{-1} x \, dx \\ &= \int_0^{\frac{1}{2}} \frac{d}{dx} \left(x \sin^{-1} x + \sqrt{1 - x^2} \right) \, dx \\ &= \left[x \sin^{-1} x + \sqrt{1 - x^2} \right]_0^{\frac{1}{2}} \\ &= \left(\frac{\pi}{12} + \frac{1}{2}\sqrt{3} - 1 \right) \text{ square units} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

9a



9b

$$\begin{aligned}A &= \int_0^{\frac{\pi}{6}} \sin y \, dy \\&= -[\cos y]_0^{\frac{\pi}{6}} \\&= \left(1 - \frac{1}{2}\sqrt{3}\right) \text{ square units}\end{aligned}$$

- 9c Note that the area formed by combined regions of the previous two questions is that of a square with area $\frac{\pi}{6} \times \frac{1}{2} = \frac{\pi}{12}$ units².

Thus, the area in this question will be $\frac{\pi}{12} - (\frac{\pi}{12} + \frac{1}{2}\sqrt{3} - 1) = 1 - \frac{1}{2}\sqrt{3}$ units²

Chapter 12 worked solutions – Further calculus

10a

$$\begin{aligned}
 & \frac{d}{dx}(\cos^{-1}(2-x)) \\
 &= \frac{-1}{\sqrt{1-(2-x)^2}} \times \frac{d}{dx}(2-x) \\
 &= \frac{-1}{\sqrt{1-(2-x)^2}} \times -1 \\
 &= \frac{1}{\sqrt{1-(4-2x+x^2)}} \\
 &= \frac{1}{\sqrt{4x-x^2-3}}
 \end{aligned}$$

10b

$$\begin{aligned}
 & \int_1^2 \frac{1}{\sqrt{4x-x^2-3}} dx \\
 &= [\cos^{-1}(2-x)]_1^2 \\
 &= \cos^{-1} 0 - \cos^{-1} 1 \\
 &= \frac{\pi}{2}
 \end{aligned}$$

11a

$$\begin{aligned}
 & \frac{d}{dx}\left(\tan^{-1}\frac{1}{2}x^3\right) \\
 &= \frac{1}{1+\left(\frac{1}{2}x^3\right)^2} \times \frac{d}{dx}\left(\frac{1}{2}x^3\right) \\
 &= \frac{1}{1+\frac{x^6}{4}} \times \frac{3}{2}x^2 \\
 &= \frac{3x^2}{2+\frac{x^6}{2}} \\
 &= \frac{6x^2}{4+x^6}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

11b

$$\begin{aligned}
 & \int \frac{x^2}{4+x^6} dx \\
 &= \frac{1}{6} \int \frac{6x^2}{4+x^6} dx \\
 &= \frac{1}{6} \int \frac{d}{dx} \left(\tan^{-1} \frac{1}{2} x^3 \right) dx \\
 &= \frac{1}{6} \tan^{-1} \frac{1}{2} x^3 + C
 \end{aligned}$$

12a

$$\begin{aligned}
 & \frac{d}{dx} (x \tan^{-1} x) \\
 &= \tan^{-1} x \times \frac{d}{dx} (x) + x \times \frac{d}{dx} (\tan^{-1} x) \\
 &= \tan^{-1} x + \frac{x}{1+x^2}
 \end{aligned}$$

12b

$$\begin{aligned}
 & \frac{d}{dx} (x \tan^{-1} x) = \tan^{-1} x + \frac{x}{1+x^2} \\
 & \tan^{-1} x = \frac{d}{dx} (x \tan^{-1} x) - \frac{x}{1+x^2} \\
 & \int_0^1 \tan^{-1} x \, dx \\
 &= \int_0^1 \left(\frac{d}{dx} (x \tan^{-1} x) - \frac{x}{1+x^2} \right) dx \\
 &= \int_0^1 \frac{d}{dx} (x \tan^{-1} x) \, dx - \int_0^1 \frac{x}{1+x^2} \, dx \\
 &= [x \tan^{-1} x]_0^1 - \left[\frac{1}{2} \ln(1+x^2) \right]_0^1 \\
 &= \frac{\pi}{4} - \frac{1}{2} (\ln 2 - 0) \\
 &= \frac{\pi}{4} - \frac{1}{2} \ln 2
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

13a 0 as this is an odd function

13b 0 as this is an odd function

13c

$$\begin{aligned} & \int_{-\frac{3}{4}}^{\frac{3}{4}} \cos^{-1} x \, dx \\ &= - \left[\frac{1}{\sqrt{1-x^2}} \right]_{-\frac{3}{4}}^{\frac{3}{4}} \\ &= \frac{3\pi}{4} \end{aligned}$$

13d 0 as this is an odd function

13e 0 as this is an odd function

13f

This is a semicircle of radius 6. Hence

$$\begin{aligned} & \int_{-6}^6 \sqrt{36-x^2} \, dx \\ &= \frac{1}{2} \times (\pi \times 6^2) \\ &= 18\pi \text{ square units} \end{aligned}$$

14a i $f(0) = \frac{0}{1+0^2} - \tan^{-1} 0 = 0 - 0 = 0$

14a ii

$$f'(x) = \frac{d}{dx} \left(\frac{x}{1+x^2} \right) - \frac{d}{dx} (\tan^{-1} x)$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \frac{(1+x^2) \times \frac{d}{dx}(x) - x \times \frac{d}{dx}(1+x^2)}{(1+x^2)^2} - \frac{1}{1+x^2} \\
 &= \frac{1+x^2 - x(2x)}{(1+x^2)^2} - \frac{1+x^2}{(1+x^2)^2} \\
 &= \frac{-2x^2}{(1+x^2)^2}
 \end{aligned}$$

14b i Firstly consider $f'(x)$

As the numerator $-2x^2 < 0$ for all $x > 0$ and the denominator $(1+x^2)^2 < 0$ for all $x \geq 0$. It follows that $f'(x) < 0$ for all $x < 0$.


Since $f(0) = 0$, and as the function is strictly decreasing for all $x > 0$, it follows that $f(x) < 0$ for all $x > 0$.

14b ii

$$\begin{aligned}
 &\int_0^1 \frac{x^2}{(1+x^2)^2} dx \\
 &= -\frac{1}{2} \int_0^1 \frac{-2x^2}{(1+x^2)^2} dx \\
 &= -\frac{1}{2} \int_0^1 f'(x) dx \\
 &= -\frac{1}{2} [f(x)]_0^1 \\
 &= -\frac{1}{2} [f(1) - f(0)] \\
 &= -\frac{1}{2} f(1) \\
 &= -\frac{1}{2} \left[\frac{1}{2} - \frac{\pi}{4} \right] \\
 &= \frac{1}{8} [\pi - 2]
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

15a,b



- 15c From the graph the domain is $-2 < x < 2$ (note that this is strictly less than) and the range is $x \geq \frac{1}{2}$. Its symmetry is even as $f(-x) = \frac{1}{\sqrt{4-(-x)^2}} = \frac{1}{\sqrt{4-x^2}} = f(x)$

15d

$$\begin{aligned}
 A &= \int_{-1}^1 \frac{1}{\sqrt{4-x^2}} dx \\
 &= \int_{-1}^1 \frac{1}{2\sqrt{1-\left(\frac{x}{2}\right)^2}} dx \\
 &= \left[\sin^{-1}\left(\frac{x}{2}\right) \right]_{-1}^1 \\
 &= \sin^{-1}\frac{1}{2} - \sin^{-1}\left(-\frac{1}{2}\right) \\
 &= \frac{\pi}{6} - \left(-\frac{\pi}{6}\right) \\
 &= \frac{\pi}{3} \text{ square units}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

15e

$$\begin{aligned}
 A &= \int_{-2}^2 \frac{1}{\sqrt{4-x^2}} dx \\
 &= \int_{-2}^2 \frac{1}{2\sqrt{1-\left(\frac{x}{2}\right)^2}} dx \\
 &= \left[\sin^{-1}\left(\frac{x}{2}\right) \right]_{-2}^2 \\
 &= \sin^{-1} 1 - \sin^{-1}(-1) \\
 &= \frac{\pi}{2} - \left(-\frac{\pi}{2}\right) \\
 &= \pi \text{ square units}
 \end{aligned}$$

16a

$$f(-x) = \frac{4}{(-x)^2 + 4} = \frac{4}{x^2 + 4} = f(x)$$

Hence the function is even and thus has an axis of symmetry about the y -axis.

16b The function is defined for all real x so the domain is all real x .

As $x^2 + 4 \geq 4$ it follows that $0 < \frac{1}{x^2+4} \leq \frac{1}{4}$ and hence $0 < \frac{4}{x^2+4} \leq 1$ thus the range is $0 < f(x) \leq 1$.

16c

$$\begin{aligned}
 f'(x) &= \frac{d}{dx}(4(x^2 + 4)^{-1}) \\
 &= -1 \times 4(x^2 + 4)^{-2} \times \frac{d}{dx}(x^2) \\
 &= -1 \times 4(x^2 + 4)^{-2} \times 2x \\
 &= -\frac{8x}{(x^2 + 4)^2}
 \end{aligned}$$

Hence

$$f'(0) = 0$$

and

$$f(0) = 1$$

Chapter 12 worked solutions – Further calculus

So there is a turning point at $(0, 1)$.


Testing for maximum

x	-1	0	1
$f'(x)$	$\frac{8}{25}$	0	$-\frac{8}{25}$

Thus the curve is increasing before the turning point decreasing afterwards.
Thus it is a maximum

16d

$$\begin{aligned}
 & \lim_{x \rightarrow \infty} f(x) \\
 &= \lim_{x \rightarrow \infty} \frac{4}{x^2 + 4} \\
 &= \lim_{x \rightarrow \infty} \frac{\frac{4}{x^2}}{1 + \frac{4}{x^2}} \\
 &= \frac{0}{1 + 0} \\
 &= 0
 \end{aligned}$$



Chapter 12 worked solutions – Further calculus

16e

$$\begin{aligned}
 A &= \int_{-2\sqrt{3}}^{2\sqrt{3}} \frac{4}{x^2 + 4} dx \\
 &= \int_{-2\sqrt{3}}^{2\sqrt{3}} \frac{1}{\left(\frac{x}{2}\right)^2 + 1} dx \\
 &= 2 \left[\tan^{-1} \frac{x}{2} \right]_{-2\sqrt{3}}^{2\sqrt{3}} \\
 &= 2 \left(\tan^{-1} \sqrt{3} - \tan^{-1} (-\sqrt{3}) \right) \\
 &= \pi \text{ square units}
 \end{aligned}$$

16f

$$\begin{aligned}
 A &= \int_{-a}^a \frac{4}{x^2 + 4} dx \\
 &= \int_{-a}^a \frac{1}{\left(\frac{x}{2}\right)^2 + 1} dx \\
 &= 2 \left[\tan^{-1} \frac{x}{2} \right]_{-a}^a \\
 &= 2 \left(\tan^{-1} \frac{a}{2} - \tan^{-1} \left(-\frac{a}{2} \right) \right) \\
 &= 4 \tan^{-1} \frac{a}{2} \text{ square units}
 \end{aligned}$$

16g

$$\begin{aligned}
 &\lim_{a \rightarrow \infty} 4 \tan^{-1} a \\
 &= 4 \lim_{a \rightarrow \infty} \tan^{-1} a \\
 &= 4 \left(\frac{\pi}{2} \right) \\
 &= 2\pi \text{ square units}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

17

Since $y = \sin^{-1} x$, $\frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}}$.

When $x = \frac{\sqrt{3}}{2}$, $y = \frac{\pi}{3}$, $\frac{dy}{dx} = 2$. Hence the tangent is

$$y - \frac{\pi}{3} = 2\left(x - \frac{\sqrt{3}}{2}\right)$$

$$y = 2x - \sqrt{3} + \frac{\pi}{3}$$

Rearranging this gives

$$x = \frac{1}{2}\left(y + \sqrt{3} - \frac{\pi}{3}\right)$$

Also note that for the tangent, when $x = 0$, $y = -\sqrt{3} + \frac{\pi}{3}$

Hence the area can be calculated by

$$\begin{aligned} A &= \int_{-\sqrt{3}+\frac{\pi}{3}}^{\frac{\pi}{3}} y_{tangent} dx - \int_0^{\frac{\pi}{3}} y_{curve} dx \\ &= \int_{-\sqrt{3}+\frac{\pi}{3}}^{\frac{\pi}{3}} \frac{1}{2}\left(y + \sqrt{3} - \frac{\pi}{3}\right) dx - \int_0^{\frac{\pi}{3}} \sin y dx \\ &= \frac{1}{2} \left[\frac{y^2}{2} + y \left(\sqrt{3} - \frac{\pi}{3} \right) \right]_{-\sqrt{3}+\frac{\pi}{3}}^{\frac{\pi}{3}} + [\cos y]_0^{\frac{\pi}{3}} \\ &= \frac{1}{4} \text{ square units} \end{aligned}$$

18a

$$\begin{aligned} I &= \int_0^1 f(x) dx \\ &\doteq \frac{b-a}{2n} (f(a) + f(b) + 2(f(x_1) + f(x_2) + \dots)) \\ &= \frac{1-0}{2(4)} (f(0) + f(1) + 2(f(0.25) + f(0.5) + f(0.75))) \\ &= \frac{1}{8} \left(1 + \frac{1}{2} + 2 \left(\frac{16}{17} + 0.8 + 0.64 \right) \right) \\ &= \frac{5323}{6800} \text{ square units} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

18b

$$I = \int_0^1 \frac{1}{1+x^2} dx = [\tan^{-1} x]_0^1 = \frac{\pi}{4}$$

Hence

$$\frac{\pi}{4} \div \frac{5323}{6800}$$

so

$$\pi \div \frac{5323}{1700}$$

19

$$\begin{aligned} & \int_{-\frac{1}{4}}^{\frac{3}{5}} \frac{1}{1+x^2} dx \\ &= [\tan^{-1} x]_{-\frac{1}{4}}^{\frac{3}{5}} \\ &= \tan^{-1} \frac{3}{5} + \tan^{-1} \left(-\frac{1}{4} \right) \end{aligned}$$

Hence

$$\begin{aligned} & \tan \int_{-\frac{1}{4}}^{\frac{3}{5}} \frac{1}{1+x^2} dx \\ &= \tan \left(\tan^{-1} \frac{3}{5} + \tan^{-1} \left(-\frac{1}{4} \right) \right) \\ &= \frac{\tan \left(\tan^{-1} \frac{3}{5} \right) + \tan \left(\tan^{-1} \left(-\frac{1}{4} \right) \right)}{1 - \tan \left(\tan^{-1} \frac{3}{5} \right) \tan \left(\tan^{-1} \left(-\frac{1}{4} \right) \right)} \\ &= \frac{\frac{3}{5} - \frac{1}{4}}{1 - \left(\frac{3}{5} \right) \left(-\frac{1}{4} \right)} \\ &= 1 \end{aligned}$$

Hence

$$\int_{-\frac{1}{4}}^{\frac{3}{5}} \frac{1}{1+x^2} dx = \frac{\pi}{4}$$

Chapter 12 worked solutions – Further calculus

20a Let $x = u^2$, $dx = 2u \, du$

$$\int \frac{1}{\sqrt{x}(1-x)} dx = \int \frac{1}{\sqrt{u^2(1+u^2)}} \times 2u \, du$$

For $u > 0$

$$\begin{aligned} & \int \frac{1}{\sqrt{x}(1-x)} dx \\ &= \int \frac{1}{u(1+u^2)} \times 2u \, du \\ &= 2 \int \frac{1}{(1+u^2)} \, du \\ &= 2 \tan^{-1} u + c \\ &= 2 \tan^{-1} \sqrt{x} + c \end{aligned}$$

For $u < 0$

$$\begin{aligned} & \int \frac{1}{\sqrt{x}(1-x)} dx \\ &= \int \frac{1}{-u(1+u^2)} \times 2u \, du \\ &= -2 \int \frac{1}{(1+u^2)} \, du \\ &= -2 \tan^{-1} u + c \\ &= -2 \tan^{-1} \sqrt{x} + c \end{aligned}$$

20b Let $u = e^x$, $du = e^x \, dx$ hence $du = u \, dx$ and so $\frac{1}{u} \, du = dx$

$$\begin{aligned} & \int_0^1 \frac{1}{e^{-x} + e^x} dx \\ &= \int_1^e \frac{1}{u^{-1} + u} \times \frac{1}{u} du \\ &= \int_1^e \frac{1}{1+u^2} du \\ &= [\tan^{-1} u]_1^e \\ &= \tan^{-1} e - \frac{\pi}{4} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

21a

$$\begin{aligned}
 & \frac{d}{dx} \left(\tan^{-1} \left(\frac{3}{2} \tan x \right) \right) \\
 &= \frac{1}{1 + \left(\frac{3}{2} \tan x \right)^2} \times \frac{d}{dx} \left(\frac{3}{2} \tan x \right) \\
 &= \frac{1}{1 + \frac{9}{4} \tan^2 x} \times \frac{3}{2} \sec^2 x \\
 &= \frac{1}{1 + \frac{9 \sin^2 x}{4 \cos^2 x}} \times \frac{3}{2} \frac{1}{\cos^2 x} \\
 &= \frac{3}{2} \times \frac{1}{\cos^2 x + \frac{9}{4} \sin^2 x} \\
 &= \frac{6}{4 \cos^2 x + 9 \sin^2 x} \\
 &= \frac{6}{4 \sin^2 x + 5(\sin^2 x + \cos^2 x)} \\
 &= \frac{6}{4 \sin^2 x + 5}
 \end{aligned}$$

21b

$$\begin{aligned}
 A &= \int_0^7 \frac{6}{4 \sin^2 x + 5} dx \\
 &= \int_0^7 \frac{d}{dx} \left(\tan^{-1} \left(\frac{3}{2} \tan x \right) \right) dx \\
 &= \left[\tan^{-1} \left(\frac{3}{2} \tan x \right) \right]_0^7 \\
 &= \tan^{-1} \left(\frac{3}{2} \tan 7 \right) - \tan^{-1} \left(\frac{3}{2} \tan 0 \right) \\
 &\doteq 0.153 \text{ square unit}
 \end{aligned}$$

- 21c The integrand is well-defined in the interval $[0, 7]$, and lies between 14 and 19, so the area lies between 74 and 79, which is much larger than the answer of 0.153 that was calculated in part (b). The primitive, however, is undefined at two values within the interval, at $x = \frac{\pi}{2}$ and at $x = \frac{3\pi}{2}$, which renders the argument completely invalid.

Chapter 12 worked solutions – Further calculus

- 22a This is a GP with $a = 1$ and $r = -t^2$. Hence $T_k = (-t)^{k-1}$, solving for $T_k = t^{4n}$

$$t^{4n} = (-t^2)^{k-1}$$

$$t^{4n} = (-1)^{k-1}(t^2)^{k-1}$$

As $4n$ is even, $k - 1$ must also be even and hence $(-1)^{k-1} = 1$. Thus

$$t^{4n} = (t^2)^{k-1}$$

$$4n = 2k - 2$$

$$k = 2n + 1$$

Hence there are $2n + 1$ terms so the sum must be

$$S_{2n+1} = \frac{a(r^n - 1)}{r - 1} = \frac{(-t^2)^{2n+1} - 1}{-t^2 - 1}$$

Since $2n + 1$ is odd

$$S_{2n+1} = \frac{-t^{4n+2} - 1}{-t^2 - 1} = \frac{t^{4n+2} + 1}{t^2 + 1}$$

Hence it follows that for positive values of t

$$\frac{1}{t^2 + 1} < \frac{t^{4n+2} + 1}{t^2 + 1}$$

And thus

$$\frac{1}{t^2 + 1} < 1 - t^2 + t^4 - \dots + t^{4n}$$

- 22b As we are adding one extra term to the previous sum, this will be S_{2n+2}

Hence there are $2n + 2$ terms so the sum must be

$$S_{2n+2} = \frac{a(r^n - 1)}{r - 1} = \frac{(-t^2)^{2n+2} - 1}{-t^2 - 1}$$

Since $2n + 2$ is even

$$S_{2n+2} = \frac{t^{4n+4} - 1}{-t^2 - 1} = \frac{-t^{4n+4} + 1}{t^2 + 1}$$

Hence it follows that for positive values of t

From part a

Chapter 12 worked solutions – Further calculus

$$\frac{1}{t^2 + 1} < 1 - t^2 + t^4 - t^6 + \cdots + t^{4n}$$

$$\frac{1}{t^2 + 1} - t^{4n+2} < 1 - t^2 + t^4 - t^6 + \cdots + t^{4n} - t^{4n+2}$$

$$\frac{1}{t^2 + 1} - t^{4n+2} < \frac{-t^{4n+4} + 1}{t^2 + 1}$$

$$\frac{1}{t^2 + 1} < \frac{-t^{4n+4} + 1}{t^2 + 1} + t^{4n+2}$$

$$\frac{1 + t^{4n+4}}{t^2 + 1} < \frac{1}{t^2 + 1} + t^{4n+2}$$

$$1 - t^2 + t^4 - t^6 + \cdots + t^{4n} < \frac{1}{t^2 + 1} + t^{4n+2}$$

$$\frac{1}{t^2 + 1} < \frac{-t^{4n+4} + 1}{t^2 + 1}$$

$$\frac{1}{t^2 + 1} < \frac{-t^{4n+4} + 1}{t^2 + 1}$$

$$\frac{1}{t^2 + 1} < \frac{1 + t^{4n+2} + t^{4n+4}}{1 + t^2}$$

$$\frac{1}{t^2 + 1} < \frac{1}{1 + t^2} + \frac{t^{4n+2} + t^{4n+4}}{1 + t^2}$$

$$\frac{1}{t^2 + 1} < \frac{1}{1 + t^2} + \frac{t^{4n+2}(1 + t^2)}{1 + t^2}$$

$$\frac{1}{t^2 + 1} < \frac{1}{1 + t^2} + t^{4n+2}$$

Chapter 12 worked solutions – Further calculus

22c The inequality

$$\frac{1}{1+t^2} < 1 - t^2 + t^4 - t^6 + \cdots + t^{4n} < \frac{1}{1+t^2} + t^{4n+2}$$

Is true for all $0 < t < x$, hence

$$\int_0^x \frac{1}{1+t^2} dt < \int_0^x (1 - t^2 + t^4 - t^6 + \cdots + t^{4n}) dt < \int_0^x \left(\frac{1}{1+t^2} + t^{4n+2} \right) dt$$

$$[\tan^{-1} x]_0^x < \left[t - \frac{t^3}{3} + \frac{t^5}{5} - \frac{t^7}{7} + \cdots + \frac{t^{4n+1}}{4n+1} \right]_0^x < \left[\tan^{-1} x + \frac{t^{4n+3}}{4n+3} \right]_0^x$$

Thus

$$\tan^{-1} x < x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \cdots + \frac{x^{4n+1}}{4n+1} < \tan^{-1} x + \frac{x^{4n+3}}{4n+3}$$

22d

For $0 \leq x \leq 1$ we know that $0 \leq x^{4n+3} \leq 1$ and as $n \rightarrow \infty$, $\frac{1}{4n+3} \rightarrow 0$, hence it follows that as $n \rightarrow \infty$, $\frac{x^{4n+3}}{4n+3} \rightarrow 0$.

Thus as $n \rightarrow \infty$

$$\tan^{-1} x < x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \cdots < \tan^{-1} x + 0$$

$$\tan^{-1} x < x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \cdots < \tan^{-1} x$$

And so

$$\tan^{-1} x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \cdots$$

22e As $\tan x$ is odd, $\tan^{-1}(-x) = -\tan^{-1}(x)$

and hence for $-1 \leq x < 0$, (where $0 < -x \leq 1$)

$$\tan^{-1} x$$

$$= \tan^{-1}(-(-x))$$

$$= -\tan^{-1}(-x)$$

$$= -\left((-1) - \frac{(-1)^3}{3} + \frac{(-1)^5}{5} - \frac{(-1)^7}{7} + \cdots \right)$$

Chapter 12 worked solutions – Further calculus

$$= x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$$

- 22f Substituting $x = 1$ into the above equation

$$\tan^{-1} 1 = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$$

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots$$

22g

$$\frac{\pi}{4} = \frac{3-1}{3 \times 5} + \frac{7-5}{5 \times 7} + \frac{11-9}{9 \times 11} \dots$$

$$\frac{\pi}{4} = \frac{2}{3 \times 5} + \frac{2}{5 \times 7} + \frac{2}{9 \times 11} \dots$$

$$\frac{\pi}{8} = \frac{1}{3 \times 5} + \frac{1}{5 \times 7} + \frac{1}{9 \times 11} \dots$$

Using 10 terms gives

$\pi \doteq 3.092$, error $\doteq 0.050$

- 23a All rectangles have the same width of 1 unit and their height is given by $\tan^{-1} n$ for the n th rectangle. This means that

$$S_n = \tan^{-1} 1 + \tan^{-1} 2 + \dots + \tan^{-1} n$$

23b

$$\frac{d}{dx}(x \tan x) = \tan x + x \times \frac{1}{1+x^2}$$

Hence

$$\int \frac{d}{dx}(x \tan x) dx = \int \tan x + x \times \frac{1}{1+x^2} dx$$

$$x \tan x = \int \tan x dx + \int \frac{x}{1+x^2} dx$$

$$x \tan x = \int \tan x dx + \frac{1}{2} \ln(1+x^2) + C$$

$$\int \tan x dx = x \tan x - \frac{1}{2} \ln(1+x^2) + C$$

23c Firstly note that since

$$\int \tan x \, dx = x \tan x - \frac{1}{2} \ln(1 + x^2) + C$$

Substituting $x = u - 1$ into the equation gives

$$\begin{aligned} & \int \tan(u-1) \, du \\ &= (u-1) \tan(u-1) - \frac{1}{2} \ln(1 + (u-1)^2) + C \\ &= (u-1) \tan^{-1}(u-1) - \frac{1}{2} \ln(1 + u^2 - 2u + 1) + C \\ &= (u-1) \tan^{-1}(u-1) - \frac{1}{2} \ln(u^2 - 2u + 2) + C \end{aligned}$$

The area in the boxes is overestimated by the area under $\tan^{-1} x$ but underestimated by the area under $\tan^{-1}(x-1)$, hence

$$\begin{aligned} & \int_1^{n+1} \tan(x-1) \, dx < S_n < \int_0^{n+1} \tan(x) \, dx \\ & \int_1^{n+1} \tan(x-1) \, dx < S_n < \int_0^{n+1} \tan(x) \, dx \\ & \left[(x-1) \tan^{-1}(x-1) - \frac{1}{2} \ln(x^2 - 2x + 2) \right]_1^{n+1} < S_n \\ & < \left[(x) \tan^{-1}(x) - \frac{1}{2} \ln(1 + x^2) \right]_1^{n+1} \\ & (n) \tan^{-1}(n) - \frac{1}{2} \ln(1 + n^2) - (0) \tan^{-1}(0) - \frac{1}{2} \ln(1) < S_n \\ & < (n+1) \tan^{-1}(n+1) - \frac{1}{2} \ln(1 + (n+1)^2) - (1) \tan^{-1}(1) - \frac{1}{2} \ln(2) \\ & (n) \tan^{-1}(n) - \frac{1}{2} \ln(1 + n^2) < S_n \\ & < (n+1) \tan^{-1}(n+1) - \frac{1}{2} \ln(1 + (n+1)^2) - \frac{\pi}{4} - \frac{1}{2} \ln(2) \\ & (n) \tan^{-1}(n) - \frac{1}{2} \ln(1 + n^2) < S_n \\ & < (n+1) \tan^{-1}(n+1) - \frac{1}{2} \ln\left(\frac{n^2 + 2n + 2}{2}\right) - \frac{\pi}{4} \\ & n \tan^{-1} n - \frac{1}{2} \ln(n^2 + 1) < S_n < (n+1) \tan^{-1}(n+1) - \frac{1}{2} \ln\left(\frac{n^2}{2} + n + 1\right) - \frac{\pi}{4} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

23d Substituting $n = 1000$ into the above equation gives

$$\begin{aligned}1000 \tan^{-1} 1000 - \frac{1}{2} \ln(1000^2 + 1) &< S_{1000} \\&< (1000 + 1) \tan^{-1}(1000 + 1) - \frac{1}{2} \ln\left(\frac{1000^2}{2} + 1000 + 1\right) - \frac{\pi}{4}\end{aligned}$$

$$1562.89 < S_{100} < 1564.02$$

$$1563 < S_{1000} < 1565$$

Hence

$$1563 < \tan^{-1} 1 + \tan^{-1} 2 + \tan^{-1} 3 + \dots + \tan^{-1} 1000 < 1565$$

Chapter 12 worked solutions – Further calculus

Solutions to Exercise 12C

1a

$$\cos 2x = \cos^2 x - \sin^2 x$$

$$\cos 2x = 1 - \sin^2 x - \sin^2 x$$

$$\cos 2x = 1 - 2 \sin^2 x$$

$$2 \sin^2 x = 1 - \cos 2x$$

$$\sin^2 x = \frac{1}{2} - \frac{1}{2} \cos 2x$$

1b

$$\cos 4x = \cos 2(2x)$$

$$\cos 4x = \cos^2 2x - \sin^2 2x$$

$$\cos 4x = \cos^2 2x - (1 - \cos^2 2x)$$

$$\cos 4x = 2 \cos^2 2x - 1$$

$$2 \cos^2 2x = 1 + \cos 4x$$

$$\cos^2 2x = \frac{1}{2} + \frac{1}{2} \cos 4x$$

1c

$$\frac{1}{2} \sin 6x = \frac{1}{2} \sin 2(3x)$$

$$= \frac{1}{2}(2 \sin 3x \cos 3x)$$

$$= \sin 3x \cos 3x$$

1d

$$\cos x = \cos 2\left(\frac{x}{2}\right) = \cos^2\left(\frac{x}{2}\right) - \sin^2\left(\frac{x}{2}\right)$$

$$\cos x = 1 - \sin^2\left(\frac{x}{2}\right) - \sin^2\left(\frac{x}{2}\right)$$

$$\cos x = 1 - 2 \sin^2\left(\frac{x}{2}\right)$$

Chapter 12 worked solutions – Further calculus

$$2 \sin^2 \left(\frac{x}{2} \right) = 1 - \cos x$$

2a $\cos^2 15^\circ$

$$\begin{aligned} &= \frac{1}{2} + \frac{1}{2} \cos 30^\circ \\ &= \frac{1}{2} + \frac{1}{2} \times \frac{\sqrt{3}}{2} \\ &= \frac{1}{4}(2 + \sqrt{3}) \end{aligned}$$

2b

$$\begin{aligned} &\sin^2 \frac{5\pi}{12} \\ &= \frac{1}{2} - \frac{1}{2} \cos \frac{5\pi}{6} \\ &= \frac{1}{2} + \frac{1}{2} \times \frac{\sqrt{3}}{2} \\ &= \frac{1}{4}(2 + \sqrt{3}) \end{aligned}$$

2c $\sin 105^\circ \cos 105^\circ$

$$\begin{aligned} &= \frac{1}{2} \sin 210^\circ \\ &= \frac{1}{2} \times \left(-\frac{1}{2} \right) \\ &= -\frac{1}{4} \end{aligned}$$

2d

$$\begin{aligned} &\sin^2 \frac{7\pi}{8} \\ &= \frac{1}{2} - \frac{1}{2} \cos \frac{7\pi}{4} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{1}{2} - \frac{1}{2} \times \left(\frac{1}{\sqrt{2}} \right)$$

$$= \frac{2 - \sqrt{2}}{4}$$

3 $\sin^2 \theta = \frac{1}{2} - \frac{1}{2} \cos 2\theta$ (see 1a)

3a

$$\begin{aligned} & \int \sin^2 x \, dx \\ &= \int \left(\frac{1}{2} - \frac{1}{2} \cos 2x \right) dx \\ &= \frac{x}{2} - \frac{\sin 2x}{4} + C \end{aligned}$$

3b

$$\begin{aligned} & \int \sin^2 2x \, dx \\ &= \int \left(\frac{1}{2} - \frac{1}{2} \cos 4x \right) dx \\ &= \frac{x}{2} - \frac{\sin 4x}{8} + C \end{aligned}$$

3c

$$\begin{aligned} & \int \sin^2 \frac{1}{4}x \, dx \\ &= \int \left(\frac{1}{2} - \frac{1}{2} \cos \frac{1}{2}x \right) dx \\ &= \frac{x}{2} - \frac{\sin \frac{1}{2}x}{2 \left(\frac{1}{2} \right)} + C \\ &= \frac{x}{2} - \sin \frac{1}{2}x + C \end{aligned}$$

Chapter 12 worked solutions – Further calculus

3d

$$\begin{aligned} & \int \sin^2 3x \, dx \\ &= \int \left(\frac{1}{2} - \frac{1}{2} \cos 6x \right) dx \\ &= \frac{x}{2} - \frac{\sin 6x}{12} + C \end{aligned}$$

4 $\cos^2 \theta = \frac{1}{2} + \frac{1}{2} \cos 2\theta$ (see 1b)

4a

$$\int \cos^2 x \, dx = \int \left(\frac{1}{2} + \frac{1}{2} \cos 2x \right) dx = \frac{x}{2} + \frac{\sin 2x}{4} + C$$

4b

$$\int \cos^2 6x \, dx = \int \left(\frac{1}{2} + \frac{1}{2} \cos 12x \right) dx = \frac{x}{2} + \frac{\sin 12x}{24} + C$$

4c

$$\int \cos^2 \frac{1}{2}x \, dx = \int \left(\frac{1}{2} + \frac{1}{2} \cos x \right) dx = \frac{x}{2} + \frac{\sin x}{2} + C$$

4d

$$\int \cos^2 10x \, dx = \int \left(\frac{1}{2} + \frac{1}{2} \cos 20x \right) dx = \frac{x}{2} + \frac{\sin 20x}{40} + C$$

5a

$$\begin{aligned} & \int_0^\pi \sin^2 x \, dx \\ &= \int_0^\pi \left(\frac{1}{2} - \frac{1}{2} \cos 2x \right) dx \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \left[\frac{x}{2} - \frac{\sin 2x}{4} \right]_0^\pi$$

$$= \frac{\pi}{2}$$

5b

$$\begin{aligned} & \int_0^{\frac{\pi}{4}} \cos^2 x \, dx \\ &= \int_0^{\frac{\pi}{4}} \left(\frac{1}{2} + \frac{1}{2} \cos 2x \right) dx \\ &= \left[\frac{x}{2} + \frac{\sin 2x}{4} \right]_0^{\frac{\pi}{4}} \\ &= \frac{1}{8}(\pi + 2) \end{aligned}$$

5c

$$\begin{aligned} & \int_0^{\frac{\pi}{6}} \sin^2 \frac{1}{2}x \, dx \\ &= \int_0^{\frac{\pi}{6}} \left(\frac{1}{2} - \frac{1}{2} \cos x \right) dx \\ &= \left[\frac{x}{2} - \frac{\sin x}{2} \right]_0^{\frac{\pi}{6}} \\ &= \frac{1}{12}(\pi - 3) \end{aligned}$$

5d

$$\begin{aligned} & \int_0^{\frac{\pi}{16}} \cos^2 2x \, dx \\ &= \int_0^{\frac{\pi}{16}} \left(\frac{1}{2} + \frac{1}{2} \cos 4x \right) dx \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \left[\frac{x}{2} + \frac{\sin 4x}{8} \right]_0^{\frac{\pi}{6}}$$

$$= \frac{1}{32}(\pi + 2\sqrt{2})$$

5e

$$\begin{aligned} & \int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} \cos^2 \left(x + \frac{\pi}{12} \right) dx \\ &= \int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} \frac{1}{2} + \frac{1}{2} \cos 2 \left(x + \frac{\pi}{12} \right) dx \\ &= \left[\frac{x}{2} + \frac{\sin 2 \left(x + \frac{\pi}{12} \right)}{4} \right]_{-\frac{\pi}{6}}^{\frac{\pi}{6}} \\ &= \frac{1}{24}(4\pi + 9) \end{aligned}$$

5f

$$\begin{aligned} & \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \sin^2 \left(x - \frac{\pi}{6} \right) dx \\ &= \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \left(\frac{1}{2} - \frac{1}{2} \cos 2 \left(x - \frac{\pi}{6} \right) \right) dx \\ &= \left[\frac{x}{2} - \frac{\sin 2 \left(x - \frac{\pi}{6} \right)}{4} \right]_{\frac{\pi}{6}}^{\frac{\pi}{3}} \\ &= \frac{1}{24}(2\pi - 3\sqrt{3}) \end{aligned}$$

Chapter 12 worked solutions – Further calculus

6a

$$\begin{aligned}
 & \int \sin 3x \cos 2x \, dx \\
 &= \int \left(\frac{1}{2} \sin(2x + 3x) + \frac{1}{2} \sin(3x - 2x) \right) dx \\
 &= \int \left(\frac{1}{2} \sin 5x + \frac{1}{2} \sin x \right) dx \\
 &= -\frac{1}{10} \cos 5x - \frac{1}{2} \cos x + C
 \end{aligned}$$

6b

$$\begin{aligned}
 & \int \cos 3x \sin x \, dx \\
 &= \int \left(\frac{1}{2} \sin 2x - \frac{1}{2} \sin 4x \right) dx \\
 &= \frac{1}{4} \cos 2x - \frac{1}{8} \cos 4x + C
 \end{aligned}$$

6c

$$\begin{aligned}
 & \int_0^{\frac{\pi}{4}} 2 \cos 2x \cos x \, dx \\
 &= \int_0^{\frac{\pi}{4}} 2 \times \frac{1}{2} (\cos x + \cos 3x) \, dx \\
 &= \int_0^{\frac{\pi}{4}} (\cos x + \cos 3x) \, dx \\
 &= \left[\sin x + \frac{1}{3} \sin 3x \right]_0^{\frac{\pi}{4}} \\
 &= \frac{2\sqrt{2}}{3}
 \end{aligned}$$


6d

$$\begin{aligned}
 & \int_0^{\frac{\pi}{3}} \sin 5x \sin 2x \, dx \\
 &= \int_0^{\frac{\pi}{3}} \frac{1}{2} (\sin 3x - \sin 7x) \, dx
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus


$$\begin{aligned}
 &= \frac{1}{2} \left[-\frac{1}{3} \cos 3x + \frac{1}{7} \cos 7x \right]_0^{\frac{\pi}{3}} \\
 &= \frac{\sqrt{3}}{28}
 \end{aligned}$$

7a



7b

$y = \frac{1}{2}(1 + \cos 2x)$ is shown in blue and $y = \frac{1}{2}(1 - \cos 2x)$ is shown in green.



7c

By observation firstly note that $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$ and that $\cos^2 x = \frac{1}{2}(1 + \cos 2x)$. By observation of the graph, we see that these two graphs always add to 1 and hence $\sin^2 x + \cos^2 x = 1$.

Chapter 12 worked solutions – Further calculus

8a Let $u = \sin x, du = \cos x dx$

$$\int \sin^3 x \cos x dx$$

$$= \int u^3 du$$
$$= \frac{u^4}{4} + C$$

$$= \frac{\sin^4 x}{4} + C$$

8b Let $u = \sin x, du = \cos x dx$

$$\int \sin^6 x \cos x dx$$

$$= \int u^6 du$$

$$= \frac{u^7}{7} + C$$

$$= \frac{\sin^7 x}{7} + C$$

8c Let $u = \cos x, du = -\sin x dx$

$$\int \cos^5 x \sin x dx$$

$$= - \int u^5 du$$

$$= -\frac{u^6}{6} + C$$

$$= -\frac{\cos^6 x}{6} + C$$

8d Let $u = \cos x, du = -\sin x$

$$\int \cos^8 x \sin x dx$$

$$= \int -u^8 du$$

Chapter 12 worked solutions – Further calculus

$$= -\frac{u^9}{9} + C$$

$$= -\frac{\cos^9 x}{9} + C$$

8e Let $u = e^x, du = e^x dx$

$$\int e^x \sin e^x dx$$

$$= \int \sin u du$$

$$= -\cos u + C$$

$$= -\cos e^x + C$$

8f Let $u = e^x, du = e^x dx$

$$\int e^x \cos 5e^x dx$$

$$= \int \cos 5u du$$

$$= \frac{1}{5} \sin 5u + C$$

$$= \frac{1}{5} \sin 5e^x + C$$

8g Let $u = \cos x, du = -\sin x$

$$\int \tan x dx$$

$$= \int \frac{\sin x}{\cos x} dx$$

$$= \int -\frac{1}{u} du$$

$$= -\ln|u| + C$$

$$= -\ln |\cos x| + c$$

Chapter 12 worked solutions – Further calculus

8h Let $u = \sin 7x, du = 7 \cos x$

$$\begin{aligned} & \int \cot 7x \, dx \\ &= \int \frac{\cos 7x}{\sin 7x} \, dx \\ &= \int \frac{1}{7u} \, du \\ &= \frac{1}{7} \ln|u| + C \\ &= \frac{1}{7} \ln |\sin 7x| + C \end{aligned}$$

9a

$$y = \cos x \sin x = \frac{1}{2} \sin 2x$$

Hence the range is

$$-\frac{1}{2} \leq y \leq \frac{1}{2}$$

9b i

$$\begin{aligned} & \int \cos x \sin x \, dx \\ &= \int \frac{1}{2} \sin 2x \, dx \\ &= -\frac{1}{4} \cos 2x + C \end{aligned}$$

9b ii

Method 1:

Let $u = \sin x, du = \cos x \, dx$

$$\begin{aligned} & \int \sin x \cos x \, dx \\ &= \int u \, du \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \frac{u^2}{2} + C \\
 &= \frac{\sin^2 x}{2} + C
 \end{aligned}$$

Method 2:

Let $u = \cos x$, $du = -\sin x dx$

$$\begin{aligned}
 &\int \sin x \cos x dx \\
 &= \int -u du \\
 &= -\frac{u^2}{2} + C \\
 &= -\frac{\cos^2 x}{2} + C
 \end{aligned}$$

- 9c Note that $\cos 2x = \cos^2 x - \sin^2 x = 2\cos^2 x - 1 = 1 - 2\sin^2 x$. Thus

$$\begin{aligned}
 &-\frac{1}{4}\cos 2x + c \\
 &= -\frac{1}{4}(2\cos^2 x - 1) \\
 &= -\frac{\cos^2 x}{2} + c_2 \\
 &= \frac{\sin^2 x}{2} + c_3
 \end{aligned}$$

Note at c, c_1 and c_2 are not the same constant.

- 10a $\sin^4 x = (\sin^2 x)^2$

$$\begin{aligned}
 &= \left(\frac{1}{2} - \frac{1}{2}\cos 2x\right)^2 \\
 &= \frac{1}{4} - \frac{1}{2}\cos 2x + \frac{1}{4}\cos^2 2x \\
 &= \frac{1}{4} - \frac{1}{2}\cos 2x + \frac{1}{4}\left(\frac{1}{2} + \cos 4x\right)
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{3}{8} - \frac{1}{2} \cos 2x + \frac{1}{8} \cos 4x$$

10b $\cos^4 x = (\cos^2 x)^2$

$$\begin{aligned} &= \left(\frac{1}{2} + \frac{1}{2} \cos 2x \right)^2 \\ &= \frac{1}{4} + \frac{1}{2} \cos 2x + \frac{1}{4} \cos^2 2x \\ &= \frac{1}{4} + \frac{1}{2} \cos 2x + \frac{1}{4} \left(\frac{1}{2} + \cos 4x \right) \\ &= \frac{3}{8} + \frac{1}{2} \cos 2x + \frac{1}{8} \cos 4x \end{aligned}$$

10c i

$$\begin{aligned} &\int_0^\pi \sin^4 x \, dx \\ &= \int_0^\pi \left(\frac{3}{8} - \frac{1}{2} \cos 2x + \frac{1}{8} \cos 4x \right) dx \\ &= \left[\frac{3}{8}x - \frac{1}{4} \sin 2x + \frac{1}{2} \sin 4x \right]_0^\pi \\ &= \frac{3\pi}{8} \end{aligned}$$

10c ii

$$\begin{aligned} &\int_0^{\frac{\pi}{4}} \cos^4 x \, dx \\ &= \int_0^{\frac{\pi}{4}} \left(\frac{3}{8} + \frac{1}{2} \cos 2x + \frac{1}{8} \cos 4x \right) dx \\ &= \left[\frac{3}{8}x + \frac{1}{4} \sin 2x + \frac{1}{2} \sin 4x \right]_0^{\frac{\pi}{4}} \\ &= \frac{1}{32}(3\pi + 8) \end{aligned}$$

Chapter 12 worked solutions – Further calculus

10d

$$\begin{aligned}
 & \int_0^{\frac{\pi}{3}} \sin^3 x \, dx \\
 &= \int_0^{\frac{\pi}{3}} \sin x (\sin^2 x) \, dx \\
 &= \int_0^{\frac{\pi}{3}} \sin x (1 - \cos^2 x) \, dx \\
 &= \int_0^{\frac{\pi}{3}} (\sin x - \sin x \cos^2 x) \, dx \\
 &= \int_0^{\frac{\pi}{3}} \sin x \, dx - \int_0^{\frac{\pi}{3}} \sin x \cos^2 x \, dx
 \end{aligned}$$

Now for $\int_0^{\frac{\pi}{3}} \sin x \cos^2 x \, dx$, let $u = \cos x$, $du = -\sin x$

$$\int_0^{\frac{\pi}{3}} \sin x \cos^2 x \, dx = \int_1^{\frac{1}{2}} u^2 \, du = \left[\frac{u^3}{3} \right]_1^{\frac{1}{2}} = \frac{1}{24}$$

Hence

$$\int_0^{\frac{\pi}{3}} \sin^3 x \, dx = [-\cos x]_0^{\frac{\pi}{3}} - \left[\frac{u^3}{3} \right]_1^{\frac{1}{2}} = \frac{5}{24}$$

11a

$$\begin{aligned}
 & \int \tan^2 2x \, dx \\
 &= \int (\sec^2 2x - 1) \, dx \\
 &= \frac{1}{2} \tan 2x - x + C
 \end{aligned}$$

11b

$$\int \cot^2 \frac{1}{2}x \, dx = \int (\operatorname{cosec}^2(\frac{1}{2}x) - 1) \, dx$$

Chapter 12 worked solutions – Further calculus

$$= -2 \cot \frac{1}{2}x - x + C$$

11c

$$\begin{aligned} & \int_{\frac{\pi}{12}}^{\frac{\pi}{9}} 3 \tan^2 3x \, dx \\ &= \int_{\frac{\pi}{12}}^{\frac{\pi}{9}} 3(\sec^2 3x + 1) \, dx \\ &= \int_{\frac{\pi}{12}}^{\frac{\pi}{9}} (3 \sec^2 3x + 3) \, dx \\ &= [\tan 3x + 3x]_{\frac{\pi}{12}}^{\frac{\pi}{9}} \\ &= \sqrt{3} - 1 - \frac{\pi}{12} \end{aligned}$$

11d

$$\begin{aligned} & \int_{\frac{\pi}{24}}^{\frac{\pi}{8}} \cot^2 4x \, dx \\ &= \int_{\frac{\pi}{24}}^{\frac{\pi}{8}} (\operatorname{cosec}^2 4x + 1) \, dx \\ &= \left[-\frac{1}{4} \cot \frac{1}{2}x + x \right]_{\frac{\pi}{24}}^{\frac{\pi}{8}} \\ &= \frac{1}{4} \sqrt{3} - \frac{\pi}{12} \end{aligned}$$

12a Let $u = \tan x, du = \sec^2 x \, dx$

$$\begin{aligned} & \int \tan x \sec^2 x \, dx \\ &= \int u \, du \\ &= \frac{u^2}{2} + C \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{\tan^2 x}{2} + C$$

12b

$$\begin{aligned} & \int \frac{\sin^2 x}{1 + \cos x} dx \\ &= \int \frac{1 - \cos^2 x}{1 + \cos x} dx \\ &= \int \frac{(1 - \cos x)(1 + \cos x)}{1 + \cos x} dx \\ &= \int (1 - \cos x) dx \\ &= x - \sin x + C \end{aligned}$$

12c

$$\begin{aligned} & \int \frac{1 + \cos^3 x}{\cos^2 x} dx \\ &= \int \left(\frac{1}{\cos^2 x} + \cos x \right) dx \\ &= \int (\sec^2 x + \cos x) dx \\ &= \tan x + \sin x + C \end{aligned}$$

13a

$$\begin{aligned} F(x) &= \int_0^x \sin^2 t dt \\ &= \int_0^x \left(\frac{1}{2} - \frac{1}{2} \cos 2t \right) dx \\ &= \left[\frac{1}{2}t - \frac{1}{4} \sin 2t \right]_0^x \\ &= \frac{1}{2}x - \frac{1}{4} \sin 2x - (0 - 0) \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{1}{2}x - \frac{1}{4}\sin 2x$$

13b

$$\begin{aligned} F'(x) &= \frac{1}{2} - \frac{1}{4} \times 2\cos 2x \\ &= \frac{1}{2} - \frac{1}{2}\cos 2x \\ &= \sin^2 x \end{aligned}$$

13b i $F(x)$ is stationary when $F'(x) = 0$ this is when $\sin^2 x = 0$, $\sin x = 0$ and hence when $x = 0, \pi, 2\pi$.

13b ii $F(x)$ is increasing when $F'(x) > 0$ this is when $\sin^2 x > 0$. This is all x such that $\sin x \neq 0$ and hence when $x \neq 0, \pi, 2\pi$.

13b iii The curve is decreasing when $F'(x) < 0$. However, $\sin^2 x \geq 0$ for all x . Hence there are no values of x for which the curve is decreasing.


13c

Since $-1 \leq \sin 2x \leq 1$, $-\frac{1}{4} \leq \frac{1}{4}\sin 2x \leq \frac{1}{4}$ and thus $\frac{1}{2}x - \frac{1}{4} \leq \frac{1}{2}x + \frac{1}{4}\sin 2x \leq \frac{1}{2}x + \frac{1}{4}$, thus $\frac{1}{2}x - \frac{1}{4} \leq F(x) \leq \frac{1}{2}x + \frac{1}{4}$. Thus $F(x)$ never differs from $\frac{1}{2}x$ by more than $\frac{1}{4}$.

13d We know that the stationary points are $x = 0, \pi, 2\pi$. As the curve is always increasing, these will all be points of inflection (if there is a maximum or minimum the slope would have to be decreasing on one side).

Chapter 12 worked solutions – Further calculus

13e



13f i

$$\int_0^k \sin^2 x \, dx = \frac{3\pi}{2}$$

$$\int_0^k \left(\frac{1}{2} - \frac{1}{2} \cos 2x \right) dx = \frac{3\pi}{2}$$

$$\left[\frac{x}{2} - \frac{\sin 2x}{4} \right]_0^k = \frac{3\pi}{2}$$

$$\frac{k}{2} - \frac{\sin 2k}{4} = \frac{3\pi}{2}$$

$$\frac{k}{2} = \frac{3\pi}{2}$$

$$k = 3\pi$$

13f ii

$$\int_0^k \sin^2 x \, dx = \frac{n\pi}{2}$$

$$\int_0^k \left(\frac{1}{2} - \frac{1}{2} \cos 2x \right) dx = \frac{n\pi}{2}$$

$$\left[\frac{x}{2} - \frac{\sin 2x}{4} \right]_0^k = \frac{n\pi}{2}$$

$$\frac{k}{2} - \frac{\sin 2k}{4} = \frac{n\pi}{2}$$

$$\frac{k}{2} = \frac{n\pi}{2}$$

Chapter 12 worked solutions – Further calculus

$$k = n\pi$$

14

$$\begin{aligned} & \lim_{R \rightarrow \infty} \left(\frac{1}{R} \int_0^R \sin^2 t \, dt \right) \\ &= \lim_{R \rightarrow \infty} \left(\frac{1}{R} \left(\frac{R}{2} - \frac{\sin 2R}{4} \right) \right) \\ &= \lim_{R \rightarrow \infty} \left(\frac{1}{2} - \frac{1}{4} \left(\frac{\sin 2R}{R} \right) \right) \end{aligned}$$

Now as $-1 \leq \sin 2R \leq 0$, then as R gets large $\frac{\sin 2R}{R} \rightarrow 0$, hence

$$\lim_{R \rightarrow \infty} \left(\frac{1}{R} \int_0^R \sin^2 t \, dt \right) = \frac{1}{2} - \frac{1}{4}(0) = \frac{1}{2}$$

Chapter 12 worked solutions – Further calculus

Solutions to Exercise 12D

1a

$$\begin{aligned} du &= \frac{d}{dx}(1 + x^2) \\ &= \frac{d}{dx}(1) + \frac{d}{dx}(x^2) \\ &= 0 + 2x \\ &= 2x \end{aligned}$$

1b

$$\begin{aligned} &\int 2x(1 + x^2)^3 dx \\ &= \int (1 + x^2)^3 (2x dx) \\ &= \int u^3 du \quad \text{where } u = 1 + x^2 \\ &= \frac{u^4}{4} + C \\ &= \frac{(1 + x^2)^4}{4} + C \end{aligned}$$

1c

$$\frac{(1 + x^2)^4}{4} + C$$

1d

$$\begin{aligned} &\frac{d}{dx}\left(\frac{(1 + x^2)^4}{4}\right) \\ &= \frac{4(2x)(1 + x^2)^3}{4} \\ &= 2x(1 + x^2)^3 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

2a Let $u = 2x + 3, du = 2 dx$

$$\int 2(2x+3)^3 dx$$

$$= \int u^3 du$$

$$= \frac{u^4}{4} + C$$

$$= \frac{(2x+3)^4}{4} + C$$

2b Let $u = 1 + x^3, du = 3x^2 dx$

$$\int 3x^2(1+x^3)^4 dx$$

$$= \int u^4 du$$

$$= \frac{u^5}{5} + C$$

$$= \frac{(1+x^3)^5}{5} + C$$

2c Let $u = 1 + x^2, du = 2x dx$

$$\int \frac{2x}{(1+x^2)^2} dx$$

$$= \int \frac{1}{u^2} du$$

$$= \int u^{-2} du$$

$$= -u^{-1} + C$$

$$= -\frac{1}{1+x^2} + C$$

Chapter 12 worked solutions – Further calculus

2d Let $u = 3x - 5, du = 3 dx$

$$\begin{aligned} & \int \frac{3}{\sqrt{3x-5}} dx \\ &= \int \frac{1}{\sqrt{u}} du \\ &= \int u^{-\frac{1}{2}} du \\ &= 2u^{\frac{1}{2}} + C \\ &= 2\sqrt{3x-5} + C \end{aligned}$$

2e Let $u = \sin x, du = \cos x dx$

$$\begin{aligned} & \int \sin^3 x \cos x dx \\ &= \int u^3 du \\ &= \frac{u^4}{4} + C \\ &= \frac{\sin^4 x}{4} + C \end{aligned}$$

2f Let $u = 1 + x^4, du = 4x^3 dx$

$$\begin{aligned} & \int \frac{4x^3}{1+x^4} dx \\ &= \int \frac{1}{u} du \\ &= \ln|u| + C \\ &= \ln(1+x^4) + C \quad (\text{as } 1+x^4 \text{ is positive}) \end{aligned}$$

3a Let $u = 1 - x^2, du = -2x dx$ hence $x dx = -\frac{1}{2}du$

3b

$$\begin{aligned} & \int \frac{x}{\sqrt{1-x^2}} dx \\ &= \int \frac{1}{\sqrt{u}} \times -\frac{1}{2} du \\ &= -\frac{1}{2} \int u^{-\frac{1}{2}} du \end{aligned}$$

3c

$$\begin{aligned} & \int \frac{x}{\sqrt{1-x^2}} dx \\ &= \int \frac{1}{\sqrt{u}} \times -\frac{1}{2} du \\ &= -\frac{1}{2} \int u^{-\frac{1}{2}} du \\ &= -u^{\frac{1}{2}} + C \\ &= -\sqrt{1-x^2} + C \end{aligned}$$

4a Let $u = x^4 + 1, du = 4x^3 dx$

$$\begin{aligned} & \int x^3(x^4 + 1)^5 dx \\ &= \int \frac{1}{4} u^5 du \\ &= \frac{u^6}{24} + C \\ &= \frac{(x^4 + 1)^6}{24} + C \end{aligned}$$

Chapter 12 worked solutions – Further calculus

4b Let $u = x^3 - 1$, $du = 3x^2 dx$

$$\int x^2 \sqrt{x^3 - 1} dx$$

$$= \int \frac{1}{3} \sqrt{u} du$$

$$= \frac{1}{3} \times \frac{2}{3} u^{\frac{3}{2}} + C$$

$$= \frac{2}{9} u^{\frac{3}{2}} + C$$

$$= \frac{2}{9} (x^3 - 1)^{\frac{3}{2}} + C$$

4c Let $u = x^3$, $du = 3x^2 dx$

$$\int x^2 e^{x^3} dx$$

$$= \int \frac{1}{3} e^u du$$

$$= \frac{e^u}{3} + C$$

$$= \frac{e^{x^3}}{3} + C$$

4d Let $u = 1 + \sqrt{x}$, hence $du = \frac{1}{2} x^{-\frac{1}{2}} dx$, hence $2 du = \frac{1}{\sqrt{x}} dx$

$$\int \frac{1}{\sqrt{x}(1 + \sqrt{x})^3} dx$$

$$= \int \frac{2}{u^3} du$$

$$= -\frac{1}{u^2} + C$$

$$= -\frac{1}{(1 + \sqrt{x})^2} + C$$

Chapter 12 worked solutions – Further calculus

4e Let $u = \tan 2x, du = 2 \sec^2 2x dx$

$$\int \tan^2 2x \sec^2 2x dx$$

$$= \int \frac{u^2}{2} du$$

$$= \frac{u^3}{6} + C$$

$$= \frac{\tan^3 2x}{6} + C$$

4f Let $u = \frac{1}{x}, du = -\frac{1}{x^2} dx$

$$\int \frac{e^{\frac{1}{x}}}{x^2} dx$$

$$= \int -e^u du$$

$$= -e^u + C$$

$$= -e^{\frac{1}{x}} + C$$

5a Let $u = 2 + x^3, du = 3x^2 dx$

When $x = 0, u = 2$

When $x = 1, u = 3$

$$\int_0^1 x^2(2 + x^3)^3 dx$$

$$= \int_2^3 \frac{1}{3} u^3 du$$

$$= \left[\frac{u^9}{9} \right]_2^3$$

$$= \frac{1}{9}(3^9 - 2^9)$$

Chapter 12 worked solutions – Further calculus

$$= \frac{65}{12}$$

5b When $u = 1 + x^4$, $du = 4x^3$

When $x = 0, u = 1$

When $x = 1, u = 2$

$$\int_0^1 \frac{2x^3}{\sqrt{1+x^4}} dx$$

$$= \frac{1}{2} \int_0^1 \frac{4x^3}{\sqrt{1+x^4}} dx$$

$$= \frac{1}{2} \int_1^2 \frac{du}{\sqrt{u}}$$

$$= \frac{1}{2} \left[2u^{\frac{1}{2}} \right]_1$$

$$= [\sqrt{u}]_1$$

$$= \sqrt{2} - 1$$

5c Let $u = \cos x, du = -\sin x$

When $x = 0, u = 1$

When $x = \frac{\pi}{2}, u = 0$

$$\int_0^{\frac{\pi}{2}} \cos^2 x \sin x dx$$

$$= \int_1^0 -u^2 du$$

$$= \int_0^1 u^2 du$$

$$= \left[\frac{u^3}{3} \right]_0^1$$

$$= \frac{1}{3}$$

Chapter 12 worked solutions – Further calculus

5d Let $u = 1 - x^2, du = 2x \, dx$

$$\text{When } x = \frac{1}{2}\sqrt{3}, u = \frac{1}{4}$$

$$\text{When } x = 1, u = 0$$

$$\int_{\frac{1}{2}\sqrt{3}}^1 x\sqrt{1-x^2} \, dx$$

$$= \int_0^{\frac{1}{4}} \frac{1}{2} \sqrt{u} \, du$$

$$= \frac{1}{2} \left[\frac{2}{3} u^{\frac{3}{2}} \right]_0^{\frac{1}{4}}$$

$$= \frac{1}{3} \left[u^{\frac{3}{2}} \right]_0^{\frac{1}{4}}$$

$$= \frac{1}{3} \left(\frac{1}{4} \right)^{\frac{3}{2}}$$

$$= \frac{1}{24}$$

5e Let $u = \ln x, du = \frac{1}{x} dx$

$$\text{When } x = 1, u = \ln 1 = 0$$

$$\text{When } x = e^2, u = \ln e^2 = 2 \ln e = 2$$

$$\int_1^{e^2} \frac{\ln x}{x} dx$$

$$= \int_0^2 u \, du$$

$$= \left[\frac{u^2}{2} \right]_0^2$$

$$= 2$$

Chapter 12 worked solutions – Further calculus

5f Let $u = \sqrt{x}$, $du = \frac{1}{2}x^{-\frac{1}{2}}dx = \frac{1}{2\sqrt{x}} dx$

$$\int_0^4 \frac{e^{\sqrt{x}}}{4\sqrt{x}} dx$$

$$= \int_0^2 \frac{e^u}{2} du$$

$$= \frac{1}{2} [e^u]_0^2$$

$$= \frac{1}{2}(e^2 - 1)$$

5g Let $u = \sin 2x$, $du = 2 \cos 2x dx$

$$\int_0^{\frac{\pi}{4}} \sin^4 2x \cos 2x dx$$

$$= \int_0^1 \frac{u^4}{2} du$$

$$= \frac{1}{2} \left[\frac{u^5}{5} \right]_0^1$$

$$= \frac{1}{10}$$

5h Let $u = \sin^{-1} x$, $du = \frac{1}{\sqrt{1-x^2}} dx$

$$\int_0^1 \frac{(\sin^{-1} x)^3}{\sqrt{1-x^2}} dx$$

$$= \int_0^{\frac{\pi}{2}} u^3 du$$

$$= \left[\frac{u^4}{4} \right]_0^{\frac{\pi}{2}}$$

$$= \frac{\left(\frac{\pi}{2} \right)^4}{4}$$

$$= \frac{\pi^4}{64}$$

Chapter 12 worked solutions – Further calculus


5i Let $u = x^2 + 2x$, $du = 2x + 2 dx$

$$\begin{aligned} & \int_0^2 \frac{x+1}{\sqrt[3]{x^2+2x}} dx \\ &= \frac{1}{2} \int_0^2 \frac{2x+2}{\sqrt[3]{x^2+2x}} dx \\ &= \frac{1}{2} \int_0^8 \frac{1}{\sqrt[3]{u}} du \\ &= \frac{1}{2} \int_0^8 u^{-\frac{1}{3}} du \\ &= \frac{1}{2} \left[\frac{3}{2} u^{\frac{2}{3}} \right]_0^8 \\ &= \frac{3}{4} \left(8^{\frac{2}{3}} \right) \\ &= 3 \end{aligned}$$

5j Let $u = \tan x$, $du = \sec^2 x$

$$\begin{aligned} & \int_{\frac{\pi}{4}}^{\frac{\pi}{3}} \frac{\sec^2 x}{\tan x} dx \\ &= \int_1^{\sqrt{3}} \frac{1}{u} du \\ &= [\ln|u|]_1^{\sqrt{3}} \\ &= \ln \sqrt{3} - \ln 1 \\ &= \frac{1}{2} \ln 3 \end{aligned}$$

6 Note that the area to be found is shown in green in the diagram below



Chapter 12 worked solutions – Further calculus

Let $u = x^3$, $du = 3x^2 dx$. The area bounded will be given by

$$\begin{aligned} A &= \int_0^1 \frac{x^2}{1+x^6} dx \\ &= \frac{1}{3} \int_0^1 \frac{3x^2}{1+x^6} dx \\ &= \frac{1}{3} \int_0^1 \frac{1}{1+u^2} du \\ &= \frac{1}{3} [\tan^{-1} u]_0^1 \\ &= \frac{\pi}{12} \text{ square units} \end{aligned}$$

- 7 For all following subsections, let $u = \sin x$, $du = \cos x dx$

7a

$$\begin{aligned} &\int_0^{\frac{\pi}{6}} \frac{\cos x}{1+\sin x} dx \\ &= \int_0^{\frac{1}{2}} \frac{1}{1+u} du \\ &= [\ln|1+u|]_0^{\frac{1}{2}} \\ &= \ln\frac{3}{2} \end{aligned}$$

7b

$$\begin{aligned} &\int_0^{\frac{\pi}{2}} \frac{\cos x}{1+\sin^2 x} dx \\ &= \int_0^1 \frac{1}{1+u^2} du \\ &= [\tan|u|]_0^1 \\ &= \frac{\pi}{4} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

7c

$$\begin{aligned}
 & \int_0^{\frac{\pi}{2}} \cos^3 x \, dx \\
 &= \int_0^{\frac{\pi}{2}} \cos x \cos^2 x \, dx \\
 &= \int_0^{\frac{\pi}{2}} \cos x (1 - \sin^2 x) \, dx \\
 &= \int_0^{\frac{\pi}{2}} \cos x \, dx - \int_0^{\frac{\pi}{2}} \cos x \sin^2 x \, dx \\
 &= [\sin x]_0^{\frac{\pi}{2}} - \int_0^1 u^2 \, du \\
 &= 1 - \left[\frac{u^3}{3} \right]_0^1 \\
 &= 1 - \frac{1}{3} \\
 &= \frac{2}{3}
 \end{aligned}$$

7d

$$\begin{aligned}
 & \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \frac{\cos^3 x}{\sin^4 x} \, dx \\
 &= \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \frac{\cos x (1 - \sin^2 x)}{\sin^4 x} \, dx \\
 &= \left(\int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \frac{\cos x}{\sin^4 x} - \frac{\cos x}{\sin^2 x} \right) dx \\
 &= \int_{\frac{1}{2}}^1 \left(\frac{1}{u^4} - \frac{1}{u^2} \right) du \\
 &= \int_{\frac{1}{2}}^1 (u^{-4} - u^{-2}) \, du
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \left[\frac{u^{-3}}{3} - \frac{u^{-1}}{1} \right]_{\frac{1}{2}}$$

$$= \frac{4}{3}$$

8a Let $u = e^{2x}$, $du = 2e^{2x} dx$

$$\begin{aligned} & \int \frac{e^{2x}}{\sqrt{1+e^{2x}}} dx \\ &= \int \frac{1}{2\sqrt{1+u}} du \\ &= \int \frac{1}{2}(1+u)^{-\frac{1}{2}} du \\ &= (1+u)^{\frac{1}{2}} + C \\ &= \sqrt{1+u} + C \\ &= \sqrt{1+e^{2x}} + C \end{aligned}$$

8b Let $u = \ln x$, $du = \frac{1}{x} dx$

$$\begin{aligned} & \int \frac{1}{x \ln x} dx \\ &= \int \frac{1}{u} du \\ &= \ln|u| + C \\ &= \ln|\ln x| + C \end{aligned}$$

8c Let $u = \ln \cos x$, $du = -\frac{\sin x}{\cos x} dx = -\tan x dx$

$$\begin{aligned} & \int \frac{\tan x}{\ln \cos x} dx \\ &= \int -\frac{1}{u} du \\ &= -\ln|u| + C \\ &= -\ln|\ln \cos x| + C \end{aligned}$$

Chapter 12 worked solutions – Further calculus

8d Let $u = \tan x, du = \sec^2 x dx$

$$\begin{aligned} & \int \tan^3 x \sec^4 x dx \\ &= \int u^3 \sec^2 x du \\ &= \int u^3(1 + \tan^2 x) du \\ &= \int u^3(1 + u^2) du \\ &= \int (u^3 + u^5) du \\ &= \frac{u^4}{4} + \frac{u^6}{6} + C \\ &= \frac{\tan^4 x}{4} + \frac{\tan^6 x}{6} + C \end{aligned}$$

9a

$$\begin{aligned} \frac{dy}{dx} &= \frac{e^{2x}}{1 + e^{4x}} \\ y &= \int \frac{e^{2x}}{1 + e^{4x}} dx \end{aligned}$$

Let $u = e^{2x}, du = 2e^{2x} dx$

$$\begin{aligned} y &= \int \frac{e^{2x}}{1 + e^{4x}} dx \\ &= \frac{1}{2} \int \frac{1}{1 + u^2} du \\ &= \frac{1}{2} \tan^{-1} u + C \\ &= \frac{1}{2} \tan^{-1} e^{2x} + C \end{aligned}$$

When $x = 0, y = \frac{\pi}{8}$

$$\frac{\pi}{8} = \frac{1}{2} \tan^{-1} e^0 + C$$

$$\frac{\pi}{8} = \frac{1}{2} \tan^{-1} 1 + C$$

$$\frac{\pi}{8} = \frac{\pi}{8} + C$$

Chapter 12 worked solutions – Further calculus

Hence $C = 0$ so the equation of the line is

$$y = \frac{1}{2} \tan^{-1} e^{2x}$$

9b

$$y'' = \frac{x}{(4 - x^2)^{\frac{3}{2}}}$$

Let $u = 4 - x^2$, $du = -2x \, dx$ hence $-\frac{1}{2} du = dx$

$$\begin{aligned} y' &= \int \frac{x}{(4 - x^2)^{\frac{3}{2}}} dx \\ &= \int \frac{1}{u^{\frac{3}{2}}} \left(-\frac{1}{2}\right) du \\ &= -\frac{1}{2} \int u^{-\frac{3}{2}} \\ &= u^{-\frac{1}{2}} + C \\ &= (4 - x^2)^{-\frac{1}{2}} + C \end{aligned}$$

When $x = 0$, $y' = 1$

$$1 = 4^{-\frac{1}{2}} + C$$

$$1 = \frac{1}{2} + C$$

$$C = \frac{1}{2}$$

So

$$y' = (4 - x^2)^{-\frac{1}{2}} + \frac{1}{2} = \frac{1}{\sqrt{4 - x^2}} + \frac{1}{2}$$

$$\begin{aligned} y &= \int \left(\frac{1}{\sqrt{4 - x^2}} + \frac{1}{2} \right) dx \\ &= \int \left(\frac{1}{2\sqrt{1 - \left(\frac{x}{2}\right)^2}} + \frac{1}{2} \right) dx \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \sin^{-1}\left(\frac{x}{2}\right) + \frac{x}{2} + D$$

$$\text{When } x = 0, y = \frac{1}{2}$$

$$\frac{1}{2} = 0 + 0 + D$$

$$D = \frac{1}{2}$$

$$y = \sin^{-1}\left(\frac{x}{2}\right) + \frac{x}{2} + \frac{1}{2}$$

10a

$$\begin{aligned} & \frac{d}{dx}(\sec x) \\ &= \frac{d}{dx}(\cos^{-1} x) \\ &= -1 \times -\sin x \times (\cos x)^{-2} \\ &= \frac{1}{\cos x} \times \frac{\sin x}{\cos x} \\ &= \sec x \tan x \end{aligned}$$

10b i Let $u = \sec x, du = \sec x \tan x dx$

$$\begin{aligned} & \int_0^{\frac{\pi}{3}} 2^{\sec x} \sec x \tan x dx \\ &= \int_1^2 2^u du \\ &= \left[\frac{2^u}{\ln 2} \right]_1^2 \\ &= \frac{4}{\ln 2} - \frac{2}{\ln 2} \\ &= \frac{2}{\ln 2} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

10b ii

$$\int_0^{\frac{\pi}{4}} \sec^5 x \tan x \, dx$$

$$= \int_1^{\sqrt{2}} u^4 \, du$$

$$= \frac{1}{5} [u^5]_1^{\sqrt{2}}$$

$$= \frac{1}{5} (4\sqrt{2} - 1)$$

11a Let $u = \sin^2 x, du = 2 \cos x \sin x \, dx = \sin 2x \, dx$

$$\int_0^{\frac{\pi}{2}} \frac{\sin 2x}{1 + \sin^2 x} \, dx$$

$$= \int_0^1 \frac{1}{1 + u^2} \, du$$

$$= [\tan^{-1} u]_0^1$$

$$= \frac{\pi}{4}$$

11b Let $u = x \ln x, du = 1 \times \ln x + x \times \frac{1}{x} \, dx = \ln x + 1 \, dx$

$$\int_1^e \frac{\ln x + 1}{(x \ln x + 1)^2} \, dx$$

$$= \int_0^e \frac{1}{(u + 1)^2} \, du$$

$$= \left[-\frac{1}{u + 1} \right]_0^e$$

$$= -\frac{1}{e + 1} - (-1)$$

$$= 1 - \frac{1}{e + 1}$$

$$= \frac{e + 1}{e + 1} - \frac{1}{e + 1}$$

$$= \frac{e}{e + 1}$$

Chapter 12 worked solutions – Further calculus

12 Let $u = \sqrt{x-1}$, $du = \frac{1}{2\sqrt{x-1}} dx$

If $u = \sqrt{x-1}$, then $u^2 = x-1$ and $x = 1+u^2$

$$\int \frac{1}{2x\sqrt{x-1}} dx$$

$$= \int \frac{1}{x} du$$

$$= \int \frac{1}{1+u^2} du$$

$$= \tan^{-1} u + C$$

$$= \tan^{-1} \sqrt{x-1} + C$$

13a Let $u = \sqrt{x}$, $du = \frac{1}{2\sqrt{x}} dx$ and $u^2 = x$

$$\int \frac{1}{\sqrt{x(1-x)}} dx$$

$$= \int \frac{2}{\sqrt{(1-x)} 2\sqrt{x}} dx$$

$$= \int \frac{2}{\sqrt{1-u^2}} du$$

$$= 2 \sin^{-1} u + C_1$$

$$= 2 \sin^{-1} \sqrt{x} + C_1$$

13b Let $u = x - \frac{1}{2}$, $du = dx$, $x = \frac{1}{2} + u$

$$\int \frac{1}{\sqrt{x(1-x)}} dx$$

$$= \int \frac{1}{\sqrt{(\frac{1}{2}+u)(\frac{1}{2}-u)}} du$$

$$= \int \frac{1}{\sqrt{\frac{1}{4}-u^2}} du$$

$$= \int \frac{2}{\sqrt{1-(2u)^2}} du$$

$$= \sin^{-1}(2u) + C_2$$

$$= \sin^{-1}(2x-1) + C_2$$

Chapter 12 worked solutions – Further calculus

13c Since

$$\begin{aligned} & \int \frac{1}{\sqrt{x(1-x)}} dx \\ &= 2 \sin^{-1} \sqrt{x} + C_1 \\ &= \sin^{-1}(2x-1) + C_2 \end{aligned}$$

It follows that

$$\sin^{-1}(2x-1) = 2 \sin^{-1} \sqrt{x} + C_3$$

Substituting $x = 0$

$$\sin^{-1}(-1) = 2 \sin^{-1} \sqrt{0} + C_3$$

$$C_3 = -\frac{\pi}{2}$$

Hence we can conclude that

$$\sin^{-1}(2x-1) = 2 \sin^{-1} \sqrt{x} - \frac{\pi}{2}$$

14 Let $u = x - \frac{1}{x}$, $du = 1 + \frac{1}{x^2} dx$,Hence $x^2 du = x^2 + 1 dx$ When $x = 1$, $u = 0$ When $x = \frac{1}{2}(\sqrt{6} + \sqrt{2})$,

$$\begin{aligned} u &= \frac{1}{2}(\sqrt{6} + \sqrt{2}) - \frac{1}{\frac{1}{2}(\sqrt{6} + \sqrt{2})} \\ &= \frac{1}{2}(\sqrt{6} + \sqrt{2}) - \frac{\sqrt{6} - \sqrt{2}}{\frac{1}{2}(6 - 2)} \\ &= \frac{1}{2}(\sqrt{6} + \sqrt{2}) - \frac{1}{2}(\sqrt{6} - \sqrt{2}) \\ &= \sqrt{2} \end{aligned}$$

$$u^2 = x^2 - 2 + \frac{1}{x^2}$$

$$u^2 = \frac{x^4 + 1}{x^2} - 2$$

$$u^2 + 2 = \frac{x^4 + 1}{x^2}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}\frac{1}{u^2 + 2} &= \frac{x^2}{1 + x^4} \\ \int_1^{\frac{1}{2}(\sqrt{6}+\sqrt{2})} \frac{1+x^2}{1+x^4} dx &= \int_0^{\sqrt{2}} \frac{x^2}{1+x^4} du \\ &= \int_0^{\sqrt{2}} \frac{1}{u^2+2} du \\ &= \frac{1}{2} \int_0^{\sqrt{2}} \frac{1}{1+\left(\frac{u}{\sqrt{2}}\right)^2} du \\ &= \frac{\sqrt{2}}{2} \left[\tan^{-1} \left(\frac{u}{\sqrt{2}} \right) \right]_0^{\sqrt{2}} \\ &= \frac{\sqrt{2}}{2} (\tan^{-1} 1 - 0) \\ &= \frac{1}{\sqrt{2}} \left(\frac{\pi}{4} \right) \\ &= \frac{\pi}{4\sqrt{2}}\end{aligned}$$

Chapter 12 worked solutions – Further calculus

Solutions to Exercise 12E

$$1a \quad dx = \frac{d}{du}(u) + \frac{d}{du}(1)du = 1du = du$$

$$1b \quad I = \int x(x-1)^5 dx$$

$$= \int (u+1)(u)^5 du$$

$$1c \quad I = \int (u+1)(u)^5 du$$

$$= \int (u^6 + u^5) du$$

$$= \frac{u^7}{7} + \frac{u^6}{6} + C$$

$$= \frac{(x-1)^7}{7} + \frac{(x-1)^6}{6} + C$$

1d

$$\frac{d}{dx} \left(\frac{(x-1)^7}{7} + \frac{(x-1)^6}{6} \right)$$

$$= (x-1)^6 + (x-1)^5$$

$$= (x-1+1)(x-1)^5$$

$$= x(x-1)^5 + C$$

2a

$$\int \frac{x}{\sqrt{x-1}} dx$$

$$= \int \frac{u+1}{\sqrt{u}} du$$

$$= \int \left(u^{\frac{1}{2}} + u^{-\frac{1}{2}} \right) du$$

$$= \frac{2}{3}u^{\frac{3}{2}} + 2u^{\frac{1}{2}} + C$$

$$= \frac{2}{3}(x-1)^{\frac{3}{2}} + 2\sqrt{x-1} + C$$

Chapter 12 worked solutions – Further calculus

2b

$$\begin{aligned}
 & \int \frac{x}{(x-1)^2} dx \\
 &= \int \frac{u+1}{u^2} du \\
 &= \int \left(\frac{1}{u} + \frac{1}{u^2} \right) du \\
 &= \ln|u| - \frac{1}{u} + C \\
 &= \ln|x-1| - \frac{1}{x-1} + C
 \end{aligned}$$

3a $dx = 2u du$

3b

$$\begin{aligned}
 J &= \int x\sqrt{x+1} dx \\
 &= \int (u^2 - 1)\sqrt{u^2 - 1 + 1} 2u du \\
 &= 2 \int (u^2 - 1)\sqrt{u^2} u du \\
 &= 2 \int (u^2 - 1)u^2 du \\
 &= 2 \int (u^4 - u^2) du
 \end{aligned}$$

3c

$$\begin{aligned}
 J &= 2 \int (u^4 - u^2) du \\
 &= 2 \left(\frac{u^5}{5} - \frac{u^3}{3} \right) + C \\
 &= 2 \left(\frac{(\sqrt{x+1})^5}{5} - \frac{(\sqrt{x+1})^3}{3} \right) + C
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

3d

$$\begin{aligned}
 & \frac{d}{dx} \left(2 \left(\frac{(\sqrt{x+1})^5}{5} - \frac{(\sqrt{x+1})^3}{3} \right) \right) \\
 &= 2 \left(\frac{\frac{5}{2}(\sqrt{x+1})^3}{5} - \frac{\frac{3}{2}(\sqrt{x+1})}{3} \right) \\
 &= (\sqrt{x+1})^3 - (\sqrt{x+1}) \\
 &= (x+1)\sqrt{x+1} - \sqrt{x+1} \\
 &= (x+1-1)\sqrt{x+1} \\
 &= x\sqrt{x+1}
 \end{aligned}$$

4a

$$\begin{aligned}
 & \int x^2 \sqrt{x+1} \, dx \\
 &= \int (u^2 - 1)^2 \sqrt{u^2} \, 2u \, du \\
 &= 2 \int (u^2 - 1)^2 u \, u \, du \\
 &= 2 \int (u^2 - 1)^2 u^2 \, du \\
 &= 2 \int (u^4 - 2u^2 + 1)u^2 \, du \\
 &= 2 \int (u^6 - 2u^4 + u^2) \, du \\
 &= 2 \left(\frac{u^7}{7} - \frac{2u^5}{5} + \frac{u^3}{3} \right) + C \\
 &= \frac{2(x+1)^{\frac{7}{2}}}{7} - \frac{4(x+1)^{\frac{5}{2}}}{5} + \frac{2(x+1)^{\frac{3}{2}}}{3} + C
 \end{aligned}$$

4b

$$\begin{aligned}
 & \int \frac{2x+3}{\sqrt{x+1}} \, dx \\
 &= \int \left(\frac{2x+2}{\sqrt{x+1}} + \frac{1}{\sqrt{x+1}} \right) \, dx \\
 &= \int \left(\frac{2(x+1)}{\sqrt{x+1}} + \frac{1}{\sqrt{x+1}} \right) \, dx
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \int \left(\frac{2(u^2)}{\sqrt{u^2}} + \frac{1}{\sqrt{u^2}} 2u \right) du \\
 &= \int (4u^2 + 2u) du \\
 &= \frac{4}{3}u^3 + u^2 + C \\
 &= \frac{4}{3}(x+1)^{\frac{3}{2}} + 2(x+1)^{\frac{1}{2}} + C
 \end{aligned}$$

5a Let $x = u - 2, dx = du$

$$\begin{aligned}
 &\int \frac{x-2}{x+2} dx \\
 &= \int \frac{u-2-2}{u-2+2} du \\
 &= \int \frac{u-4}{u} du \\
 &= \int \left(1 - \frac{4}{u} \right) du \\
 &= u - 4 \ln|u| + C \\
 &= x + 2 - 4 \ln|x+2| + C
 \end{aligned}$$

5b Let $x = \frac{1}{2}(u+1), dx = \frac{1}{2}du, u = 2x - 1$

$$\begin{aligned}
 &\int \frac{2x+1}{\sqrt{2x-1}} dx \\
 &= \int \frac{2\left(\frac{1}{2}(u+1)\right)+1}{\sqrt{2\left(\frac{1}{2}(u+1)\right)-1}} \frac{1}{2} du \\
 &= \int \frac{u+1+1}{\sqrt{u+1-1}} \frac{1}{2} du \\
 &= \frac{1}{2} \int \frac{u+2}{\sqrt{u}} du \\
 &= \frac{1}{2} \int \left(\sqrt{u} + \frac{2}{\sqrt{u}} \right) du \\
 &= \frac{1}{2} \left(\frac{2}{3}u^{\frac{3}{2}} + 4\sqrt{u} \right) + C
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \frac{1}{3}u^{\frac{3}{2}} + 2\sqrt{u} + C \\
 &= \frac{1}{3}(2x-1)^{\frac{3}{2}} + 2\sqrt{2x-1} + C
 \end{aligned}$$

5c $x = \frac{1}{4}(u^2 + 5)$, $dx = \frac{1}{2}u du$, $u^2 = 4x - 5$, $u = \sqrt{4x-5}$

$$\begin{aligned}
 &\int 3x\sqrt{4x-5} dx \\
 &= 3 \int \frac{1}{4}(u^2 + 5) \sqrt{4\left(\frac{1}{4}(u^2 + 5)\right) - 5} \times \frac{1}{2}u du \\
 &= 3 \int \frac{1}{4}(u^2 + 5) \sqrt{u^2 + 5 - 5} \times \frac{1}{2}u du \\
 &= 3 \int \frac{1}{4}(u^2 + 5) \sqrt{u^2} \times \frac{1}{2}u du \\
 &= 3 \int \frac{1}{8}(u^2 + 5)u^2 du \\
 &= \frac{3}{8} \int (u^4 + 5u^2) du \\
 &= \frac{3}{8} \left(\frac{u^5}{5} + \frac{5u^3}{3} \right) + C \\
 &= \frac{3}{8} \left(\frac{u^5}{5} + \frac{5u^3}{3} \right) + C \\
 &= \frac{3}{8} \left(\frac{(\sqrt{4x-5})^5}{5} + \frac{5(\sqrt{4x-5})^3}{3} \right) + C
 \end{aligned}$$

5d $x = (u-1)^2$, $dx = 2(u-1) du$

Hence $\sqrt{x} = u-1$, $u = 1 + \sqrt{x}$

$$\begin{aligned}
 &\int \frac{1}{1+\sqrt{x}} dx \\
 &= \int \frac{1}{u} \times 2(u-1) du \\
 &= 2 \int \left(1 - \frac{1}{u}\right) du \\
 &= 2(u - \ln|u|) \\
 &= 2((1 + \sqrt{x}) - \ln(1 + \sqrt{x})) + C \quad (\text{as } \sqrt{x} \text{ positive}) \\
 &= 2(1 + \sqrt{x}) - 2 \ln(1 + \sqrt{x}) + C
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

6a $x = u - 1, dx = du$

$$\begin{aligned} & \int_0^1 x(x+1)^3 dx \\ &= \int_1^2 (u-1)u^3 du \\ &= \int_1^2 (u^4 - u^3) du \\ &= \left[\frac{u^5}{5} - \frac{u^4}{4} \right]_1^2 \\ &= \frac{49}{20} \end{aligned}$$

6b $x = 1 - u, dx = -du$

$$\begin{aligned} & \int_0^{\frac{1}{2}} \frac{1+x}{1-x} dx \\ &= \int_1^{\frac{1}{2}} \frac{2-u}{u} (-du) \\ &= \int_{\frac{1}{2}}^1 \left(\frac{2}{u} - 1 \right) du \\ &= [2 \ln|u| - u]_{\frac{1}{2}}^1 \\ &= 2 \ln 2 - \frac{1}{2} \end{aligned}$$

6c Let $x = \frac{1}{3}(u-1), dx = \frac{1}{3}du, u = 3x + 1$

$$\begin{aligned} & \int_0^1 \frac{3x}{\sqrt{3x+1}} dx \\ &= \int_1^4 \frac{u-1}{\sqrt{u}} du \\ &= \int_1^4 \left(\sqrt{u} - u^{-\frac{1}{2}} \right) du \\ &= \left[\frac{2}{3}u^{\frac{3}{2}} - 2u^{\frac{1}{2}} \right]_1^4 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{8}{9}$$

6d Let $x = u - 2, dx = du, u = x + 2$

$$\begin{aligned} & \int_0^1 \frac{2-x}{(2+x)^3} dx \\ &= \int_2^3 \frac{4-u}{(u)^3} du \\ &= \int_2^3 \left(\frac{4}{u^3} - \frac{1}{u^2} \right) du \\ &= \left[-\frac{2}{u^2} + \frac{1}{u} \right]_2^3 \\ &= \frac{1}{9} \end{aligned}$$

6e Let $x = 4 - u^2, dx = -2u du, u = \sqrt{4-x}$

$$\begin{aligned} & \int_0^4 x \sqrt{4-x} dx \\ &= \int_2^0 (4-u^2)u (-2u du) \\ &= \int_0^2 (4-u^2)u (2u du) \\ &= \int_0^2 (8u^2 - 2u^4) du \\ &= \left[\frac{8u^3}{3} - \frac{2u^5}{5} \right]_0^2 \\ &= \frac{128}{15} \end{aligned}$$

6f $x = \frac{1}{2}(u^2 + 1), dx = u du, u^2 = 2x - 1, u = \sqrt{2x-1}$

$$\int_1^5 \frac{x}{(2x-1)^{\frac{3}{2}}} dx$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \int_1^5 \frac{x}{(2x-1)^{\frac{3}{2}}} dx \\
 &= \int_1^3 \frac{\frac{1}{2}(u^2+1)}{(u^2)^{\frac{3}{2}}} u du \\
 &= \frac{1}{2} \int_1^3 \frac{(u^2+1)}{u^3} u du \\
 &= \frac{1}{2} \int_1^3 \frac{(u^2+1)}{u^2} du \\
 &= \frac{1}{2} \int_1^3 \left(1 + \frac{1}{u^2}\right) du \\
 &= \frac{1}{2} \left[u - \frac{1}{u}\right]_1^3 \\
 &= \frac{4}{3}
 \end{aligned}$$

6g $x = (u-3)^2, dx = 2(u-3) du, u = \sqrt{x} + 3$

$$\begin{aligned}
 &\int_0^4 \frac{1}{3+\sqrt{x}} dx \\
 &= \int_3^5 \frac{1}{u} 2(u-3) du \\
 &= \int_3^5 2 - \frac{6}{u} du \\
 &= [2u - 6 \ln|u|]_3^5 \\
 &= 4 - 6 \ln \frac{5}{3}
 \end{aligned}$$

6h $x = u^3 - 1, dx = 3u^2 du, u = \sqrt[3]{x+1}$

$$\begin{aligned}
 &\int_0^7 \frac{x^2}{\sqrt[3]{x+1}} dx \\
 &= \int_1^2 \frac{(u^3-1)^2}{u} 3u^2 du \\
 &= \int_1^2 3u(u^3-1)^2 du
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \int_1^2 3u(u^6 - 2u^3 + 1)du \\
 &= \int_1^2 (3u^7 - 6u^4 + 3u) du \\
 &= \left[\frac{3u^8}{8} - \frac{6u^5}{5} + \frac{3u^2}{2} \right]_1^2 \\
 &= \frac{2517}{40}
 \end{aligned}$$

7a Let $x = u - 2, dx = du$

$$\begin{aligned}
 I &= \int \frac{1}{\sqrt{5 - 4x - x^2}} dx \\
 &= \int \frac{1}{\sqrt{5 - 4(u - 2) - (u - 2)^2}} du \\
 &= \int \frac{1}{\sqrt{5 - 4u + 8 - (u^2 - 4u + 4)}} \\
 &= \int \frac{1}{\sqrt{9 - u^2}} \\
 &= \int \frac{1}{3\sqrt{1 - \left(\frac{u}{3}\right)^2}} du \\
 &= \sin^{-1}\left(\frac{u}{3}\right) + C \\
 &= \sin^{-1}\left(\frac{x+2}{3}\right) + C
 \end{aligned}$$

7b i Let $x = u - 1, dx = du$

$$\begin{aligned}
 &\int \frac{1}{x^2 + 2x + 4} dx \\
 &= \int \frac{1}{(u - 1)^2 + 2(u - 1) + 4} du \\
 &= \int \frac{1}{u^2 - 2u + 1 + 2u - 2 + 4} du \\
 &= \int \frac{1}{u^2 + 3} du \\
 &= \frac{1}{3} \int \frac{1}{1 + \left(\frac{u}{\sqrt{3}}\right)^2} du
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \frac{\sqrt{3}}{3} \tan^{-1} \left(\frac{u}{\sqrt{3}} \right) + C \\
 &= \frac{1}{\sqrt{3}} \tan^{-1} \frac{x+1}{\sqrt{3}} + C
 \end{aligned}$$

7b ii Let $x = u - 1, dx = du$

$$\begin{aligned}
 &\int \frac{1}{\sqrt{4 - 2x - x^2}} dx \\
 &= \int \frac{1}{\sqrt{4 - 2(u-1) - (u-1)^2}} du \\
 &= \int \frac{1}{\sqrt{4 - 2u + 2 - (u^2 - 2u + 1)}} du \\
 &= \int \frac{1}{\sqrt{5 - u^2}} du \\
 &= \int \frac{1}{\sqrt{5} \sqrt{1 - \left(\frac{u}{\sqrt{5}}\right)^2}} du \\
 &= \sin^{-1} \left(\frac{u}{\sqrt{5}} \right) + C \\
 &= \sin^{-1} \frac{x+1}{\sqrt{5}} + C
 \end{aligned}$$

7b iii Let $x = u + 1, dx = du$

$$\begin{aligned}
 &\int_1^2 \frac{1}{\sqrt{3 + 2x - x^2}} dx \\
 &= \int_0^1 \frac{1}{\sqrt{3 + 2(u+1) - (u+1)^2}} du \\
 &= \int_0^1 \frac{1}{\sqrt{3 + 2u + 2 - (u^2 + 2u + 1)}} du \\
 &= \int_0^1 \frac{1}{\sqrt{4 - u^2}} du \\
 &= \left[\sin^{-1} \frac{u}{2} \right]_0^1 \\
 &= \frac{\pi}{6}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

7b iv Let $x = u + 3, dx = du$

$$\begin{aligned}
 & \int_3^7 \frac{1}{x^2 - 6x + 25} dx \\
 &= \int_0^4 \frac{1}{(u+3)^2 - 6(u+3) + 25} du \\
 &= \int_0^4 \frac{1}{(u+3)^2 - 6(u+3) + 25} du \\
 &= \int_0^4 \frac{1}{u^2 + 6u + 9 - 6u - 18 + 25} du \\
 &= \int_0^4 \frac{1}{u^2 + 16} du \\
 &= \frac{1}{4} \left[\tan^{-1} \frac{u}{4} \right]_0^4 \\
 &= \frac{\pi}{16}
 \end{aligned}$$

8a Let $x = 2 \sin \theta, dx = 2 \cos \theta d\theta$

Hence

$$\begin{aligned}
 J &= \int \frac{1}{\sqrt{4-x^2}} dx \\
 &= \int \frac{1}{\sqrt{4-(2 \sin \theta)^2}} \times 2 \cos \theta d\theta \\
 &= \int \frac{1}{\sqrt{4-4 \sin^2 \theta}} \times 2 \cos \theta d\theta \\
 &= \int \frac{1}{2\sqrt{1-\sin^2 \theta}} \times 2 \cos \theta d\theta \\
 &= \int \frac{1}{2\sqrt{\cos^2 \theta}} \times 2 \cos \theta d\theta \\
 &= \int \frac{1}{2 \cos \theta} \times 2 \cos \theta d\theta \\
 &= \int 1 d\theta \\
 &= \theta + C
 \end{aligned}$$

Since

$$x = 2 \sin \theta$$

$$\sin \theta = \frac{x}{2}$$

Hence

Chapter 12 worked solutions – Further calculus

$$\theta = \sin^{-1} \frac{x}{2}$$

Thus

$$J = \sin^{-1} \frac{x}{2} + C$$

8b i Let $x = 3 \tan \theta$, $dx = 3 \sec^2 \theta d\theta$ and $\theta = \tan^{-1} \frac{x}{3}$

$$\begin{aligned} & \int \frac{1}{9+x^2} dx \\ &= \int \frac{1}{9+(3\tan\theta)^2} 3\sec^2\theta d\theta \\ &= \int \frac{1}{9+9\tan^2\theta} 3\sec^2\theta d\theta \\ &= \frac{1}{3} \int \frac{\sec^2\theta}{1+\tan^2\theta} d\theta \\ &= \frac{1}{3} \int \frac{\sec^2\theta}{\sec^2\theta} d\theta \\ &= \frac{1}{3} \int 1 d\theta \\ &= \frac{1}{3}\theta + C \\ &= \frac{1}{3}\tan^{-1}\frac{x}{3} + C \end{aligned}$$

8b ii Let $x = \sqrt{3} \cos \theta$, $dx = -\sqrt{3} \sin \theta d\theta$, $\theta = \cos^{-1} \frac{x}{\sqrt{3}}$

$$\begin{aligned} & \int -\frac{1}{\sqrt{3-x^2}} dx \\ &= \int -\frac{1}{\sqrt{3-(\sqrt{3}\cos\theta)^2}} \times -\sqrt{3}\sin\theta d\theta \\ &= \int \frac{\sqrt{3}\sin\theta}{\sqrt{3\sqrt{1-\cos^2\theta}}} d\theta \\ &= \int \frac{\sqrt{3}\sin\theta}{\sqrt{3\sin^2\theta}} d\theta \\ &= \int \frac{\sqrt{3}\sin\theta}{\sqrt{3}\sin\theta} d\theta \\ &= \int 1 d\theta \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \theta + C \\
 &= \cos^{-1} \frac{x}{\sqrt{3}} + C
 \end{aligned}$$

8b iii Let $x = \frac{1}{2} \sin \theta$, $dx = \frac{1}{2} \cos \theta \ d\theta$ and $\theta = \sin^{-1} 2x$

$$\begin{aligned}
 &\int \frac{1}{\sqrt{1 - 4x^2}} dx \\
 &= \int \frac{1}{\sqrt{1 - 4\left(\frac{1}{2} \sin \theta\right)^2}} \frac{1}{2} \cos \theta \ d\theta \\
 &= \frac{1}{2} \int \frac{\cos \theta}{\sqrt{1 - \sin^2 \theta}} d\theta \\
 &= \frac{1}{2} \int \frac{\cos \theta}{\sqrt{\cos^2 \theta}} d\theta \\
 &= \frac{1}{2} \int \frac{\cos \theta}{\cos \theta} d\theta \\
 &= \frac{1}{2} \int 1 d\theta \\
 &= \frac{1}{2} \theta + C \\
 &= \frac{1}{2} \sin^{-1} 2x + C
 \end{aligned}$$

8b iv Let $x = \frac{1}{4} \tan \theta$, $dx = \frac{1}{4} \sec^2 \theta \ d\theta$ and $\theta = \tan^{-1} 4x$

$$\begin{aligned}
 &\int \frac{1}{1 + 16x^2} dx \\
 &= \int \frac{1}{1 + 16\left(\frac{1}{4} \tan \theta\right)^2} \frac{1}{4} \sec^2 \theta \ d\theta \\
 &= \frac{1}{4} \int \frac{\sec^2 \theta}{1 + \tan^2 \theta} d\theta \\
 &= \frac{1}{4} \int \frac{\sec^2 \theta}{\sec^2 \theta} d\theta \\
 &= \frac{1}{4} \int 1 d\theta \\
 &= \frac{1}{4} \theta + C \\
 &= \frac{1}{4} \tan^{-1} 4x + C
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

8bv Let $x = 6 \sin \theta$, $dx = 6 \cos \theta d\theta$ and $\theta = \sin^{-1} \frac{x}{6}$.

When $x = 0, \theta = 0$

When $x = 3, \theta = \sin^{-1} \frac{1}{2} = \frac{\pi}{6}$

$$\begin{aligned} & \int_0^3 \frac{1}{\sqrt{36 - x^2}} dx \\ &= \int_0^{\frac{\pi}{6}} \frac{1}{\sqrt{36 - (6 \sin \theta)^2}} 6 \cos \theta d\theta \\ &= \int_0^{\frac{\pi}{6}} \frac{1}{6\sqrt{1 - \sin^2 \theta}} 6 \cos \theta d\theta \\ &= \int_0^{\frac{\pi}{6}} \frac{6 \cos \theta}{6\sqrt{\cos^2 \theta}} d\theta \\ &= \int_0^{\frac{\pi}{6}} \frac{6 \cos \theta}{6 \cos \theta} d\theta \\ &= \int_0^{\frac{\pi}{6}} 1 d\theta \\ &= [\theta]_0^{\frac{\pi}{6}} \\ &= \frac{\pi}{6} \end{aligned}$$

8b vi

Let $x = \frac{2}{3} \tan \theta$, $dx = \frac{2}{3} \sec^2 \theta d\theta$ and $\theta = \tan^{-1} \frac{3}{2}x$

When $x = 0, \theta = 0$

When $x = \frac{2}{3}, \theta = \tan^{-1} 1 = \frac{\pi}{4}$

$$\begin{aligned} & \int_0^{\frac{2}{3}} \frac{1}{4 + 9x^2} dx \\ &= \int_0^{\frac{\pi}{4}} \frac{1}{4 + 9\left(\frac{2}{3} \tan \theta\right)^2} \frac{2}{3} \sec^2 \theta d\theta \\ &= \int_0^{\frac{\pi}{4}} \frac{1}{4 + 4 \tan^2 \theta} \frac{2}{3} \sec^2 \theta d\theta \\ &= \frac{1}{6} \int_0^{\frac{\pi}{4}} \frac{\sec^2 \theta}{\sec^2 \theta} d\theta \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \frac{1}{6} \int_0^{\frac{\pi}{4}} 1 d\theta \\
 &= \frac{1}{6} [\theta]_0^{\frac{\pi}{4}} \\
 &= \frac{\pi}{24}
 \end{aligned}$$

9a Let $x = \sin \theta$, $dx = \cos \theta d\theta$ and $\theta = \sin^{-1} x$

$$\begin{aligned}
 I &= \int \frac{1}{(1-x^2)^{\frac{3}{2}}} dx \\
 &= \int \frac{1}{(1-\sin^2 \theta)^{\frac{3}{2}}} \cos \theta d\theta \\
 &= \int \frac{1}{(\cos^2 \theta)^{\frac{3}{2}}} \cos \theta d\theta \\
 &= \int \frac{1}{\cos^3 \theta} \cos \theta d\theta \\
 &= \int \frac{1}{\cos^2 \theta} d\theta \\
 &= \int \sec^2 \theta d\theta \\
 &= \tan \theta + C
 \end{aligned}$$

Now since $\sin \theta = x = \frac{x}{1}$, it follows, from the Pythagorean theorem that

$$\tan \theta = \frac{x}{\sqrt{1-x^2}}$$

Hence

$$I = \frac{x}{\sqrt{1-x^2}} + C$$

9b i Let $x = 2 \tan \theta$, $dx = 2 \sec^2 \theta d\theta$

$$\begin{aligned}
 &\int \frac{1}{(4+x^2)^{\frac{3}{2}}} dx \\
 &= \int \frac{1}{(4+4\tan^2 \theta)^{\frac{3}{2}}} 2 \sec^2 \theta d\theta \\
 &= \int \frac{1}{(4\sec^2 \theta)^{\frac{3}{2}}} 2 \sec^2 \theta d\theta
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \int \frac{1}{8 \sec^3 \theta} 2 \sec^2 \theta d\theta \\
 &= \frac{1}{4} \int \frac{1}{\sec \theta} d\theta \\
 &= \frac{1}{4} \int \cos \theta d\theta \\
 &= \frac{1}{4} \sin \theta + C \\
 &= \frac{1}{4} \frac{x}{\sqrt{4+x^2}} + C
 \end{aligned}$$

9b ii Let $x = \sin \theta, dx = \cos \theta d\theta$

$$\begin{aligned}
 &\int_0^{\frac{1}{2}} \frac{x^2}{\sqrt{1-x^2}} dx \\
 &= \int_0^{\frac{\pi}{6}} \frac{\sin^2 \theta}{\sqrt{1-\sin^2 \theta}} \cos \theta d\theta \\
 &= \int_0^{\frac{\pi}{6}} \frac{\sin^2 \theta}{\sqrt{\cos^2 \theta}} \cos \theta d\theta \\
 &= \int_0^{\frac{\pi}{6}} \frac{\sin^2 \theta}{\cos \theta} \cos \theta d\theta \\
 &= \int_0^{\frac{\pi}{6}} \sin^2 \theta d\theta \\
 &= \frac{1}{2} \int_0^{\frac{\pi}{6}} (1 - \cos 2\theta) d\theta \\
 &= \frac{1}{2} \left[\theta - \frac{\sin 2\theta}{2} \right]_0^{\frac{\pi}{6}} \\
 &= \frac{\pi}{12} - \frac{\sqrt{3}}{8}
 \end{aligned}$$

9b iii Let $x = 2 \sin \theta, dx = 2 \cos \theta d\theta$

$$\begin{aligned}
 &\int_0^2 \sqrt{4-x^2} dx \\
 &= \int_0^{\frac{\pi}{2}} \sqrt{4-(2 \sin \theta)^2} 2 \cos \theta d\theta \\
 &= \int_0^{\frac{\pi}{2}} 2\sqrt{1-\sin^2 \theta} 2 \cos \theta d\theta
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= 4 \int_0^{\frac{\pi}{2}} \cos^2 \theta \, d\theta \\
 &= 2 \int_0^{\frac{\pi}{2}} (1 + \cos 2\theta) \, d\theta \\
 &= 2 \left[\theta + \frac{\sin 2\theta}{2} \right]_0^{\frac{\pi}{2}} \\
 &= \pi
 \end{aligned}$$

9b iv Let $x = 5 \cos \theta, dx = -5 \sin \theta \, d\theta$

$$\begin{aligned}
 &\int \frac{1}{x^2 \sqrt{25 - x^2}} \, dx \\
 &= \int \frac{1}{(5 \cos \theta)^2 \sqrt{25 - (5 \cos \theta)^2}} \times -5 \sin \theta \, d\theta \\
 &= \int \frac{1}{25 \cos^2 \theta 5 \sqrt{1 - \cos^2 \theta}} \times -5 \sin \theta \, d\theta \\
 &= \int \frac{1}{25 \cos^2 \theta 5 \sin \theta} \times -5 \sin \theta \, d\theta \\
 &= -\frac{1}{25} \int \frac{1}{\cos^2 \theta} \, d\theta \\
 &= -\frac{1}{25} \int \sec^2 \theta \, d\theta \\
 &= -\frac{1}{25} \tan \theta + C \\
 &= -\frac{1}{25} \times \frac{\sqrt{25 - x^2}}{x} + C \\
 &= -\frac{\sqrt{25 - x^2}}{25x} + C
 \end{aligned}$$

9b v Let $x = 3 \tan \theta, dx = 3 \sec^2 \theta \, d\theta$

$$\begin{aligned}
 &\int \frac{1}{x^2 \sqrt{9 + x^2}} \, dx \\
 &= \int \frac{1}{(3 \tan \theta)^2 \sqrt{9 + (3 \tan \theta)^2}} 3 \sec^2 \theta \, d\theta \\
 &= \int \frac{1}{(3 \tan \theta)^2 3 \sqrt{1 + \tan^2 \theta}} 3 \sec^2 \theta \, d\theta \\
 &= \int \frac{1}{(3 \tan \theta)^2 3 \sec \theta} 3 \sec^2 \theta \, d\theta
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \int \frac{\sec \theta}{(3 \tan \theta)^2} d\theta \\
 &= \frac{1}{9} \int \frac{\cos^2 \theta}{\sin^2 \theta} \times \frac{1}{\cos \theta} d\theta \\
 &= \frac{1}{9} \int \frac{\cos \theta}{\sin \theta} \times \frac{1}{\sin \theta} d\theta \\
 &= \frac{1}{9} \int \cot \theta \cosec \theta d\theta \\
 &= -\frac{1}{9} \cosec \theta + C \\
 &= -\frac{1}{9} \times \frac{\sqrt{9+x^2}}{x} + C \\
 &= -\frac{\sqrt{9+x^2}}{9x} + C
 \end{aligned}$$

9b vi Let $x = 2 \sec \theta$, $dx = 2 \sec \theta \tan \theta d\theta$

$$\begin{aligned}
 &\int_2^4 \frac{1}{x^2 \sqrt{x^2 - 4}} dx \\
 &= \int_0^{\frac{\pi}{3}} \frac{1}{(2 \sec \theta)^2 \sqrt{(2 \sec \theta)^2 - 4}} 2 \sec \theta \tan \theta d\theta \\
 &= \int_0^{\frac{\pi}{3}} \frac{1}{(2 \sec \theta)^2 2 \sqrt{\sec^2 \theta - 1}} 2 \sec \theta \tan \theta d\theta \\
 &= \int_0^{\frac{\pi}{3}} \frac{1}{(2 \sec \theta)^2 2 \sqrt{\tan^2 \theta}} 2 \sec \theta \tan \theta d\theta \\
 &= \int_0^{\frac{\pi}{3}} \frac{1}{(2 \sec \theta)^2 2 \tan \theta} 2 \sec \theta \tan \theta d\theta \\
 &= \int_0^{\frac{\pi}{3}} \frac{1}{4 \sec \theta} d\theta \\
 &= \frac{1}{4} \int_0^{\frac{\pi}{3}} \cos \theta d\theta \\
 &= \frac{1}{4} [\sin \theta]_0^{\frac{\pi}{3}} \\
 &= \frac{\sqrt{3}}{8}
 \end{aligned}$$

10 Let $x = 3 \sec \theta$, $dx = 3 \sec \theta \tan \theta d\theta$ and $\theta = \sec^{-1} \frac{x}{3}$
 $f(x)$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \int f'(x) dx \\
 &= \int \frac{\sqrt{x^2 - 9}}{x} dx \\
 &= \int \frac{\sqrt{9 \sec^2 \theta - 9}}{3 \sec \theta} 3 \sec \theta \tan \theta d\theta \\
 &= \int \frac{3 \tan \theta}{3 \sec \theta} 3 \sec \theta \tan \theta d\theta \\
 &= 3 \int \tan^2 \theta d\theta \\
 &= 3 \int (\sec^2 \theta - 1) d\theta \\
 &= 3(\tan \theta - \theta) + C \\
 &= \sqrt{x^2 - 9} - 3 \sec^{-1} \frac{x}{3} + C
 \end{aligned}$$

Now note that

$$\sqrt{x^2 - 9} = \sqrt{9 \sec^2 \theta - 9} = 3 \tan \theta$$

$$\therefore \theta = \tan^{-1} \frac{\sqrt{x^2 - 9}}{3}$$

$$f(3) = 0$$

$$\sqrt{3^2 - 9} - 3 \sec^{-1} \frac{3}{3} + C = 0$$

$$C = 3$$

Hence,

$$\begin{aligned}
 f(x) &= 3[\tan \theta - \theta] + C \\
 &= \sqrt{x^2 - 9} - 3 \tan^{-1} \frac{\sqrt{x^2 - 9}}{3}
 \end{aligned}$$

- 11 Let $x = \sqrt{3} \sin \theta$, $dx = \sqrt{3} \cos \theta d\theta$

$$\begin{aligned}
 A &= \int_0^1 \frac{x^3}{\sqrt{3 - x^2}} dx \\
 &= \int_0^{\frac{\pi}{6}} \frac{3\sqrt{3} \sin^3 \theta}{\sqrt{3 - 3 \sin^2 \theta}} \sqrt{3} \cos \theta d\theta \\
 &= \int_0^{\frac{\pi}{6}} \frac{3\sqrt{3} \sin^3 \theta}{\sqrt{3} \cos \theta} \sqrt{3} \cos \theta d\theta
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \int_0^{\frac{\pi}{6}} 3\sqrt{3} \sin^3 \theta \, d\theta$$

Let $u = \cos \theta, du = -\sin \theta \, d\theta$

$$\begin{aligned} A &= \int_1^{\frac{\sqrt{3}}{2}} -3\sqrt{3} \sin^2 \theta \, du \\ &= 3\sqrt{3} \int_{\frac{\sqrt{3}}{2}}^1 1 - \cos^2 \theta \, du \\ &= 3\sqrt{3} \int_{\frac{\sqrt{3}}{2}}^1 1 - u^2 \, du \\ &= 3\sqrt{3} \left[u - \frac{u^3}{3} \right]_{\frac{\sqrt{3}}{2}}^1 \\ &= \frac{1}{3}(6\sqrt{3} - 7\sqrt{2}) \text{ square units} \end{aligned}$$

12a Let $x = r \sin \theta, dx = r \cos \theta \, d\theta$

$$\begin{aligned} A &= 2 \int_{-r}^r \sqrt{r^2 - x^2} \, dx \\ &= 2 \int_{-r}^r \sqrt{r^2 - x^2} \, dx \\ &= 2 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sqrt{r^2 - (r \sin \theta)^2} r \cos \theta \, d\theta \\ &= 2 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} r \sqrt{1 - \sin^2 \theta} r \cos \theta \, d\theta \\ &= 2r^2 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^2 \theta \, d\theta \\ &= r^2 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} (1 - \cos 2\theta) \, d\theta \\ &= r^2 \left[\theta - \frac{\sin 2\theta}{2} \right]_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \\ &= \pi r^2 \end{aligned}$$

12b i Let P be the x -position of chord AB

Chapter 12 worked solutions – Further calculus

$$\sin \frac{\alpha}{2} = \frac{P}{OA} = \frac{P}{r}$$

Hence

$$P = r \sin \frac{\alpha}{2}$$

The area is twice the area from P to r under the semicircle

$$A = 2 \int_P^r \sqrt{r^2 - x^2} dx = 2 \int_{r \sin \frac{\alpha}{2}}^r \sqrt{r^2 - x^2} dx$$

12b ii Let $x = r \cos \theta$, $dx = -r \sin \theta d\theta$

$$\begin{aligned} A &= 2 \int_{r \sin \frac{\alpha}{2}}^r \sqrt{r^2 - x^2} dx \\ &= 2 \int_{r \sin \frac{\alpha}{2}}^r \sqrt{r^2 - x^2} dx \\ &= 2 \int_{r \sin \frac{\alpha}{2}}^{\frac{\pi}{2}} -\sqrt{r^2 - (r \cos \theta)^2} r \sin \theta d\theta \\ &= 2 \int_{\frac{1}{2}\alpha}^0 -r \sqrt{1 - \cos^2 \theta} r \sin \theta d\theta \\ &= 2 \int_{\frac{1}{2}\alpha}^0 -r \sin \theta r \sin \theta d\theta \\ &= -2r^2 \int_{\frac{1}{2}\alpha}^0 \sin^2 \theta d\theta \end{aligned}$$

12b iii

$$\begin{aligned} A &= 2 \int_{r \sin \frac{\alpha}{2}}^r \sqrt{r^2 - x^2} dx \\ &= -2r^2 \int_{\frac{1}{2}\alpha}^0 \sin^2 \theta d\theta \\ &= 2r^2 \int_0^{\frac{1}{2}\alpha} \sin^2 \theta d\theta \\ &= 2r^2 \int_0^{\frac{1}{2}\alpha} (1 - \cos 2\theta) d\theta \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= 2r^2 \left[\theta - \frac{1}{2} \sin 2\theta \right]_0^{\frac{1}{2}\alpha} \\
 &= \frac{1}{2} r^2 (\alpha - \sin \alpha)
 \end{aligned}$$

12c For an ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\frac{y^2}{b^2} = 1 - \frac{x^2}{a^2}$$

$$y = b \sqrt{1 - \frac{x^2}{a^2}}$$

Let $x = a \sin \theta$, $dx = a \cos \theta d\theta$

The area will be given by

$$\begin{aligned}
 A &= 2 \int_{-a}^a b \sqrt{1 - \frac{x^2}{a^2}} dx \\
 &= 2 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} b \sqrt{1 - \frac{a^2 \sin^2 \theta}{a^2}} a \cos \theta d\theta \\
 &= 2ab \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sqrt{1 - \sin^2 \theta} \cos \theta d\theta \\
 &= 2ab \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^2 \theta d\theta \\
 &= 2ab \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos^2 \theta d\theta \\
 &= ab \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} 1 + \cos 2\theta d\theta \\
 &= ab \left[\theta + \frac{1}{2} \sin 2\theta \right]_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \\
 &= \pi ab
 \end{aligned}$$

The unit circle has area π , dilating it a units horizontally causes it to become $a\pi$ and then dilating it a further b units vertically causes the area to become πab .

Chapter 12 worked solutions – Further calculus

13a Let $x = -u, dx = -du$

$$\begin{aligned}
 & \int_{-2}^2 \frac{x^2}{e^x + 1} dx \\
 &= \int_2^{-2} \frac{(-u)^2}{e^{-u} + 1} (-du) \quad (\text{using when } x = -2, u = 2 \text{ and when } x = 2, u = -2) \\
 &= \int_{-2}^2 \frac{u^2}{e^{-u} + 1} du \\
 &= \int_{-2}^2 \frac{u^2 e^u}{1 + e^u} du \\
 &= \int_{-2}^2 \frac{u^2 e^u}{e^u + 1} du
 \end{aligned}$$

Now let $x = u$

$$\int_{-2}^2 \frac{u^2 e^u}{e^u + 1} du = \int_{-2}^2 \frac{x^2 e^x}{e^x + 1} dx$$

Hence

$$\int_{-2}^2 \frac{x^2}{e^x + 1} dx = \int_{-2}^2 \frac{x^2 e^x}{e^x + 1} dx$$

13b

$$\begin{aligned}
 & \int_{-2}^2 \frac{x^2}{e^x + 1} dx \\
 &= \frac{1}{2} \left(\int_{-2}^2 \frac{x^2}{e^x + 1} dx + \int_{-2}^2 \frac{x^2}{e^x + 1} dx \right) \\
 &= \frac{1}{2} \left(\int_{-2}^2 \frac{x^2}{e^x + 1} dx + \int_{-2}^2 \frac{x^2 e^x}{e^x + 1} dx \right) \\
 &= \frac{1}{2} \left(\int_{-2}^2 \frac{x^2 + x^2 e^x}{e^x + 1} dx \right) \\
 &= \frac{1}{2} \left(\int_{-2}^2 \frac{x^2 (1 + e^x)}{e^x + 1} dx \right) \\
 &= \frac{1}{2} \left(\int_{-2}^2 x^2 dx \right) \\
 &= \frac{1}{2} \left[\frac{x^3}{3} \right]_{-2}^2
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus


$$= \frac{1}{2} \left(\frac{8}{3} - \left(-\frac{8}{3} \right) \right)$$

$$= \frac{8}{3}$$

Uncorrected proofs

Solutions to Exercise 12F

1a



1b The radius will be $r = 3(3) = 9$ units and the height will be 3 units.

Hence


$$\begin{aligned} V &= \frac{1}{3}\pi r^2 \\ &= \frac{1}{3}\pi(9)^2(3) \\ &= 81\pi \text{ cubic units} \end{aligned}$$

1c

$$\begin{aligned} &\pi \int_0^3 9x^2 dx \\ &= 3\pi[x^3]_0^3 \\ &= 81\pi \text{ cubic units} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

2a



2b This is a sphere of radius 3, hence the volume is

$$\begin{aligned}V &= \frac{4}{3}\pi r^3 \\&= \frac{4}{3}\pi(3)^3 \\&= 36\pi \text{ cubic units}\end{aligned}$$

2c

$$\begin{aligned}\pi \int_{-3}^3 (9 - x^2) dx \\&= \pi \left[9x - \frac{x^3}{3} \right]_{-3}^3 \\&= 36\pi \text{ cubic units}\end{aligned}$$

3a This is a cylinder of radius 2, height 4. Hence

$$\begin{aligned}V &= \pi r^2 h \\&= \pi(2)^2 \times 4 \\&= 16\pi \text{ cubic units}\end{aligned}$$

3b This is a cone of radius 3 and height 3. Hence

$$\begin{aligned}V &= \frac{1}{3}\pi r^2 h \\&= \frac{1}{3}\pi(3)^2 \times 3\end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= 9\pi \text{ cubic units}$$

3c

$$\begin{aligned} V &= \pi \int_0^2 y^2 dx \\ &= \pi \int_0^2 x^4 dx \\ &= \pi \left[\frac{x^5}{5} \right]_0 \\ &= \frac{32\pi}{5} \text{ cubic units} \end{aligned}$$

3d

$$\begin{aligned} V &= \pi \int_2^4 y^2 dx \\ &= \pi \int_2^4 x dx \\ &= \frac{\pi}{2} [x^2]_2^4 \\ &= 6\pi \text{ cubic units} \end{aligned}$$

3e

$$\begin{aligned} V &= \pi \int_0^2 y^2 dx \\ &= \int_0^2 (4 - x^2) dx \\ &= \left[4x - \frac{x^3}{3} \right]_0 \\ &= \frac{16\pi}{3} \text{ cubic units} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

3f

$$\begin{aligned}V &= \pi \int_{-1}^0 y^2 dx \\&= \pi \int_{-1}^0 x^6 dx \\&= \pi \left[\frac{x^7}{7} \right]_{-1}^0 \\&= \frac{\pi}{7} \text{ cubic units}\end{aligned}$$

3g

$$\begin{aligned}V &= \pi \int_{-5}^{-2} y^2 dx \\&= \pi \int_{-5}^{-2} (x + 2)^2 dx \\&= \pi \int_{-5}^{-2} (x^2 + 4x + 4) dx \\&= \pi \left[\frac{x^3}{3} + 2x^2 + 4x \right]_{-5}^{-2} \\&= 9\pi \text{ cubic units}\end{aligned}$$

3h

$$\begin{aligned}V &= \pi \int_{-3}^3 y^2 dx \\&= \pi \int_{-3}^3 \left(4 - \frac{4}{9}x^2 \right) dx \\&= \pi \left[4x - \frac{4}{27}x^3 \right]_{-3}^3 \\&= \frac{16\pi}{3} \text{ cubic units}\end{aligned}$$

Chapter 12 worked solutions – Further calculus

4a This is a cylinder of radius 1, height 3, hence

$$\begin{aligned}V &= \pi r^2 h \\&= \pi(1)^2(3) \\&= 3\pi \text{ cubic units}\end{aligned}$$

4b This is a cone with radius 2, height 4, hence

$$\begin{aligned}V &= \frac{1}{3}\pi r^2 h \\&= \frac{1}{3}\pi(4)^2(2) \\&= \frac{256}{3}\pi \text{ cubic units}\end{aligned}$$

4c

$$\begin{aligned}V &= \pi \int_2^5 x^2 dy \\&= \pi \int_2^5 y^4 dy \\&= \left[\frac{\pi y^5}{5} \right]_2^5 \\&= \frac{3093\pi}{5} \text{ cubic units}\end{aligned}$$

4d

$$\begin{aligned}V &= \pi \int_0^1 x^2 dy \\&= \pi \int_0^1 y dy \\&= \pi \left[\frac{y^2}{2} \right]_0^1 \\&= \frac{\pi}{2} \text{ cubic units}\end{aligned}$$

Chapter 12 worked solutions – Further calculus

4e This is a sphere of radius 4, hence

$$\begin{aligned}V &= \frac{4}{3}\pi r^3 \\&= \frac{4}{3}\pi(4)^3 \\&= \frac{256}{3}\pi \text{ cubic units}\end{aligned}$$

4f

$$\begin{aligned}V &= \pi \int_{-3}^0 x^2 dy \\&= \pi \int_{-3}^0 (-y^2)^2 dy \\&= \pi \int_{-3}^0 y^4 dy \\&= \pi \left[\frac{y^5}{5} \right]_{-3}^0 \\&= \frac{243\pi}{5} \text{ cubic units}\end{aligned}$$

4g

$$\begin{aligned}V &= \pi \int_0^2 x^2 dy \\&= \pi \int_0^2 (2y - y^2)^2 dy \\&= \pi \int_0^2 (4y^2 - 4y^3 + y^4) dy \\&= \pi \left[\frac{4}{3}y^3 - y^4 + \frac{y^5}{5} \right]_0^2 \\&= \frac{16\pi}{15} \text{ cubic units}\end{aligned}$$

Chapter 12 worked solutions – Further calculus

4h

$$\begin{aligned}V &= \int_{-1}^1 x^2 dy \\&= \int_{-1}^1 (4 - 4y^2) dy \\&= \left[4y - \frac{4y^3}{3} \right]_{-1}^1 \\&= \frac{16\pi}{3} \text{ cubic units}\end{aligned}$$

5

$$\begin{aligned}V &= \pi \int_0^1 y^2 dx \\&= \pi \int_0^1 e^{2x} dx \\&= \frac{\pi}{2} [e^{2x}]_0^1 \\&= \frac{\pi}{2} (e^2 - 1) \text{ cubic units}\end{aligned}$$

6

$$\begin{aligned}V &= \pi \int_2^4 y^2 dx \\&= \pi \int_2^4 \frac{1}{x} dx \\&= \pi [\ln x]_2^4 \\&= \pi [\ln 4 - \ln 2] \\&= \pi \ln 2 \text{ cubic units}\end{aligned}$$

Chapter 12 worked solutions – Further calculus

7

$$\begin{aligned}
 V &= \pi \int_1^6 x^2 dy \\
 &= \pi \int_1^6 \frac{1}{y} dy \\
 &= \pi [\ln y]_1^6 \\
 &= \pi \ln 6 \text{ cubic units}
 \end{aligned}$$

8a $\tan^2 x = \sec^2 x - 1$

8b

$$\begin{aligned}
 V &= \pi \int_1^{\frac{\pi}{3}} y^2 dx \\
 &= \pi \int_1^{\frac{\pi}{3}} \tan^2 x dx \\
 &= \pi \int_1^{\frac{\pi}{3}} (\sec^2 x - 1) dx \\
 &= \pi [\tan x - x]_1^{\frac{\pi}{3}} \\
 &= \pi \left(\sqrt{3} - \frac{\pi}{3} \right) \text{ cubic units}
 \end{aligned}$$

9a $\sin^2 x = \frac{1}{2} - \frac{1}{2} \cos 2x$

9b


$$\begin{aligned}
 V &= \pi \int_0^{\frac{\pi}{2}} y^2 dx \\
 &= \pi \int_0^{\frac{\pi}{2}} \sin^2 x dx \\
 &= \pi \int_0^{\frac{\pi}{2}} \sin^2 x dx \\
 &= \pi \int_0^{\frac{\pi}{2}} \left(\frac{1}{2} - \frac{1}{2} \cos 2x \right) dx
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned} &= \pi \left[\frac{1}{2}x - \frac{1}{4}\sin 2x \right]_0^{\frac{\pi}{2}} \\ &= \frac{\pi^2}{4} \text{ cubic units} \end{aligned}$$

10a


$$\begin{aligned} V &= \pi \int_3^5 y^2 dx \\ &= \int_3^5 (x+3)^2 dx \\ &= \pi \left[\frac{(x+3)^3}{3} \right]_3^5 \\ &= \frac{296\pi}{3} \text{ cubic units} \end{aligned}$$



Chapter 12 worked solutions – Further calculus

10b


$$\begin{aligned}
 V &= \pi \int_1^4 y^2 dx \\
 &= \pi \int_1^4 (1 + \sqrt{x})^2 dx \\
 &= \pi \int_1^4 1 + 2\sqrt{x} + x dx \\
 &= \pi \int_1^4 1 + 2x^{\frac{1}{2}} + x dx \\
 &= \pi \left[x + \frac{4}{3}x^{\frac{3}{2}} + \frac{x^2}{2} \right]_1^4 \\
 &= \frac{119\pi}{6} \text{ cubic units}
 \end{aligned}$$



10c


$$\begin{aligned}
 V &= \pi \int_0^5 y^2 dx \\
 &= \pi \int_0^5 (5x - x^2)^2 dx \\
 &= \pi \int_0^5 (25x^2 - 10x^3 + x^4) dx \\
 &= \pi \left[\frac{25x^3}{3} - \frac{10x^4}{4} + \frac{x^5}{5} \right]_0^5 \\
 &= \frac{625\pi}{6} \text{ cubic units}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus



10d

$$\begin{aligned}
 V &= \pi \int_{-1}^1 y^2 dx \\
 &= \pi \int_{-1}^1 (1 - x^2)^2 dx \\
 &= \pi \int_{-1}^1 (1 - 2x^2 + x^4) dx \\
 &= \pi \left[x - \frac{2x^3}{3} + \frac{x^5}{5} \right]_{-1}^1 \\
 &= \frac{16\pi}{15} \text{ cubic units}
 \end{aligned}$$




11a

$$\begin{aligned}
 V &= \pi \int_1^2 x^2 dy \\
 &= \pi \int_1^2 (y - 2)^2 dy \\
 &= \pi \int_1^2 (y^2 - 4y + 4) dy
 \end{aligned}$$


Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \pi \left[\frac{y^3}{3} - 2y^2 + 4y \right]_1^2 \\
 &= \frac{\pi}{3} \text{ cubic units}
 \end{aligned}$$



11b


$$\begin{aligned}
 V &= \pi \int_0^1 x^2 dy \\
 &= \pi \int_0^1 (y^2 + 1)^2 dy \\
 &= \pi \int_0^1 (y^4 + 2y^2 + 1) dy \\
 &= \pi \left[\frac{y^5}{5} + \frac{2y^3}{3} + y \right]_0^1 \\
 &= \frac{28\pi}{15} \text{ cubic units}
 \end{aligned}$$



Chapter 12 worked solutions – Further calculus

11c

$$\begin{aligned}
 V &= \pi \int_0^2 x^2 dy \\
 &= \pi \int_0^2 y^2(y-3)^2 dy \\
 &= \pi \int_0^2 y^2(y^2 - 6y + 9) dy \\
 &= \pi \int_0^2 (y^4 - 6y^3 + 9y^2) dy \\
 &= \pi \left[\frac{y^5}{5} - \frac{6y^4}{4} + 3y^3 \right]_0^2 \\
 &= \frac{81\pi}{10} \text{ cubic units}
 \end{aligned}$$




11d

$$\begin{aligned}
 V &= \pi \int_0^1 x^2 dy \\
 &= \pi \int_0^1 (1-y) dy \\
 &= \pi \left[y - \frac{y^2}{2} \right]_0^1
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{\pi}{2} \text{ cubic units}$$



12

$$\begin{aligned} V &= \pi \int_{\frac{1}{2}}^3 y^2 dx \\ &= \pi \int_{\frac{1}{2}}^3 \left(1 + \frac{1}{x}\right)^2 dx \\ &= \pi \int_{\frac{1}{2}}^3 \left(1 + \frac{2}{x} + \frac{1}{x^2}\right) dx \\ &= \pi \left[x + 2 \ln|x| - \frac{1}{x} \right]_{\frac{1}{2}}^3 \\ &= \frac{\pi}{6} (25 + 12 \ln 6) \text{ cubic units} \end{aligned}$$

13

$$\begin{aligned} V &= \pi \int_0^{\frac{1}{2}} y^2 dx \\ &= \pi \int_0^{\frac{1}{2}} (e^x - e^{-x})^2 dx \\ &= \pi \int_0^{\frac{1}{2}} (e^{2x} - 2 + e^{-2x}) dx \\ &= \pi \left[\frac{e^{2x}}{2} - 2x - \frac{e^{-2x}}{2} \right]_0^{\frac{1}{2}} \\ &= \frac{\pi}{2} (e - 2 - e^{-1}) \text{ cubic units} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

14

$$\begin{aligned}
 V &= \pi \int_{4\pi}^{6\pi} y^2 dx \\
 &= \pi \int_{4\pi}^{6\pi} \left(4 + 4 \sin \frac{x}{4}\right)^2 dx \\
 &= 16\pi \int_{4\pi}^{6\pi} \left(1 + 2 \sin \frac{x}{4} + \sin^2 \frac{x}{4}\right) dx \\
 &= 16\pi \int_{4\pi}^{6\pi} \left(1 + 2 \sin \frac{x}{4} + \frac{1}{2}(1 - \cos \frac{x}{2})\right) dx \\
 &= 16\pi \int_{4\pi}^{6\pi} \left(\frac{3}{2} + 2 \sin \frac{x}{4} - \cos \frac{x}{2}\right) dx \\
 &= 16\pi \left[x - 8 \cos \frac{x}{4} - 2 \sin \frac{x}{2}\right]_{4\pi}^{6\pi} \\
 &= 71.62 \text{ mL}
 \end{aligned}$$

15a

$$\begin{aligned}
 V &= \pi \int_0^{16} y^2 dx \\
 &= \pi \int_0^{16} 4^2 dx \\
 &= \pi [4^2 x]_0^{16} \\
 &= 256\pi \text{ cubic units}
 \end{aligned}$$

15b

$$\begin{aligned}
 V &= \pi \int_0^{16} x^2 dy \\
 &= \pi \int_0^{16} y dy \\
 &= \pi \left[\frac{y^2}{2}\right]_0^{16} \\
 &= 128\pi \text{ cubic units}
 \end{aligned}$$

15c $V = 256\pi - 128\pi = 128\pi \text{ cubic units}$

Chapter 12 worked solutions – Further calculus

16a i

$$\begin{aligned}V &= \pi \int_0^2 y^2 dx \\&= \pi \int_0^2 x^4 dx \\&= \pi \left[\frac{x^5}{5} \right]_0 \\&= \frac{32\pi}{5} \text{ cubic units}\end{aligned}$$

16a ii

$$\begin{aligned}V &= \pi \int_0^4 x^2 dy \\&= \pi \int_0^4 y dy \\&= \pi \left[\frac{y^2}{2} \right] \\&= 8\pi \text{ cubic units}\end{aligned}$$

16b i

$$\begin{aligned}V &= \pi \times 5^2 \times 1 - \pi \int_0^1 y^2 dx \\&= \pi \times 5^2 \times 1 - \pi \int_0^1 25x^2 dx \\&= \pi \times 5^2 \times 1 - \pi \left[\frac{25x^3}{3} \right]_0 \\&= \frac{50\pi}{3} \text{ cubic units}\end{aligned}$$

Chapter 12 worked solutions – Further calculus

16b ii

$$\begin{aligned}
 V &= \pi \int_0^5 x^2 dy \\
 &= \pi \int_0^5 \left(\frac{y}{5}\right)^2 dy \\
 &= \frac{\pi}{25} \int_0^5 y^2 dy \\
 &= \frac{\pi}{25} \left[\frac{y^3}{3}\right]_0^5 \\
 &= \frac{5\pi}{3} \text{ cubic units}
 \end{aligned}$$

16c i

$$\begin{aligned}
 V &= \pi \int_0^4 y^2 dx \\
 &= \pi \int_0^4 x dx \\
 &= \pi \left[\frac{x^2}{2}\right]_0^4 \\
 &= 8\pi \text{ cubic units}
 \end{aligned}$$

16c ii

$$\begin{aligned}
 V &= \pi \times 4^2 \times 2 - \pi \int_0^2 x^2 dy \\
 &= \pi \times 4^2 \times 2 - \pi \int_0^2 y^4 dy \\
 &= \pi \times 4^2 \times 2 - \pi \left[\frac{y^5}{5}\right]_0^2 \\
 &= \frac{128\pi}{5} \text{ cubic units}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus


16d i

$$\begin{aligned}
 V &= \pi \times 4^2 \times 1 - \int_0^1 y^2 \, dx \\
 &= 16\pi - \pi \int_0^1 (x^2 + 3)^2 \, dx \\
 &= 16\pi - \pi \int_0^1 (x^4 + 6x^2 + 9) \, dx \\
 &= 16\pi - \pi \left[\frac{x^5}{5} + 2x^3 + 9x \right]_0^1 \\
 &= \frac{24\pi}{5} \text{ cubic units}
 \end{aligned}$$

16d ii

$$\begin{aligned}
 V &= \pi \int_3^4 x^2 \, dy \\
 &= \pi \int_3^4 (y - 3) \, dy \\
 &= \pi \left[\frac{y^2}{2} - 3y \right]_3^4 \\
 &= \frac{\pi}{2} \text{ cubic units}
 \end{aligned}$$

17a



Chapter 12 worked solutions – Further calculus


17b i

$$\begin{aligned}
 V &= \pi \int_0^1 y_1^2 dx - \pi \int_0^1 y_2^2 dx \\
 &= \pi \int_0^1 x^4 dx - \pi \int_0^1 x^6 dx \\
 &= \pi \int_0^1 (x^4 - x^6) dx \\
 &= \pi \left[\frac{x^5}{5} - \frac{x^7}{7} \right]_0^1 \\
 &= \frac{2\pi}{35} \text{ cubic units}
 \end{aligned}$$

17b ii

$$\begin{aligned}
 V &= \pi \int_0^1 x_1^2 dy - \pi \int_0^1 x_2^2 dy \\
 &= \pi \int_0^1 y^1 dy - \pi \int_0^1 y^{\frac{3}{2}} dy \\
 &= \pi \int_0^1 \left(y^1 - y^{\frac{3}{2}} \right) dy \\
 &= \pi \left[\frac{y^2}{2} - \frac{2}{5} y^{\frac{5}{2}} \right]_0^1 \\
 &= \frac{\pi}{10} \text{ cubic units}
 \end{aligned}$$

18a




Chapter 12 worked solutions – Further calculus

18b

$$\begin{aligned}
 V &= \pi \int_1^5 (y_1^2 - y_2^2) dx \\
 &= \pi \int_1^5 \left((6-x)^2 - \left(\frac{5}{x}\right)^2 \right) dx \\
 &= \pi \int_1^5 \left(36 - 12x + x^2 - \frac{25}{x^2} \right) dx \\
 &= \pi \left[36x - 6x^2 + \frac{x^3}{3} + \frac{25}{x} \right]_1^5 \\
 &= \frac{64\pi}{3} \text{ cubic units}
 \end{aligned}$$

19



$$\begin{aligned}
 V &= \pi \int_1^2 (y_1^2 - y_2^2) dx \\
 &= \pi \int_1^2 \left(1^2 - \left(2 - \frac{2}{x} \right)^2 \right) dx \\
 &= \pi \int_1^2 \left(1 - \frac{4}{x^2} - 4 + \frac{8}{x} \right) dx \\
 &= \pi \int_1^2 \left(-3 + \frac{4}{x^2} + \frac{8}{x} \right) dx \\
 &= \pi \left[-3x + \frac{4}{x} + 8 \ln x \right]_1^2 \\
 &= \pi(8 \ln 2 - 5) \text{ cubic units}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

20a $y = x^3 + 2$


$$\frac{dy}{dx} = 3x^2$$

When $x = 1, y = 3$ and $\frac{dy}{dx} = 3$. Hence the equation of the tangent is

$$y - 3 = 3(x - 1)$$

$$y = 3x$$

20b



Chapter 12 worked solutions – Further calculus

20c i

$$\begin{aligned}
 V &= \pi \int_{-2}^1 (y_1^2 - y_2^2) dx \\
 &= \pi \int_{-2}^1 (x^6 - 9x^2) dx \\
 &= \pi \left[\frac{x^7}{7} - 3x^3 \right]_{-2}^1 \\
 &= \frac{15\pi}{7} \text{ cubic units}
 \end{aligned}$$

20c ii

$$\begin{aligned}
 V &= \pi \int_{-6}^3 (x_1^2 - x_2^2) dy \\
 &= \pi \int_{-6}^3 \left(y^{\frac{2}{3}} - \frac{y^2}{9} \right) dy \\
 &= \pi \left[\frac{3}{2} y^{\frac{2}{3}} - \frac{y^3}{27} \right]_{-6}^3 \\
 &= \frac{2\pi}{5} \text{ cubic units}
 \end{aligned}$$

21

$$\begin{aligned}
 V &= \pi \times 2^2 \times \ln 2 - \pi \int_0^{\ln 2} x^2 dy \\
 &= 4\pi \ln 2 - \pi \int_0^{\ln 2} (e^y)^2 dy \\
 &= 4\pi \ln 2 - \pi \int_0^{\ln 2} e^{2y} dy \\
 &= 4\pi \ln 2 - \pi \left[\frac{e^{2y}}{2} \right]_0^{\ln 2} \\
 &= 4\pi \ln 2 - \pi \left(\frac{e^{2\ln 2}}{2} - \frac{e^0}{2} \right) \\
 &= 4\pi \ln 2 - \pi \left(\frac{e^{\ln 4}}{2} - \frac{e^0}{2} \right)
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned} &= 4\pi \ln 2 - \pi \left(\frac{4}{2} - \frac{1}{2} \right) \\ &= 4\pi \ln 2 - \pi \left(\frac{3}{2} \right) \\ &= \frac{\pi}{2} (8 \ln 2 - 3) \text{ cubic units} \end{aligned}$$

22a

$$\begin{aligned} V &= \pi \int_0^h y^2 dx \\ &= \pi \int_0^h \left(\frac{rx}{h} \right)^2 dx \\ &= \pi \int_0^h \frac{r^2 x^2}{h^2} dx \\ &= \pi \left[\frac{r^2 x^3}{3h^2} \right]_0^h \\ &= \pi \left[\frac{r^2 h^3}{3h^2} \right] \\ &= \frac{1}{3} \pi r^2 h \end{aligned}$$

22b

$$\begin{aligned} V &= \pi \int_0^h y^2 dx \\ &= \pi \int_0^h r^2 dx \\ &= \pi [r^2 x]_0^h \\ &= \pi r^2 h \end{aligned}$$

Chapter 12 worked solutions – Further calculus

22c i

$$\begin{aligned}
 V &= \pi \int_{-r}^r y^2 dx \\
 &= \pi \int_{-r}^r (r^2 - x^2) dx \\
 &= \pi \left[r^2x - \frac{x^3}{3} \right]_{-r}^r \\
 &= \pi \left[\left(r^2r - \frac{r^3}{3} \right) - \left(r^2(-r) - \frac{(-r)^3}{3} \right) \right] \\
 &= \frac{4}{3}\pi r^3
 \end{aligned}$$

22c ii

$$\begin{aligned}
 V &= \pi \int_{r-h}^r y^2 dx \\
 &= \pi \int_{r-h}^r r^2 - x^2 dx \\
 &= \pi \left[r^2x - \frac{x^3}{3} \right]_{r-h}^r \\
 &= \pi \left[\left(r^2r - \frac{r^3}{3} \right) - \left(r^2(r-h) - \frac{(r-h)^3}{3} \right) \right] \\
 &= \pi \left[\left(r^2r - \frac{r^3}{3} \right) - \left(r^3 - r^2h - \frac{r^3 - 3r^2h + 3rh^2 - h^3}{3} \right) \right] \\
 &= \frac{1}{3}\pi h^2(3r - h)
 \end{aligned}$$

23a $\sec \theta$

$$\begin{aligned}
 &= \sec \theta \times \frac{\sec \theta + \tan \theta}{\sec \theta + \tan \theta} \\
 &= \frac{\sec^2 \theta + \sec \theta \tan \theta}{\sec \theta + \tan \theta} \\
 &= \frac{\sec \theta \tan \theta + \sec^2 \theta}{\sec \theta + \tan \theta}
 \end{aligned}$$

$$\begin{aligned}
 &\int \sec \theta d\theta \\
 &= \int \frac{\sec \theta \tan \theta + \sec^2 \theta}{\sec \theta + \tan \theta} d\theta
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \ln(\sec \theta + \tan \theta) + C$$

23b

$$y = \frac{x}{\sqrt{x^2 + 16}}$$

$$y^2 = \frac{x^2}{x^2 + 16}$$

$$x^2 + 16 = \frac{x^2}{y^2}$$

$$16 = \frac{x^2}{y^2} - x^2$$

$$16 = x^2 \left(\frac{1}{y^2} - 1 \right)$$

$$x^2 = \frac{16}{\left(\frac{1}{y^2} - 1 \right)}$$

$$= \frac{16}{\left(\frac{1}{y^2} - \frac{y^2}{y^2} \right)}$$

$$= \frac{16}{\left(\frac{1 - y^2}{y^2} \right)}$$

$$= \frac{16y^2}{1 - y^2}$$

$$V = \pi \int_0^{\frac{1}{\sqrt{2}}} x^2 dy$$

$$= \pi \int_0^{\frac{1}{\sqrt{2}}} \frac{16y^2}{1 - y^2} dy$$

Let $y = \sin \theta$, $dy = \cos \theta d\theta$

$$\begin{aligned} V &= \pi \int_0^{\frac{1}{\sqrt{2}}} x^2 dy \\ &= \pi \int_0^{\frac{\pi}{4}} \frac{16 \sin^2 \theta}{1 - \sin^2 \theta} \cos \theta d\theta \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \pi \int_0^{\frac{\pi}{4}} \frac{16 \sin^2 \theta}{\cos^2 \theta} \cos \theta \, d\theta \\
 &= \pi \int_0^{\frac{\pi}{4}} \frac{16 \sin^2 \theta}{\cos \theta} \, d\theta \\
 &= \pi \int_0^{\frac{\pi}{4}} \frac{16(1 - \cos^2 \theta)}{\cos \theta} \, d\theta \\
 &= 16\pi \int_0^{\frac{\pi}{4}} (\sec \theta - \cos \theta) \, d\theta \\
 &= 16\pi [\ln(\sec \theta + \tan \theta) + \sin \theta]_0^{\frac{\pi}{4}} \\
 &= 16\pi(\sqrt{2} - \ln(\sqrt{2} + 1)) \text{ cubic units}
 \end{aligned}$$

24a

$$x^n = x^{n+1}$$

$$x^n - x^{n+1} = 0$$

$$x^n(1 - x) = 0$$

Hence $x = 0$ or 1 . Substituting back into the equations gives the points of intersection as $(0, 0)$ and $(1, 1)$.

24b

$$\begin{aligned}
 V &= \pi \int_0^1 f(x)^2 \, dx - \pi \int_0^1 f(x)^2 \, dx \\
 &= \pi \int_0^1 (x^n)^2 \, dx - \pi \int_0^1 (x^{n+1})^2 \, dx \\
 &= \pi \int_0^1 x^{2n} \, dx - \pi \int_0^1 x^{2n+2} \, dx \\
 &= \pi \left[\frac{x^{2n+1}}{2n+1} \right]_0^1 - \pi \left[\frac{x^{2n+3}}{2n+3} \right]_0^1 \\
 &= \pi \left(\frac{1}{2n+1} - \frac{1}{2n+3} \right)
 \end{aligned}$$

- 24c It is the cone formed by rotating the line $y = x$ from $x = 0$ to $x = 1$ about the x -axis.

Chapter 12 worked solutions – Further calculus


24d

$$\begin{aligned}
 & \lim_{n \rightarrow \infty} (V_1 + V_2 + V_3 + \cdots + V_n) \\
 &= \lim_{n \rightarrow \infty} \left(\pi \left(\frac{1}{2+1} - \frac{1}{2+3} \right) + \pi \left(\frac{1}{2(2)+1} - \frac{1}{2(2)+3} \right) + \pi \left(\frac{1}{2(3)+1} - \frac{1}{2(3)+3} \right) \right. \\
 &\quad \left. + \cdots + \pi \left(\frac{1}{2(n)+1} - \frac{1}{2(n)+3} \right) \right) \\
 &= \pi \lim_{n \rightarrow \infty} \left(\frac{1}{3} - \frac{1}{5} + \frac{1}{5} - \frac{1}{7} + \frac{1}{7} - \frac{1}{9} + \cdots + \frac{1}{2n+1} - \frac{1}{2n+3} \right) \\
 &= \pi \lim_{n \rightarrow \infty} \left(\frac{1}{3} - \frac{1}{2n+3} \right) \\
 &= \pi \left(\frac{1}{3} - 0 \right) \\
 &= \frac{\pi}{3} \text{ cubic units}
 \end{aligned}$$

24e

$$\begin{aligned}
 & \frac{1}{3 \times 5} + \frac{1}{5 \times 7} + \frac{1}{7 \times 9} + \cdots \\
 &= \frac{1}{2} \left(\frac{1}{3} - \frac{1}{5} \right) + \frac{1}{2} \left(\frac{1}{5} - \frac{1}{7} \right) + \frac{1}{2} \left(\frac{1}{7} - \frac{1}{9} \right) + \cdots \\
 &= \frac{1}{2} \left(\frac{1}{3} - \frac{1}{5} + \frac{1}{5} - \frac{1}{7} + \frac{1}{7} - \frac{1}{9} + \cdots \right) \\
 &= \frac{1}{2} \left(\frac{1}{3} \right) \\
 &= \frac{1}{6}
 \end{aligned}$$

25a



25b $y = -x^2 + 6x - 8$

$y = -x^2 + 6x - 9 + 1$

$y = -(x^2 - 6x + 9) + 1$

Chapter 12 worked solutions – Further calculus

$$y = -(x - 3)^2 + 1$$

$$(x - 3)^2 = 1 - y$$

$x - 3 = \sqrt{1 - y}$ for $3 \leq x \leq 4$ and $x - 3 = -\sqrt{1 - y}$ for $2 \leq x \leq 3$

Hence

$$x = 3 + \sqrt{1 - y} \text{ for } 3 \leq x \leq 4 \text{ and } x = 3 - \sqrt{1 - y} \text{ for } 2 \leq x \leq 3$$

25c

$$\begin{aligned} V &= \pi \int_0^1 (3 + \sqrt{1 - y})^2 dy - \pi \int_0^1 (3 - \sqrt{1 - y})^2 dy \\ &= \pi \int_0^1 \left((3 + \sqrt{1 - y})^2 - (3 - \sqrt{1 - y})^2 \right) dy \\ &= \pi \int_0^1 \left((9 + 6\sqrt{1 - y} + (1 - y)) - (9 - 6\sqrt{1 - y} + (1 - y)) \right) dy \\ &= \pi \int_0^1 (12\sqrt{1 - y}) dy \\ &= \int_0^1 (12\pi\sqrt{1 - y}) dy \\ &= \int_0^1 \left(12\pi(1 - y)^{\frac{1}{2}} \right) dy \\ &= 12\pi \left[-\frac{2}{3}(1 - y)^{\frac{3}{2}} \right]_0^1 \\ &= 12\pi \left(-\frac{2}{3} \times 0 + \frac{2}{3} \times 1 \right) \\ &= 8\pi \text{ cubic units} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

Solutions to Chapter review

1a

$$\begin{aligned} & \frac{d}{dx}(\sin^{-1} 3x) \\ &= \frac{1}{\sqrt{1 - (3x)^2}} \times \frac{d}{dx}(3x) \\ &= \frac{3}{\sqrt{1 - 9x^2}} \end{aligned}$$

1b

$$\begin{aligned} & \frac{d}{dx}\left(\tan^{-1}\left(\frac{x}{3}\right)\right) \\ &= \frac{1}{1 + \left(\frac{x}{3}\right)^2} \times \frac{d}{dx}\left(\frac{x}{3}\right) \\ &= \frac{1}{1 + \frac{x^2}{9}} \times \frac{1}{3} \\ &= \frac{3}{9 + x^2} \end{aligned}$$

1c

$$\begin{aligned} & \frac{d}{dx}(\cos^{-1}(1-x)) \\ &= \frac{-1}{\sqrt{1 - (1-x)^2}} \times \frac{d}{dx}(1-x) \\ &= \frac{-1}{\sqrt{1 - (1-2x+x^2)}} \times -1 \\ &= \frac{1}{\sqrt{2x-x^2}} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

1d

$$\begin{aligned}
 & \frac{d}{dx}(x^2 \tan^{-1} x) \\
 &= (\tan^{-1} x) \times \frac{d}{dx}(x^2) + x^2 \times \frac{d}{dx}(\tan^{-1} x) \\
 &= 2x \tan^{-1} x + x^2 \times \frac{1}{1+x^2} \\
 &= \frac{x^2}{1+x^2} + 2x \tan^{-1} x
 \end{aligned}$$

1e

$$\begin{aligned}
 & \frac{d}{dx}\left(\tan^{-1}\left(\frac{1}{2}x + 1\right)\right) \\
 &= \frac{1}{1+\left(\frac{1}{2}x+1\right)^2} \times \frac{d}{dx}\left(\frac{1}{2}x+1\right) \\
 &= \frac{1}{1+\frac{x^2}{4}+x+1} \times \frac{1}{2} \\
 &= \frac{1}{\frac{x^2}{4}+x+2} \times \frac{1}{2} \\
 &= \frac{4}{x^2+4x+8} \times \frac{1}{2} \\
 &= \frac{2}{x^2+4x+8}
 \end{aligned}$$

1f

$$\begin{aligned}
 & \frac{d}{dx}\left(\sin^{-1}\frac{1}{x}\right) \\
 &= \frac{1}{\sqrt{1-\left(\frac{1}{x}\right)^2}} \times \frac{d}{dx}\left(\frac{1}{x}\right) \\
 &= \frac{1}{\sqrt{1-\frac{1}{x^2}}} \times \left(-\frac{1}{x^2}\right) \\
 &= -\frac{1}{\sqrt{x^4}\sqrt{1-\frac{1}{x^2}}} \\
 &= -\frac{1}{\sqrt{x^4-x^2}}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

2a $y = \tan^{-1} x$

$$y' = \frac{1}{1+x^2} = (1+x^2)^{-1}$$

$$y'' = -1 \times (1+x^2)^{-2} \times 2x = -\frac{2x}{(1+x^2)^2}$$

Substituting for y' and y'' :

$$\begin{aligned} & (1+x^2)y'' + 2xy' \\ &= (1+x^2) \times -\frac{2x}{(1+x^2)^2} + 2x \times \frac{1}{(1+x^2)} \\ &= -\frac{2x}{(1+x^2)} + \frac{2x}{(1+x^2)} \\ &= 0 \end{aligned}$$

3a

$$\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}}$$

$$\begin{aligned} & \frac{d}{dx}(\sin^{-1} \sqrt{1-x^2}) \\ &= \frac{1}{\sqrt{1-(\sqrt{1-x^2})^2}} \times \frac{d}{dx}(\sqrt{1-x^2}) \\ &= \frac{1}{\sqrt{1-(1-x^2)}} \times \frac{1}{2} \times (1-x^2)^{-\frac{1}{2}} \times -2x \\ &= \frac{1}{\sqrt{x^2}} \times -\frac{x}{\sqrt{1-x^2}} \\ &= -\frac{1}{\sqrt{x^2}} \times \frac{x}{\sqrt{1-x^2}} \end{aligned}$$

For $0 \leq x \leq 1$, $\sqrt{x^2} = x$

Hence

$$\begin{aligned} & \frac{d}{dx}(\sin^{-1} \sqrt{1-x^2}) \\ &= -\frac{1}{x} \times \frac{x}{\sqrt{1-x^2}} \\ &= -\frac{1}{\sqrt{1-x^2}} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{d}{dx} (\cos^{-1} x)$$

- 3b The functions differ by at most a constant.

This constant is zero, so $\cos^{-1} x = \sin^{-1} \sqrt{1 - x^2}$ for $0 \leq x \leq 1$.

4a

$$\begin{aligned} & \int \frac{1}{3+x^2} dx \\ &= \frac{1}{3} \int \frac{1}{1 + \left(\frac{x}{\sqrt{3}}\right)^2} dx \\ &= \frac{\sqrt{3}}{3} \tan^{-1} \frac{x}{\sqrt{3}} + C \end{aligned}$$

4b

$$\begin{aligned} & \int \frac{1}{\sqrt{3-x^2}} dx \\ &= \int \frac{1}{\sqrt{3}} \frac{1}{\sqrt{1 - \left(\frac{x}{\sqrt{3}}\right)^2}} dx \\ &= \sin^{-1} \frac{x}{\sqrt{3}} + C \end{aligned}$$

4c

$$\begin{aligned} & \int \frac{1}{9+4x^2} dx \\ &= \frac{1}{4} \int \frac{1}{\frac{9}{4} + \left(\frac{x}{\frac{3}{2}}\right)^2} dx \\ &= \frac{1}{9} \int \frac{1}{1 + \left(\frac{2x}{3}\right)^2} dx \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{1}{6} \tan^{-1} \frac{2x}{3} + C$$

4d

$$\begin{aligned} & \int -\frac{1}{\sqrt{16 - 9x^2}} dx \\ &= \int -\frac{1}{4\sqrt{1 - \left(\frac{3x}{4}\right)^2}} dx \\ &= \frac{1}{3} \cos^{-1} \frac{3x}{4} + C \end{aligned}$$

5a

$$\begin{aligned} & \int_{\frac{1}{3}}^{\frac{1}{\sqrt{3}}} \frac{1}{1 + 9x^2} dx \\ &= \int_{\frac{1}{3}}^{\frac{1}{\sqrt{3}}} \frac{1}{1 + (3x)^2} dx \\ &= \frac{1}{3} [\tan^{-1} 3x]_{\frac{1}{3}}^{\frac{1}{\sqrt{3}}} \\ &= \frac{\pi}{36} \end{aligned}$$

5b

$$\begin{aligned} & \int_{-\frac{3}{4}}^{\frac{3}{4}} \frac{1}{\sqrt{3 - 4x^2}} dx \\ &= \int_{-\frac{3}{4}}^{\frac{3}{4}} \frac{1}{\sqrt{3}} \frac{1}{\sqrt{1 - \left(\frac{2x}{\sqrt{3}}\right)^2}} dx \\ &= \frac{1}{\sqrt{3}} \times \frac{\sqrt{3}}{2} \left[\sin^{-1} \frac{2x}{\sqrt{3}} \right]_{-\frac{3}{4}}^{\frac{3}{4}} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \frac{1}{2} \left[\sin^{-1} \frac{2x}{\sqrt{3}} \right]_{-\frac{3}{4}}^{\frac{3}{4}} \\
 &= \frac{\pi}{3}
 \end{aligned}$$

6a

$$\begin{aligned}
 &\int \cos^2 x \, dx \\
 &= \int \left(\frac{1}{2} + \frac{1}{2} \cos 2x \right) dx \\
 &= \frac{1}{2}x + \frac{1}{4} \sin 2x + C
 \end{aligned}$$

6b

$$\begin{aligned}
 &\int \sin^2 x \, dx \\
 &= \int \left(\frac{1}{2} - \frac{1}{2} \cos 2x \right) dx \\
 &= \frac{1}{2}x - \frac{1}{4} \sin 2x + C
 \end{aligned}$$

6c

$$\begin{aligned}
 &\int \cos^2 2x \, dx \\
 &= \int \left(\frac{1}{2} + \frac{1}{2} \cos 4x \right) dx \\
 &= \frac{1}{2}x + \frac{1}{8} \sin 4x + C
 \end{aligned}$$

6d

$$\begin{aligned}
 &\int \sin^2 4x \, dx \\
 &= \int \left(\frac{1}{2} - \frac{1}{2} \cos 8x \right) dx
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{1}{2}x - \frac{1}{16}\sin 4x + C$$

7a

$$\begin{aligned} & \int_0^{\frac{\pi}{3}} \sin^2 3x \, dx \\ &= \int_0^{\frac{\pi}{3}} \left(\frac{1}{2} - \frac{1}{2}\cos 6x \right) dx \\ &= \left[\frac{1}{2}x - \frac{1}{16}\sin 4x \right]_0^{\frac{\pi}{3}} \\ &= \frac{\pi}{6} \end{aligned}$$

7b

$$\begin{aligned} & \int_0^{\frac{\pi}{6}} \cos^2 \frac{1}{2}x \, dx \\ &= \int_0^{\frac{\pi}{6}} \left(\frac{1}{2} + \frac{1}{2}\sin x \right) dx \\ &= \left[\frac{1}{2}x - \frac{1}{2}\cos x \right]_0^{\frac{\pi}{6}} \\ &= \frac{1}{12}(\pi + 3) \end{aligned}$$

8

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \cos^2 x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1}{2} + \frac{1}{2}\cos 2x \right) dx = \left[\frac{1}{2}x + \frac{1}{4}\sin 2x \right]_0^{\frac{\pi}{2}} = \frac{\pi}{4} \\ \int_0^{\frac{\pi}{2}} \cos^2 2x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1}{2} + \frac{1}{2}\cos 4x \right) dx = \left[\frac{1}{2}x + \frac{1}{8}\sin 4x \right]_0^{\frac{\pi}{2}} = \frac{\pi}{4} \\ \int_0^{\frac{\pi}{2}} \cos^2 4x \, dx &= \int_0^{\frac{\pi}{2}} \left(\frac{1}{2} + \frac{1}{2}\cos 8x \right) dx = \left[\frac{1}{2}x + \frac{1}{16}\sin 8x \right]_0^{\frac{\pi}{2}} = \frac{\pi}{4} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

9a Let $u = 5x - 1, du = 5 dx$

$$\begin{aligned} & \int 5(5x - 1)^5 \\ &= \int u^5 du \\ &= \frac{u^6}{6} + C \\ &= \frac{(5x - 1)^6}{6} + C \end{aligned}$$

9b Let $u = x^2 + 2$

$$\begin{aligned} du &= 2x dx \\ \int 2x(x^2 + 2)^2 dx & \\ &= \int u^2 du \\ &= \frac{u^3}{3} + C \\ &= \frac{(x^2 + 2)^3}{3} + C \end{aligned}$$

9c Let $u = x^4 + 1, du = 4x^3$

$$\begin{aligned} \int \frac{4x^3}{(x^4 + 1)^2} dx & \\ &= \int \frac{1}{u^2} du \\ &= \int u^{-2} du \\ &= -u^{-1} + C \\ &= \frac{1}{x^4 + 1} + C \end{aligned}$$

Chapter 12 worked solutions – Further calculus

9d Let $u = 4x + 3, du = 4 dx$

$$\begin{aligned} & \int \frac{1}{\sqrt{4x+3}} dx \\ &= \int \frac{1}{\sqrt{u}} \times \frac{1}{4} du \\ &= \frac{1}{4} \int u^{-\frac{1}{2}} du \\ &= \frac{1}{4} \times 2u^{\frac{1}{2}} + C \\ &= \frac{1}{2} \sqrt{4x+3} + C \end{aligned}$$

9e Let $u = \sin x, du = \cos x dx$

$$\begin{aligned} & \int \sin^2 x \cos x dx \\ &= \int u^2 du \\ &= \frac{u^3}{3} + C \\ &= \frac{\sin^3 x}{3} + C \end{aligned}$$

9f Let $u = \tan x, du = \sec^2 x dx$

$$\begin{aligned} & \int \tan^3 x \sec^2 x dx \\ &= \int u^3 du \\ &= \frac{u^4}{4} + C \\ &= \frac{\tan^4 x}{4} + C \end{aligned}$$

Chapter 12 worked solutions – Further calculus

10a Let $u = 1 + x^3$, $du = 3x^2 dx$ and $\frac{1}{3}du = x^2 dx$

$$\int_{-1}^0 x^2(1+x^3)^4 dx$$

$$= \frac{1}{3} \int_0^1 u^4 du$$

$$= \frac{1}{3} \left[\frac{u^5}{5} \right]_0^1$$

$$= \frac{1}{15}$$

10b Let $u = \cos x$, $du = -\sin x$

$$\int_0^{\frac{\pi}{2}} \cos^3 x \sin x dx$$

$$= \frac{1}{4} [-u^4]_1^0$$

$$= -\frac{1}{4} (0^3 - 1^2)$$

$$= \frac{1}{4}$$

10c Let $u = x^2 - 1$, $du = 2x dx$ and $\frac{1}{2}du = x dx$

$$\int_1^{\sqrt{2}} x \sqrt{x^2 - 1} dx$$

$$= \frac{1}{2} \int_0^1 \sqrt{u} du$$

$$= \frac{1}{2} \int_0^1 u^{\frac{1}{2}} du$$

$$= \frac{1}{2} \left[\frac{2}{3} u^{\frac{3}{2}} \right]_0^1$$

$$= \frac{1}{2} \left(\frac{2}{3} - 0 \right)$$

Chapter 12 worked solutions – Further calculus

$$= \frac{1}{3}$$

10d Let $u = \ln x, du = \frac{1}{x} dx$

$$\int_1^e \frac{(\ln x)^2}{x} dx$$

$$= \int_0^1 u^2 du$$

$$= \left[\frac{1}{3} u^3 \right]_0^1$$

$$= \frac{1}{3}$$

10e Let $u = \frac{1}{x}, du = -\frac{1}{x^2} dx$

$$\int_{\frac{1}{2}}^1 \frac{e^{\frac{1}{x}}}{x^2} dx$$

$$= \int_2^1 -e^u du$$

$$= \int_1^2 e^u du$$

$$= [e^u]_1^2$$

$$= e^2 - e$$

10f Let $u = \tan 2x, du = 2 \sec^2 2x dx$

$$\int_0^{\frac{\pi}{8}} \frac{\sec^2 2x}{1 + \tan 2x} dx$$

$$= \frac{1}{2} \int_0^1 \frac{1}{1+u} du$$

$$= \frac{1}{2} [\ln|1+u|]_0^1$$

Chapter 12 worked solutions – Further calculus

$$= \frac{1}{2}(\ln 2 - \ln 1)$$

$$= \frac{1}{2}\ln 2$$

11a Let $x = u + 1, du = dx$

$$\begin{aligned} & \int \frac{x}{x-1} dx \\ &= \int \frac{u+1}{u} du \\ &= \int \left(1 + \frac{1}{u}\right) du \\ &= u + \ln|u| + C \\ &= x - 1 + \ln|x-1| + C \end{aligned}$$

11b Let $x = u - 2, du = dx$

$$\begin{aligned} & \int \frac{x-1}{\sqrt{x+2}} dx \\ &= \int \frac{u-3}{\sqrt{u}} du \\ &= \int (u^{\frac{1}{2}} - 3u^{-\frac{1}{2}}) du \\ &= \frac{2}{3}u^{\frac{3}{2}} - 6u^{\frac{1}{2}} + C \\ &= \frac{2}{3}(x+2)^{\frac{3}{2}} - 6\sqrt{x+2} + C \end{aligned}$$

11c Let $x = \frac{1}{2}u^2 - \frac{1}{2}, dx = u du$

Note that $2x + 1 = u^2$

$$\begin{aligned} & \int x\sqrt{2x+1} dx \\ &= \int \left(\frac{1}{2}u^2 - \frac{1}{2}\right) (\sqrt{u^2}) u du \\ &= \frac{1}{2} \int (u^4 - u^2) du \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \frac{1}{2} \left(\frac{u^5}{5} - \frac{u^3}{3} \right) + C \\
 &= \frac{1}{2} \left(\frac{(2x+1)^{\frac{5}{2}}}{5} - \frac{(2x+1)^{\frac{3}{2}}}{3} \right) + C \\
 &= \frac{1}{10} (2x+1)^{\frac{5}{2}} - \frac{1}{6} (2x+1)^{\frac{3}{2}} + C
 \end{aligned}$$

11d Let $x = (u-4)^2$, $dx = 2(u-4) du$

$$\begin{aligned}
 &\int \frac{1}{4+\sqrt{x}} dx \\
 &= \int \frac{2(u-4)}{4+u-4} du \\
 &= \int \left(2 - \frac{8}{u} \right) du \\
 &= 2u - 8 \ln|u| + C \\
 &= 2(\sqrt{x}+4) - 8 \ln(\sqrt{x}+4) + C
 \end{aligned}$$

12a Let $x = u+1$, $dx = du$

$$\begin{aligned}
 &\int_1^2 x(x-1)^4 dx \\
 &= \int_0^1 (u+1)u^4 du \\
 &= \int_0^1 u^5 + u^4 du \\
 &= \left[\frac{u^6}{6} + \frac{u^5}{5} \right]_0^1 \\
 &= \frac{1}{6} + \frac{1}{5} \\
 &= \frac{11}{30}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

12b Let $x = u - 3, dx = du$

$$\begin{aligned} & \int_1^5 \frac{x}{x+3} dx \\ &= \int_4^8 \frac{u-3}{u} du \\ &= \int_4^8 \left(1 - \frac{3}{u}\right) du \\ &= [u - 3 \ln|u|]_4^8 \\ &= (8 - 3 \ln 8) - (4 - 3 \ln 4) \\ &= 4 - 3 \ln 2 \end{aligned}$$

12c Let $x = u^2 - 1, dx = 2u du$

$$\begin{aligned} & \int_0^{15} \frac{x}{\sqrt{x+1}} dx \\ &= \int_1^4 \frac{u^2 - 1}{\sqrt{u^2}} \times 2u du \\ &= \int_1^4 (2u^2 - 2) du \\ &= \left[\frac{2}{3}u^3 - 2u \right]_1^4 \\ &= 36 \end{aligned}$$

12d Let $x = u^2 + 2, dx = 2u du$


$$\begin{aligned} & \int_2^3 \frac{1}{2}x\sqrt{x-2} dx \\ &= \int_0^1 \frac{1}{2}(u^2 + 2)\sqrt{u^2} \times 2u du \\ &= \int_0^1 (u^4 + 2u^2) du \\ &= \left[\frac{u^5}{5} + \frac{2}{3}u^3 \right]_0^1 \\ &= \frac{1}{5} + \frac{2}{3} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$= \frac{13}{15}$$

- 13a The domain is all x such that $9 - x \geq 0$ and hence the domain is $x \leq 9$. As the square root function always returns a positive number, the range is $y \geq 0$.

13b



13c

$$\begin{aligned} A &= \int_0^9 \sqrt{9-x} dx \\ &= \int_0^9 (9-x)^{\frac{1}{2}} dx \\ &= -\frac{4}{3} \left[(9-x)^{\frac{3}{2}} \right]_0^9 \\ &= 18 \text{ square units} \end{aligned}$$

13d i

$$\begin{aligned} V &= \pi \int_0^9 y^2 dx \\ &= \pi \int_0^9 (9-x) dx \\ &= \pi \left[9x - \frac{x^2}{2} \right]_0^9 \\ &= \frac{81\pi}{2} \text{ cubic units} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

13d ii

$$\begin{aligned}
 V &= \pi \int_0^3 x^2 dy \\
 &= \pi \int_0^3 (y^2 - 9)^2 dy \\
 &= \pi \int_0^3 (y^4 - 18y^2 + 81) dy \\
 &= \pi \left[\frac{y^5}{5} - 6y^3 + 81y \right]_0^3 \\
 &= \frac{648\pi}{5} \text{ cubic units}
 \end{aligned}$$

14

$$\begin{aligned}
 V &= \int_0^{3/4} \pi y^2 dy \\
 &= \int_0^{3/4} \pi \left(\frac{1}{\sqrt{4-x}} \right)^2 dx \\
 &= \int_0^{3/4} \frac{\pi}{4-x} dx \\
 &= -\pi [\ln|4-x|]_0^{3/4} \\
 &= 4\pi \ln 2 \text{ cubic units}
 \end{aligned}$$

15 $y^2 = 18(x - 6)$

$$x = \frac{y^2}{18} + 6$$

$$\begin{aligned}
 V &= \pi \int_{-6}^6 x^2 dy \\
 &= \pi \int_{-6}^6 \left(\frac{y^2}{18} + 6 \right)^2 dy \\
 &= \pi \int_{-6}^6 \left(\frac{y^4}{18^2} + \frac{12}{18} y^2 + 36 \right) dy
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

$$\begin{aligned}
 &= \pi \left[\frac{y^5}{5 \times 18^2} + \frac{12}{18 \times 3} y^3 + 36y \right]_{-6}^6 \\
 &= \frac{2688\pi}{5} \text{ cubic units}
 \end{aligned}$$

16 $\cos^2 2x = \frac{1}{2} + \frac{1}{2} \cos 4x$

16b


$$\begin{aligned}
 V &= \pi \int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} y^2 dx \\
 &= \pi \int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} \cos^2 2x dx \\
 &= \pi \int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} \left(\frac{1}{2} + \frac{1}{2} \cos 4x \right) dx \\
 &= \pi \left[\frac{1}{2}x + \frac{1}{8} \sin 4x \right]_{-\frac{\pi}{6}}^{\frac{\pi}{6}} \\
 &= \frac{\pi}{24} (4\pi + 3\sqrt{3}) \text{ cubic units}
 \end{aligned}$$

17

$$\begin{aligned}
 V &= \pi \int_1^3 y^2 dx \\
 &= \pi \int_1^3 (1 + e^{-x})^2 dx \\
 &= \pi \int_1^3 (1 + 2e^{-x} + e^{-2x}) dx \\
 &= \pi \left[x - 2e^{-x} - \frac{1}{2}e^{-2x} \right]_1^3 \\
 &= \pi \left(\left(3 - 2e^{-3} - \frac{1}{2}e^{-6} \right) - \left(1 - 2e^{-1} - \frac{1}{2}e^{-2} \right) \right) \\
 &\doteq 8.49 \text{ cubic units}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

18a



18b

$$\begin{aligned}
 V &= \pi \int_0^{\frac{\pi}{4}} y^2 dx \\
 &= \pi \int_0^{\frac{\pi}{4}} (1 - \tan x)^2 dx \\
 &= \pi \int_0^{\frac{\pi}{4}} (1 - 2 \tan x + \tan^2 x) dx \\
 &= \pi \int_0^{\frac{\pi}{4}} (1 - 2 \tan x + \sec^2 x - 1) dx \\
 &= \pi \int_0^{\frac{\pi}{4}} \left(-\frac{2 \sin x}{\cos x} + \sec^2 x \right) dx \\
 &= \pi [2 \ln |\cos x| + \tan x]_0^{\frac{\pi}{4}} \\
 &= \pi \left(\left(2 \ln \left(2^{-\frac{1}{2}} \right) + 1 \right) - (-2 \ln 1 + 0) \right) \\
 &= \pi ((-\ln 2 + 1) - (0 + 0)) \\
 &= \pi (1 - \ln 2) \text{ cubic units}
 \end{aligned}$$

Chapter 12 worked solutions – Further calculus

19a

$$\frac{dy}{dx} = 1 - \frac{1}{x^2}$$

$$\frac{d^2y}{dx^2} = \frac{2}{x^3}$$

Stationary points occur when $\frac{dy}{dx} = 0$

$$1 - \frac{1}{x^2} = 0$$

$$1 = \frac{1}{x^2}$$


$$x^2 = 1$$

$$x = \pm 1$$

When $x = -1, y = -2, \frac{d^2y}{dx^2} = -2 < 0$, hence the curve is concave down and this is a maximum.

When $x = 1, y = 2, \frac{d^2y}{dx^2} = 2 > 0$, hence the curve is concave up and this is a minimum.

Hence there is a minimum at $(1, 2)$ and a maximum at $(-1, -2)$.



Chapter 12 worked solutions – Further calculus

19b

$$\frac{5}{2} = x + \frac{1}{x}$$

$$5 = 2x + \frac{2}{x}$$

$$5x = 2x^2 + 2$$

$$2x^2 - 5x + 2 = 0$$

$$(2x - 1)(x - 2) = 0$$

Hence


$$x = \frac{1}{2} \text{ or } 2$$

19c

$$\begin{aligned} V &= \pi \int_{\frac{1}{2}}^2 \left(\frac{5}{2}\right)^2 dx - \pi \int_{\frac{1}{2}}^2 \left(x + \frac{1}{x}\right)^2 dx \\ &= \pi \int_{\frac{1}{2}}^2 \left(\frac{25}{4} - x^2 - 2 - \frac{1}{x^2}\right) dx \\ &= \pi \int_{\frac{1}{2}}^2 \left(\frac{17}{4} - x^2 - \frac{1}{x^2}\right) dx \\ &= \pi \left[\frac{17}{4}x - \frac{x^3}{3} + \frac{1}{x} \right]_{\frac{1}{2}}^2 \\ &= \pi \left(\left(\frac{17}{2} - \frac{8}{3} + \frac{1}{2}\right) - \left(\frac{17}{8} - \frac{1}{24} + 2\right) \right) \\ &= \frac{9\pi}{4} \text{ cubic units} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

20a <To come>



21a

$$\begin{aligned} V &= \pi \int_1^3 y^2 dx \\ &= \pi \int_1^3 (2^{x+1})^2 dx \\ &= \pi \int_1^3 2^{2x+2} dx \end{aligned}$$

21b

$$\begin{aligned} V &\doteq \pi \times \frac{3-1}{2(4)} (f(1) + f(3) + 2(f(1.5) + f(2) + f(2.5))) \\ &= \frac{\pi}{4} (2^2 + 2^8 + 2(2^5 + 2^6 + 2^7)) \\ &= 180\pi \text{ cubic units} \end{aligned}$$

Chapter 12 worked solutions – Further calculus

21c

$$\begin{aligned}V &= \pi \int_1^3 y^2 dx \\&= \pi \int_1^3 (2^{x+1})^2 dx \\&= \pi \int_1^3 2^{2x+2} dx \\&= \pi \int_1^3 e^{\ln 2^{2x+2}} dx \\&= \pi \int_1^3 e^{(2x+2)\ln 2} dx \\&= \frac{\pi}{2 \ln 2} [e^{(2x+2)\ln 2}]_1^3 \\&= \frac{120\pi}{\ln 2} \\&\doteq 173\pi \text{ cubic units}\end{aligned}$$

The exact answer is smaller than the trapezoidal-rule approximation because the curve is concave up.