

Lecture 1: Finding Solutions

Introduction to Data Analysis Techniques

1 Data Analysis in Science

Definition 1.1 (Analytical Method and Analytical Solution). [def:analytical-method] When we use *analytical methods* to solve a problem, we are manipulating the mathematical expressions until unknowns are being isolated into a standard mathematical form.

An *analytical solution* is a solution for a problem such that the unknowns are expressed in a closed form.

Example 1.1 (Analytical Solution of a Quadratic Equation). [eg:quadratic-equation] Consider the quadratic equation

$$ax^2 + bx + c = 0, \quad (1)$$

where a , b , and c are constants. The analytic solution for x can be found using the quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}. \quad (2)$$

Definition 1.2 (Numerical Method and Numerical Solution). [def:numerical-method] *Numerical methods* are methods to obtain numerical solutions.

A *numerical solution* is a solution for a problem such that the unknowns are expressed in an numerical form.

Example 1.2 (Numerical Solution of a Trigonometric Equation). [eg:trig-equation] Consider the trigonometric equation

$$x = \beta \cos x \quad (3)$$

where β is a constant.

It is very difficult to find an analytical solution for this equation using algebraic manipulations. However, we can use numerical methods to find an approximate solution for specific β values.

The numerical solutions can be

$$\beta = 1, \quad x \approx 0.73909, \quad \beta = 0.5, \quad x \approx 0.4508$$

Remark 1.1 (Advantages and Disadvantages of Numerical Methods). [rem:numerical-methods]

1. For numerical methods, we must have all parameters of the problem known.
2. If parameters change, solution must be re-done.
3. Solutions may be not exact.
4. Numerical solutions often considered inferior to analytic solutions, if exist. *ie, we should use analytical methods whenever possible or too difficult.*

2 Using Taylor Series to Approximate Functions

Definition 2.1 (Taylor Series). [def:taylor-series] For a function $f(x)$ that is infinitely differentiable at a point a , then:

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

Remark 2.1 (Taylor Series Approximation). [rem:taylor-series-approx] It is impossible to compute the infinite sum in practice. When we find the analytical solution of a problem, we can use the first few terms of the Taylor Series to approximate the function.

The more terms we use, the better the approximation.

Example 2.1 (Electric Field of a Dipole for $x \gg a$). [[eg:dipole-field](#)] Consider the electric field on the axis of a dipole with charges $+q$ at $x = a$ and $-q$ at $x = -a$:

$$E = \frac{q}{4\pi\epsilon_0} \left[\frac{1}{(x-a)^2} - \frac{1}{(x+a)^2} \right]$$

Take the Taylor series expansion of E for $x \gg a$:

$$E = \frac{q}{4\pi\epsilon_0 x^2} \left[4\frac{a}{x} + 8\left(\frac{a}{x}\right)^3 + 12\left(\frac{a}{x}\right)^5 + \dots \right]$$

Keeping the first two terms gives an approximation for $x \gg a$:

$$E \approx \frac{q}{4\pi\epsilon_0 x^2} \left[4\frac{a}{x} + 8\left(\frac{a}{x}\right)^3 \right]$$

Which means, for $x \gg a$, $E \propto x^{-3}$.

3 Fixed Point Iteration Method

Definition 3.1 (Fixed Point Iteration Method). [[def:fixed-point-iteration-method](#)] The idea for *fixed point iteration method* is to first find a function $g(x)$ such that root r is a fixed point of $g(x)$, i.e., $g(r) = r$

Convention 3.1 (Procedure of Fixed Point Iteration Method). [[conv:fixed-point-iteration](#)] The classical procedure of fixed point iteration method is as follows:

1. Find a function $g(x)$ such that root r is a fixed point of $g(x)$, i.e., $g(r) = r$.
2. Choose an initial guess x_0 .
3. Choose a maximum number of iterations N
4. Iterate $x_{n+1} = g(x_n)$ for N times

Remark 3.1 (Error Analysis of Fixed Point Iteration Method). [[rem:fixed-point-iteration-error](#)] By using this method, we usually believe that the root, r , is close to the initial guess, x_0 . Also, it should be within x_n and x_{n+1} for all n .

Which means, the error of the n -th iteration is $|x_n - r| \leq |x_{n+1} - x_n|$.

Pf. Convergence of Fixed Point Iteration Method. □

Convention 3.2 (Optimized Fixed Point Iteration). [[conv:optimized-fpi](#)] We may want to set an error tolerance, tol , and stop the iteration when $|x_{n+1} - x_n| < \text{tol}$.

1. Find a function $g(x)$ such that root r is a fixed point of $g(x)$, i.e., $g(r) = r$.
2. Choose an initial guess x_0 .
3. Choose a maximum number of iterations N
4. Set an error tolerance, tol
5. Iterate $x_{n+1} = g(x_n)$
6. Check if $|x_{n+1} - x_n| < \text{tol}$, if yes, stop the iteration and return the value.
7. If the maximum number of iterations is reached, stop the iteration and return the value.
(For some algorithms, we may want to return an error message instead.)

```
def find_fixed_point(
    g,                # g(x): fixed point r satisfies g(r) = r
    x0: float,        # initial guess
    tol: float = 1e-5, # stop when |x_new - x| < tol
    max_iter: int = 100, # safety cap on iterations
```

```

) -> float:
    x = x0
    for i in range(max_iter):
        x_new = g(x)
        print(f"Iteration {i+1}: x = {x_new}, g(x) = {g(x_new)}")
        if abs(x_new - x) < tol:
            print(f"Converged after {i+1} iterations.")
            return x_new
        x = x_new
    raise ValueError("Fixed point not found within max_iter.")

```

Example 3.1 (Convergent Case: Fixed Point Iteration on $x - \cos x = 0$). [[eg:cos-fixed-point](#)] Apply Optimized Fixed Point Iteration to $f(x) = x - \cos x = 0$.

1. Rewrite $f(x) = 0$ as $x = \cos x$, so $g(x) = \cos x$ and $g(r) = r$.
2. Initial guess: $x_0 = 1$.
3. Maximum iterations: $N = 100$.
4. Error tolerance: $\text{tol} \in \{5 \times 10^{-2}, 1 \times 10^{-3}\}$ (two runs for comparison).
5. Iterate $x_{n+1} = \cos x_n$.
6. Stop when $|x_{n+1} - x_n| < \text{tol}$ and return x_{n+1} .
7. If n reaches N , stop and return the last x_{n+1} .

In Python:

```

from math import cos
from find_fixed_point import find_fixed_point

# f(x) = x - cos(x)
# We want to solve f(x) = 0, i.e., find the root of f(x).
# So our equation is x - cos(x) = 0
# Which means that we can rewrite it as x = cos(x)
# So we want to find the fixed point of the function g(x) = cos(x)
def g(x):
    return cos(x)

# Initial guess: x = 1
x = 1

# Find the fixed point of g(x) = cos(x) using the fixed-point iteration method
r = find_fixed_point(g, x, tol=5e-2)
print(r)


# Find the fixed point of g(x) = cos(x) using the fixed-point iteration method with a
# smaller tolerance
r = find_fixed_point(g, x, tol=1e-3)
print(r)

```


Both runs spiral into the fixed point $r \approx 0.739$; a tighter tolerance just adds more steps near r . The top row shows the iteration geometrically on the (x, y) plane; the bottom row shows the same iterates plotted against their step number.

- Figure 3.1a: x-axis is x , y-axis is y ; shows $y = g(x) = \cos x$ (blue), the diagonal $y = x$ (dashed), and an orange path that traces the iteration
- Figure 3.1b: same axes as (a); tighter $\text{tol} = 1 \times 10^{-3}$ makes the orange path pack more segments around the fixed point r .
- Figure 3.1c: x-axis is the iteration index n , y-axis is the iterate x_n ; shows the sequence of iterates as damped oscillation for $\text{tol} = 5 \times 10^{-2}$, with the dashed line marking the final x_n .
- Figure 3.1d: same axes as (c); tighter run $\text{tol} = 1 \times 10^{-3}$ settles visibly onto $r \approx 0.739$.


Remark 3.2 (Convergence not Guaranteed). [[rem:fixed-point-iteration-convergence](#)] The convergence of fixed point iteration is not guaranteed.




(a) Trace Iteration, $\text{tol} = 5 \times 10^{-2}$.



(b) Trace Iteration, $\text{tol} = 1 \times 10^{-3}$.



(c) x_n vs. n , $\text{tol} = 5 \times 10^{-2}$.



(d) x_n vs. n , $\text{tol} = 1 \times 10^{-3}$.

Figure 3.1: Convergent case: fixed point iteration on $x - \cos x = 0$ from $x_0 = 1$.

Example 3.2 (Divergent Case: $x - 1.5 \cos x = 0$). [eg:divergent-fpi] Same procedure as example 3.1, but with $g(x) = 1.5 \cos x$ and $x_0 = 1$. It diverges and oscillates between ≈ 0.123 and ≈ 1.489 ; both runs hit $N = 100$ without ever satisfying the stop condition.

Remark 3.3 (Advantage and Disadvantage of Fixed Point Iteration Method). [rem:fpi-evaluation]

1. Advantage: It is easy to implement.
2. Disadvantage: The convergence is not guaranteed.


Usually used for *good initial guess* and *good $g(x)$ function*.

4 Bisection Method


Definition 4.1 (Bisection Method). [def:bisection-method] The idea of *bisection method* is when we know that a root r is in the interval $[a, b]$, then we can find the midpoint of the range and evaluate which part of the interval contains the root. We can then repeat this process until we find a sufficiently accurate approximation of the root.

Convention 4.1 (Procedure of Bisection Method). [conv:bisection-method-procedure] The procedure of bisection method is as follows:


1. Find an interval $[a, b]$ such that $\exists r \in [a, b]$ st $f(r) = 0$ and $f(a)f(b) < 0$.




(a) Trace Iteration, $\text{tol} = 5 \times 10^{-2}$.



(b) Trace Iteration, $\text{tol} = 1 \times 10^{-3}$.



(c) x_n vs. n , $\text{tol} = 5 \times 10^{-2}$.



(d) x_n vs. n , $\text{tol} = 1 \times 10^{-3}$.

Figure 3.2: Divergent case: fixed point iteration on $x - 1.5 \cos x = 0$ from $x_0 = 1$.

2. Set the maximum number of iterations, N .
3. Find the midpoint of the interval, $c = \frac{a+b}{2}$.
4. Evaluate $f(c)$.
5. If $f(a)$ and $f(c)$ have opposite signs, then set $b = c$; otherwise, set $a = c$.
6. If the maximum number of iterations is reached, stop the iteration and return the value. Otherwise goto step 3.

Pf. Bisection Method Convergence. Continuous Function, IVT. TODO □

Remark 4.1 (Error Analysis of Bisection Method). [rem:bisection-method-error] By using this method, we usually believe that the root, r , is in the interval $[a, b]$.

Which means, we can have a one to one map between the number of iterations and the error tolerance, tol ,

$$\text{tol} = |a - b| \times 2^{-n}$$

5 Newton-Raphson Method

Definition 5.1 (Newton-Raphson Method). [def:newton-raphson-method] (newton-raphson-method) The idea of *Newton-Raphson method* is to use the slope of the function at a point to find the root of the function.

Convention 5.1 (Procedure of Newton-Raphson Method). [conv:newton-raphson-method-procedure]

1. Take an initial guess x_0 .
2. Set the maximum number of iterations, N .
3. Calculate the derivative of the function f at the chosen point, x_n , call it as t_n .
4. Define $y = t_n(x_n - x_{n-1}) + f(x_{n-1})$. *Intuition: This is the equation of the tangent line at x_{n-1} .*
5. Find the root of the tangent line, x_n , by solving $y = 0$.
6. If the maximum number of iterations is reached, stop the iteration and return the value. Otherwise goto step 3.

Remark 5.1 (Error Analysis of Newton-Raphson Method). [rem:newton-raphson-method-error] Let the r to be the root, ie, $f(r) = 0$. So, the error of each step is $e_n = x_n - r$.