Lecture 4: Limit of Sequence

Mathematical Analysis

1 Convergent Sequences

Definition 1.1 (Convergent Sequence)**.** [def:convergent-sequence] A sequence $a : \mathbb{N}_1 \to F$, where F is a ordered field, is said to be *convergent* if

$$\exists \ell \in F,\, \forall \varepsilon \in F_+,\, \exists N \in \mathbb{N}_1,\, \forall n \in \mathbb{N}_1,\, n \geq N,\, d(a_n, \ell) < \varepsilon$$

Remark 1.1 (The ε-N Challenge). [rem:epsilon-n-challenge] A value ℓ passes the ε-N challenge if for any proposed $\varepsilon > 0$, there exists an index $N \in \mathbb{N}_1$ such that for all $n \in \mathbb{N}_1$ with $n \geq N$, $d(a_n, \ell) < \varepsilon$

Remark 1.2. In the The ε-N Challenge, the value of N may depend on the choice of ε. And the value N may be not unique (ie, $N' > N$ also works)

Theorem 1.1 (Uniqueness of Limit)**.** [thm:limit-uniqueness] *Let F be an ordered field, $a : \mathbb{N}_1 \to F$, and $\ell_1, \ell_2 \in F$. Then*

$$\begin{gathered}\big(\forall \varepsilon \in F_+,\, \exists N \in \mathbb{N}_1,\, \forall n \in \mathbb{N}_1,\, n \geq N,\, d(a_n, \ell_1) < \varepsilon\big) \\ \wedge \\ \big(\forall \varepsilon \in F_+,\, \exists N \in \mathbb{N}_1,\, \forall n \in \mathbb{N}_1,\, n \geq N,\, d(a_n, \ell_2) < \varepsilon\big) \\ \implies \ell_1 = \ell_2\end{gathered}$$

Proof. Assume for contradiction that $\ell_1 \neq \ell_2$. Set

$$\varepsilon = \frac{d(\ell_1, \ell_2)}{2}.$$

Since $\ell_1 \neq \ell_2$, we have $\varepsilon > 0$
By the hypothesis for ℓ_1, there exists $N_1 \in \mathbb{N}_1$ such that for all $n \geq N_1$, $d(a_n, \ell_1) < \varepsilon$. Similarly, there exists $N_2 \in \mathbb{N}_1$ such that for all $n \geq N_2$, $d(a_n, \ell_2) < \varepsilon$
Let $N = \max(N_1, N_2)$. For any $n \geq N$, both hypotheses hold:

$$d(a_n, \ell_1) < \varepsilon \quad \text{and} \quad d(a_n, \ell_2) < \varepsilon$$

By the triangle inequality for distance applied via the intermediate point a_n,

$$d(\ell_1, \ell_2) \leq d(\ell_1, a_n) + d(a_n, \ell_2) = d(a_n, \ell_1) + d(a_n, \ell_2),$$

where the last equality uses symmetry of distance. Combining,

$$d(\ell_1, \ell_2) \leq d(a_n, \ell_1) + d(a_n, \ell_2) < 2\varepsilon = d(\ell_1, \ell_2),$$

a contradiction. Hence $\ell_1 = \ell_2$ □

Remark 1.3 (Compact Form Using $\exists!$). [rem:limit-uniqueness-compact] Combined with the existence from the definition of convergent sequence, this theorem states: if a is convergent, then there exists a unique such ℓ:

$$\exists! \ell \in F,\, \forall \varepsilon \in F_+,\, \exists N \in \mathbb{N}_1,\, \forall n \in \mathbb{N}_1,\, n \geq N,\, d(a_n, \ell) < \varepsilon.$$

Here $\exists!$ is shorthand for "existence and uniqueness"; it should be formally introduced in `lec01` as a FOL convention before being used

2 Convergence Challenge Functions

Definition 2.1 (Convergence Challenge Function)**.** [def:convergence-challenge-function] Let F be an ordered field. A function $g : F \to F$ is a *convergence challenge function* if it satisfies two conditions:

- *Positivity on positives*: $\forall \gamma \in F_+,\ g(\gamma) > 0$
- *Arbitrarily small values*: $\forall \delta \in F_+,\ \exists \gamma \in F_+,\ g(\gamma) < \delta$

Remark 2.1 (Intuition: Covering Arbitrarily Small Positives)*.* [rem:challenge-intuition]

Theorem 2.1 (Convergence Challenge Replacement)**.** [thm:convergence-challenge-replacement] *Let F be an ordered field, $a : \mathbb{N}_1 \to F$, and let g be a convergence challenge function. Then*

$$\big(\exists \ell \in F,\ \forall \varepsilon \in F_+,\ \exists N \in \mathbb{N}_1,\ \forall n \in \mathbb{N}_1,\ n \geq N,\ d(a_n, \ell) < \varepsilon\big)$$

$$\iff$$

$$\big(\exists \ell \in F,\ \forall \varepsilon \in F_+,\ \exists N \in \mathbb{N}_1,\ \forall n \in \mathbb{N}_1,\ n \geq N,\ d(a_n, \ell) < g(\varepsilon)\big)$$

Proof. Fix the witness $\ell \in F$ in either direction
($\Rightarrow$) For $\varepsilon > 0$, the positivity-on-positives condition gives $g(\varepsilon) > 0$. Apply the definition of convergent sequence at the challenge $g(\varepsilon)$
($\Leftarrow$) For $\varepsilon > 0$, the challenge-function property of g gives $\gamma > 0$ with $g(\gamma) < \varepsilon$. The hypothesis at γ gives N such that

$$d(a_n, \ell) < g(\gamma) < \varepsilon \quad \text{for all } n \geq N$$

□

Lemma 2.1 (Restricting to Small Tolerances)**.** [lem:restricting-to-small-tolerances] *Let F be an ordered field and fix $r \in F_+$. The function*

$$g : F \to F \equiv \gamma \mapsto \min(\gamma, r/2)$$

is a convergence challenge function

Proof. For every $\gamma > 0$, $0 < g(\gamma) \leq r/2 < r$. For every $\delta > 0$, choose $\gamma \equiv \delta/2 > 0$; then $g(\gamma) \leq \delta/2 < \delta$ □

Remark 2.2 (Restricting the Tolerance)*.* [rem:restricting-the-tolerance] Since $0 < g(\gamma) < r$ for $\gamma > 0$, challenge replacement lets us check $\forall\, 0 < \varepsilon < r$ instead of $\forall \varepsilon > 0$, for any fixed $r > 0$, eg, $\forall\, 0 < \varepsilon < 1$

Lemma 2.2 (Linear Scaling)**.** [lem:linear-is-challenge] *For any $c \in F$ with $c > 0$, the function $g(\gamma) = c\gamma$ is a convergence challenge function*

Proof. We verify both conditions
Positivity on positives. For any $\gamma \in F$ with $\gamma > 0$, both $c > 0$ and $\gamma > 0$, so $g(\gamma) = c\gamma > 0$
Arbitrarily small values. Fix any $\delta \in F$ with $\delta > 0$. We seek $\gamma \in F$ with $\gamma > 0$ and $g(\gamma) < \delta$
Take $\gamma \equiv \delta/(c+1)$, which is derived in remark 2.3. Since $\delta > 0$ and $c + 1 > 0$ (because $c > 0$), we have $\gamma > 0$
Now $g(\gamma) = c\gamma = c\delta/(c+1)$. Since $c < c + 1$, we have $c/(c+1) < 1$, hence $c\delta/(c+1) < \delta$ □

Remark 2.3 (Deriving the Witness for Linear Scaling)*.* [rem:linear-scaling-witness-derivation] To find the witness γ used in the proof of lemma 2.2, we start from the target inequality $g(\gamma) < \delta$ and manipulate it to isolate γ:

$$g(\gamma) < \delta \iff c\gamma < \delta \iff \gamma < \frac{\delta}{c}.$$

Any γ in the open interval $\left(0, \frac{\delta}{c}\right)$ will therefore satisfy $g(\gamma) < \delta$. To pick a concrete such γ, we use the fact that $c + 1 > c$, which gives

$$\frac{\delta}{c+1} < \frac{\delta}{c}.$$

Hence we may take $\gamma \equiv \delta/(c+1)$ as the witness

Remark 2.4 (Why $c \leq 0$ is Excluded). [rem:linear-scaling-cnonpos] If $c \leq 0$, then for any $\gamma \in F$ with $\gamma > 0$, $g(\gamma) = c\gamma \leq 0$. This violates the positivity-on-positives condition, so g is not a convergence challenge function

Lemma 2.3 (Power). [lem:power-is-challenge] *For any positive integer k, the function $g(\gamma) = \gamma^k$ is a convergence challenge function*

Proof. Both conditions follow by induction on k. Positivity: $\gamma^k > 0$ for $\gamma > 0$. For arbitrarily small values, take $\gamma \equiv \min(\delta/2,\, 1/2)$, so $\gamma > 0$ and $\gamma \leq 1$; induction gives $\gamma^k \leq \gamma \leq \delta/2 < \delta$ □

Lemma 2.4 (Summation). [lem:sum-is-challenge] *Let f, g be convergence challenge functions, and suppose both are monotone increasing (i.e. $x \leq y \implies f(x) \leq f(y)$, and similarly for g). Then $f + g$, defined by $(f + g)(\gamma) \equiv f(\gamma) + g(\gamma)$, is a convergence challenge function*

Proof. Positivity is immediate from $f(\gamma), g(\gamma) > 0$. For $\delta > 0$, pick $\gamma_1, \gamma_2 > 0$ with $f(\gamma_1), g(\gamma_2) < \delta/2$; set $\gamma \equiv \min(\gamma_1, \gamma_2)$. Monotonicity gives $(f + g)(\gamma) \leq f(\gamma_1) + g(\gamma_2) < \delta$ □

Lemma 2.5 (Composition). [lem:challenge-composition] *Let f, g be convergence challenge functions, and suppose f is monotone increasing (i.e. $x \leq y \implies f(x) \leq f(y)$). Then $f \circ g$ is a convergence challenge function*

Proof. Positivity: $f(g(\gamma)) > 0$ since $g(\gamma) > 0$ and f preserves positivity. For $\delta > 0$, pick $\eta > 0$ with $f(\eta) < \delta$, then $\gamma > 0$ with $g(\gamma) < \eta$. Monotonicity of f gives $(f \circ g)(\gamma) \leq f(\eta) < \delta$ □

Theorem 2.2 (Closed Form of Convergence). [thm:convergence-closed-form] *Let F be an ordered field, let $a : \mathbb{N}_1 \to F$. Then*

$$\left(\exists \ell \in F,\, \forall \varepsilon \in F_+,\, \exists N \in \mathbb{N}_1,\, \forall n \in \mathbb{N}_1,\, n \geq N,\, d(a_n, \ell) < \varepsilon\right)$$

$$\iff$$

$$\left(\exists \ell \in F,\, \forall \varepsilon \in F_+,\, \exists N \in \mathbb{N}_1,\, \forall n \in \mathbb{N}_1,\, n \geq N,\, d(a_n, \ell) \leq \varepsilon\right)$$

Proof. ($\Rightarrow$) If $d(a_n, \ell) < \varepsilon$, then $d(a_n, \ell) \leq \varepsilon$
($\Leftarrow$) By lemma 2.2 with $c = 2$, $g(\varepsilon) = 2\varepsilon$ is a convergence challenge function. From the closed-form hypothesis, for every $\varepsilon > 0$ there exists N with $d(a_n, \ell) \leq \varepsilon$ for all $n \geq N$. Since $\varepsilon > 0 \implies \varepsilon < 2\varepsilon$, we obtain $d(a_n, \ell) \leq \varepsilon < 2\varepsilon$, hence

$$\exists \ell \in F,\, \forall \varepsilon \in F_+,\, \exists N \in \mathbb{N}_1,\, \forall n \in \mathbb{N}_1,\, n \geq N,\, d(a_n, \ell) < 2\varepsilon. \tag{$*$}$$

For any $\varepsilon > 0$, applying $(*)$ at $\varepsilon/2 > 0$ gives N such that $d(a_n, \ell) < 2 \cdot (\varepsilon/2) = \varepsilon$ for all $n \geq N$ □

3 Limit of a Sequence

Definition 3.1 (Limit of a Sequence). [def:limit] Let F be an ordered field. We say the sequence $a : \mathbb{N}_1 \to F$ *has a limit of* $\ell \in F$ if

$$\forall \varepsilon \in F_+,\, \exists N \in \mathbb{N}_1,\, \forall n \in \mathbb{N}_1,\, n \geq N,\, d(a_n, \ell) < \varepsilon$$

Redirect (Converges To) [def:converges-to]
Moved to definition 3.1.

Definition 3.3 (Set of Convergent Sequences)**.** [def:convergent-sequences-set] Let F be an ordered field. The *set of convergent sequences in* F is

$$\mathrm{Conv}(F) \equiv \{a : \mathbb{N}_1 \to F \mid a \text{ is convergent}\}$$

LaTeX: \Conv (F)

Definition 3.4 (Limit Operator)**.** [def:limit-op] Let F be an ordered field. The *limit operator* is the function

$$\lim : \mathrm{Conv}(F) \to F \equiv a \mapsto \text{the unique } \ell \in F \text{ such that } a\text{has a limit of}\ell$$

We write

$$\lim(a) = \ell$$

The input is the whole sequence, rather than an individual term

LaTeX: \seqlim (a)

Redirect (Limit of a Sequence) [def:limit-of-sequence]
Moved to definition 3.4.

Remark 3.1 (Well-Definedness of the Limit)*.* [rem:limit-well-defined] The function lim in definition 3.4 is well-defined for two reasons:

- *Every input has an output*: for every $a \in \mathrm{Conv}(F)$, the definition of convergent sequence guarantees some $\ell \in F$ with $a$has a limit ofℓ
- *Every input has exactly one output*: by theorem 1.1, this ℓ is unique

Theorem 3.1 (Addition of the Limit)**.** [thm:limit-addition] *Let F be an ordered field, and let $a, b \in \mathrm{Conv}(F)$. Then*

$$\lim(n \mapsto a_n + b_n) = \lim(a) + \lim(b)$$

Proof. We want to prove

$$\forall \varepsilon \in F_+,\ \exists N \in \mathbb{N}_1,\ \forall n \in \mathbb{N}_1,\ n \geq N \implies$$
$$|a_n + b_n - \lim(a) - \lim(b)| < \varepsilon$$

According to the definition of a limit, for every $\varepsilon \in F_+$, there exist $N_a, N_b \in \mathbb{N}_1$ such that, setting $N_M \equiv \max(N_a, N_b)$, both estimates hold for $n \geq N_M$:

$$|a_n - \lim(a)| < \varepsilon \quad \wedge \quad |b_n - \lim(b)| < \varepsilon$$

Therefore, by the triangle inequality for distance,

$$\begin{aligned} |a_n + b_n - \lim(a) - \lim(b)| &\leq |a_n - \lim(a)| + |b_n - \lim(b)| \\ &< \varepsilon + \varepsilon = 2 \cdot \varepsilon \end{aligned}$$

Notice that

$$g : F \to F \equiv \varepsilon \mapsto 2 \cdot \varepsilon$$

is a convergence challenge function by linear scaling. Hence the challenge replacement theorem gives

$$\lim(n \mapsto a_n + b_n) = \lim(a) + \lim(b)$$

□

Theorem 3.2 (Multiplication of the Limit). [thm:limit-multiplication] *Let F be an ordered field, and let $a, b \in \mathrm{Conv}(F)$. Then*

$$\lim(n \mapsto a_n \cdot b_n) = \lim(a) \cdot \lim(b)$$

Proof. For $n \mapsto a_n \cdot b_n$, we want to show that $\lim(a) \cdot \lim(b)$ is its limit, ie,

$$\begin{gathered}\forall \varepsilon \in F_+,\, \exists N \in \mathbb{N}_1,\, \forall n \in \mathbb{N}_1,\, n \geq N \implies \\ |a_n \cdot b_n - \lim(a) \cdot \lim(b)| < \varepsilon\end{gathered}$$

According to the definition of a limit, for every $\varepsilon \in F_+$, there exist $N_a, N_b \in \mathbb{N}_1$ such that

$$\begin{aligned} n \geq N_a &\implies |a_n - \lim(a)| < \varepsilon, \\ n \geq N_b &\implies |b_n - \lim(b)| < \varepsilon \end{aligned}$$

Let $N_M \equiv \max(N_a, N_b)$. Then, for every $n \geq N_M$, both inequalities hold
Adding and subtracting $\lim(a) \cdot b_n$, we have

$$\begin{aligned} &|a_n \cdot b_n - \lim(a) \cdot \lim(b)| \\ &\quad = |a_n \cdot b_n - \lim(a) \cdot b_n + \lim(a) \cdot b_n - \lim(a) \cdot \lim(b)| \\ &\quad \leq |a_n \cdot b_n - \lim(a) \cdot b_n| + |\lim(a) \cdot b_n - \lim(a) \cdot \lim(b)| \\ &\quad = |b_n| \cdot |a_n - \lim(a)| + |\lim(a)| \cdot |b_n - \lim(b)| \end{aligned}$$

The inequality uses the triangle inequality for distance
Therefore, when $n \geq N_M$,

$$\begin{aligned} |a_n \cdot b_n - \lim(a) \cdot \lim(b)| &\leq |b_n| \cdot \varepsilon + |\lim(a)| \cdot \varepsilon \\ &= \varepsilon \cdot \big(|b_n| + |\lim(a)|\big) \end{aligned}$$

Since all convergent sequences are bounded, choose a strict bound $M' \in F$ such that $\forall n \in \mathbb{N}_1$, $|b_n| < M'$, and let $M \equiv \max(M', 1)$. This bound is independent of ε. Hence $M \geq 1$ and

$$|a_n \cdot b_n - \lim(a) \cdot \lim(b)| < \varepsilon \cdot \big(M + |\lim(a)|\big)$$

Notice that

$$g : F \to F \equiv \varepsilon \mapsto \varepsilon \cdot \big(M + |\lim(a)|\big)$$

is a convergence challenge function by linear scaling, since $M + |\lim(a)| > 0$. We have shown that

$$\begin{gathered}\forall \varepsilon \in F_+,\, \exists N_M \in \mathbb{N}_1,\, \forall n \in \mathbb{N}_1,\, n \geq N_M \implies \\ |a_n \cdot b_n - \lim(a) \cdot \lim(b)| < \varepsilon \cdot \big(M + |\lim(a)|\big)\end{gathered}$$

Therefore, by the challenge replacement theorem,

$$\lim(n \mapsto a_n \cdot b_n) = \lim(a) \cdot \lim(b)$$

□

Theorem 3.3 (Scalar Multiplication of the Limit). [thm:limit-scalar-multiplication] *Let F be an ordered field, let $a \in \mathrm{Conv}(F)$, and let $c \in F$. Then*

$$\lim(n \mapsto c \cdot a_n) = c \cdot \lim(a)$$

Proof. Define the constant sequence

$$p : \mathbb{N}_1 \to F \equiv n \mapsto c$$

For every $\varepsilon \in F_+$ and every $n \in \mathbb{N}_1$,

$$|p_n - c| = 0 < \varepsilon$$

Hence $p \in \mathrm{Conv}(F)$ and $\lim(p) = c$
By the definition of p,

$$\forall n \in \mathbb{N}_1, \quad c \cdot a_n = p_n \cdot a_n$$

By multiplication of limits, the sequence $n \mapsto p_n \cdot a_n$ is convergent, and

$$\begin{aligned} \lim(n \mapsto c \cdot a_n) &= \lim(n \mapsto p_n \cdot a_n) \\ &= \lim(p) \cdot \lim(a) \\ &= c \cdot \lim(a) \end{aligned}$$

□

Theorem 3.4 (Linearity of the Limit)**.** [thm:limit-linearity] *Let F be an ordered field, let $a, b \in \mathrm{Conv}(F)$, and let $c, d \in F$. Then*

$$\lim(n \mapsto c \cdot a_n + d \cdot b_n) = c \cdot \lim(a) + d \cdot \lim(b)$$

Proof. This follows from addition of limits and scalar multiplication of limits □

Lemma 3.1 (Reciprocal of the Limit)**.** [lem:limit-reciprocal] *Let F be an ordered field, and let $b \in \mathrm{Conv}(F)$ with $\forall n \in \mathbb{N}_1$, $b_n \neq 0$ and $\lim(b) \neq 0$. Then*

$$\lim\left(n \mapsto \frac{1}{b_n}\right) = \frac{1}{\lim(b)}$$

Proof. Define

$$p : \mathbb{N}_1 \to F \equiv n \mapsto \frac{1}{b_n}$$

We check whether $1/\lim(b)$ is a limit of p
For $n \in \mathbb{N}_1$,

$$\begin{aligned} |p_n - \frac{1}{\lim(b)}| &= |\frac{1}{b_n} - \frac{1}{\lim(b)}| \\ &= |\frac{\lim(b) - b_n}{b_n \cdot \lim(b)}| \\ &= \frac{|\lim(b) - b_n|}{|b_n \cdot \lim(b)|} \end{aligned}$$

To bound the denominator below, set $M \equiv |\lim(b)|/2 > 0$. By the convergence of b, choose $N_0 \in \mathbb{N}_1$ such that $n \geq N_0 \implies |b_n - \lim(b)| < M$. For $n \geq N_0$, the triangle inequality gives

$$|\lim(b)| \leq |\lim(b) - b_n| + |b_n| < M + |b_n|$$

Since $|\lim(b)| = 2 \cdot M$, we have $|b_n| > M$. So

$$\exists M \in F_+, \exists N_0 \in \mathbb{N}_1, \forall n \in \mathbb{N}_1, n \geq N_0 \implies |b_n| > M$$

Fix $\varepsilon \in F_+$ and choose $N_b \in \mathbb{N}_1$ so that $n \geq N_b \implies |b_n - \lim(b)| < \varepsilon$. Set $N \equiv \max(N_0, N_b)$.
For $n \geq N$,

$$|b_n \cdot \lim(b)| > M \cdot |\lim(b)|, \qquad |\lim(b) - b_n| < \varepsilon$$

Therefore,

$$\frac{|\lim(b) - b_n|}{|b_n \cdot \lim(b)|} < \frac{\varepsilon}{M \cdot |\lim(b)|}$$

Since $M \cdot |\lim(b)| > 0$, the function

$$g : F \to F \equiv \varepsilon \mapsto \frac{\varepsilon}{M \cdot |\lim(b)|}$$

is a convergence challenge function by linear scaling. Hence

$$\forall \varepsilon \in F_+, \exists N \in \mathbb{N}_1, \forall n \in \mathbb{N}_1, n \geq N \implies |p_n - \frac{1}{\lim(b)}| < g(\varepsilon)$$

By challenge replacement, $p \in \mathrm{Conv}(F)$ and

$$\lim(p) = \frac{1}{\lim(b)}$$

□

Theorem 3.5 (Division of the Limit). [thm:limit-division] *Let F be an ordered field, and let $a, b \in \mathrm{Conv}(F)$ with $\forall n \in \mathbb{N}_1, b_n \neq 0$ and $\lim(b) \neq 0$. Then*

$$\lim\left(n \mapsto \frac{a_n}{b_n}\right) = \frac{\lim(a)}{\lim(b)}$$

Proof. Define

$$p : \mathbb{N}_1 \to F \equiv n \mapsto \frac{1}{b_n}$$

By the reciprocal lemma, $p \in \mathrm{Conv}(F)$ and $\lim(p) = 1/\lim(b)$. Since $a_n/b_n = a_n \cdot p_n$ for every $n \in \mathbb{N}_1$, multiplication of limits gives

$$\begin{aligned}\lim\left(n \mapsto \frac{a_n}{b_n}\right) &= \lim(n \mapsto a_n \cdot p_n) \\ &= \lim(a) \cdot \lim(p) \\ &= \frac{\lim(a)}{\lim(b)}\end{aligned}$$

□

Theorem 3.6 (Order Preservation of Limits). [thm:limit-order-preservation] *Let F be an ordered field, and let $a, b \in \mathrm{Conv}(F)$ with $\forall n \in \mathbb{N}_1$, $a_n \leq b_n$. Then*

$$\lim(a) \leq \lim(b)$$

4 Bounded Sequences

Definition 4.1 (Bounded Sequence). [def:bounded-sequence] Let F be an ordered field. A sequence $a : \mathbb{N}_1 \to F$ is *bounded* if

$$\exists M \in F, \forall n \in \mathbb{N}_1, \quad d(a(n), 0) \leq M$$

Definition 4.2 (Set of Bounded Sequences). [def:bounded-sequences-set] Let F be an ordered field. The set of bounded sequences in F is

$$\mathrm{Bdd}(F) \equiv \{a : \mathbb{N}_1 \to F \mid a \text{ is bounded}\}$$

`LaTeX: \Bdd (F)`

Theorem 4.1 (Convergent Sequences are Bounded). [thm:convergent-sequences-are-bounded] *Let F be an ordered field and let $a : \mathrm{Conv}(F)$. Then a is bounded, so $a \in \mathrm{Bdd}(F)$*

Proof. Suppose the convergent sequence a, $\lim a = \ell$
Take $\varepsilon = 1$, then there exists $N \in \mathbb{N}_1$ such that for all $n \in \mathbb{N}_1$ with $n > N$, $d(a_n, \ell) < 1$
According to the triangle inequality for distance, and $d(a_n, \ell) < 1$, we have

$$d(0, a_n) \leq d(0, \ell) + d(\ell, a_n) < d(0, \ell) + 1$$

Which means for all $n \in \mathbb{N}_1$ with $n > N$, $d(0, a_n) < d(0, \ell) + 1$
And for $n \in \mathbb{N}_1$ with $n \leq N$, we have $d(0, a_n) \leq \max\{d(0, a_1), d(0, a_2), ..., d(0, a_N)\}$
So, we can take $M = \max\{\max\{d(0, a_1), d(0, a_2), ..., d(0, a_N)\}, d(0, \ell) + 1\}$. Hence M exists for all convergent sequences a □

Remark 4.1. Although Convergent $\implies$ Bounded, the converse is not true. A bounded sequence may not be convergent
For example, the sequence $a_n = (-1)^n$ is bounded (since $d(a_n, 0) = 1$ for all n), but not convergent

5 Monotone Convergence Theorem

Definition 5.1 (Increasing Sequence). [def:increasing-sequence] A sequence $a : \mathbb{N}_1 \to F$ is said to be *increasing* if

$$\forall n \in \mathbb{N}_1, a_n \leq a_{n+1}$$

Definition 5.2 (Strictly Increasing Sequence). [def:strictly-increasing-sequence] A sequence $a : \mathbb{N}_1 \to F$ is said to be *strictly increasing* if

$$\forall n \in \mathbb{N}_1, a_n < a_{n+1}$$

Definition 5.3 (Decreasing Sequence). [def:decreasing-sequence] A sequence $a : \mathbb{N}_1 \to F$ is said to be *decreasing* if

$$\forall n \in \mathbb{N}_1, a_n \geq a_{n+1}$$

Definition 5.4 (Strictly Decreasing Sequence). [def:strictly-decreasing-sequence] A sequence $a : \mathbb{N}_1 \to F$ is said to be *strictly decreasing* if

$$\forall n \in \mathbb{N}_1, a_n > a_{n+1}$$

Lemma 5.1 (Strictly Increasing Extends to All Indices). [lem:strictly-increasing-extends] *Let $a : \mathbb{N}_1 \to F$ be strictly increasing. Then $\forall n, m \in \mathbb{N}_1$, $n > m \implies a_n > a_m$*

Lemma 5.2 (Increasing Extends to All Indices). [lem:increasing-extends] *Let $a : \mathbb{N}_1 \to F$ be increasing. Then $\forall n, m \in \mathbb{N}_1$, $n \geq m \implies a_n \geq a_m$*

Lemma 5.3 (Strictly Decreasing Extends to All Indices). [lem:strictly-decreasing-extends] *Let $a : \mathbb{N}_1 \to F$ be strictly decreasing. Then $\forall n, m \in \mathbb{N}_1$, $n > m \implies a_n < a_m$*

Lemma 5.4 (Decreasing Extends to All Indices). [lem:decreasing-extends] *Let $a : \mathbb{N}_1 \to F$ be decreasing. Then $\forall n, m \in \mathbb{N}_1$, $n \geq m \implies a_n \leq a_m$*

Definition 5.5 (Monotone Sequence). [def:monotone-sequence] A sequence $a : \mathbb{N}_1 \to F$ is said to be *monotone* if it is either increasing or decreasing

Theorem 5.1 (Monotone Convergence Theorem). [thm:mct] *Let F be a complete ordered field, and let $a : \mathrm{Bdd}(F)$ be a monotone sequence. Then a is convergent*

Proof. Suppose a is a bounded sequence in F and increasing sequence, which means exist $x = \sup\{a_n : n \in \mathbb{N}_1\}$ (by the Axiom of Completeness, applied to the nonempty set $\{a_n : n \in \mathbb{N}_1\}$ bounded above in F), and $\forall n, m \in \mathbb{N}_1,\ n \leq m \implies a_n \leq a_m$
According to the ε criterion for supremum, $\forall n \in \mathbb{N}_1,\ a_n \leq x$ and $\forall \varepsilon \in F_+,\ \exists N \in \mathbb{N}_1,\ a_N > x - \varepsilon$
Adding the increasing property, ie

$$N \leq n \implies a_N \leq a_n \leq x \implies (a_N > x - \varepsilon \implies a_n \geq a_N > x - \varepsilon)$$

We can have

$$\forall \varepsilon \in F_+,\ \exists N \in \mathbb{N}_1,\ \forall n \in \mathbb{N}_1 \geq N,\ a_n > x - \varepsilon$$

Also,

$$a_n \leq x \ \wedge\ \varepsilon > 0 \implies a_n < x + \varepsilon$$

$$a_n > x - \varepsilon \ \wedge\ a_n < x + \varepsilon \implies x - \varepsilon < a_n < x + \varepsilon \implies d(a_n, x) < \varepsilon$$

Notice that such x is the supremum of the upper bounded sequence a, it exists.

$$\exists x \in F,\ \forall \varepsilon \in F_+,\ \exists N \in \mathbb{N}_1,\ \forall n \in \mathbb{N}_1 \geq N,\ d(a_n, x) < \varepsilon$$

Which means it is a convergent sequence and its limit is x
If instead a is decreasing, the same argument applies with $x \equiv \inf\{a_n : n \in \mathbb{N}_1\}$ (which exists by the Axiom of Completeness, since $\{a_n : n \in \mathbb{N}_1\}$ is nonempty and bounded below), using the dual ε-criterion for the infimum □

6 Comparison of Sequences

Theorem 6.1 (Comparison for Monotone Sequences)**.** [thm:monotone-comparison-increasing] *Let F be a complete ordered field, and let $s, t : \mathbb{N}_1 \to F$ be increasing with $s_n \leq t_n$ for all $n \in \mathbb{N}_1$. If t is convergent, then*

$$s \text{ is convergent and } \lim(s) \leq \lim(t)$$

Proof. **TODO** □

Theorem 6.2 (Comparison for Decreasing Sequences)**.** [thm:monotone-comparison-decreasing] *Let F be a complete ordered field, and let $s, t : \mathbb{N}_1 \to F$ be decreasing with $s_n \leq t_n$ for all $n \in \mathbb{N}_1$. If s is convergent, then*

$$t \text{ is convergent and } \lim(s) \leq \lim(t)$$

Proof. Apply theorem 6.1 to the increasing sequences $n \mapsto -t_n$ and $n \mapsto -s_n$ □

7 Squeeze Theorem

Theorem 7.1 (Squeeze Theorem)**.** [thm:squeeze-theorem] *Let F be an ordered field, and let $a, x, b : \mathbb{N}_1 \to F$ with $a_n \leq x_n \leq b_n$ for all $n \in \mathbb{N}_1$. If a and b are convergent with $\lim(a) = \lim(b) = L$, then x is convergent and $\lim(x) = L$*

Proof. **TODO** □