

Lecture 1: Incompleteness of Rational Numbers

Mathematical Analysis

1 Numbers

Definition 1.1 (Natural Numbers). [def:natural-number] Let S denote the *successor*. The *natural numbers* are inductively generated by

$$\mathbb{N} := 0 \mid S(n : \mathbb{N}).$$

These constructors exhaust $\mathbb{N}$.

Theorem 1.1 (Natural Number Induction). [thm:natural-number-induction] Let $P : \mathbb{N} \rightarrow \text{Prop}$. If

$$P(0) \quad \text{and} \quad \forall n : \mathbb{N}, P(n) \implies P(S(n)),$$

then $P(n)$ holds for every $n : \mathbb{N}$.

Proof. Let $A = \{n : \mathbb{N} \mid P(n)\}$. The hypotheses say that A contains 0 and is closed under S . Because the constructors in definition 1.1 exhaust $\mathbb{N}$, we have $A = \mathbb{N}$. $\square$

Remark 1.1 (Number Systems as Types and Sets). [rem:number-systems-types-and-sets] Throughout these notes, a symbol denoting a number system may represent either a type of numbers or the set of all numbers of that type. Thus $n : \mathbb{N}$ is the type-theoretic form, whereas $n \in \mathbb{N}$ is the corresponding set-theoretic form. The intended interpretation is determined by context.

Convention 1.1 (Positive Natural Numbers). [conv:positive-natural-numbers] We write

$$\mathbb{N}_1 = \mathbb{N} \setminus \{0\}$$

when zero must be excluded, such as for one-based indices or denominators.

Definition 1.2 (Integers). [def:integer] The *integers* are the whole numbers, including the positive integers, the negative integers, and zero:

$$\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}.$$

Definition 1.3 (Rational Numbers). [def:rational-numbers] A real number is called *rational* if it can be expressed as a quotient of two integers. The set of rational numbers is

$$\mathbb{Q} = \left\{ \frac{p}{q} : p, q \in \mathbb{Z}, q \neq 0 \right\}.$$

Proposition 1.1 (Unique Reduced Form). [prop:unique-reduced-rational-form] For every $x \in \mathbb{Q}$, there exists a unique pair $p, q \in \mathbb{Z}$ such that

$$x = \frac{p}{q}, \quad q > 0, \quad \gcd(|p|, q) = 1.$$

Remark 1.2. [rem:number-set-inclusions] Under the usual identifications, $\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q}$.

Definition 1.4 (Floor). [def:floor] For $x \in \mathbb{R}$, the *floor* $\lfloor x \rfloor$ is the unique $m \in \mathbb{Z}$ satisfying

$$m \leq x < m + 1.$$

Definition 1.5 (Ceiling). [def:ceiling] For $x \in \mathbb{R}$, the *ceiling* $\lceil x \rceil$ is the unique $m \in \mathbb{Z}$ satisfying

$$m - 1 < x \leq m.$$

Theorem 1.2 (Well-Definedness of Floor and Ceiling). [thm:floor-ceiling-well-defined] For every $x \in \mathbb{R}$:

- there exists a unique $m \in \mathbb{Z}$ satisfying $m \leq x < m + 1$, namely $\lfloor x \rfloor$;
- there exists a unique $m \in \mathbb{Z}$ satisfying $m - 1 < x \leq m$, namely $\lceil x \rceil$.

Proof. Existence and uniqueness are deferred. Sketch: existence follows from the Archimedean property combined with the well-ordering of $\mathbb{N}$ (equivalent to Natural Number Induction); uniqueness is immediate from the strict inequalities. A rigorous proof belongs in abstract algebra / order-theory prerequisites. $\square$

2 Fields

Definition 2.1 (Closed Operation). [def:closed-operation] An operation $*$ is said to be *closed* on a set A if for all $a, b \in A$, we have $a * b \in A$.

Definition 2.2 (Field). [def:field] A *field* is a set F equipped with addition and multiplication such that both operations are

- closed operations on F ,
- associative, meaning for all $a, b, c \in F$, we have $(a + b) + c = a + (b + c)$ and $(a \cdot b) \cdot c = a \cdot (b \cdot c)$,
- commutative, meaning for all $a, b \in F$, we have $a + b = b + a$ and $a \cdot b = b \cdot a$,
- have distinct identity elements $0, 1 \in F$, meaning $a + 0 = a$, $a \cdot 1 = a$, and $0 \neq 1$,
- each element has an additive inverse, such that for each $a \in F$, there exists $-a \in F$ with $a + (-a) = 0$,
- each non-zero element has a multiplicative inverse, such that for each $a \in F$ with $a \neq 0$, there exists $a^{-1} \in F$ with $a \cdot a^{-1} = 1$, and
- multiplication distributes over addition, meaning $a \cdot (b + c) = a \cdot b + a \cdot c$.

Proposition 2.1 ($\mathbb{Q}$ Is a Field). [prop:rational-are-a-field] The rational numbers with their usual addition and multiplication form a field.

Remark 2.1 ($\mathbb{N}$ and $\mathbb{Z}$ Are Not Fields). [rem:natural-and-integers-not-fields] The natural numbers and the integers are not fields because they do not have multiplicative inverses for every non-zero element.

For a counter example, consider the integer 2. There is no integer x such that $2 \cdot x = 1$

3 Ordered Set

Definition 3.1 (Transitive). [def:transitive] An ordering on a set is *transitive* if

$$\forall x, y, z \in F, \text{ if } x \leq y \text{ and } y \leq z, \text{ then } x \leq z$$

Definition 3.2 (Reflexive). [def:reflexive] An ordering on a set is *reflexive* if

$$\forall x \in F, x \leq x$$

Definition 3.3 (Antisymmetric). [def:antisymmetric] An ordering on a set is *antisymmetric* if

$$\forall x, y \in F, (x \leq y) \wedge (y \leq x) \implies x = y$$

Definition 3.4 (Partial Order). [def:partial-order] A binary relation $\leq$ on a set F is a *partial order* if it is reflexive, antisymmetric, and transitive.

Definition 3.5 (Totality). [def:totality] An ordering on a set F is *total* if every pair of elements is comparable:

$$\forall x, y \in F, \quad x \leq y \vee y \leq x.$$

Definition 3.6 (Total Order). [def:total-order] A partial order $\leq$ on F is a *total order* if it also satisfies totality.

Convention 3.1 (Order Notation). [conv:order-notation] For $a, b \in F$, define

$$a \geq b \iff b \leq a, \quad a < b \iff a \leq b \wedge a \neq b, \quad a > b \iff b < a.$$

Definition 3.7 (Binary Maximum). [def:binary-max] Let X be equipped with a total order $\leq$. For $a, b \in X$, define

$$\max(a, b) := \begin{cases} a & \text{if } a \geq b, \\ b & \text{otherwise.} \end{cases}$$

Definition 3.8 (Binary Minimum). [def:binary-min] Let X be equipped with a total order $\leq$. For $a, b \in X$, define

$$\min(a, b) := \begin{cases} a & \text{if } a \leq b, \\ b & \text{otherwise.} \end{cases}$$

Definition 3.9 (Maximum of a Set). [def:set-maximum] Let X be equipped with a total order $\leq$, and let $A \subseteq X$ be nonempty. An element $m \in A$ is a *maximum* of A , written $m = \max A$, if

$$\forall x \in A, m \geq x.$$

Definition 3.10 (Minimum of a Set). [def:set-minimum] Let X be equipped with a total order $\leq$, and let $A \subseteq X$ be nonempty. An element $m \in A$ is a *minimum* of A , written $m = \min A$, if

$$\forall x \in A, m \leq x.$$

Proposition 3.1 (Finite Nonempty Families Have a Maximum and Minimum). [prop:finite-max-min-inductive] *Let X be a totally ordered set. For any nonempty finite family $(a_1, \dots, a_n)$ in X , the maximum $\max\{a_1, \dots, a_n\}$ and minimum $\min\{a_1, \dots, a_n\}$ exist. They may be computed inductively:*

$$\max\{a_1\} = a_1, \quad \max\{a_1, \dots, a_k\} = \max(\max\{a_1, \dots, a_{k-1}\}, a_k) \text{ for } k \geq 2,$$

and symmetrically for min.

Definition 3.11 (Ordered Field). [def:ordered-field] An *ordered field* is a field F equipped with a total order $\leq$ such that for all $x, y, z \in F$,

- if $x \leq y$, then $x + z \leq y + z$; and
- if $0 \leq x$ and $0 \leq y$, then $0 \leq xy$.

Definition 3.12 (Absolute Value). [def:absolute-value] Let F be an ordered field. For $a \in F$, the *absolute value* of a is

$$|a| := \begin{cases} a, & a \geq 0, \\ -a, & a < 0. \end{cases}$$

Lemma 3.1 (Difference Characterization). [lem:difference-characterization] *Let F be an ordered field. For all $a, b \in F$,*

$$a < b \iff 0 < b - a.$$

Proof. Since addition preserves order and is cancellative in an ordered field,

$$a < b \iff a - a < b - a \iff 0 < b - a.$$

Here $a - a = 0$ by the additive-inverse axiom in the definition of a field. $\square$

Lemma 3.2 (Positive Perturbation). [lem:positive-perturbation] *Let F be an ordered field. For all $s, \varepsilon \in F$,*

$$\varepsilon > 0 \implies s - \varepsilon < s < s + \varepsilon.$$

Proof. If $\varepsilon > 0$, then

$$s - (s - \varepsilon) = \varepsilon > 0 \quad \text{and} \quad (s + \varepsilon) - s = \varepsilon > 0.$$

Both inequalities follow from the difference characterization. $\square$

4 Indexed Family

Definition 4.1 (Indexed Family). [def:indexed-family] Let I be an index set or type, and let X be a set or type. An *I -indexed family in X* is a mapping

$$a : I \rightarrow X.$$

Definition 4.2 (Term). [def:term] Let $a : I \rightarrow X$ be an indexed family. For each $i \in I$, the value

$$a_i := a(i)$$

is called the *term indexed by i* . For a sequence, a_n is called the *n -th term*.

Convention 4.1 (Indexed Family Notation). [conv:indexed-family-notation] We denote an indexed family $a : I \rightarrow X$ by

$$(a_i)_{i \in I}.$$

We may write simply (a_i) when the index set is clear.

Definition 4.3 (Sequence). [def:sequence] Let $I = \mathbb{N}$ or $I = \mathbb{N}_1$. An I -indexed family $a : I \rightarrow X$ is called a *sequence in X* . For a sequence, we conventionally use n as the index and write $(a_n)_{n \in I}$.

Example 4.1. $(\frac{1}{n})_{n \in \mathbb{N}_1}$ is a sequence.

5 Finite Sums and Products

Definition 5.1 (Finite Summation). [def:finite-summation] Let F be a field, $a : \mathbb{N}_1 \rightarrow F$, and $m, n \in \mathbb{N}_1$. The *finite summation* $\sum_{k=m}^n a_k$ is defined recursively:

$$\sum_{k=m}^n a_k = \begin{cases} 0 & \text{if } m > n, \\ \left(\sum_{k=m}^{n-1} a_k \right) + a_n & \text{if } m \leq n. \end{cases}$$

Definition 5.2 (Finite Product). [def:finite-product] Let F be a field, $a : \mathbb{N}_1 \rightarrow F$, and $m, n \in \mathbb{N}_1$. The *finite product* $\prod_{k=m}^n a_k$ is defined recursively:

$$\prod_{k=m}^n a_k = \begin{cases} 1 & \text{if } m > n, \\ \left(\prod_{k=m}^{n-1} a_k \right) \cdot a_n & \text{if } m \leq n. \end{cases}$$

Proposition 5.1 (Linearity of Summation). [prop:summation-linearity] Let F be a field, $a, b : \mathbb{N}_1 \rightarrow F$, $c, d \in F$, and $m, n \in \mathbb{N}_1$. Then

$$\sum_{k=m}^n (c \cdot a_k + d \cdot b_k) = c \cdot \sum_{k=m}^n a_k + d \cdot \sum_{k=m}^n b_k.$$

Proposition 5.2 (Index Shift for Summation). [prop:summation-shift] Let F be a field, $a : \mathbb{N}_1 \rightarrow F$, and $m, n, p \in \mathbb{N}_1$ with $m + p, n + p \in \mathbb{N}_1$. Then

$$\sum_{k=m}^n a_k = \sum_{k=m+p}^{n+p} a_{k-p}.$$

Proposition 5.3 (Splitting of Summation). [prop:summation-split] Let F be a field, $a : \mathbb{N}_1 \rightarrow F$, and $m, p, n \in \mathbb{N}_1$ with $m \leq p \leq n$. Then

$$\sum_{k=m}^n a_k = \sum_{k=m}^p a_k + \sum_{k=p+1}^n a_k.$$

Proposition 5.4 (Multiplicativity of Product). [prop:product-multiplicativity] Let F be a field, $a, b : \mathbb{N}_1 \rightarrow F$, and $m, n \in \mathbb{N}_1$. Then

$$\prod_{k=m}^n (a_k \cdot b_k) = \left(\prod_{k=m}^n a_k \right) \cdot \left(\prod_{k=m}^n b_k \right).$$

Proposition 5.5 (Index Shift for Product). [prop:product-shift] Let F be a field, $a : \mathbb{N}_1 \rightarrow F$, and $m, n, p \in \mathbb{N}_1$ with $m + p, n + p \in \mathbb{N}_1$. Then

$$\prod_{k=m}^n a_k = \prod_{k=m+p}^{n+p} a_{k-p}.$$

Proposition 5.6 (Splitting of Product). [prop:product-split] Let F be a field, $a : \mathbb{N}_1 \rightarrow F$, and $m, p, n \in \mathbb{N}_1$ with $m \leq p \leq n$. Then

$$\prod_{k=m}^n a_k = \left(\prod_{k=m}^p a_k \right) \cdot \left(\prod_{k=p+1}^n a_k \right).$$

6 Sets

Remark 6.1 (Membership and Classical Logic). [rem:membership-classical-logic] For every set A and every x in the chosen universe, exactly one of $x \in A$ and $x \notin A$ holds.

Proof. Let P be the proposition $x \in A$. By the law of excluded middle, $P \vee \neg P$, while the law of non-contradiction gives $\neg(P \wedge \neg P)$. Hence exactly one of $x \in A$ and $x \notin A$ holds. $\square$

Definition 6.1 (Intersection). [def:intersection] For sets A and B ,

$$x \in A \cap B \iff x \in A \text{ and } x \in B.$$

Definition 6.2 (Indexed Intersection). [def:indexed-intersection] Let I be a nonempty index type and let

$$A : I \rightarrow \mathbf{Set}$$

be an indexed family of sets. The *indexed intersection* is characterized by

$$x \in \bigcap_{i:I} A_i \iff \forall i : I, \quad x \in A_i.$$

In particular, for $a, b \in \mathbb{N}$ with $a \leq b$,

$$\bigcap_{n=a}^b A_n \equiv \bigcap_{n \in \{m \in \mathbb{N} : a \leq m \leq b\}} A_n, \quad \bigcap_{n=0}^{\infty} A_n \equiv \bigcap_{n:\mathbb{N}} A_n.$$

Definition 6.3 (Union). [def:union] For sets A and B ,

$$x \in A \cup B \iff x \in A \text{ or } x \in B.$$

Definition 6.4 (Indexed Union). [def:indexed-union] Let I be an index type and let

$$A : I \rightarrow \mathbf{Set}$$

be an indexed family of sets. The *indexed union* is characterized by

$$x \in \bigcup_{i:I} A_i \iff \exists i : I, \quad x \in A_i.$$

In particular, for $a, b \in \mathbb{N}$ with $a \leq b$,

$$\bigcup_{n=a}^b A_n \equiv \bigcup_{n \in \{m \in \mathbb{N} : a \leq m \leq b\}} A_n, \quad \bigcup_{n=0}^{\infty} A_n \equiv \bigcup_{n:\mathbb{N}} A_n.$$

If I is empty, then $\bigcup_{i:I} A_i = \emptyset$.

Definition 6.5 (Complement). [def:complement] Given a universal set U and $A \subseteq U$, the complement of A is $A^c = U \setminus A$; equivalently,

$$x \in A^c \iff x \notin A.$$

Definition 6.6 (Set Difference). [def:set-difference] For sets A and B ,

$$x \in A \setminus B \iff x \in A \text{ and } x \notin B.$$

Definition 6.7 (Subset). [def:subset] For sets A and B ,

$$A \subseteq B \iff \text{for every } x, \quad x \in A \implies x \in B.$$

Definition 6.8 (Set Equality). [def:set-equality] For sets A and B ,

$$A = B \iff A \subseteq B \text{ and } B \subseteq A.$$

Theorem 6.1 (De Morgan's Laws). [thm:de-morgan-laws] For sets A and B in a universal set U ,

$$(A \cap B)^c = A^c \cup B^c \quad \text{and} \quad (A \cup B)^c = A^c \cap B^c.$$

Also,

$$\left(\bigcap_{n=1}^{\infty} A_n \right)^c = \bigcup_{n=1}^{\infty} (A_n)^c \quad \text{and} \quad \left(\bigcup_{n=1}^{\infty} A_n \right)^c = \bigcap_{n=1}^{\infty} (A_n)^c$$

Example 6.1 (Nested Sequence). [[eg:nested-sequence](#)] Let $A_n = \{n, n+1, n+2, \dots\}, n \in \mathbb{N}_1$ ie, $A_1 = \{1, 2, 3, \dots\}, A_5 = \{5, 6, 7, \dots\}$ Trivially we can get $A_1 \subset A_2 \subset \dots$. This is called a nested sequence. Where

$$\bigcup_{n=1}^{\infty} A_n = A_1 \quad \bigcap_{n=1}^{\infty} = \emptyset$$

The intersection is $\emptyset$ due to if a number, let's say for any natural number p is in the set, then A_{p+1} does not include that number. So in the universe $\mathbb{N}$,

$$\forall p \notin \bigcap_{n=1}^{\infty} \implies \bigcap_{n=1}^{\infty} = \emptyset$$

7 Mapping

Definition 7.1 (Mapping). [[def:mapping](#)] A mapping $f : A \rightarrow B$ is a set of rule st.:

$$\forall x \in A, \exists! y = f(x) \in B \leftrightarrow \forall x \in A, \exists! y \in B \text{ st. } (x, y) \in f$$

Definition 7.2 (Domain). [[def:domain](#)] The *domain* of a mapping $f : A \rightarrow B$ is the set A .

Definition 7.3 (Codomain). [[def:codomain](#)] The *codomain* of a mapping $f : A \rightarrow B$ is the set B .

Definition 7.4 (Range). [[def:range](#)] The *range* of a mapping $f : A \rightarrow B$ is a set C such that:

$$C = \{y : \exists x \in A, \text{ st. } y = f(x)\}$$

Remark 7.1 (Range and codomains are different). [[rem:range-and-codomains-are-different](#)] The range is a subset of codomain. Notice that some of the values in the codomain is not reachable. For example,

$$f = x^2 : \mathbb{R} \rightarrow \mathbb{R}$$

The codomain is $\mathbb{R}$ but the range is $\mathbb{R}^+ \cup \{0\}$.

Let $\mathbb{B} := \{\text{tt}, \text{ff}\}$ denote the set of Boolean values.

Definition 7.5 (Injective Mapping/One to One Mapping). [[def:injective-mapping](#)] A mapping $f : A \rightarrow B$ is called *injective* or *one to one* if

$$f(x_1) = f(x_2) \implies x_1 = x_2$$

equivalently,

$$x_1 \neq x_2 \implies f(x_1) \neq f(x_2).$$

Example 7.1. The following is an injective mapping:

$$f : \mathbb{B} \rightarrow \mathbb{Z}, \quad \text{tt} \mapsto 0, \quad \text{ff} \mapsto 1.$$

Definition 7.6 (Surjective Mapping/onto Mapping). [[def:surjective-mapping](#)] A mapping $f : A \rightarrow B$ is called *surjective* or *onto* if

$$\forall y \in B, \exists x \in A, \text{ st. } f(x) = y$$

equivalently,

$$\text{Range}(f) = B$$

Example 7.2. The following is a surjective mapping:

$$g : \mathbb{Z} \rightarrow \mathbb{B}, \quad g(n) = \begin{cases} \text{tt}, & n = 0, \\ \text{ff}, & n \neq 0. \end{cases}$$

Definition 7.7 (Bijective Mapping). [def:bijective-mapping] A mapping $f : A \rightarrow B$ is called *bijective* if it is both injective and surjective.

Example 7.3. The following is a bijective mapping, so it is also injective and surjective:

$$h : \{0, 1\} \rightarrow \mathbb{B}, \quad 0 \mapsto \text{tt}, \quad 1 \mapsto \text{ff}.$$

8 Distance

Definition 8.1 (Distance). [def:distance] Let F be an ordered field. For $x, y \in F$, the *distance* between x and y is

$$d(x, y) := |x - y|.$$

Lemma 8.1 (Symmetry of Distance). [lem:distance-symmetry] Let F be an ordered field. For all $x, y \in F$,

$$d(x, y) = d(y, x).$$

Proof. By definition, $d(x, y) = |x - y|$ and $d(y, x) = |y - x| = |-(x - y)|$. It suffices to show $|-(x - y)| = |x - y|$; we split on the sign of $x - y$.

Case 1: $x - y \geq 0$. Then $|x - y| = x - y$. Also $-(x - y) \leq 0$, so $|-(x - y)| = -(-(x - y)) = x - y = |x - y|$.

Case 2: $x - y < 0$. Then $|x - y| = -(x - y)$. Also $-(x - y) > 0$, so $|-(x - y)| = -(x - y) = |x - y|$. $\square$

Lemma 8.2 (Triangle Inequality for Distance). [lem:triangle-inequality-distance] Let F be an ordered field. For all $x, y, z \in F$,

$$d(x, z) \leq d(x, y) + d(y, z).$$

Proof. Let $a = x - y$ and $b = y - z$. Then $x - z = a + b$, so $d(x, z) = |a + b|$. We split on the sign of $a + b$.

Case 1: $a + b \geq 0$. Then $|a + b| = a + b$. Since $a \leq |a|$ and $b \leq |b|$,

$$|a + b| = a + b \leq |a| + |b| = d(x, y) + d(y, z).$$

Case 2: $a + b < 0$. Then $|a + b| = -(a + b) = -a - b$. Since $-a \leq |a|$ and $-b \leq |b|$,

$$|a + b| = -a - b \leq |a| + |b| = d(x, y) + d(y, z).$$

$\square$

Lemma 8.3 (ε -Criterion for Equality). [lem:epsilon-equality] Let F be an ordered field. For $a, b \in F$,

$$a = b \iff \forall \varepsilon \in F, \varepsilon > 0, |a - b| < \varepsilon.$$

Proof. In order to proof lem:epsilon-equality, we need to prove both directions.

(a): $a = b \implies \forall \varepsilon > 0, |a - b| < \varepsilon$

Suppose $a = b$. Then $|a - b| = 0$. For any $\varepsilon > 0$, we have $|a - b| = 0 < \varepsilon$.

(b): $\forall \varepsilon > 0, |a - b| < \varepsilon \implies a = b$

Assume $a \neq b$.

Then, this means that

$$\exists \varepsilon_0 > 0, |a - b| = \varepsilon_0$$

Taking $\varepsilon = \varepsilon_0$ gives $\varepsilon_0 = |a - b| < \varepsilon_0$, a contradiction.

So, we have $\forall \varepsilon > 0, |a - b| < \varepsilon \implies \neg a \neq b$

Since $\neg a \neq b \iff a = b$, So, $\forall \varepsilon > 0, |a - b| < \varepsilon \implies a = b$

Therefore, both implications hold, and hence proven. $\square$

9 Bounds

Definition 9.1 (Bounded Above). [def:bounded-above] A set A is *bounded above* in the universe U if

$$\exists b \in U, \forall a \in A, a \leq b$$

Such b is called an *upper bound* of A .

Definition 9.2 (Bounded Below). [def:bounded-below] A set A is *bounded below* in the universe U if

$$\exists b \in U, \forall a \in A, a \geq b$$

Such b is called a *lower bound* of A .

Definition 9.3 (Least Upper Bound/Supremum). [def:supremum] s is the *least upper bound* or *supremum* of a set A if

1. s is an upper bound of A
2. $\forall b$ is an upper bound of A , $s \leq b$

Proposition 9.1 (ε -Criterion for the Supremum). [prop:epsilon-supremum] Let F be an ordered field, let $A \subseteq F$ be nonempty, and let $s \in F$. Then $s = \sup A$ if and only if

1. s is an upper bound of A ; and
2. for every $\varepsilon \in F$ with $\varepsilon > 0$, $s - \varepsilon$ is not an upper bound of A .

Equivalently,

$$s = \sup A \iff \begin{cases} \forall x \in A, & x \leq s, \\ \forall \varepsilon \in F, & \varepsilon > 0 \implies \exists x \in A \text{ s.t. } s - \varepsilon < x. \end{cases}$$

Proof. In order to proof ε -Characterization of the Supremum, we need to proof both directions are true.

(a):

$$s = \sup A \implies \begin{cases} \forall x \in A, & x \leq s, \\ \forall \varepsilon > 0, & \exists x \in A \text{ s.t. } s - \varepsilon < x. \end{cases}$$

According to the definition of supremum, we know that s is an upper bound of A , ie, $\forall x \in A, x \leq s$. Take arbitrary $\varepsilon \in F$ with $\varepsilon > 0$. By lemma 3.2, $s' := s - \varepsilon$, $s' < s$.

Again, according to the definition of supremum, we know $\forall b$ is an upper bound of A , $s \leq b$. Since $s' < s$, we have that s' is not an upper bound of A .

Therefore, $s' = s - \varepsilon$ is not an upper bound of A . According to the definition of upper bound, $\exists x \in A, x > s' \implies \exists x \in A, x > s - \varepsilon$.

So, (a) part is proven.

(b):

$$\begin{cases} \forall x \in A, & x \leq s, \\ \forall \varepsilon > 0, & \exists x \in A \text{ s.t. } s - \varepsilon < x. \end{cases} \implies s = \sup A$$

Assume $\forall x \in A, x \leq s$ and $\forall \varepsilon > 0, \exists x \in A \text{ s.t. } s - \varepsilon < x$, and suppose for contradiction that $s \neq \sup A$.

From $\forall x \in A, x \leq s$, it follows that s is an upper bound of A .

If $s \neq \sup A$, then there exists an upper bound b of A such that $b < s$. Let $\varepsilon := s - b > 0$. By the assumption, there exists $x \in A$ such that $s - \varepsilon < x$. However, $s - \varepsilon = s - (s - b) = b$, so $b < x$, contradicting that b is an upper bound of A .

So (b) is proven. □

Definition 9.4 (Greatest Lower Bound/Infimum). [def:infimum] i is the *greatest lower bound* or *infimum* of a set A if

1. i is a lower bound of A
2. $\forall b$ is a lower bound of A , $i \geq b$

Proposition 9.2 (ε -Criterion for the Infimum). [prop:epsilon-infimum] *Let F be an ordered field, let $A \subseteq F$ be nonempty, and let $i \in F$. Then $i = \inf A$ if and only if*

1. i is a lower bound of A ; and
2. for every $\varepsilon \in F$ with $\varepsilon > 0$, $i + \varepsilon$ is not a lower bound of A .

Equivalently,

$$i = \inf A \iff \begin{cases} \forall x \in A, & i \leq x, \\ \forall \varepsilon \in F, & \varepsilon > 0 \implies \exists x \in A \text{ s.t. } x < i + \varepsilon. \end{cases}$$

Convention 9.1. We use the notation $\sup A$ to denote the supremum of a set A and $\inf A$ to denote the infimum of a set A .

Remark 9.1 (Supremums and Infimums are unique). [rem:supremum-infimum-uniqueness] If a set A has a supremum, then it is unique. If a set A has an infimum, then it is unique.

Pf. Supremums are unique. Assume there are 2 supremums for set A , s_1, s_2 . Where $s_1 \neq s_2$.

Since $\forall s$ is an upper bound of A , $s_1 \leq s$, also $s_2 \leq s$.

Notice that s_1 is an upper bound of A , and s_2 is a least upper bound, so $s_2 \leq s_1$.

Also, s_2 is an upper bound of A , and s_1 is a least upper bound, so $s_1 \leq s_2$.

Trivially, $s_1 = s_2$. Contradicts with $s_1 \neq s_2$.

So there are no multiple supremum. Supremums are unique for each set. □

Definition 9.5 (Bounded Set). [def:bounded-set] A set A is *bounded* in the universe U if it is both bounded above and bounded below.

10 Axiom of Completeness

Definition 10.1 (Complete Ordered Field). [def:complete-ordered-field] A *complete ordered field* is an ordered field F such that every nonempty set $A \subseteq F$ that is bounded above has a least upper bound in F .

Remark 10.1 (Hierarchy of Order Structures). [rem:order-structure-hierarchy] Just now discussed the four ordered structures, each strictly extends the previous one and unlocks one new capability. When stating a theorem, use the weakest tier that suffices.

- Partial order: comparison via $\leq$ (reflexive, antisymmetric, transitive). Minimal and maximal elements make sense when they exist.
- Total order: adds totality, so any two elements are comparable. This is the minimum needed to define intervals.
- Ordered field: adds the field operations $+$, $-$, $\cdot$ compatible with $\leq$. Enables absolute value, distance, and ε -arguments.
- Complete ordered field: adds the Axiom of Completeness. Suprema and infima of bounded sets exist, and limit-existence results (e.g. NIP) require this tier.

Axiom 10.1 (Axiom of Completeness). [axm:aoc] *Every nonempty set $A \subseteq \mathbb{R}$ that is bounded above has a least upper bound in $\mathbb{R}$*

Lemma 10.1 (Supremum–Infimum Equivalence). [lem:supremum-infimum-equivalence] *Let F be an ordered field. The following statements are equivalent:*

1. Every nonempty set $A \subseteq F$ that is bounded above in F has a supremum in F .
2. Every nonempty set $A \subseteq F$ that is bounded below in F has an infimum in F .

Consequently, the definition of a complete ordered field may equivalently use the infimum property. In particular, the Axiom of Completeness is equivalent to the assertion that every nonempty subset of $\mathbb{R}$ bounded below has an infimum in $\mathbb{R}$.

Proof. First assume the supremum property, and let $A \subseteq \mathbb{R}$ be nonempty and bounded below. Define

$$-A = \{-a : a \in A\}.$$

Then $-A$ is nonempty and bounded above, so let $s = \sup(-A)$. For every $a \in A$, we have $-a \leq s$, and hence $-s \leq a$. Thus $-s$ is a lower bound of A . If ℓ is any lower bound of A , then $-\ell$ is an upper bound of $-A$. Therefore $s \leq -\ell$, so $\ell \leq -s$. Hence $-s = \inf A$.

Conversely, assume the infimum property, and let $A \subseteq \mathbb{R}$ be nonempty and bounded above. Then $-A$ is nonempty and bounded below, so let $i = \inf(-A)$. For every $a \in A$, we have $i \leq -a$, and hence $a \leq -i$. Thus $-i$ is an upper bound of A . If u is any upper bound of A , then $-u$ is a lower bound of $-A$. Therefore $-u \leq i$, so $-i \leq u$. Hence $-i = \sup A$. $\square$

Remark 10.2 (Integers and Natural Numbers). [rem:integers-naturals-not-fields] Neither $\mathbb{Z}$ nor $\mathbb{N}$ is a field, so neither can be called a complete ordered field. As ordered sets, however, their nonempty subsets that are bounded above have greatest elements.

Example 10.1. If $A = \{\frac{1}{n} : n \in \mathbb{N}_1\}$, then $\sup A = 1$ and $\inf A = 0$. Although it has a infimum, it does not have a minimum since $0 \notin A$.

Corollary 10.1 (Supremum and Infimum Existence in $\mathbb{R}$). [cor:supremum-infimum-existence] Every nonempty set $A \subseteq \mathbb{R}$ that is bounded above has a least upper bound in $\mathbb{R}$ and every nonempty set $A \subseteq \mathbb{R}$ that is bounded below has a greatest lower bound in $\mathbb{R}$.

Proof. The supremum statement is the Axiom of Completeness, and the infimum statement follows from lemma 10.1. $\square$

Remark 10.3 (Intuition for Completeness). [rem:intuition-for-completeness] We can have a set, let's say,

$$S = \{r \in \mathbb{Q} : r^2 < 3\}, \text{ where } r \in \mathbb{Q} \subset \mathbb{R}$$

Although the set S is bounded above in $\mathbb{Q}$, it does not have a supremum in $\mathbb{Q}$. According to the AoC, we know that $\exists s \in \mathbb{R}, s = \sup S$. This means we can find a number that can be described as the least upper bound a subset of rational numbers which is not in the rational numbers.

11 Irrational Numbers

Lemma 11.1 (Parity of a Square). [lem:square-parity] For every $n \in \mathbb{Z}$,

$$n^2 \text{ is even} \iff n \text{ is even},$$

and equivalently,

$$n^2 \text{ is odd} \iff n \text{ is odd}.$$

Proposition 11.1 (No Rational Square Root of 2). [prop:no-rational-square-root-of-two] There is no $q \in \mathbb{Q}$ such that $q^2 = 2$.

Proof. Suppose, for contradiction, that some $q \in \mathbb{Q}$ satisfies $q^2 = 2$. According to the proposition 1.1, and $q \in \mathbb{Q}$

$$\exists p, r \in \mathbb{Z}, q = \frac{p}{r}, r > 0, \gcd(|p|, r) = 1$$

Then we have

$$\left(\frac{p}{r}\right)^2 = 2 \implies p^2 = 2r^2$$

Since p^2 is even, the square-parity lemma implies that p is even. Therefore,

$$\exists k \in \mathbb{Z}, \quad p = 2k.$$

So,

$$(2k)^2 = 2r^2 \implies 2k^2 = r^2$$

By the same argument, r is also even. Meaning that both p and r are even, which contradicts the assumption that $\gcd(|p|, r) = 1$.

So, it contradicts the assumption that $q \in \mathbb{Q}$ satisfies $q^2 = 2$. Therefore, there is no $q \in \mathbb{Q}$ such that $q^2 = 2$. $\square$

Remark 11.1 (Larger Number System). [rem:sqrt-two-larger-number-system] Within $\mathbb{Q}$, the proposition only says that the equation $q^2 = 2$ has no solution $q \in \mathbb{Q}$. Writing $\sqrt{2} \notin \mathbb{Q}$ presupposes a larger number system in which $\sqrt{2}$ exists.

Remark 11.2 (Completeness). [rem:rational-incompleteness] The existence, in a larger number system, of some $x \notin \mathbb{Q}$ satisfying $x^2 = 2$ does not by itself prove that $\mathbb{Q}$ is incomplete. Incompleteness must be demonstrated internally by finding a nonempty subset of $\mathbb{Q}$ that is bounded above in $\mathbb{Q}$ but has no supremum in $\mathbb{Q}$, as in Proposition 11.3.

Convention 11.1 (Irrational Numbers). [conv:irrational-numbers] We use $\mathbb{I}$ to denote the set of *irrational numbers*

Remark 11.3 (Number Sets). [rem:number-sets-inclusions-real] The set of real numbers $\mathbb{R}$ is the union of the set of rational numbers $\mathbb{Q}$ and the set of irrational numbers $\mathbb{I}$:

$$\mathbb{R} = \mathbb{Q} \cup \mathbb{I}$$

Also, in normal conventions, we have:

$$\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R}$$

Proposition 11.2 (Integer–Irrational Root Dichotomy). [prop:nth-root-dichotomy] For $x, n \in \mathbb{N}$ with $n > 1$, $\sqrt[n]{x} \in \mathbb{Z}$ if x is a perfect n -th power, and otherwise $\sqrt[n]{x} \in \mathbb{I}$.

Remark 11.4 (Sources of Irrational). [rem:sources-of-irrational]

- By proposition 11.2, $\sqrt[n]{x}$ is irrational whenever $x \in \mathbb{N}$, $n > 1$, and x is not a perfect n -th power.
- Taking logarithms
- Constants defined by taking limits, such as π and e .

Remark 11.5 (Irrational Numbers Are Not Closed). [rem:irrational-not-closed] The irrational numbers are not closed under addition, multiplication, or exponentiation. For example,

$$\sqrt{2} + (-\sqrt{2}) = 0, \quad \sqrt{2} \cdot \sqrt{2} = 2, \quad (\sqrt{3})^{\log_3 4} = 2,$$

where all the inputs are irrational. Although $\pi, e \in \mathbb{I}$, it is not known whether either $\pi + e$ or πe is irrational; both are expected to be transcendental. By contrast, e^π has been proved transcendental, and is therefore irrational.

Proposition 11.3 (The Rational Numbers Are Not Complete). [prop:rational-not-complete] $\mathbb{Q}$ is not a complete ordered field. In fact, the set

$$A = \{q \in \mathbb{Q} : q^2 < 2\}$$

is nonempty and bounded above in $\mathbb{Q}$, but it has no supremum in $\mathbb{Q}$.

Proof. The set A is nonempty because $1 \in A$. It is bounded above in $\mathbb{Q}$ by 2: if some $q \in A$ satisfied $q > 2$, then $q^2 > 4 > 2$, contradicting $q^2 < 2$. Suppose, for contradiction, that A has a supremum $s \in \mathbb{Q}$. Since $1 \in A$, we have $s \geq 1$. First suppose $s^2 < 2$, and define

$$h = \frac{2 - s^2}{2s + 2} \in \mathbb{Q}.$$

Then $h > 0$; moreover, $s \geq 1$ implies $h < 2$. Hence

$$\begin{aligned} (s + h)^2 &= s^2 + h(2s + h) \\ &< s^2 + h(2s + 2) \\ &= 2. \end{aligned}$$

Thus $s + h \in A$, but $s + h > s$, contradicting that s is an upper bound of A . Therefore $s^2 \not< 2$. Now suppose $s^2 > 2$, and define

$$h = \frac{s^2 - 2}{2s} \in \mathbb{Q}, \quad t = s - h.$$

Then $h > 0$, so $t < s$, and

$$t = \frac{s^2 + 2}{2s} > 0, \quad t^2 = (s - h)^2 = 2 + h^2 > 2.$$

If some $q \in A$ satisfied $q > t$, then $q > t > 0$ and

$$q^2 - t^2 = (q - t)(q + t) > 0,$$

so $q^2 > t^2 > 2$, contradicting $q \in A$. Consequently every $q \in A$ satisfies $q \leq t$, making t an upper bound of A . This contradicts the minimality of s , because $t < s$. Therefore $s^2 \not> 2$.

By totality, $s^2 = 2$. This contradicts the fact that no rational number has square 2. Hence A has no supremum in $\mathbb{Q}$, so $\mathbb{Q}$ is not complete. $\square$