Luego pienso en el título Va a estar chilero

Hernández, Giovanna¹*

¹Escuela de Ciencias Físicas y Matemáticas, Universidad de San Carlos de Guatemala

Abstract

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Praesent porttitor arcu luctus, imperdiet urna iaculis, mattis eros. Pellentesque iaculis odio vel nisl ullamcorper, nec faucibus ipsum molestie. Sed dictum nisl non aliquet porttitor. Etiam vulputate arcu dignissim, finibus sem et, viverra nisl. Aenean luctus congue massa, ut laoreet metus ornare in. Nunc fermentum nisi imperdiet lectus tincidunt vestibulum at ac elit. Nulla mattis nisl eu malesuada suscipit. Aliquam arcu turpis, ultrices sed luctus ac, vehicula id metus. Morbi eu feugiat velit, et tempus augue. Proin ac mattis tortor. Donec tincidunt, ante rhoncus luctus semper, arcu lorem lobortis justo, nec convallis ante quam quis lectus. Aenean tincidunt sodales massa, et hendrerit tellus mattis ac. Sed non pretium nibh. Donec cursus maximus luctus. Vivamus lobortis eros et massa porta porttitor.

Introduction

Near-Earth Objects (NEOs) are comets and asteroids that have been nudged by the gravitational attraction of nearby planets into orbits that allow them to enter the Earth's neighborhood. Composed mostly of water ice with embedded dust particles, comets originally formed in the cold outer planetary system while most of the rocky asteroids formed in the warmer inner solar system between the orbits of Mars and Jupiter.¹

A meteoroid is generally defined as an asteroid or comet fragment that orbits the Sun and has an approximate size between ten microns and a meter or so. Meteors, or "shooting stars", are the visible paths of meteoroids that have entered the Earth's atmosphere at high velocities. A fireball is an unusually bright meteor that reaches a visual magnitude of -3 or brighter when seen at the observer's zenith.²

Near-Earth Asteroids (NEAs) are small bodies of the Solar System with perihelion distance q 1.3 AU (Astronomical Units) and aphelion distances Q 0.983 AU, whose orbits approach or intersect Earth orbit. The NEAs are classified into three main classes: Apollo, Amor and Aten on the basis of derived orbital parameters.³

Potentially Hazardous Asteroids (PHAs) are a special subset of NEAs that, according to The Center for Near-Earth Object Studies (CNEOS), have an absolute magnitude (*H*) of 22.0 or less that can come close to the Earth and are large enough to cause significant

*Corresponding author: gioreneeha@gmail.com Received: May 19, 2024 damage in the event of an impact.4

Sentry is a highly automated collision monitoring system that continually scans the most current asteroid catalog for possibilities of future impact with Earth over the next 100 years. Whenever a potential impact is detected it will be analyzed and the results immediately published here, except in unusual cases where we seek independent confirmation.⁵

Data and Methods

The data about fireballs, NEOs, NEAs and impact probabilities have been collected from the database of The Center for Near-Earth Object Studies (CNEOS) and its monitoring system Sentry. The parameters studied include absolute visual magnitude (*H*), impact probability, impact energy (*kt*) and geographic location of fireball objects.

To determine if the impact energy (kt) of fireballs is consistent with some type of distribution it was decided to use the logarithm of the data and then a histogram was made with the counts of the impact energy (log(kt)) in intervals of 0.2. With these data, some distribution fit were applied to confirm wich one was more accurate.

At first, the covariance matrix was used to find the linear bond between NEOs' absolute magnitude (*H*) and impact probability, but due to the correlation not being linear, it was discarded. Then proceeded to use Pearson's and Spearman's correlation coefficients to analyze better the data and determine if a correlation existed and which type it was.




Figure 1: Fireballs reported by US government sensors from April 15, 1988, to May 15, 2024. The size and color of each circle were determined by the logarithm of the impact energy (kt), and the location was determined by the altitude and longitude. All this data was obtained from the database published by NASA's CNEOS at JPL.

At last, using Gnuplot and Python with various libraries such as pandas, numpy, plotly, a graphic representation of geographic locations with their respective impact energy as the size of the reported events was made.




Figure 2: Histogram of the logarithm of fireballs' impact energy (kt) reported by US government sensors from April 15, 1988, to May 15, 2024. The red line indicates the fit of a Gaussian distribution that was applied to the data. The larger impact energy was of $6 \log(kt)$.

Results

Here there goes the results




Figure 3: Plot between NEOs' impact probability and absolute magnitude (H) obtained from the database published by NASA's CNEOS at JPL. The olive line indicates an exponential fit and the yellow one indicates a logarithmic fit, both were applied to the data.

Discussion

References

- [1] CNEOS, NEO Basics cneos.jpl.nasa.gov, https://cneos.jpl.nasa.gov/about/basics.html, [Accessed 19-05-2024].
- [2] CNEOS, Fireballs cneos.jpl.nasa.gov, https://cneos.jpl.nasa.gov/fireballs/intro.html, [Accessed 19-05-2024].
- [3] J. Rukmini, G. Raghavendra, S. A. Ahmed, D. S. Priya, and S. A. Unnisa, *Statistical survey and analysis of photometric and spectroscopic data on neas*, 2016.
- [4] Y. F. Zhou, H. Li, Z. Li, and L. Zhou, *Martians* (mars2020, tianwen and so on) would see more potentially hazardous asteroids than earthlings, 2024.
- [5] CNEOS, Sentry: Earth Impact Monitoring—
 cneos.jpl.nasa.gov, https://cneos.jpl.nasa.
 gov/sentry/, [Accessed 19-05-2024].