

A Spectral Approach To Meshes

Eli Griffiths

University of California, Irvine

November 22, 2024


- 1 Basics of Graphs
- 2 Matrix Representations
- 3 Computational Geometry

Definition (Graph)

A graph is a pair $G = (V, E)$ where E is comprised of two element subsets of V . The elements of V are called *vertices* and the elements of E *edges*

Definition (Graph)

A graph is a pair $G = (V, E)$ where E is comprised of two element subsets of V . The elements of V are called *vertices* and the elements of E *edges*



Graphs at their core encode connectivity

Graphs at their core encode connectivity


- Friend networks on social media platforms



Graphs at their core encode connectivity

- Friend networks on social media platforms
- Molecules and bonds between atoms in structural chemistry


Graphs at their core encode connectivity

- Friend networks on social media platforms
- Molecules and bonds between atoms in structural chemistry
- Connectivity of meshes in computational geometry






	v_1	v_2	v_3	v_4	v_5	v_6	v_7
v_1							
v_2							
v_3							
v_4							
v_5							
v_6							
v_7							



	v_1	v_2	v_3	v_4	v_5	v_6	v_7
v_1		✓	✓	✓	✓	✓	✓
v_2	✓		✓		✓	✓	✓
v_3	✓	✓					✓
v_4	✓				✓	✓	✓
v_5	✓	✓		✓		✓	
v_6	✓	✓		✓	✓		
v_7	✓	✓	✓	✓			



	v_1	v_2	v_3	v_4	v_5	v_6	v_7
v_1		✓	✓	✓	✓	✓	✓
v_2	✓		✓		✓	✓	✓
v_3	✓	✓					✓
v_4	✓				✓	✓	✓
v_5	✓	✓		✓		✓	
v_6	✓	✓		✓	✓		
v_7	✓	✓	✓	✓			



$$\begin{pmatrix}
 0 & 1 & 1 & 1 & 1 & 1 & 1 \\
 1 & 0 & 1 & 0 & 1 & 1 & 1 \\
 1 & 1 & 0 & 0 & 0 & 0 & 1 \\
 1 & 0 & 0 & 0 & 1 & 1 & 1 \\
 1 & 1 & 0 & 1 & 0 & 1 & 0 \\
 1 & 1 & 0 & 1 & 1 & 0 & 0 \\
 1 & 1 & 1 & 1 & 0 & 0 & 0
 \end{pmatrix}$$

Adjacency Matrix A

Definition (Degree Matrix)

The degree matrix of a graph G with n vertices is the $n \times n$ matrix D such that

$$(D)_{ij} = \begin{cases} \deg(v_i) & i = j \\ 0 & i \neq j \end{cases}.$$

Definition (Laplacian Matrix)

The Laplacian matrix of a graph G is

$$L := D - A.$$

Theorem


A graph has m connected components if and only if zero has algebraic multiplicity m for L

Theorem

The eigenvector associated with the smallest positive eigenvalue of L (called the “Fiedler Vector”) defines a well behaved partitioning scheme of a graph.

Definition (Triangular Mesh)

A **triangular mesh** is a triple $K = (V, E, F)$ such that




Definition (Triangular Mesh)

A **triangular mesh** is a triple

$K = (V, E, F)$ such that

- $V \subseteq \mathbb{R}^3$ is the set of vertices




Definition (Triangular Mesh)

A **triangular mesh** is a triple

$K = (V, E, F)$ such that

- $V \subseteq \mathbb{R}^3$ is the set of vertices
- $E \subseteq [V]^2$ is the set of representing non-intersecting edges



Definition (Triangular Mesh)

A **triangular mesh** is a triple

$K = (V, E, F)$ such that

- $V \subseteq \mathbb{R}^3$ is the set of vertices
- $E \subseteq [V]^2$ is the set of representing non-intersecting edges
- $F \subseteq [E]^3$ is the set of faces such that for any $f = \{e_1, e_2, e_3\} \in F$,

$$e_1 \cap e_2 = \{v_1\}$$

$$e_2 \cap e_3 = \{v_2\}$$

$$e_3 \cap e_1 = \{v_3\}$$

for $v_1 \neq v_2 \neq v_3$.

