

Math 121A: Linear Algebra

Eli Griffiths

October 18, 2023

Table of Contents

Introduction	2
1.1 Fields	2
1.2 Vector Spaces	2
1.2.1 Subspaces	3
1.3 Span and Independence	4
1.4 Basis and Dimension	5
Linear Maps	7
2.1 Linearity	7
List of Theorems	9
List of Definitions	9

Introduction

1.1 Fields

A field is an algebraic structure intended to capture the properties of the rational and real numbers.

Definition 1.1 (Field). A field \mathbb{F} is a set \mathbb{F} equipped with an addition and multiplication operator such that

1. Elements commute under both operations
2. Associativity holds for both operators
3. $\exists 1, 0 \in \mathbb{F}$ such that $\lambda + 0 = \lambda$ and $\lambda 1 = \lambda$ for all $\lambda \in \mathbb{F}$
4. $\forall \lambda \in \mathbb{F}, \exists ! \gamma \in \mathbb{F}$ such that $\lambda + \gamma = 0$
5. $\forall \lambda \in \mathbb{F}$ where $\lambda \neq 0, \exists ! \gamma \in \mathbb{F}$ such that $\lambda \gamma = 0$

Consider a prime number p . It is then possible to construct the field \mathbb{F}_p where

$$\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$$

Consider the simplest example, $\mathbb{F}_2 = \{0, 1\}$. It is clear that this makes up a field as 1 has a multiplicative and additive inverse. In the case of $\mathbb{F}_4 = \{0, 1, 2, 3\}$, 2 has no multiplicative inverse and hence isn't a field. In general, if $a, b \neq 0$ and $ab = 0$, then a and b do not have inverses.

Proof. Let $a, b \in \mathbb{F}$. Assume towards contradiction that $ab = 0$ and that a or b have inverses. WLOG, assume that there is a $c \in \mathbb{F}$ such that $ac = 1$. ■

1.2 Vector Spaces

Definition 1.2 (Vector Space). Let \mathbb{F} be a field. A vector space V over a field \mathbb{F} is a set V equipped with addition and scalar multiplication such that

1. $(V, +)$ is an abelian group
2. (V, \cdot) is associative and distributive, that is $\forall a, b \in \mathbb{F}$ and $\forall u, v \in V$
 - (a) $a(u + v) = au + av$
 - (b) $(a + b)v = av + bv$
3. $\exists ! 1 \in \mathbb{F}$ such that $\forall v \in V, 1v = v$

Example 1.1. Let \mathbb{F} be a field and $n \in \mathbb{N}$. Let

$$\mathbb{F}^n = \{(a_1, a_2, a_3, \dots) : a_i \in \mathbb{F}\}$$

where

$$(a_1, a_2, a_3, \dots) + (b_1, b_2, b_3, \dots) = (a_1 + b_1, a_2 + b_2, a_3 + b_3, \dots) \\ e \cdot (a_1, a_2, a_3, \dots) = (ea_1, ea_2, ea_3, \dots)$$

Then $(\mathbb{F}^n, +, \cdot)$ is a vector space. This gives rise to all the familiar vector spaces $\mathbb{C}^n, \mathbb{R}^n, \mathbb{Q}^n, \dots$

Remark. If $K \supset F$ are fields, then $(K, +, \cdot)$ is a vector space over F

Theorem 1.1. Let V be a vector space and $u, v, w \in V$. Then $u + w = v + w$ implies $u = v$.

Proof. Im too lazy :P ■

There are more exotic examples of vector spaces. Consider the set of continuous, real value functions over the interval $[0, 1]$.

Example 1.2.

$$C = \{f : [0, 1] \rightarrow \mathbb{R} \mid f \text{ is continuous}\}$$

This set can be turned into a vector space by defining the following operations

$$+ \Rightarrow (f + g)(x) = f(x) + g(x) \\ \cdot \Rightarrow (cf)(x) = cf(x)$$

1.2.1 Subspaces

Definition 1.3 (Subspace). Let V be a vector space over a field \mathbb{F} . A subset $W \subset V$ is called a subspace if W is also a vector space over \mathbb{F} .

Theorem 1.2. Let $W \subset V$ be a non-empty subset of V . W is a subspace of V if and only if

1. W is closed under $+$

$$w_1 + w_2 \in W, \forall w_1, w_2 \in W$$

2. W is closed under \cdot

$$cw \in W, \forall w \in W, c \in \mathbb{F}$$

An equivalent formulation of the conditions is that W is closed under linear combination, or symbolically

$$c_1w_1 + c_2w_2 \in W, \forall c_1, c_2 \in \mathbb{F}, w_1, w_2 \in W$$

Consider the vector space \mathbb{R}^2 . After some time, it becomes clear that the only subspaces of \mathbb{R}^2 are $\{\vec{0}\}$ and $\{c(a, b) | c \in \mathbb{R}\}$ where $a, b \in \mathbb{R}$ (aka all lines that go through the origin). If a line does not go through the origin of \mathbb{R}^2 , then it's clear it fails to be a subspace. Examine the line $y + x = 1$. Since it does not contain the zero vector, it fails to be closed under scalar multiplication as any vector in $y + x = 1$ will become the zero vector.

Example 1.3 (Polynomial Space). Let $\mathcal{P}_n(\mathbb{F}) := \{a_n x^n + a_{n-1} x^{n-1} + \dots + a_0 : a_i \in \mathbb{F}\}$. $\mathcal{P}_n(\mathbb{F})$ is a vector space $\forall n \in \mathbb{N}_0$. Additionally, $\mathcal{P}_{n-1}(\mathbb{F})$ is a subspace of $\mathcal{P}_n(\mathbb{F})$.

Example 1.4 (Matrix Space). Let $M_{n \times m}(\mathbb{F}) := \left\{ \begin{pmatrix} a_{11} & \dots & a_{1m} \\ \vdots & & \vdots \\ a_{n1} & \dots & a_{nm} \end{pmatrix} : a_{ij} \in \mathbb{F} \right\}$. Then $M_{n \times m}(\mathbb{F})$ is a vector space. An example subspace of $M_{n \times m}(\mathbb{F})$ is the set of upper triangular matrices.

Theorem 1.3 (Trivial Subspaces). For every non-zero vector space V , it has at least two subspaces $\{\vec{0}\}$ and V .

Theorem 1.4 (Subspace Construction). Let V_1, V_2 be subspaces of V over \mathbb{F} . Then

1. $V_1 \cap V_2$ is a subspace
2. $V_1 + V_2 = \{v_1 + v_2 : v_1 \in V_1, v_2 \in V_2\}$ is a subspace
3. $V_1 \cup V_2$ is a subspace if and only if $V_1 \subset V_2$ or $V_2 \subset V_1$

1.3 Span and Independence

Definition 1.4 (Linear Combination). Let V be a vector space and $v_1, v_2, \dots, v_n \in V$. A linear combination of $\{v_1, v_2, \dots, v_n\}$ is a vector of the form

$$c_1 v_1 + c_2 v_2 + \dots + c_n v_n$$

where $c_i \in \mathbb{F}$.

Definition 1.5 (Span). The span of a set of vectors $\{v_1, v_2, \dots, v_n\}$ is the set of all linear combinations of those vectors. It is denoted by $\text{span}\{v_1, v_2, \dots, v_n\}$. For an infinite set of vectors, an alternative definition helps

$$\text{span}(S) = \bigcap_{\substack{W \supseteq S \\ W \leq V}} W = \text{Smallest subspace that contains } S$$

Definition 1.6 (Subset Independence). Let S be a subset of a vector space V .

1. If every non-empty finite subset of S is linearly independent, S is called indepen-

dent.

2. If there exists a non-empty finite subset of S that is linearly dependent, S is called dependent.

1.4 Basis and Dimension

Definition 1.7. A maximal independent set of vectors is a independent subset of a vector space that, if any other vector is added from the vector space, would be become dependent. That is, given a maximally independent subset $W = \{v_1, v_2, v_3, \dots\}$, $u \in \text{span}(W)$.

Definition 1.8 (Basis). Let $\{u_1, u_2, u_3, \dots\} \subset V$. Then $\{u_1, u_2, u_3, \dots\}$ is a basis of V if every vector $v \in V$ can be uniquely written as $v = c_1u_1 + c_2u_2 + \dots$ with $c_i \in \mathbb{F}$. Equivalently, it is a basis if

1. $V = \text{span}\{u_1, u_2, u_3, \dots\}$
2. $\{u_1, u_2, u_3, \dots\}$ is independent

Theorem 1.5. Let $V = \text{span}\{v_1, v_2, \dots, v_n\}$ and $W = \{v_1, v_2, \dots, v_r\}$ be a maximally independent subset of V . Then W is a basis for V .

Proof. Since W is maximally independent, $v_i \in W$ for $1 \leq i \leq n$. Therefore $V = \text{span}(W)$. Since W is independent and spans V , it is a basis for V . ■

Corollary 1.1. Let $V = \{v_1, v_2, \dots, v_n\}$. Any maximally independent subset of V forms a basis of V over \mathbb{F} .

Theorem 1.6. If S is an independent subset of a vector space V , it can be extended to a basis of V (by the Axiom of Choice).

Theorem 1.7. Let $R = \{v_1, v_2, \dots, v_n\}$ and $W = \{w_1, w_2, \dots, w_m\}$ be bases of V over \mathbb{F} . Then $m = n$.

Proof. Since R is a basis, $w_i = \sum_{j=1}^n a_{ij}v_j$. This can be expressed in matrix form

$$\begin{pmatrix} w_1 \\ \vdots \\ w_m \end{pmatrix} = \begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{pmatrix} \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix}$$

Since W is also a basis, $v_i = \sum_{j=1}^m b_j w_j$ meaning

$$\begin{pmatrix} w_1 \\ \vdots \\ w_m \end{pmatrix} = \underbrace{\begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{pmatrix}}_A \underbrace{\begin{pmatrix} b_{11} & \cdots & b_{1m} \\ \vdots & & \vdots \\ b_{m1} & \cdots & b_{nm} \end{pmatrix}}_B \begin{pmatrix} w_1 \\ \vdots \\ w_n \end{pmatrix}$$

This means that $AB = I$. By starting with W instead of R , it follows that $BA = I$. Therefore A and B are invertible meaning they must be square, hence $n = m$. ■

Definition 1.9 (Dimension). The dimension of a vector space V is the number of vectors in a basis of V .

Theorem 1.8. Let W be a subspace of V such that $\dim V = n < \infty$. If $\dim W = \dim V$, $W = V$.

Proof. ■

Linear Maps

2.1 Linearity

Definition 2.10 (Linear Map). A map $T : V \rightarrow W$ is linear if $T(au+bv) = aT(u)+bT(v)$ for all $a, b \in \mathbb{F}$ and $u, v \in V$.

Theorem 2.9. If $\{v_1, v_2, \dots, v_n\}$ is a basis of V ,

$$T(a_1v_1 + a_2v_2 + \dots + a_nv_n) = a_1T(v_1) + a_2T(v_2) + \dots + a_nT(v_n)$$

A linear map can be defined just by declaring the images of a vector spaces basis vectors as the map has to obey linearity over a basis. This leads to a natural formulation of a linear map as a matrix where the columns are the images of the basis vectors under T .

Definition 2.11 (Null Space and Range). Let $T : V \rightarrow W$ be \mathbb{F} -linear. Then

1. $N(T) = \{v \in V : T(v) = 0\}$ is a subspace of V
2. $R(T) = T(V) = \{T(v) : v \in V\}$ is a subspace of W

Theorem 2.10. Let $T : V \rightarrow W$ be \mathbb{F} -linear. Then

$$\begin{aligned}\dim V &= \dim(N(T)) + \dim(R(T)) \\ &= \text{nullity } T + \text{rank } T\end{aligned}$$

Proof. Let S be a basis of $N(T)$. Then $\#S = \text{nullity } T$. S can be extended to be a basis of V with some S' such that $S \cup S'$ is a basis. Therefore

$$\dim V = \#(S \cup S') = \#S + \#S' = \text{nullity } T + \#S$$

■

Theorem 2.11. $T : V \rightarrow W$ is one-to-one if and only if $N(T) = \{0\}$

Proof. The forward direction follows by considering $T(v_1) = T(v_2)$ and setting v_2 to zero. Consider the backwards direction. If $T(v_1) = T(v_2)$, then $T(v_1 - v_2) = 0$. Therefore $v_1 - v_2 \in N(T) = \{0\}$ meaning $v_1 - v_2 = 0$. Hence $v_1 = v_2$ meaning T is one-to-one. ■

Theorem 2.12. Let $T : V \rightarrow V$ be a linear map where $\dim V = n < \infty$. Then T is bijective.

Proof. By the dimension formula,

$$n = \dim V = \text{nullity } T + \text{rank } T$$

If T is injective, then

$$\begin{aligned}\text{nullity } T = 0 &\implies \text{rank } T \\ &\implies \text{rank } T = n \\ &\implies R(T) = V\end{aligned}$$

■

List of Theorems

1.3 Theorem (Trivial Subspaces) 4

1.4 Theorem (Subspace Construction) 4

List of Definitions

1.1 Definition (Field) 2

1.2 Definition (Vector Space) 2

1.3 Definition (Subspace) 3

1.4 Definition (Linear Combination) 4

1.5 Definition (Span) 4

1.6 Definition (Subset Independence) 4

1.8 Definition (Basis) 5

1.9 Definition (Dimension) 6

2.10 Definition (Linear Map) 7

2.11 Definition (Null Space and Range) 7