

# A Spectral Approach To Meshes

Eli Griffiths

November 12, 2024

## Definition (Triangular Mesh)

A **triangular mesh** is a triple  $K = (V, E, F)$  such that


- $V \subseteq \mathbb{R}^3$  is a finite set representing the vertices
- $E \subseteq [V]^2$  is a set representing non-intersecting edges
- $F \subseteq [E]^3$  is the set of faces such that for any  $f = \{e_1, e_2, e_3\} \in F$ ,

$$e_1 \cap e_2 = \{v_1\}$$

$$e_2 \cap e_3 = \{v_2\}$$

$$e_3 \cap e_1 = \{v_3\}$$

for  $v_1 \neq v_2 \neq v_3$ .



## Definition (Triangular Mesh)

A **triangular mesh** is a triple  $K = (V, E, F)$  such that

- $V \subseteq \mathbb{R}^3$  is a finite set representing the vertices
- $E \subseteq [V]^2$  is a set representing non-intersecting edges
- $F \subseteq [E]^3$  is the set of faces such that for any  $f = \{e_1, e_2, e_3\} \in F$ ,

$$e_1 \cap e_2 = \{v_1\}$$

$$e_2 \cap e_3 = \{v_2\}$$

$$e_3 \cap e_1 = \{v_3\}$$

for  $v_1 \neq v_2 \neq v_3$ .



## Definition (Triangular Mesh)

A **triangular mesh** is a triple  $K = (V, E, F)$  such that


- $V \subseteq \mathbb{R}^3$  is a finite set representing the vertices
- $E \subseteq [V]^2$  is a set representing non-intersecting edges
- $F \subseteq [E]^3$  is the set of faces such that for any  $f = \{e_1, e_2, e_3\} \in F$ ,

$$e_1 \cap e_2 = \{v_1\}$$

$$e_2 \cap e_3 = \{v_2\}$$

$$e_3 \cap e_1 = \{v_3\}$$

for  $v_1 \neq v_2 \neq v_3$ .



## Definition (Triangular Mesh)

A **triangular mesh** is a triple  $K = (V, E, F)$  such that

- $V \subseteq \mathbb{R}^3$  is a finite set representing the vertices
- $E \subseteq [V]^2$  is a set representing non-intersecting edges
- $F \subseteq [E]^3$  is the set of faces such that for any  $f = \{e_1, e_2, e_3\} \in F$ ,

$$e_1 \cap e_2 = \{v_1\}$$

$$e_2 \cap e_3 = \{v_2\}$$

$$e_3 \cap e_1 = \{v_3\}$$

for  $v_1 \neq v_2 \neq v_3$ .

