

# Directed C–H Functionalization of C3-Aldehyde, Ketone, and Acid/Ester-Substituted Free (NH) Indoles with Iodoarenes *via* a Palladium Catalyst System

Yunus Taskesenligil, Murat Aslan, Tuba Cogurcu, and Nurullah Saracoglu\*

Cite This: *J. Org. Chem.* 2023, 88, 1299–1318

Read Online

ACCESS |



Metrics &amp; More




Article Recommendations



Supporting Information


**ABSTRACT:** Pd(II)-catalyzed C–H arylations of free (NH) indoles including different carbonyl directing groups on C3-position with aryl iodides are demonstrated. Importantly, the reactions are carried out using the same catalyst system without any additional transient directing group (TDG). In this study, the formyl group as a directing group gave the C4-arylated indoles versus C2-arylation. Using this catalyst system, C–H functionalization of 3-acetylindoles provided domino C4-arylation/3,2-carbonyl migration products. This transformation involves the unusual migration of the acetyl group to the C2-position following C4-arylation in one pot. Meanwhile, migration of the acetyl group could be simply controlled and *N*-protected 3-acetylindoles afforded C4-arylation products without migration of the acetyl group. Functionalization of indole-3-carboxylic acid (or methyl ester) with aryl iodides using the present Pd(II)-catalyst system resulted in decarboxylation followed by the formation of C2-arylated indoles. Based on the control experiments and the literature, plausible mechanisms are proposed. The synthetic utilities of these acetylindole derivatives have also been demonstrated. Remarkably, C4-arylated acetylindoles have allowed the construction of functionalized pityacitrin (a natural product).



## INTRODUCTION

Transition-metal (TM)-catalyzed functionalization reactions through directing group-assisted C–H activation have emerged as a powerful tool for C–C and C–X bond-forming reactions, providing an atom- and step-economical strategy for organic synthesis.<sup>1</sup> The C–H activation, which represents a paradigmatic change in the field of synthesis of complex heterocyclic and carbocyclic molecules, allows not only specifically functionalizes of the inert C–H bonds but also provides the formation of various compounds by coupling the introduced functional groups.<sup>2</sup>

Indole core is found in numerous natural products, pharmaceuticals, materials chemistry, and other bio-relevant compounds and has been recognized as a privileged structure scaffold.<sup>3</sup> Because of this, enormous efforts have been devoted to the construction of functionalized indoles. Due to the high nucleophilic activity of the pyrrole ring of the indole, most C–H activation reactions primarily occur at the C2 or C3 position of pyrrole moiety (Figure 1a).<sup>4</sup> As opposed to these positions, C–H functionalization of the low active C4–C7 positions in the benzenoid ring is less scrutinized and remains a long-standing challenge (Figure 1a). In recent years, transition-metal-catalyzed directed C–H activation has been studied as a powerful synthetic tool to access the functionalized indoles at C4–C7 positions.<sup>5</sup> Especially, Shi,<sup>6</sup> You,<sup>7</sup> Yu,<sup>8</sup> Ackermann,<sup>9</sup>




**Figure 1.** C–H functionalization of indoles and directed C–H bond activation mode examples.

and others<sup>10</sup> pioneered the developments in benzenoid functionalization. C–H activation through C3 at the pyrrole

Received: March 29, 2022

Published: May 24, 2022




Scheme 1. Transition-Metal-Catalyzed Arylation of Indoles *via* C–H Functionalization

ring of indole has two directing modes. The first is vicinal activation, in which the directing group (DG) and the target C–H bond are at vicinal positions of the indole (Figure 1b).<sup>11</sup> With this activation mode for a suitable DG at the C3-position, a metallacycle is formed after C–H activation, which leads to C2-functionalization *via* the resulting metal–C bond. The second directing mode for DG on the indole C3-position involves nonvicinal activation to form a metallacycle for C4-functionalization (Figure 1b).<sup>7,11a,b</sup> Similarly, by installing a directing group at the indole nitrogen, both C2- and C7-activation have also been enabled.<sup>5j,11c</sup> However, both C5 and C6 positions lie even more remote from a directing group, and therefore, used strategies have mimicked those used for remote meta functionalization. These strategies include template-controlled palladium-catalyzed C–H alkenylation,<sup>8</sup> C–H alkylation *via*  $\sigma$ -activation,<sup>12</sup> and copper-catalyzed arylation using diaryliodonium salts as the arylating agent.<sup>13</sup>

Due to the remarkable importance of C4-functionalized indoles in natural products and medicinal chemistry, researchers have concentrated on the direct diversification of relatively less explored C-4 position of indoles *via* C–H

activation strategy. Therefore, alkenylation,<sup>10a,14</sup> acylation,<sup>10c</sup> amidation,<sup>7b,15</sup> allylation,<sup>16</sup> alkylation,<sup>17</sup> borylation,<sup>5f</sup> cyclization,<sup>18</sup> fluoroalkylation,<sup>6e,19</sup> and halogenation<sup>20</sup> of indoles at C4-positions have been demonstrated using different transition-metal catalysts (such as Rh, Ir, Pd, Co, and Ru) in the presence of appropriate directing groups. Despite the importance of biaryl compounds, a limited number of arylations at the C4-position through C–H activation have been performed so far. In 2017, Shi's group installed an elegant Pd(0)-catalyzed strategy for the C4-arylation of *N*-benzylindoles using a pivaloyl directing group (Scheme 1a).<sup>13a</sup> Almost simultaneously, Yu and Zhang's group showed a single example (in a yield of 82%) of Pd(II)-C4-arylation of *N*-tosyl-3-formylindole with methyl 4-iodobenzoate, in which the formyl group was installed in the C3 position of the indole (Scheme 1a).<sup>21</sup> In this work, 2-amino-2-methylpropanoic acid was used as a transient directing group. Later, Maiti and Volla's group reported a similar transformation for the C4-arylation of unprotected 3-formylindole (single example) and *N*-protected (methyl, tosyl, and benzyl) 3-formylindoles with aryl iodides using glycine as an inexpensive transient directing group

Table 1. Optimization of Reaction Conditions<sup>a</sup>


| entry | catalyst   | oxidant                                | temp (°C)  | solvent              | additive                   | time (h)   | yield (3aa, %) <sup>b</sup> |
|-------|--|--|------------|----------------------|----------------------------|------------|-----------------------------|
| 1–7   | Pd(OAc) <sub>2</sub>                               | AgOAc                                  | 100        | solvent <sup>c</sup> |                            | 10–12      | trace                       |
| 8     | Pd(OAc) <sub>2</sub>                               | AgOAc                                  | 100        | TFA                  |                            | 10         | 23                          |
| 9     | Pd(OAc) <sub>2</sub>                               | AgOAc                                  | 100        | HFIP                 | HOAc                       | 10         | 47                          |
| 10    | Pd(OAc) <sub>2</sub>                               | AgOAc                                  | 100        | HFIP                 | TFA                        | 10         | 57                          |
| 11–15 | Pd(OAc) <sub>2</sub>                               | AgOAc                                  | 100        | HFIP                 | other solvent <sup>d</sup> | 5          | trace                       |
| 16    | Pd(OAc) <sub>2</sub>                               | AgOAc                                  | 100        | HFIP                 | TFA                        | 5          | 70                          |
| 17    | <b>Pd(OAc)<sub>2</sub></b>                         | <b>AgOAc</b>                           | <b>100</b> | <b>HFIP</b>          | <b>TFA</b>                 | <b>3.5</b> | <b>87</b>                   |
| 18    | Pd(OAc) <sub>2</sub>                               | AgOAc                                  | 120        | HFIP                 | TFA                        | 3.5        | 72                          |
| 19    | Pd(OAc) <sub>2</sub>                               | AgOAc                                  | 65         | HFIP                 | TFA                        | 15         | 81                          |
| 20    | Pd(OAc) <sub>2</sub>                               | Ag <sub>2</sub> CO <sub>3</sub>        | 100        | HFIP                 | TFA                        | 3.5        | 45                          |
| 21    | Pd(OAc) <sub>2</sub>                               | Ag <sub>2</sub> O                      | 100        | HFIP                 | TFA                        | 3.5        | 70                          |
| 22    | Pd(OAc) <sub>2</sub>                               | Cu(OAc) <sub>2</sub> ·H <sub>2</sub> O | 100        | HFIP                 | TFA                        | 3.5        | trace                       |
| 23    | Pd(OAc) <sub>2</sub>                               | AgTFA                                  | 100        | HFIP                 |                            | 5          | trace                       |
| 24    | PdCl <sub>2</sub>                                  | AgOAc                                  | 100        | HFIP                 | TFA                        | 3.5        | 71                          |
| 25    | Pd(PPh <sub>3</sub> ) <sub>2</sub> Cl <sub>2</sub> | AgOAc                                  | 100        | HFIP                 | TFA                        | 3.5        | 67                          |
| 26    | Pd(TFA) <sub>2</sub>                               | AgOAc                                  | 100        | HFIP                 |                            | 3.5        | trace                       |
| 27    | Pd(TFA) <sub>2</sub>                               | AgTFA                                  | 100        | HFIP                 |                            | 36         | 74                          |
| 28    | <b>Pd(TFA)<sub>2</sub></b>                         | <b>AgOAc</b>                           | <b>100</b> | <b>HFIP</b>          | <b>TFA</b>                 | <b>3.5</b> | <b>87</b>                   |

<sup>a</sup>Reaction conditions: **1a** (0.40 mmol), Pd(OAc)<sub>2</sub> (10 mol %), **2a** (0.80 mmol), oxidant (0.80 mmol), solvent (1 mL), additive (1 mL). <sup>b</sup>Isolated yield. <sup>c</sup>Solvent: HFIP, HOAc, DCE, DMA, DMF, Tol, or TFE. <sup>d</sup>Other solvent: DCE, DMA, DMF, Tol, or TFE.


(Scheme 1a).<sup>10d</sup> Yang and You's group disclosed an iridium-catalyzed C2/C4-regioselective C–H heteroarylation of indoles with the help of a pivaloyl group at the C3-position (Scheme 1a/b).<sup>7a</sup> The oxidants Cu(OAc)<sub>2</sub>·H<sub>2</sub>O and Ag<sub>2</sub>O have been demonstrated to play a vital role in the C2/C4-regioselectivity. Recently, Punniyamurthy's group reported the palladium-catalyzed weak chelation-assisted regioselective C4-arylation of indoles utilizing arenes as the aryl source *via* a twofold C–H activation/C–C bond formation (Scheme 1a).<sup>22</sup> When using carboxylic acid as the directing group, *N*-heterocyclic carbene (NHC) and abnormal NHC (*a*NHC)-based Pd-catalyzed arylation reactions of *N*-alkylindole-2-carboxylic acids with aryl bromides and aryl chlorides as the coupling partners resulted in decarboxylative C2-arylation (Scheme 1c).<sup>23</sup> Synthesis of 2-arylindoles was also reported *via* a Pd-catalyzed decarboxylative strategy in water without base, oxidant, and ligand using diaryliodonium salts as the aryl partners (Scheme 1c).<sup>24</sup> Despite the above background, the examples of the directed C–H arylation of unprotected indoles remain extremely limited and include transient directing group strategies. At this point, it is important to note that An, Li, and Yang's group recently reported C4-arylation or domino C4-arylation/3,2-carbonyl migration (*via* migration of acetyl substituent from the C3- to C2-indole position) of 3-acetyl indoles *via* the different pathways by tuning either the Pd(I)–Pd(II) pathway or Pd(II) catalysis.<sup>25</sup> Our simultaneous results complete this research, and also expand its scope, and include new findings. Herein, we now wish to report the arylations for unprotected indoles including formyl, acetyl, carboxylic acid, and methyl ester groups as a directing group at the C3-position with aryl iodides and without a transient directing group using

the single Pd(II) catalyst system. With different directing groups, different pathways were observed (Scheme 1d).

## RESULTS AND DISCUSSION

In continuation of our research interest in C–H arylation reactions,<sup>26</sup> we started our investigation, focusing on the C–H arylation reactions of unprotected indole **1a** involving the formyl group at the C3-position as a directing group without a transient directing group. Initially, 1*H*-indole-3-carbaldehyde (**1a**) and iodobenzene (**2a**) were selected as model substrates to screen the reaction conditions (Table 1). When 10 mol % Pd(OAc)<sub>2</sub> was used as the catalyst and AgOAc (2 equiv) was used as the oxidant at 100 °C, among the employed solvents such as 1,1,1,3,3,3-hexafluoroisopropanol (HFIP), HOAc, 1,2-dichloroethane (DCE), *N,N*-dimethylformamide (DMF), toluene (Tol), trifluoroacetic acid (TFA), and 2,2,2-trifluoroethanol (TFE) (Table 1, entries 1–8), TFA gave the desired product **3aa** in 23% yield (Table 1, entry 8). The yields were improved to 47 and 58% when HOAc and TFA were used as the additive, respectively (Table 1, entries 9 and 10). Entries 9 and 10 showed that the use of TFA as a co-solvent was also effective. To test the effect of the co-solvents such as DCE, *N,N*-dimethylacetamide (DMA), DMF, Tol, or TFE, further optimization was conducted (Table 1, entries 11–15). Under these conditions, the formation of **3aa** was observed as trace amounts. To our delight, decreasing the reaction time to 3.5 h, we obtained the desired product **3aa** with an 87% isolated yield (Table 1, entries 16 and 17). When the reaction was carried out at 120 °C for 3.5 h, the yield decreased (Table 1, entry 18). The reaction at 65 °C for 15 h did not give the expected increase in yield (Table 1, entry 19). Other oxidants such as


Scheme 2. Substrate Scope of Iodoarenes and 3-Formylindoles



<sup>a</sup>Reaction conditions: **1a** (0.40 mmol), Pd(OAc)<sub>2</sub> (10 mol %), **2** (0.80 mmol), AgOAc (0.80 mmol), HFIP (1 mL), TFA (1 mL). <sup>b</sup>At 65 °C. <sup>c</sup>At 120 °C.

Ag<sub>2</sub>CO<sub>3</sub>, Ag<sub>2</sub>O, Cu(OAc)<sub>2</sub>·H<sub>2</sub>O, and AgTFA were also screened, but none of them achieved the same effect as silver acetate (Table 1, entries 20–23). Further experiments revealed that other palladium catalysts such as PdCl<sub>2</sub> and Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>, and Pd(TFA)<sub>2</sub> in HFIP (with and without TFA) were inferior to Pd(OAc)<sub>2</sub> (entry 24–27). Also, when the Pd(TFA)<sub>2</sub>/AgOAc catalyst system in HFIP/TFA was used, the C4-arylation proceeded cleanly, and **3aa** was isolated in 87% yield (entry 28). Based on the screening conditions described above, the use of Pd(OAc)<sub>2</sub> as a catalyst due to inexpensive AgOAc as an oxidant and TFA as an additive in HFIP at 100 °C was determined to be the optimal reaction conditions (entry 17).

With the optimized condition in hand, the substrate scope of this C–H arylation reaction was investigated, and the results are depicted in Scheme 2. Initially, both various electron-rich (Me, *t*-Bu, OMe) and electron-poor (Br, CO<sub>2</sub>Me, COMe, and CF<sub>3</sub>) aryl iodides **2a-I** were subjected to C4 arylation with 1*H*-indole-3-carbaldehyde (**1a**) under the standard conditions. Generally, reactions of aryl iodides possessing diverse substituents at different positions of the phenyl ring proceeded smoothly, and the desired C4 arylated products **3aa-ai** and **3al** were obtained in good to excellent yields (Scheme 2a). *ortho*-Methyl group at the iodoarene is not tolerated, suggesting that a sterically crowded intermediate is being formed during the coupling (Scheme 2a). Using 1-iodo-1,3-dimethylbenzene

Scheme 3. Substrate Scope of Iodoarenes and 3-Acetylindoles<sup>d</sup>


<sup>a</sup>Reaction conditions: **1a** (0.40 mmol), Pd(OAc)<sub>2</sub> (10 mol %), **2** (0.80 mmol), AgOAc (0.80 mmol), HFIP (1 mL), TFA (1 mL). <sup>b</sup>At 130 °C. <sup>c</sup>At 100 °C. <sup>d</sup>At 110 °C.

(**2k**) gave a reaction mixture that could not be purified. Then, the influence of the substitution pattern at the indole ring was investigated. The coupling of iodobenzene (**2a**) with a variety of substituted 1*H*-indole-3-carbaldehydes **1b-j** was tested (Scheme 2b). No arylation products **3ea-ga** were obtained from 5-substituted indoles, indicating that the reaction is very sensitive to steric hindrance at this position. Conversely, substituted indoles at C2, C7, or N1 positions readily couple with iodobenzene to give the corresponding C4 arylated products **3ba-da**, **3ha**, and **3ia** in excellent yields. Particularly noteworthy are the 7-halogen substrates in which the presence of the fluorine and bromine substituents at C7 does not hamper the arylation in the C4 position. The structures of C–H arylation products were confirmed by <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy and high-resolution mass spectrometry (HRMS). The arylation at C4-position in **1a** was also assigned by a

nuclear Overhauser effect (NOE) study. NOE correlations between the C2 hydrogen atom and *ortho*-hydrogen atoms of the C4 phenyl substituent of the aldehyde hydrogen atom within **3a** (see blue arrows) support the relative arylation depicted (Scheme 2a).

Notably, 3-acetylindole (**4a**) was subjected to reaction with **2a** under the determined optimized reaction conditions for indole-3-aldehydes, and the desired product **5aa** was obtained in 83% yields (Scheme 3a). Recently, An, Li, and Yang's group researched the same reaction and reported this domino C4-arylation/3,2-carbonyl migration and structures of corresponding products.<sup>25</sup> The structure of products was assigned by NMR spectra and HRMS. At the <sup>1</sup>H NMR spectrum, the chemical shift of the C2-H (low-field) and C3-H (high-field) protons for indoles is a characteristic indication. The C3-H resonance ( $\delta$  7.34 ppm) of **5aa** appeared at a lower field than




Scheme 4. Substrate Scope of Iodoarenes and 3-Acetyl-*N*-Benzylindoles<sup>c</sup>

<sup>a</sup>Reaction conditions: **1a** (0.20 mmol), Pd(OAc)<sub>2</sub> (10 mol %), **2** (0.40 mmol), AgOAc (0.40 mmol), HFIP (1 mL), TFA (1 mL). <sup>b</sup>At 110 °C. <sup>c</sup>At 120 °C. <sup>d</sup>At 90 °C. <sup>e</sup>At 75 °C.

the C2-H resonance ( $\delta$  7.87 ppm) of **4a**. Comparison of the NMR spectra of **4a** and **5aa**, the disappearance of characteristic C2-H resonance, and the appearance of a new C3-H resonance at high-field indicated the formation of **5aa**, which reveals that the ketone group is migrated under these conditions. Additionally, for domino C4-arylation/1,2-carbonyl migration product **5aa**, structure assignment was confirmed according to NOE signals (see blue arrows) between acetyl methyl protons and C3-H/N-H protons (Scheme 3a). The evaluation of the substrate scope for this transformation is depicted in Scheme 3a. A wide variety of aryl iodides **2a-l** (except **2j**) were also well tolerated by the palladium catalyst to deliver domino C4-arylation/1,2-carbonyl migration products **5ab-al** (except **5aj**). For example, aryl iodides **2b-d** with the electron-donating

groups at the para and meta positions, such as methyl and *tert*-butyl, promoted this transformation smoothly. The corresponding products (**5ab-ad**) were obtained in excellent yields (74–81%). *para*-Methoxy-substituted iodobenzene gives the product **5af** in a high yield (70%). In contrast, aryl iodides **2e** and **2g-i** substituted with electron-withdrawing groups, such as –Br, –COOMe, –COMe, and –CF<sub>3</sub>, also reacted with **4a** smoothly, giving the corresponding products **5ae** and **5ag-ai** in 53–80% yields. When *ortho*-methyl-substituted iodobenzene **5j** was employed under standard conditions, no corresponding product **5aj** was determined. This entry indicated that the steric hindrance of *ortho*-substitution had a significant effect on the progress of the reaction. In addition, product **5al** was obtained in a low yield (53%) when the –NO<sub>2</sub> substitution as

Scheme 5. Substrate Scope of Iodoarenes and 1*H*-Indole-3-Carboxylic Acid or Methyl 1*H*-Indole-3-Carboxylate<sup>c</sup>

<sup>a</sup>Reaction conditions: 10a (or 10b) (0.40 mmol), Pd(OAc)<sub>2</sub> (10 mol %), 2 (0.80 mmol), AgOAc (0.80 mmol), HFIP (1 mL), TFA (1 mL). <sup>b</sup>For R = H. <sup>c</sup>For R = Me.

a strong electron-withdrawing substituent exists on the meta position. When 3,4-dimethyliodobenzene (2k) was used as a substrate, the corresponding product 5ak was obtained in good yield (73%). To further widen the scope of this domino C4-arylation/3,2-carbonyl migration strategy, 3-acetylindoles 4b–d with iodobenzene (2a) were subjected to optimized reaction conditions (Scheme 3b). Also, 3-acetyl-7-fluoro-1*H*-indole (4b) worked well to yield 5ba in good yield (73%). Interestingly, 3-acetyl-7-bromo-1*H*-indole (4c) afforded C4-arylated product 6ca (76% yield) in which the acetyl group did not migrate at 120 °C for 12 h, while domino C4-arylation/3,2-carbonyl migration product 5ca was obtained when the reaction was performed at 120 °C for 24 h. In the case of C2-protected indole, 3-acetyl-2-methyl-1*H*-indole (4e) provided C4-arylated product 7da with directing group removal.





Shi's group reported C4/C5-arylation (single examples) in a lower yield of an *N*-Bn-protected indole bearing directing groups at the C3 position, such as formyl, acetyl, and isobutyryl substituents.<sup>13</sup> To confirm the importance of the NH-unprotected indoles for this unusual DG migration, we investigated the reaction between 3-acetyl-*N*-benzylindole 8a with a variety of aryl iodides 2a–l containing a wide variety of electron-donating or electron-withdrawing substituents under the same optimized reaction conditions (Scheme 4a). No significant electronic effect on the reaction progress was observed. Aryl iodides with substituents on meta, para positions delivered corresponding C4-arylation products 9aa–al in good to excellent yields. But 2-iodotoluene (9j) led to a trace amount of product 9aj. Substrates 8b and 8c containing fluoro and bromo participated in C4-arylation smoothly in excellent yields (Scheme 4b). Compared to 3-acetyl-*N*-benzylindole (8a), 3-acetyl-*N*-methylindole (8e) gave C4 arylation products 9ea and 9eg in almost similar yields. However, the presence of the methyl group on the C2-position of indole led to an unpurified reaction mixture (Scheme 4b, for 9d).

With the catalyst system in hand, we next examined the scope of acid/ester-directed C–H arylation with iodoarenes using *N*-unprotected 1*H*-indole-3-carboxylic acid (10a) and methyl 1*H*-indole-3-carboxylate (10b) (Scheme 5). First, the



indole-3-carboxylic acid (10a) without substituents on position N1 with aryl iodides (2a–e and 2k) was tested. All gave the corresponding decarboxylative C2-arylation products 11a–e and 11k in high yields (75–87%) with complete site selectivity.<sup>27–30</sup> However, the methyl 1*H*-indole-3-carboxylate (11b) was effective and gave the same products 11a–e and 11k successfully, but in moderate yields (60–66%). Since this catalytic process yields decarboxylative C2-arylation products, we have limited the scope of aryl iodides.

To explore the practical utility of this C(sp<sup>2</sup>)–H arylation reaction, a gram-scale reaction of 3-formylindole (1a) was carried out under the standard conditions (Scheme 6a). The desired arylation product 3aa could be obtained in 69% yield. Also, to confirm the synthetic utility of this domino C4-arylation/1,2-carbonyl migration process, a gram-scale reaction of 3-acetylindole (4a, 6.3 mmol) and iodobenzene (2a, 25.1 mmol) was conducted under the standard conditions. A total of 0.98 g of compound 5aa was obtained in a satisfying yield (65%), which was comparable with the small-scale reaction (Scheme 6b). Furthermore, the potential applications of both domino C4-arylation/1,2-carbonyl migration products and C4-arylation products as useful synthetic blocks are illustrated (Scheme 6c–e). To perform structure confirmation as well as synthetic diversification, the benzylation reaction of 5aa gave the N1-benzylated indole derivative 12 in good yields (Scheme 6c). Comparison of the NMR spectra of 12 and 9aa is important to confirm that the acetyl group does not migrate during the arylation of 3-acetyl-*N*-benzylindole 8a under the standard conditions. Besides C–H arylation on the C4-position of the indole ring, the bromine group on the 7-position of 5ca has a potential derivatization site. The classical palladium-catalyzed Suzuki–Miyaura coupling of 5ca with thiophene-2-boronic acid provided 13 in excellent yield (89%) (Scheme 6d).<sup>31</sup> Pityiacitrin (14) is a 1-indolyl- $\beta$ -carboline alkaloid isolated from various sources and has also been tested for various biological activities.<sup>32</sup> To demonstrate utility in the synthesis of the functionalized natural product of C4-arylated acetylindoles, the synthesis of the substituted pityiacitrin starting from 9ad was carried out successfully (Scheme 6e). For this, 9ad and 5-methoxytryptamine (15) were reacted with

## Scheme 6. Gram-Scale Reaction and Synthetic Applications

a) Gram-scale reaction for **3aa**b) Gram-scale reaction for **5aa**c) N-Benzylation reaction of **5aa**d) Cross-coupling of 2-thienyl boronic acid and **5ca**

## e) Construction of functionalized natural product



1.0 equiv of  $\text{I}_2$  and 1.5 equiv of  $\text{H}_2\text{O}_2$  in dimethyl sulfoxide (DMSO) at 100 °C for 24 h to form substituted pityiacitrin **16**. The protocol involves Kornblum oxidation to form indolyl-

glyoxal, Pictet–Spengler condensation with tryptamine to yield dihydro- $\beta$ -carboline, and finally, aromatization of dihydro- $\beta$ -carboline to provide the desired product **16**. This one-pot




process successfully led to the synthesis of structurally related analogue **17** of pityiacitrin from **5aa** (Scheme 6e).


To further understand both structure characterization and the progress of these reactions, several control experiments were performed. During arylation of **4c**, product **6ca** was also isolated in 76% yield (Scheme 3b). Under the standard conditions, **6ca** itself provided 3,2-carbonyl migration product **5ca** in 74% yield (Scheme 7a). In fact, the C–H arylation

### Scheme 7. Control Experiments


(a) 3,2-Carbonyl migration from **6ca**




(b) Reaction under standard conditions of **10a** without **2a**




(c) Reaction under standard conditions of **19a** and **2a**



(d) Reaction under standard conditions of **10c** without **2a** / with **2a**



(e) Reaction under standard conditions of **19b** and **2a**





reaction of *N*-unprotected 3-acetylindoles **4a–d** can follow two paths: respectively, migration/arylation or arylation/migration. This result strongly supports both the migration of the acetyl group after the arylation reaction and the reported results of An, Li, and Yang's group.<sup>25</sup> To understand the mechanism of the decarboxylative arylation reaction, several control experiments were performed (Scheme 7b–e). First, indole-3-carboxylic acid (**10a**) (or methylindole-3-carboxylate (**10b**)) was subjected to standard reaction conditions without using

substrate **2a**, wherein we isolated 3-acylation product **18** in 75% (or 70%) yield as a sole product (Scheme 7b).<sup>33</sup> We first assume that indole-3-carboxylic acid yields indole (**19a**) via palladium-catalyzed decarboxylation. Then, product **18** is formed from the acylation of indole (**19a**) with TFA. In addition, when indole (**19a**) and **2a** were reacted under the standard reaction conditions, 2-phenylindole (**11a**) was obtained in 70% yield (Scheme 7c). These results indicate that indole (**19a**) might be the key intermediate for the decarboxylative arylation reaction. Furthermore, the reaction of 2-methylindole-3-carboxylic acid (**10c**)<sup>34</sup> without **2a**/with **2a** under standard reaction conditions gave 2-methylindole (**19b**) via decarboxylation (Scheme 7d). But the treatment of **19b** with **2a** did not provide any product (Scheme 7e). These experiments showed that C–H arylation could not proceed via C2 protection under our standard conditions.

To probe the role of TFA in C4-arylation, optimization experiments were carried out (Table 1, especially entries 17 and 28). The role of TFA can be categorized into two aspects: (i) Pd(OAc)<sub>2</sub> could be readily converted to Pd(TFA)<sub>2</sub> to catalyze reactions (Table 1, entries 8, 10, 16–21, 24, and 25); (ii) it may affect the reaction rate or by increasing the solubility of the reaction mixture (Table 1, entries 26–28). Based on previous reports,<sup>25,35</sup> our findings, and control experiments, possible catalytic mechanisms are illustrated in Scheme 8. Initially, the active catalytic species **A** is formed from Pd(II) catalyst (Pd(TFA)<sub>2</sub>, a catalyst formed *in situ* from Pd(OAc)<sub>2</sub>

### Scheme 8. Proposed Pd<sup>II</sup>–Pd<sup>IV</sup> Catalytic Cycle for C4-Arylation and Migration Mechanism



Scheme 9. Proposed Pd<sup>II</sup>-Pd<sup>IV</sup> Catalytic Cycle for C2-Arylation

and TFA) and 3-formylindole (**1a**) (or 3-acetylindole) (or **4a**) in the presence of AgOAc, followed by the C–H bond activation to form the cyclometalated Pd(II) intermediate **B**. The intermediate **B** further underwent oxidative addition with aryl iodide **2** to produce diaryl Pd(IV) species **C**. Reductive elimination of **C** produced the desired C4-arylated product **3** (or **6**) and regeneration of the active Pd(II) species by AgOAc to regenerate the catalytic cycle. To explain carbonyl migration, An, Li, and Yang's group conducted many independent experiments and mechanism studies and proposed a plausible reaction pathway.<sup>25</sup> Accordingly, after the formation of Pd(II)-catalyzed C4-arylation product **6**, the 3,2-carbonyl migration (or Friedel–Crafts acyl rearrangement) process takes place. The first step would be a reaction between **6** and TFA, yielding the product **D** and mixed (or unsymmetrical) acid anhydride **E** via protonation and reverse Friedel–Crafts process. Next, TFA (or Pd(TFA)<sub>2</sub>)-promoted intermolecular Friedel–Crafts reaction takes place between **D** and **E** to yield selectively migration product **5** through acylation rearomatization. This unusual Friedel–Crafts acyl rearrangement is not reversible. We speculate that C3-acetylated product **6** was the kinetically controlled product, whereas C2-acetylated product **5** was the thermodynamically controlled product. NH-free 3-acetylindoles underwent unusual migration of the acetyl group to the C2-position following C4-arylation in one pot, whereas *N*-alkylated 3-acetylindoles showed C4-arylations without migration. These results indicate that an alkyl group at the nitrogen atom of the indoles plays a crucial role to prevent the migration of the acetyl group. We believe the existence of iminium intermediate **F** via a hydrogen-bond interaction between the CF<sub>3</sub>COO<sup>−</sup> and the NH-free indole, which could promote both reverse Friedel–Crafts process and 1,2-acetyl migration process.

Two different pathways for the formation of decarboxylative C2-arylated indoles **11**, through either the C–H arylation/decarboxylation process (path 1) or the decarboxylation/C–H arylation process (path 2), could be evaluated (Scheme 9). Based on the above control results and literature reports,<sup>26,36</sup> a plausible reaction pathway is path 2. In this context, the

mechanism involves initial palladation at C3 followed by palladium migration to C2 via the Pd(II)/Pd(IV) pathway. Accordingly, this reaction is progressing to yield aryl-palladium intermediate indole cations **G** and **H**, followed by oxidative addition of palladium(II) to the ArI. This aryl-palladium intermediate **I** undergoes sequential reductive elimination to afford the C2-arylated product **11** and PdIOAc or PdI<sub>2</sub> (from 2 turnovers). Finally, the catalytically active Pd(OAc)<sub>2</sub> (or Pd(TFA)<sub>2</sub>) for the cycle is regenerated from inactive PdIOAc or PdI<sub>2</sub> via silver salt and TFA.

## CONCLUSIONS

In summary, we have reported Pd(II)-catalyzed protocol for accessing arylated indole scaffolds utilizing iodoarenes as the aryl source via the C–H bond activation of *N*-unprotected indoles with the aid of readily accessible carbonyl directing groups (aldehyde, acetyl, carboxylic acid, and methyl ester) at the C3 position. The protocol is operationally simple and utilizes Pd(OAc)<sub>2</sub> as a catalyst, AgOAc as an oxidant, and TFA as an additive in HFIP at 65–120 °C. The substrate scope is broad and displays excellent selectivity (C4-arylation for 3-formylindoles and *N*-protected 3-acetylindoles, C4-arylation including 3,2-carbonyl migration for 3-acetylindoles, C2-arylation via decarboxylation/arylation steps for both indole-3-carboxylic acid and methylindole-3-carboxylate). Based on the control experiments and the literature, plausible mechanisms are proposed via Pd(II)/Pd(IV) catalytic cycles. To show the synthetic utility of both our catalytic system and the arylation products, gram-scale reaction and synthetic applications were performed. In this context, C4-arylated acetylindoles allowed us to construct the functionalized and structural-related analogues of ptyiacitrin.

## EXPERIMENTAL SECTION

**General Information.** Unless otherwise mentioned, all reagents and solvents from commercial sources were used without further purification. NMR spectra were recorded in CDCl<sub>3</sub>, DMSO-*d*<sub>6</sub>, or acetone-*d*<sub>6</sub> solvents at 400 MHz (<sup>1</sup>H) and 100 MHz (<sup>13</sup>C), respectively. Chemical shifts (δ) are reported in parts per million (ppm), using the residual solvent peak in CDCl<sub>3</sub> (δ = 7.26 ppm for

$^1\text{H}$  NMR and  $\delta = 77.0$  ppm for  $^{13}\text{C}$  NMR), DMSO- $d_6$  ( $\delta = 2.50$  ppm for  $^1\text{H}$  NMR and  $\delta = 39.4$  ppm for  $^{13}\text{C}$  NMR), and acetone- $d_6$  ( $\delta = 2.05$  ppm for  $^1\text{H}$  NMR and  $\delta = 29.8$  ppm for  $^{13}\text{C}$  NMR) as an internal standard, and coupling constants ( $J$ ) are indicated in hertz (Hz). Signal multiplicities are abbreviated as s = singlet, d = doublet, t = triplet, q = quartet, dd = doublet of doublets, m = multiplet, and br = broad. High-resolution mass spectrometry (HRMS) of all compounds was performed using a quadrupole time-of-flight (QTOF) spectrometry device. Column chromatography was performed using silica gel (70–230 mesh).

**General Procedures.** *General Procedure A: Preparation of 1H-Indole-3-carbaldehydes (1b–g).*<sup>37</sup> 1H-Indole-3-carbaldehydes **1b–g** were prepared according to the reported literature method.<sup>37</sup> Pyrophosphoryl chloride (0.9 mL, 9.4 mmol, 1.1 equiv) was added dropwise to a stirred mixture of *N,N*-dimethylformamide (2.8 mL, 36.7 mmol, 4.3 equiv) at 10–20 °C. To this mixture, a solution of indole substrate (8.5 mmol) in *N,N*-dimethylformamide (1.67 mL, 5 M for indole substrate) was added slowly, keeping the temperature at 20–30 °C. The mixture was then stirred in a preheated oil bath at 35 °C for 45 min. After completion of the reaction, the mixture was cooled to room temperature and carefully quenched with crushed ice (3.5 g). The mixture was stirred vigorously, and further crushed ice (3.5 g) was added, followed by a solution of NaOH (3.77 g, 94.3 mmol, 11 equiv) in water (10 mL). Then, the mixture was heated under reflux for 15 min, and the precipitate was filtered, washed with water ( $3 \times 20$  mL), and dried in vacuo to afford 1H-indole-3-carbaldehydes **1b–g**.

*General Procedure B: Preparation of 3-Acetyl-1H-indoles (4a–d).*<sup>38</sup> 3-Acetyl-1H-indoles **4a–d** were prepared according to the reported literature method.<sup>38</sup>  $\text{SnCl}_4$  (0.75 mL, 6 mmol, 1.2 equiv) was added dropwise into a stirred solution of 1H-indole derivative (5 mmol) in dichloromethane (10 mL, 2 M for 1H-indole derivative) at 0 °C in an inert atmosphere of  $\text{N}_2$ . The resultant mixture was warmed to room temperature and stirred for 30 min, then acetic anhydride (510 mg, 0.5 mL, 5 mmol, 1 equiv) was added, followed by nitromethane (7.5 mL), and stirred for 2 h at room temperature. After the completion of the reaction, the reaction was quenched by the addition of ice and water (20 mL), then extracted with dichloromethane ( $3 \times 30$  mL), and dried over anhydrous  $\text{Na}_2\text{SO}_4$ . The solvent was evaporated, and the crude product was then purified by silica gel chromatography to give 3-acetyl-1H-indoles.

*General Procedure C: Preparation of N-Alkyl Indoles (1h–i and 8a–d).*<sup>10a</sup> To a suspension of NaH (1.1 mmol, 1.1 equiv, 60% dispersion in mineral oil) in THF at 0 °C, a solution of 1H-3-acetylindole (or 1H-indole-3-carbaldehyde) (1 mmol) in THF (5 mL, 0.2 M for 1H-3-acetylindole or 1H-indole-3-carbaldehyde) was added dropwise. Benzyl bromide (methyl iodide for methylation) (1.1 mmol, 1.1 equiv) was then added dropwise to this solution and stirred for 12 h at room temperature. After completion, the reaction was quenched with water and extracted by EtOAc ( $2 \times 30$  mL). The combined organic phase was dried over  $\text{Na}_2\text{SO}_4$ , then concentrated under reduced pressure, and the residue was purified on silica gel column chromatography to provide the corresponding *N*-alkyl 3-acetyl indoles (or *N*-alkyl 3-carbaldehydes) indoles.

*General Procedure D: C–H Arylation.* The indole material **1** (or **4**, or **8**, or **10**) (0.4 mmol),  $\text{Pd}(\text{OAc})_2$  (9 mg, 40  $\mu\text{mol}$ , 10 mol %), and  $\text{AgOAc}$  (133 mg, 0.8 mmol, 2 equiv) were weighed in air and placed in a sealed tube (15 mL) with a magnetic stir bar. To the reaction mixture, aryl iodide **2** (0.8 mmol, 2 equiv) and HFIP/TFA (2 mL, 1:1, v/v, 0.2 M for the indole material **1** (or **4**, or **8**, or **10**)) were added. The reaction mixture was then stirred in a preheated oil bath at 65–130 °C for 3–24 h. Upon completion, the reaction mixture was cooled to room temperature, the solvents were removed under reduced pressure, and the resulting mixture was purified by a silica gel column chromatography column to obtain the corresponding C–H arylation product using hexane/EtOAc as the eluent.

**Spectral Data of Starting Materials.** *7-Fluoro-1H-indole-3-carbaldehyde (1b).*<sup>39</sup> Off-white solid, mp: 140–141 °C;  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ ):  $\delta$  12.70 (bs, NH, 1H), 9.97 (s, CHO, 1H), 8.37 (s, CH, 1H), 7.91 (d,  $J = 7.8$  Hz, CH, 1H), 7.24–7.16 (m, CH,

1H), 7.13–7.07 (m, CH, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz, DMSO- $d_6$ ):  $\delta$  185.2, 149.0 (d,  $J = 245.2$  Hz), 139.0 (s), 127.7 (d,  $J = 4.5$  Hz), 124.7 (d,  $J = 13.2$  Hz), 122.8 (d,  $J = 6.0$  Hz), 118.7, 116.9 (d,  $J = 3.6$  Hz), 108.5 (d,  $J = 15.9$  Hz).

*7-Bromo-1H-indole-3-carbaldehyde (1c).*<sup>40</sup> Yellow solid, mp: 166–167 °C;  $^1\text{H}$  NMR (400 MHz, Acetone- $d_6$ ):  $\delta$  11.32 (bs, NH, 1H), 10.06 (s, CHO, 1H), 8.29 (s, CH, 1H), 8.23 (d,  $J = 8.2$  Hz, CH, 1H), 7.51–7.49 (m, CH, 1H), 7.20 (t,  $J = 7.8$  Hz, CH, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz, Acetone- $d_6$ ):  $\delta$  185.7, 138.5, 136.7, 127.2, 127.0, 124.5, 121.7, 120.9, 105.5.

*2-Methyl-1H-indole-3-carbaldehyde (1d).*<sup>41</sup> White solid, mp: 200–201 °C;  $^1\text{H}$  NMR (400 MHz, Acetone- $d_6$ ):  $\delta$  10.94 (bs, NH, 1H), 10.18 (s, CHO, 1H), 8.29–8.09 (m, CH, 1H), 7.52–7.34 (m, CH, 1H), 7.27–7.09 (m, CH, 2H), 2.75 (s, CH<sub>3</sub>, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz, Acetone- $d_6$ ):  $\delta$  184.6, 148.5, 136.5, 127.1, 123.6, 122.8, 121.3, 115.3, 112.0, 11.8.

*5-Bromo-1H-indole-3-carbaldehyde (1e).*<sup>40</sup> White solid, mp: 204–205 °C;  $^1\text{H}$  NMR (400 MHz, Acetone- $d_6$ ):  $\delta$  11.31 (bs, NH, 1H), 10.02 (s, CHO, 1H), 8.39 (d,  $J = 1.9$  Hz, CH, 1H), 8.26 (s, CH, 1H), 7.53 (d,  $J = 8.6$  Hz, CH, 1H), 7.40 (dd,  $J = 8.6$ , 2.0 Hz, CH, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz, Acetone- $d_6$ ):  $\delta$  185.4, 139.0, 137.0, 127.3, 127.2, 124.6, 119.4, 116.1, 115.0.

*5-(Benzyloxy)-1H-indole-3-carbaldehyde (1f).*<sup>42</sup> Yellow solid, mp: 108–109 °C;  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ ):  $\delta$  12.10 (bs, NH, 1H), 9.89 (s, CHO, 1H), 8.22 (s, CH, 1H), 7.70 (d,  $J = 2.5$  Hz, CH, 1H), 7.52–7.46 (m, CH, 2H), 7.46–7.37 (m, CH, 3H), 7.35–7.29 (m, CH, 1H), 6.97 (dd,  $J = 8.7$ , 2.5 Hz, CH, 1H), 5.12 (s, CH<sub>2</sub>, 2H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz, DMSO- $d_6$ ):  $\delta$  184.8, 154.6, 138.5, 137.4, 132.0, 128.3, 127.6, 127.6, 124.8, 118.0, 113.8, 113.2, 104.0, 69.6.

*5-Nitro-1H-indole-3-carbaldehyde (1g).*<sup>40</sup> Yellow solid, mp: 292–293 °C;  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ ):  $\delta$  10.02 (s, CHO, 1H), 8.93 (s, CH, 1H), 8.56 (s, CH, 1H), 8.14 (d,  $J = 8.2$  Hz, CH, 1H), 7.71 (d,  $J = 8.2$  Hz, CH, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz, DMSO- $d_6$ ):  $\delta$  185.4, 142.7, 141.6, 140.3, 123.5, 119.0, 118.6, 117.0, 113.2.

*1-Benzyl-1H-indole-3-carbaldehyde (1h).*<sup>40</sup> White solid, mp: 107–108 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  9.99 (s, CHO, 1H), 8.46–8.23 (m, CH, 1H), 7.70 (s, CH, 1H), 7.40–7.28 (m, CH, 6H), 7.21–7.16 (m, CH, 2H), 5.35 (s, CH<sub>2</sub>, 2H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  184.6, 138.6, 137.5, 135.3, 129.1, 128.4, 127.2, 125.5, 124.2, 123.1, 122.2, 118.5, 110.4, 50.9.

*1-Methyl-1H-indole-3-carbaldehyde (1i).*<sup>40</sup> Pale-brown solid, mp: 68–69 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  9.94 (s, CHO, 1H), 8.34–8.30 (m, CH, 1H), 7.60 (s, CH, 1H), 7.41–7.28 (m, CH, 3H), 3.80 (s, CH<sub>3</sub>, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  184.5, 139.5, 137.9, 125.2, 124.0, 122.9, 121.9, 117.9, 110.0, 33.6.

*1-Tosyl-1H-indole-3-carbaldehyde (1j).*<sup>43</sup> 1-Tosyl-1H-indole-3-carbaldehyde (**1j**) were prepared according to the reported literature method.<sup>41</sup> Purple solid, mp: 139–140 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.09 (s, CHO, 1H), 8.25 (d,  $J = 7.3$  Hz, =CH, 1H), 8.23 (s, =CH, 1H), 7.95 (d,  $J = 7.9$  Hz, =CH, 1H), 7.87–7.82 (m, AA' part of AA'/BB' system, =CH, 2H), 7.44–7.32 (m, =CH, 2H), 7.31–7.26 (m, BB' part of AA'/BB' system, =CH, 2H), 2.36 (s, CH<sub>3</sub>, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  185.4, 146.2 (2C), 136.2, 135.2, 134.4, 130.3, 127.2, 126.3, 125.1, 122.6, 122.4, 113.3, 21.6.

*1-(1H-indol-3-yl)ethan-1-one (4a).*<sup>10a</sup> White solid, mp: 189–190 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.93 (bs, NH 1H), 8.43–8.37 (m, CH, 1H), 7.87 (d,  $J = 3.0$  Hz, CH, 1H), 7.48–7.39 (m, CH, 1H), 7.35–7.23 (m, CH, 2H), 2.56 (s, CH<sub>3</sub>, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  194.2, 136.6, 132.3, 125.4, 123.6, 122.6, 122.2, 118.2, 111.7, 27.6.

*1-(7-Fluoro-1H-indol-3-yl)ethan-1-one (4b).* White solid, mp: 195–196 °C;  $^1\text{H}$  NMR (400 MHz, Acetone- $d_6$ ):  $\delta$  11.41 (bs, NH, 1H), 8.28 (d,  $J = 3.0$  Hz, CH, 1H), 8.12 (d,  $J = 8.0$  Hz, CH, 1H), 7.20–7.14 (m, CH, 1H), 7.01 (dd,  $J = 11.5$ , 8.0 Hz, CH, 1H), 2.50 (s, CH<sub>3</sub>, 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz, Acetone- $d_6$ ):  $\delta$  193.2 (d,  $J = 0.9$  Hz), 150.3 (d,  $J = 243.8$  Hz), 134.5 (d,  $J = 17.1$  Hz), 130.4 (d,  $J = 4.4$  Hz), 125.8 (d,  $J = 13.4$  Hz), 123.2 (d,  $J = 6.0$  Hz), 119.3 (d,  $J = 1.6$  Hz), 118.9 (d,  $J = 3.7$  Hz), 108.6 (d,  $J = 15.8$  Hz), 27.5. HRMS (ESI-TOF)  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{10}\text{H}_9\text{FNO}$ : 178.0663; found: 178.0659.



**1-(7-Bromo-1H-indol-3-yl)ethan-1-one (4c).**<sup>44</sup> White solid, mp: 191–192 °C; <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 12.16 (bs, NH, 1H), 8.35 (s, CH, 1H), 8.19 (d, *J* = 7.7 Hz, CH, 1H), 7.44 (d, *J* = 7.7 Hz, CH, 1H), 7.12 (t, *J* = 7.7 Hz, CH, 1H), 2.48 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, DMSO-*d*<sub>6</sub>): δ 193.0, 135.1, 135.0, 126.9, 125.4, 123.1, 120.7, 117.6, 104.6, 27.4.

**1-(2-Methyl-1H-indol-3-yl)ethan-1-one (4d).**<sup>45</sup> Off-white solid, mp: 201–202 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.70 (bs, NH, 1H), 8.03 (d, *J* = 7.8 Hz, CH, 1H), 7.37–7.33 (m, CH, 1H), 7.29–7.19 (m, CH, 2H), 2.76 (s, CH<sub>3</sub>, 3H), 2.67 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 194.8, 143.7, 134.5, 127.0, 122.4, 122.1, 120.9, 114.7, 110.8, 31.3, 15.5.

**1-(1-Benzyl-1H-indol-3-yl)ethan-1-one (8a).**<sup>10a</sup> White solid; mp 114–115 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.36–8.28 (m, CH, 1H), 7.64 (s, CH, 1H), 7.29–7.13 (m, CH, 6H), 7.08–7.04 (m, CH, 2H), 5.23 (s, CH<sub>2</sub>, 2H), 2.41 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 193.0, 137.1, 135.8, 135.0, 129.1, 128.2, 127.0, 126.5, 123.5, 122.7 (2C), 117.5, 110.2, 50.7, 27.7.

**1-(1-Benzyl-7-fluoro-1H-indol-3-yl)ethan-1-one (8b).** White solid; mp 132–133 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.19 (d, *J* = 8.0 Hz, CH, 1H), 7.69 (s, CH, 1H), 7.41–7.29 (m, CH, 3H), 7.22–7.14 (m, CH, 3H), 6.95 (dd, *J* = 12.6, 7.9 Hz, CH, 1H), 5.50 (s, CH<sub>2</sub>, 2H), 2.50 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 192.9, 149.8 (d, *J* = 245.5 Hz), 136.6, 136.1, 130.1 (d, *J* = 4.1 Hz), 129.0, 128.2, 127.0, 124.9 (d, *J* = 9.1 Hz), 123.1 (d, *J* = 6.7 Hz), 118.5 (d, *J* = 3.9 Hz), 118.0, 109.4 (d, *J* = 17.7 Hz), 53.0 (d, *J* = 6.1 Hz), 27.7. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>15</sub>FNO: 268.1132; found: 268.1130.

**1-(1-Benzyl-7-bromo-1H-indol-3-yl)ethan-1-one (8c).** White solid; mp 154–153 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.47 (dd, *J* = 8.0, 0.9 Hz, CH, 1H), 7.70 (s, CH, 1H), 7.43 (dd, *J* = 7.6, 0.8 Hz, CH, 1H), 7.36–7.29 (m, CH, 3H), 7.13 (t, *J* = 7.9 Hz, CH, 1H), 7.06–7.01 (m, CH, 2H), 5.85 (s, CH<sub>2</sub>, 2H), 2.49 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 192.8, 137.8, 137.4, 133.6, 129.5, 129.1, 129.0, 127.9, 126.3, 123.9, 122.2, 117.1, 104.0, 52.2, 27.7. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>15</sub>BrNO: 328.0332; found: 328.0329.

**1-(1-Benzyl-2-methyl-1H-indol-3-yl)ethan-1-one (8d).**<sup>10a</sup> White solid; mp 96–97 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.12–8.08 (m, CH, 1H), 7.39–7.31 (m, CH, 5H), 7.29 (dd, *J* = 7.0, 1.4 Hz, CH, 1H), 7.06 (d, *J* = 6.5 Hz, CH, 2H), 5.43 (s, CH<sub>2</sub>, 2H), 2.80 (s, CH<sub>3</sub>, 3H), 2.79 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 194.8, 144.9, 136.5, 136.1, 129.0, 127.8, 126.5, 125.9, 122.3, 122.1, 120.8, 114.7, 110.0, 46.4, 31.8, 12.7.

**1-(1-Benzyl-2-methyl-1H-indol-3-yl)ethan-1-one (8e).**<sup>10a</sup> White solid; mp 101–102 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.43–8.31 (m, CH, 1H), 7.67 (s, CH, 1H), 7.40–7.24 (m, CH, 3H), 3.82 (s, CH<sub>3</sub>, 3H), 2.51 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 192.9, 137.5, 135.8, 126.2, 123.3, 122.5 (2C), 116.9, 109.6, 33.5, 27.6.

**2-Methyl-1H-indole-3-carboxylic Acid (10c).**<sup>34</sup> (Pink solid; mp: 174–175 °C); <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): 11.86 (bs, NH, OH, 2H), 7.94–7.89 (m, CH, 1H), 7.37–7.31 (m, CH, 1H), 7.11–7.05 (m, CH, 2H), 2.64 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, DMSO-*d*<sub>6</sub>): δ 166.7, 144.3, 134.7, 127.2, 121.3, 120.6, 120.4, 111.0, 103.2, 13.7.

**Spectral Data for C–H Arylation Products.** **4-Phenyl-1H-indole-3-carbaldehyde (3aa).** Compound 3aa was synthesized by following general procedure D using 1H-indole-3-carbaldehyde (1a, 58 mg, 0.4 mmol) and iodobenzene (2a, 90 μL, 0.8 mmol) at 100 °C for 3.5 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): 3aa (76 mg, 87%, a brown solid, mp: 151–152 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 10.05 (bs, NH, 1H), 9.52 (s, CHO, 1H), 8.01 (d, *J* = 2.6 Hz, CH, 1H), 7.57–7.51 (m, CH, 2H), 7.50–7.39 (m, CH, 4H), 7.33 (t, *J* = 7.5 Hz, CH, 1H), 7.20 (d, *J* = 7.5 Hz, CH, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 187.2, 141.9, 137.1, 135.6, 131.8, 129.0, 128.6, 127.7, 124.2, 123.9, 123.4, 119.2, 111.5. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>15</sub>H<sub>12</sub>NO: 222.0913; found: 222.0914.

**4-(p-Tolyl)-1H-indole-3-carbaldehyde (3ab).** Compound 3ab was synthesized by following general procedure D using 1H-indole-3-

carbaldehyde (1a, 58 mg, 0.4 mmol) and 1-iodo-4-methylbenzene (2b, 175 mg, 0.8 mmol) at 65 °C for 15 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): 3ab (76 mg, 81%, a brown solid, mp: 207–208 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.52 (bs, NH, 1H), 9.02 (s, CHO, 1H), 8.02 (d, *J* = 3.1 Hz, CH, 1H), 7.46–7.39 (m, CH, 3H), 7.33 (t, *J* = 7.3 Hz, CH, 1H), 7.29–7.25 (m, BB' part of AA'BB' system, CH, 2H), 7.18 (dd, *J* = 7.1, 0.7 Hz, CH, 1H), 2.42 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 187.3, 138.9, 137.5, 137.0, 135.7, 131.2, 129.3, 128.8, 124.3, 123.9, 123.4, 119.3, 111.1, 21.3. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>14</sub>NO: 236.1070; found: 236.1071.

**4-(m-Tolyl)-1H-indole-3-carbaldehyde (3ac).** Compound 3ac was synthesized by following general procedure D using 1H-indole-3-carbaldehyde (1a, 58 mg, 0.4 mmol) and 1-iodo-3-methylbenzene (2c, 103 μL, 0.8 mmol) at 65 °C for 20 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): 3ac (65 mg, 70%, a dark brown solid, mp: 87–88 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.78 (bs, NH, 1H), 9.51 (s, CHO, 1H), 8.01 (d, *J* = 3.0 Hz, CH, 1H), 7.42 (d, *J* = 8.0 Hz, CH, 1H), 7.38–7.28 (m, CH, 4H), 7.22 (d, *J* = 7.0 Hz, CH, 1H), 7.18 (d, *J* = 6.9 Hz, CH, 1H), 2.40 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 187.4, 138.3, 137.0, 135.7, 131.3, 129.7, 128.5, 128.4, 126.0, 123.8, 123.4, 119.2, 111.3, 21.5. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>14</sub>NO: 236.1070; found: 236.1070.

**4-(4-(tert-Butyl)phenyl)-1H-indole-3-carbaldehyde (3ad).** Compound 3ad was synthesized by following general procedure D using 1H-indole-3-carbaldehyde (1a, 58 mg, 0.4 mmol) and 1-(tert-butyl)-4-iodobenzene (2d, 141 μL, 0.8 mmol) at 100 °C for 3 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): 3ad (86 mg, 78%, a dark brown solid, mp: 153–154 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.79 (bs, NH, 1H), 9.56 (s, CHO, 1H), 8.00 (d, *J* = 3.0 Hz, CH, 1H), 7.52–7.44 (m, CH, 4H), 7.41 (d, *J* = 8.0 Hz, CH, 1H), 7.30 (t, *J* = 7.2 Hz, CH, 1H), 7.18 (d, *J* = 7.2 Hz, CH, 1H), 1.37 (s, CH<sub>3</sub>, 9H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 187.5, 150.7, 144.6, 138.8, 137.0, 135.6, 131.2, 128.6, 125.5, 124.3, 124.0, 123.4, 119.3, 117.7, 111.1, 34.6, 31.4. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>19</sub>H<sub>20</sub>NO: 278.1539; found: 278.1541.

**4-(4-Bromophenyl)-1H-indole-3-carbaldehyde (3ae).** Compound 3ae was synthesized by following general procedure D using 1H-indole-3-carbaldehyde (1a, 58 mg, 0.4 mmol) and 1-bromo-4-iodobenzene (2e, 226 mg, 0.8 mmol) at 100 °C for 6 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): 3ae (102 mg, 86%, a dark brown solid, mp: >300 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.55 (bs, NH, 1H), 9.04 (s, CHO, 1H), 8.03 (d, *J* = 2.7 Hz, CH, 1H), 7.64–7.56 (m, AA' part of AA'BB' system, CH, 2H), 7.51–7.44 (m, CH, 1H), 7.42–7.37 (m, BB' part of AA'BB' system, CH, 2H), 7.34 (t, *J* = 7.5 Hz, CH, 1H), 7.16 (d, *J* = 7.5 Hz, CH, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 186.2, 140.8, 137.0, 134.4, 131.7, 131.6, 130.7, 123.9, 123.7, 123.6, 121.9, 119.4, 111.5. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>15</sub>H<sub>11</sub>BrNO: 300.0019; found: 300.0018.

**4-(4-Methoxyphenyl)-1H-indole-3-carbaldehyde (3af).** Compound 3af was synthesized by following general procedure D using 1H-indole-3-carbaldehyde (1a, 58 mg, 0.4 mmol) and 1-iodo-4-methoxybenzene (2f, 187 mg, 0.8 mmol) at 100 °C for 3 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): 3af (88 mg, 79%, a dark brown solid, mp: 111–112 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.55 (bs, NH, 1H), 9.21 (s, CHO, 1H), 8.02 (d, *J* = 3.2 Hz, CH, 1H), 7.48–7.40 (m, CH, 3H), 7.32 (t, *J* = 7.3 Hz, CH, 1H), 7.19–7.14 (m, CH, 1H), 7.03–6.97 (m, BB' part of AA'BB' system, CH, 2H), 3.87 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 187.2, 159.2, 137.0, 135.4, 134.2, 131.2, 130.1, 124.4, 123.9, 123.4, 119.4, 114.0, 111.0, 55.3. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>14</sub>NO<sub>2</sub>: 252.1019; found: 252.1019.

**Methyl 4-(3-Formyl-1H-indol-4-yl)benzoate (3ag).** Compound 3ag was synthesized by following general procedure D using 1H-indole-3-carbaldehyde (1a, 58 mg, 0.4 mmol) and methyl 4-iodobenzoate (2g, 210 mg, 0.8 mmol) at 100 °C for 12 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): 3ag (97 mg, 87%, a brown solid, mp: 190–191 °C); <sup>1</sup>H

NMR (400 MHz, Acetone- $d_6$ ):  $\delta$  11.46 (bs, NH, 1H), 9.56 (s, CHO, 1H), 8.20 (d,  $J$  = 3.3 Hz, CH, 1H), 8.15–8.06 (m, AA' part of AA'BB' system, CH, 2H), 7.70–7.58 (m, CH, 3H), 7.36 (t,  $J$  = 7.4 Hz, CH, 1H), 7.17 (dd,  $J$  = 7.5, 0.8 Hz, CH, 1H), 3.92 (s,  $\text{CH}_3$ , 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz, Acetone- $d_6$ ):  $\delta$  184.6, 167.2, 148.2, 138.8, 135.2, 134.7, 130.2 (2C), 130.0, 124.2, 124.1, 124.0, 119.9, 113.1, 52.4. HRMS (ESI-TOF)  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{17}\text{H}_{14}\text{NO}_3$ : 280.0968; found: 280.0968.

**4-(4-Acetylphenyl)-1H-indole-3-carbaldehyde (3ah).** Compound **3ah** was synthesized by following general procedure D using 1H-indole-3-carbaldehyde (**1a**, 58 mg, 0.4 mmol) and 1-(4-iodophenyl)ethan-1-one (**2h**, 196 mg, 0.8 mmol) at 100 °C for 3 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): **3ah** (96 mg, 92%, a dirty white solid, mp: 220–221 °C);  $^1\text{H}$  NMR (400 MHz, Acetone- $d_6$ ):  $\delta$  11.45 (bs, NH, 1H), 9.56 (s, CHO, 1H), 8.19 (d,  $J$  = 3.2 Hz, CH, 1H), 8.14–8.03 (m, AA' part of AA'BB' system, CH, 2H), 7.70–7.60 (m, CH, 3H), 7.36 (t,  $J$  = 7.4 Hz, CH, 1H), 7.19–7.14 (m, CH, 1H), 2.65 (s,  $\text{CH}_3$ , 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz, Acetone- $d_6$ ):  $\delta$  197.7, 184.7, 148.2, 138.8, 137.1, 135.2, 134.5, 130.2, 129.1, 124.2, 124.1, 124.0, 119.8, 113.1, 26.8. HRMS (ESI-TOF)  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{17}\text{H}_{14}\text{NO}_2$ : 264.1019; found: 264.1024.

**4-(4-(Trifluoromethyl)phenyl)-1H-indole-3-carbaldehyde (3ai).** Compound **3ai** was synthesized by following general procedure D using 1H-indole-3-carbaldehyde (**1a**, 58 mg, 0.4 mmol) and 1-iodo-4-(trifluoromethyl)benzene (**2i**, 117  $\mu\text{L}$ , 0.8 mmol) at 100 °C for 15 h and purified by silica gel column chromatography (70:30 hexane/ethyl acetate): **3ai** (95 mg, 83%, a dark brown solid, mp: 190–191 °C);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  9.53 (bs, NH, 1H), 9.03 (s, CHO, 1H), 8.04 (d,  $J$  = 3.1 Hz, CH, 1H), 7.79–7.70 (m, AA' part of AA'BB' system, CH, 2H), 7.66–7.62 (m, BB' part of AA'BB' system, CH, 2H), 7.50 (d,  $J$  = 7.6 Hz, CH, 1H), 7.37 (t,  $J$  = 7.6 Hz, CH, 1H), 7.19 (d,  $J$  = 7.6 Hz, CH, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  185.9, 145.6, 137.1, 134.2, 132.4, 129.8 (q,  $J$  = 32.6 Hz), 129.4, 126.9 (q,  $J$  = 27.2 Hz), 125.4 (q,  $J$  = 3.4 Hz), 124.1, 123.6, 123.5, 119.2, 111.9. HRMS (ESI-TOF)  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{16}\text{H}_{11}\text{F}_3\text{NO}$ : 290.0787; found: 290.0792.

**4-(3-Nitrophenyl)-1H-indole-3-carbaldehyde (3al).** Compound **3al** was synthesized by following general procedure D using 1H-indole-3-carbaldehyde (**1a**, 58 mg, 0.4 mmol) and 1-iodo-3-nitrobenzene (**2l**, 199 mg, 0.8 mmol) at 120 °C for 16 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): **3al** (72 mg, 68%, a yellow solid, mp: 187–188 °C);  $^1\text{H}$  NMR (400 MHz, Acetone- $d_6$ ):  $\delta$  11.54 (bs, NH, 1H), 9.65 (s, CHO, 1H), 8.30 (d,  $J$  = 3.3 Hz, CH, 1H), 8.29–8.24 (m, CH, 2H), 7.94–7.81 (m, CH, 1H), 7.72 (t,  $J$  = 7.9 Hz, CH, 1H), 7.69–7.66 (m, CH, 1H), 7.40 (t,  $J$  = 7.4 Hz, CH, 1H), 7.27–7.21 (m, CH, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz, Acetone- $d_6$ ):  $\delta$  183.8, 148.7, 145.0, 139.3, 138.1, 136.2, 134.2, 129.8, 124.9, 124.8, 124.3, 123.2, 122.6, 120.1, 113.4. HRMS (ESI-TOF)  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{15}\text{H}_{11}\text{N}_2\text{O}_3$ : 267.0764; found: 267.0765.

**7-Fluoro-4-phenyl-1H-indole-3-carbaldehyde (3ba).** Compound **3ba** was synthesized by following general procedure D using 7-fluoro-1H-indole-3-carbaldehyde (**1b**, 65 mg, 0.4 mmol) and iodobenzene (**2a**, 90  $\mu\text{L}$ , 0.8 mmol) at 100 °C for 5 h and purified by silica gel column chromatography (70:30 hexane/ethyl acetate): **3ba** (85 mg, 90%, a purple solid, mp: 114–115 °C);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  9.48 (bs, NH, 1H), 9.21 (s, CHO, 1H), 8.03 (d,  $J$  = 3.0 Hz, C, 1H), 7.51–7.40 (m, CH, 5H), 7.12–7.02 (m, CH, 2H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  187.1, 149.2 (d,  $J$  = 246.3 Hz), 141.1, 132.2, 131.5 (d,  $J$  = 3.8 Hz), 129.1, 128.6, 127.8, 127.3 (d,  $J$  = 4.3 Hz), 125.2 (d,  $J$  = 13.8 Hz), 124.0 (d,  $J$  = 6.2 Hz), 119.7 (d,  $J$  = 1.1 Hz), 108.1 (d,  $J$  = 16.0 Hz). HRMS (ESI-TOF)  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{15}\text{H}_{11}\text{FNO}$ : 240.0819; found: 240.0820.

**7-Bromo-4-phenyl-1H-indole-3-carbaldehyde (3ca).** Compound **3ca** was synthesized by following general procedure D using 7-bromo-1H-indole-3-carbaldehyde (**1c**, 45 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc) $_2$  (5 mg, 20  $\mu\text{mol}$ , 10 mol %), and iodobenzene (**2a**, 90  $\mu\text{L}$ , 0.8 mmol) at 100 °C for 5 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): **3ca** (50 mg, 84%, a dark brown solid, mp: >300 °C);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$

9.46 (bs, NH, 1H), 9.15 (s, CHO, 1H), 8.07 (d,  $J$  = 2.9 Hz, CH, 1H), 7.58–7.37 (m, CH, 6H), 7.07 (d,  $J$  = 7.8 Hz, B part of AB system, CH, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  186.7, 140.8, 135.3, 135.2, 130.9, 128.9, 128.7, 128.0, 125.7, 125.2, 124.8, 120.5, 104.6. HRMS (ESI-TOF)  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{15}\text{H}_{11}\text{BrNO}$ : 300.0019; found: 300.0018.

**2-Methyl-4-phenyl-1H-indole-3-carbaldehyde (3da).** Compound **3da** was synthesized by following general procedure D using 2-methyl-1H-indole-3-carbaldehyde (**1d**, 58 mg, 0.4 mmol) and iodobenzene (**2a**, 90  $\mu\text{L}$ , 0.8 mmol) at 100 °C for 3.5 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): **3da** (77 mg, 82%, a pale-brown solid, mp: 190–191 °C);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  9.54 (bs, NH, 1H), 8.66 (s, CHO, 1H), 7.54–7.49 (m, CH, 2H), 7.48–7.34 (m, CH, 3H), 7.31–7.27 (m, CH, 2H), 7.13 (dd,  $J$  = 7.4, 0.7 Hz, CH, 1H), 2.78 (s,  $\text{CH}_3$ , 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  187.9, 145.4, 142.2, 135.1, 134.8, 128.9, 128.6, 127.6, 125.7, 123.8, 122.6, 114.4, 110.4, 29.7. HRMS (ESI-TOF)  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{16}\text{H}_{14}\text{NO}$ : 236.1070; found: 236.1070.

**1-Benzyl-4-phenyl-1H-indole-3-carbaldehyde (3ha).** Compound **3ha** was synthesized by following general procedure D using 1-benzyl-1H-indole-3-carbaldehyde (**1h**, 47 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc) $_2$  (5 mg, 20  $\mu\text{mol}$ , 10 mol %), and iodobenzene (**2a**, 45  $\mu\text{L}$ , 0.4 mmol) at 100 °C for 3 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): **3ha** (52 mg, 84%, a pale-brown solid, mp: 171–172 °C);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  9.45 (s, CHO, 1H), 7.96 (s, CH, 1H), 7.53 (d,  $J$  = 7.0 Hz, CH, 2H), 7.50–7.40 (m, CH, 3H), 7.39–7.28 (m, CH, 5H), 7.25–7.16 (m, CH, 3H), 5.38 (s,  $\text{CH}_2$ , 2H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  186.3, 141.9, 137.5, 136.0, 135.3, 134.2, 129.2, 129.1, 128.6, 128.4, 127.8, 127.4, 125.3, 123.9, 123.2, 118.4, 110.0, 51.2. HRMS (ESI-TOF)  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{22}\text{H}_{18}\text{NO}$ : 312.1383; found: 312.1382.

**1-Methyl-4-phenyl-1H-indole-3-carbaldehyde (3ia).** Compound **3ia** was synthesized by following general procedure D using 1-methyl-1H-indole-3-carbaldehyde (**1i**, 64 mg, 0.4 mmol) and iodobenzene (**2a**, 90  $\mu\text{L}$ , 0.8 mmol) at 65 °C for 3 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): **3ia** (76 mg, 81%, a yellow oil);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  9.35 (s, CHO, 1H), 7.82 (s, CH, 1H), 7.42 (d,  $J$  = 7.4 Hz, CH, 2H), 7.37 (t,  $J$  = 7.4 Hz, CH, 2H), 7.34–7.27 (m, CH, 3H), 7.11 (dd,  $J$  = 6.0, 2.0 Hz, CH, 1H), 3.80 (s,  $\text{CH}_3$ , 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  186.2, 141.9, 137.9, 135.8, 134.8, 129.0, 128.6, 127.7, 125.0, 123.8, 123.0, 118.0, 109.4, 33.9. HRMS (ESI-TOF)  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{16}\text{H}_{14}\text{NO}$ : 236.1070; found: 236.1070.

**1-(4-Phenyl-1H-indol-2-yl)ethan-1-one (5aa).** Compound **5aa** was synthesized by following general procedure D using 1-(1H-indol-3-yl)ethan-1-one (**4a**, 64 mg, 0.4 mmol) and iodobenzene (**2a**, 90  $\mu\text{L}$ , 0.8 mmol) at 120 °C for 7 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **5aa** (78 mg, 83%, a yellow solid, mp: 188–189 °C);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  9.14 (bs, NH, 1H), 7.69–7.65 (m, CH, 2H), 7.53 (t,  $J$  = 7.6 Hz, CH, 2H), 7.46–7.39 (m, CH, 3H), 7.34 (d,  $J$  = 1.8 Hz, CH, 1H), 7.22 (dd,  $J$  = 5.9, 2.2 Hz, CH, 1H), 2.58 (s,  $\text{CH}_3$ , 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  190.8, 140.4, 138.0, 137.0, 135.6, 128.8, 128.7, 127.5, 126.6, 126.2, 120.6, 111.4, 109.7, 25.9. HRMS (ESI-TOF)  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{16}\text{H}_{14}\text{NO}$ : 236.1070; found: 236.1070.

**1-(4-(p-Tolyl)-1H-indol-2-yl)ethan-1-one (5ab).** Compound **5ab** was synthesized by following general procedure D using 1-(1H-indol-3-yl)ethan-1-one (**4a**, 64 mg, 0.4 mmol) and 1-iodo-4-methylbenzene (**2b**, 175, 0.8 mmol) at 130 °C for 7 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **5ab** (81 mg, 81%, a dirty white solid, mp: 185–186 °C);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  9.46 (bs, NH, 1H), 7.62–7.56 (m, AA' part of AA'BB' system, CH, 2H), 7.45–7.39 (m, BB' part of AA'BB' system, 2H), 7.38–7.32 (m, CH, 3H), 7.20 (t,  $J$  = 4.1 Hz, CH, 1H), 2.59 (s,  $\text{CH}_3$ , 3H), 2.47 (s,  $\text{CH}_3$ , 3H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  190.6, 138.0, 137.5, 137.3, 137.0, 135.5, 129.5, 128.6, 126.7, 126.2, 120.5, 111.1, 109.7, 25.9, 21.3. HRMS (ESI-TOF)  $m/z$ :  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{17}\text{H}_{16}\text{NO}$ : 250.1226; found: 250.1226.



**1-(4-(*m*-Tolyl)-1*H*-indol-2-yl)ethan-1-one (5ac).** Compound **5ac** was synthesized by following general procedure D using 1-(1*H*-indol-3-yl)ethan-1-one (**4a**, 64 mg, 0.4 mmol) and 1-iodo-3-methylbenzene (**2c**, 103  $\mu$ L, 0.8 mmol) at 120 °C for 7 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **5ac** (74 mg, 74%, a pale-yellow solid, mp: 110–111 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.59 (bs, NH, 1H), 7.51–7.47 (m, CH, 2H), 7.45–7.38 (m, CH, 3H), 7.35 (d, *J* = 1.9 Hz, CH, 1H), 7.25 (d, *J* = 7.4 Hz, CH, 1H), 7.20 (dd, *J* = 6.3, 1.9 Hz, CH, 1H), 2.60 (s, CH<sub>3</sub>, 3H), 2.47 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  190.7, 140.4, 138.4, 138.0, 137.2, 135.6, 129.5, 128.6, 128.3, 126.6, 126.2, 125.9, 120.6, 111.3, 109.7, 26.0, 21.6. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>16</sub>NO: 250.1226; found: 250.1226.

**1-(4-(4-(*tert*-Butyl)phenyl)-1*H*-indol-2-yl)ethan-1-one (5ad).** Compound **5ad** was synthesized by following general procedure D using 1-(1*H*-indol-3-yl)ethan-1-one (**4a**, 64 mg, 0.4 mmol) and 1-(*tert*-butyl)-4-iodobenzene (**2d**, 142  $\mu$ L, 0.8 mmol) at 120 °C for 8 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **5ad** (89 mg, 76%, a brown solid, mp: 209–210 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.38 (bs, NH, 1H), 7.67–7.60 (m, AA' part of AA'BB' system, CH, 2H), 7.59–7.54 (m, BB' part of AA'BB' system, 2H), 7.45–7.39 (m, CH, 3H), 7.22 (dd, *J* = 4.6, 3.5 Hz, CH, 1H), 2.60 (s, CH<sub>3</sub>, 3H), 1.43 (s, CH<sub>3</sub>, 9H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  190.6, 150.5, 137.9, 137.4, 136.9, 135.5, 128.4, 126.7, 126.2, 125.7, 120.5, 111.0, 109.8, 34.7, 31.4, 25.9. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>20</sub>H<sub>22</sub>NO: 292.1696; found: 292.1696.

**1-(4-(4-Bromophenyl)-1*H*-indol-2-yl)ethan-1-one (5ae).** Compound **5ae** was synthesized by following general procedure D using 1-(1*H*-indol-3-yl)ethan-1-one (**4a**, 64 mg, 0.4 mmol) and 1-bromo-4-iodobenzene (**2e**, 227 mg, 0.8 mmol) at 100 °C for 13 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **5ae** (101 mg, 80%, a white solid, mp: 198–199 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.14 (bs, NH, 1H), 7.68–7.61 (m, AA' part of AA'BB' system, CH, 2H), 7.56–7.51 (m, BB' part of AA'BB' system, CH, 2H), 7.45–7.40 (m, CH, 2H), 7.27 (d, *J* = 1.9 Hz, CH, 1H), 7.18 (dd, *J* = 4.7, 3.4 Hz, CH, 1H), 2.58 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  190.5, 139.3, 137.8, 135.8, 135.7, 131.9, 130.3, 126.6, 125.9, 121.7, 120.6, 111.7, 109.0, 25.9. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>13</sub>BrNO: 314.0175; found: 314.0174.

**1-(4-(4-Methoxyphenyl)-1*H*-indol-2-yl)ethan-1-one (5af).** Compound **5af** was synthesized by following general procedure D using 1-(1*H*-indol-3-yl)ethan-1-one (**4a**, 64 mg, 0.4 mmol) and 1-iodo-4-methoxybenzene (**2f**, 188 mg, 0.8 mmol) at 120 °C for 5 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **5af** (74 mg, 70%, a pale-yellow solid, mp: 196–197 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.34 (bs, NH, 1H), 7.66–7.56 (m, AA' part of AA'BB' system, CH, 2H), 7.46–7.37 (m, CH, 2H), 7.34 (d, *J* = 1.5 Hz, CH, 1H), 7.17 (dd, *J* = 5.8, 2.2 Hz, CH, 1H), 7.14–7.03 (m, CH, BB' part of AA'BB' system, 2H), 3.90 (s, CH<sub>3</sub>, 3H), 2.59 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  190.6, 159.2, 138.0, 136.7, 135.5, 132.9, 129.8, 126.7, 126.2, 120.3, 114.2, 110.8, 109.7, 55.4, 25.9. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>16</sub>NO<sub>2</sub>: 266.1176; found: 266.1175.

**Methyl 4-(2-Acetyl-1*H*-indol-4-yl)benzoate (5ag).** Compound **5ag** was synthesized by following general procedure D using 1-(1*H*-indol-3-yl)ethan-1-one (**4a**, 64 mg, 0.4 mmol) and methyl 4-iodobenzoate (**2g**, 210 mg, 0.8 mmol) at 110 °C for 15 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **5ag** (82 mg, 70%, a white solid, mp: 190–191 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.53 (bs, NH, 1H), 8.42–7.96 (m, AA' part of AA'BB' system, CH, 2H), 7.95–7.67 (m, BB' part of AA'BB' system, CH, 2H), 7.52–7.40 (m, CH, 2H), 7.32 (s, CH, 1H), 7.24 (d, *J* = 6.8 Hz, CH, 1H), 3.98 (s, CH<sub>3</sub>, 3H), 2.60 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  190.6, 167.0, 145.0, 137.9, 135.8, 135.8, 130.0, 129.1, 128.7, 126.6, 126.0, 120.9, 112.1, 109.0, 52.2, 25.9. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>18</sub>H<sub>16</sub>NO<sub>3</sub>: 294.1125; found: 294.1124.

**1-(4-(2-Acetyl-1*H*-indol-4-yl)phenyl)ethan-1-one (5ah).** Compound **5ah** was synthesized by following general procedure D using

1-(1*H*-indol-3-yl)ethan-1-one (**4a**, 64 mg, 0.4 mmol) and 1-(4-iodophenyl)ethan-1-one (**2h**, 198 mg, 0.8 mmol) at 120 °C for 7 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **5ah** (59 mg, 53%, a white solid, mp: 223–224 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.53 (bs, NH, 1H), 8.42–7.96 (m, AA' part of AA'BB' system, CH, 2H), 7.95–7.67 (m, BB' part of AA'BB' system, CH, 2H), 7.52–7.40 (m, CH, 2H), 7.32 (s, CH, 1H), 7.24 (d, *J* = 6.8 Hz, CH, 1H), 3.98 (s, CH<sub>3</sub>, 3H), 2.60 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  190.6, 167.0, 145.0, 137.9, 135.8, 135.8, 130.0, 129.1, 128.7, 126.6, 126.0, 120.9, 112.1, 109.0, 52.2, 25.9. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>18</sub>H<sub>16</sub>NO<sub>2</sub>: 278.1176; found: 278.1182.

**1-(4-(4-(Trifluoromethyl)phenyl)-1*H*-indol-2-yl)ethan-1-one (5ai).** Compound **5ai** was synthesized by following general procedure D using 1-(1*H*-indol-3-yl)ethan-1-one (**4a**, 64 mg, 0.4 mmol) and 1-iodo-4-(trifluoromethyl)benzene (**2i**, 118  $\mu$ L, 0.8 mmol) at 110 °C for 20 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **5ai** (83 mg, 68%, a white solid, mp: 216–217 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.56 (bs, NH, 1H), 7.82–7.75 (m, CH, 4H), 7.50 (d, *J* = 7.5 Hz, CH, 1H), 7.45 (t, *J* = 7.5 Hz, CH, 1H), 7.30 (d, *J* = 1.1 Hz, CH, 1H), 7.23 (d, *J* = 7.5 Hz, CH, 1H), 2.61 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  190.6, 144.0, 137.9, 135.9, 135.4, 129.5 (q, *J* = 32.4 Hz), 129.0, 126.6, 126.0, 125.7 (q, *J* = 3.8 Hz), 121.6 (q, *J* = 272.4 Hz), 120.9, 112.2, 108.8, 25.9. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>13</sub>F<sub>3</sub>NO: 304.0944; found: 304.0944.

**1-(4-(3,4-Dimethylphenyl)-1*H*-indol-2-yl)ethan-1-one (5ak).** Compound **5aa** was synthesized by following general procedure D using 1-(1*H*-indol-3-yl)ethan-1-one (**4a**, 64 mg, 0.4 mmol) and 4-iodo-1,2-dimethylbenzene (**2k**, 114  $\mu$ L, 0.8 mmol) at 120 °C for 6 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **5ak** (77 mg, 73%, a brown solid, mp: 180–181 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.41 (bs, NH, 1H), 7.47–7.40 (m, CH, 4H), 7.37 (d, *J* = 2.0 Hz, CH, 1H), 7.31 (d, *J* = 7.6 Hz, CH, 1H), 7.23–7.18 (m, CH, 1H), 2.60 (s, CH<sub>3</sub>, 3H), 2.40 (s, CH<sub>3</sub>, 3H), 2.38 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  190.6, 138.0, 137.9, 137.2, 137.0, 136.0, 135.5, 129.98, 129.96, 126.6, 126.2, 126.2, 120.4, 110.9, 109.7, 25.9, 20.0, 19.6. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>18</sub>H<sub>18</sub>NO: 264.1383; found: 264.1382.

**1-(4-(3-Nitrophenyl)-1*H*-indol-2-yl)ethan-1-one (5al).** Compound **5al** was synthesized by following general procedure D using 1-(1*H*-indol-3-yl)ethan-1-one (**4a**, 64 mg, 0.4 mmol) and 1-iodo-3-nitrobenzene (**2l**, 200 mg, 0.8 mmol) at 100 °C for 15 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **5al** (59 mg, 53%, a yellow solid, mp: 210–211 °C); <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  12.02 (bs, NH, 1H), 8.42 (s, CH, 1H), 8.28 (d, *J* = 7.8 Hz, CH, 1H), 8.16 (d, *J* = 7.8 Hz, CH, 1H), 7.83 (t, *J* = 7.8 Hz, CH, 1H), 7.54 (d, *J* = 7.7 Hz, CH, 1H), 7.48 (s, CH, 1H), 7.42 (t, *J* = 7.7 Hz, CH, 1H), 7.28 (d, *J* = 7.7 Hz, CH, 1H), 2.56 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  190.3, 148.3, 141.4, 138.2, 136.5, 134.8, 133.1, 130.4, 125.7, 124.8, 122.7, 122.1, 120.3, 113.1, 107.7, 26.2. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>13</sub>N<sub>2</sub>O<sub>3</sub>: 281.0921; found: 281.0920.

**1-(7-Fluoro-4-phenyl-1*H*-indol-2-yl)ethan-1-one (5ba).** Compound **5ba** was synthesized by following general procedure D using 1-(7-fluoro-1*H*-indol-3-yl)ethan-1-one (**4b**, 71 mg, 0.4 mmol) and iodobenzene (**2a**, 90  $\mu$ L, 0.8 mmol) at 120 °C for 18 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **5ba** (74 mg, 73%, an orange solid, mp: 164–165 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.31 (bs, NH, 1H), 7.62 (d, *J* = 7.2 Hz, CH, 2H), 7.52 (t, *J* = 7.6 Hz, CH, 2H), 7.46–7.40 (m, CH, 1H), 7.34–7.31 (m, CH, 1H), 7.14–7.10 (m, CH, 2H), 2.59 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  190.2, 149.2 (d, *J* = 247.0 Hz), 139.7, 136.2, 133.0 (d, *J* = 4.0 Hz), 129.2 (d, *J* = 5.2 Hz), 128.8, 128.7, 127.5, 126.4 (d, *J* = 14.8 Hz), 120.6 (d, *J* = 5.6 Hz), 110.8 (d, *J* = 16.0 Hz), 109.7, 26.0. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>13</sub>FNO: 254.0976; found: 254.0976.

**1-(7-Bromo-4-phenyl-1*H*-indol-2-yl)ethan-1-one (5ca).** Compound **5ca** was synthesized by following general procedure D using 1-(7-bromo-1*H*-indol-3-yl)ethan-1-one (**4c**, 71 mg, 0.3 mmol),

AgOAc (100 mg, 0.6 mmol), Pd(OAc)<sub>2</sub> (7 mg, 30 μmol, 10 mol %), and iodobenzene (**2a**, 60 μL, 0.6 mmol) at 120 °C for 24 h and purified by silica gel column chromatography (85:15 hexane/ethyl acetate): **5ca** (69 mg, 74%, a pale-brown solid, mp: 125–126 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.21 (bs, NH, 1H), 7.64 (d, *J* = 8.0 Hz, CH, 2H), 7.59–7.50 (m, CH, 3H), 7.48–7.42 (m, CH, 1H), 7.38 (d, *J* = 1.7 Hz, CH, 1H), 7.14–7.08 (m, CH, 1H), 2.59 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 190.2, 139.5, 136.4, 136.4, 135.9, 128.9, 128.7, 128.7, 127.8, 127.0, 121.7, 110.1, 104.5, 25.9. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>13</sub>BrNO: 314.0175; found: 314.0175.

**1-(7-Bromo-4-phenyl-1H-indol-3-yl)ethan-1-one (6ca)**. Compound **6ca** was synthesized by following general procedure D using 1-(7-bromo-1H-indol-3-yl)ethan-1-one (**4c**, 71 mg, 0.3 mmol), AgOAc (100 mg, 0.6 mmol), Pd(OAc)<sub>2</sub> (7 mg, 30 μmol, 10 mol %), and iodobenzene (**2a**, 60 μL, 0.6 mmol) at 120 °C for 20 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **6ca** (71 mg, 76%, a brown solid, mp: 162–163 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.01 (bs, NH, 1H), 7.79 (d, *J* = 2.9 Hz, CH, 1H), 7.49 (d, *J* = 7.9 Hz, A part of AB system, CH, 1H), 7.45–7.33 (m, CH, 5H), 7.09 (d, *J* = 7.9 Hz, B part of AB system, CH, 1H), 2.07 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 194.2, 142.0, 136.0, 135.8, 131.2, 128.6, 128.1, 127.2, 125.9, 125.1, 123.6, 121.9, 104.1, 29.1. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>16</sub>H<sub>13</sub>BrNO: 314.0175; found: 314.0175.

**2-Methyl-4-phenyl-1H-indole (7da)**. Compound **7da** was synthesized by following general procedure D using 1-(2-methyl-1H-indol-3-yl)ethan-1-one (**4d**, 69 mg, 0.4 mmol) and iodobenzene (**2a**, 90 μL, 0.8 mmol) at 120 °C for 13 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **7da** (66 mg, 80%, a brown solid, mp: 96–97 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.78 (bs, NH, 1H), 7.64–7.57 (m, CH, 2H), 7.38 (t, *J* = 7.6 Hz, CH, 2H), 7.30–7.23 (m, CH, 1H), 7.18–7.03 (m, CH, 3H), 6.32 (s, CH, 1H), 2.32 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 141.5, 136.5, 135.5, 133.4, 128.8, 128.4, 127.3, 126.8, 121.3, 119.6, 109.5, 100.0, 13.8. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>15</sub>H<sub>14</sub>N: 208.1121; found: 208.1120.

**1-(1-Benzyl-4-phenyl-1H-indol-3-yl)ethan-1-one (9aa)**. Compound **9aa** was synthesized by following general procedure D using 1-(1-benzyl-1H-indol-3-yl)ethan-1-one (**8a**, 50 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and iodobenzene (**2a**, 45 μL, 0.4 mmol) at 100 °C for 5 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9aa** (58 mg, 89%, a white solid, mp: 132–133 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.71 (s, CH, 1H), 7.47–7.42 (m, CH, 4H), 7.42–7.29 (m, CH, 6H), 7.24–7.18 (m, CH, 3H), 5.37 (s, CH<sub>2</sub>, 2H), 2.00 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 194.0, 143.0, 138.0, 136.8, 135.8, 134.6, 129.1, 128.7, 128.2, 128.0, 127.1, 127.0, 124.1, 123.6, 123.3, 119.9, 109.3, 50.8, 29.1. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>23</sub>H<sub>20</sub>NO: 326.1539; found: 326.1539.

**1-(1-Benzyl-4-phenyl-1H-indol-3-yl)ethan-1-one (9ab)**. Compound **9ab** was synthesized by following general procedure D using 1-(1-benzyl-1H-indol-3-yl)ethan-1-one (**8a**, 50 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and 1-iodo-4-methylbenzene (**2b**, 87 mg, 0.4 mmol) at 100 °C for 5.5 h and purified by silica gel column chromatography (85:15 hexane/ethyl acetate): **9ab** (59 mg, 87%, a white solid, mp: 125–126 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.68 (s, CH, 1H), 7.36–7.31 (m, CH, 5H), 7.30–7.27 (m, CH, 2H), 7.24–7.16 (m, CH, 5H), 5.34 (s, CH<sub>2</sub>, 2H), 2.40 (s, CH<sub>3</sub>, 3H), 1.98 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 194.3, 140.1, 138.0, 136.8, 136.6, 135.8, 134.5, 129.1, 128.8, 128.6, 128.2, 127.1, 124.1, 123.6, 123.3, 120.0, 109.1, 50.8, 29.3, 21.3. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>24</sub>H<sub>22</sub>NO: 340.1696; found: 340.1695.

**1-(1-Benzyl-4-(*m*-tolyl)-1H-indol-3-yl)ethan-1-one (9ac)**. Compound **9ac** was synthesized by following general procedure D using 1-(1-benzyl-1H-indol-3-yl)ethan-1-one (**8a**, 50 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and 1-iodo-3-methylbenzene (**2c**, 51 μL, 0.4 mmol) at 100 °C for 5.5

h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9ac** (57 mg, 85%, a pale-brown solid, mp: 115–116 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.60 (s, 1H), 7.29–7.19 (m, 7H), 7.17–7.08 (m, 5H), 5.27 (s, 2H), 2.31 (s, 3H), 1.88 (s, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 194.4, 142.8, 137.9, 137.7, 136.8, 135.8, 134.3, 129.3, 129.0, 128.2, 128.0, 127.8, 127.1, 125.8, 123.9, 123.6, 123.3, 120.0, 109.2, 50.8, 29.2, 21.5. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>24</sub>H<sub>22</sub>NO: 340.1696; found: 340.1696.

**1-(1-Benzyl-4-(4-(*tert*-butyl)phenyl)-1H-indol-3-yl)ethan-1-one (9ad)**. Compound **9ad** was synthesized by following general procedure D using 1-(1-benzyl-1H-indol-3-yl)ethan-1-one (**8a**, 50 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and 1-(*tert*-butyl)-4-iodobenzene (**2d**, 71 μL, 0.4 mmol) at 100 °C for 5.5 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9ad** (68 mg, 90%, a white solid, mp: 166–167 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.66 (s, CH, 1H), 7.47–7.40 (m, CH, 4H), 7.39–7.29 (m, CH, 5H), 7.24–7.19 (m, CH, 3H), 5.35 (s, CH<sub>2</sub>, 2H), 1.79 (s, CH<sub>3</sub>, 3H), 1.38 (s, CH<sub>3</sub>, 9H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 195.9, 150.2, 139.9, 137.9, 136.4, 135.8, 133.8, 129.0, 128.4, 128.2, 127.2, 125.3, 123.8, 123.7, 123.2, 120.5, 109.2, 50.8, 34.6, 31.4, 29.6. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>27</sub>H<sub>28</sub>NO: 382.2165; found: 382.2165.

**1-(1-Benzyl-4-(4-bromophenyl)-1H-indol-3-yl)ethan-1-one (9ae)**. Compound **9ae** was synthesized by following general procedure D using 1-(1-benzyl-1H-indol-3-yl)ethan-1-one (**8a**, 50 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and 1-bromo-4-iodobenzene (**2e**, 113 mg, 0.4 mmol) at 100 °C for 6 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9ae** (64 mg, 79%, a pale-brown solid, mp: 145–146 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.75 (s, CH, 1H), 7.55–7.50 (m, AA' part of AA'BB' system, CH, 2H), 7.39–7.24 (m, CH, 7H), 7.20–7.13 (m, CH, 3H), 5.39 (s, CH<sub>2</sub>, 2H), 2.17 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 192.7, 142.0, 138.1, 135.7, 135.7, 135.5, 130.9, 130.3, 129.1, 128.3, 127.0, 124.5, 123.5, 123.2, 120.9, 119.2, 109.6, 50.8, 28.8. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>23</sub>H<sub>19</sub>BrNO: 404.0645; found: 404.0644.

**1-(1-Benzyl-4-(4-methoxyphenyl)-1H-indol-3-yl)ethan-1-one (9af)**. Compound **9af** was synthesized by following general procedure D using 1-(1-benzyl-1H-indol-3-yl)ethan-1-one (**8a**, 50 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and 1-iodo-4-methoxybenzene (**2f**, 113 mg, 0.4 mmol) at 100 °C for 5.5 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9af** (57 mg, 80%, a brown solid, mp: 132–133 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.70 (s, CH, 1H), 7.40–7.37 (m, AA' part of AA'BB' system, CH, 2H), 7.36–7.28 (m, CH, 5H), 7.22–7.16 (m, CH, 3H), 7.05–6.87 (m, BB' part of AA'BB' system, CH, 2H), 5.36 (s, CH<sub>2</sub>, 2H), 3.86 (s, CH<sub>3</sub>, 3H), 2.01 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 194.6, 158.8, 138.0, 136.4, 135.8, 135.5, 134.4, 129.7, 129.1, 128.2, 127.1, 124.0, 123.7, 123.3, 120.1, 113.6, 109.0, 55.2, 50.8, 29.4. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>24</sub>H<sub>22</sub>NO<sub>2</sub>: 356.1645; found: 356.1645.

**Methyl 4-(3-Acetyl-1-benzyl-1H-indol-4-yl)benzoate (9ag)**. Compound **9ag** was synthesized by following general procedure D using 1-(1-benzyl-1H-indol-3-yl)ethan-1-one (**8a**, 50 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and methyl 4-iodobenzoate (**2g**, 105 mg, 0.4 mmol) at 110 °C for 10 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9ag** (62 mg, 81%, a brown solid, mp: 165–166 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.18–8.04 (m, AA' part of AA'BB' system, CH, 2H), 7.78 (s, CH, 1H), 7.55–7.41 (m, BB' part of AA'BB' system, CH, 2H), 7.42–7.27 (m, CH, 5H), 7.25–7.15 (m, CH, 3H), 5.39 (s, CH<sub>2</sub>, 2H), 3.93 (s, CH<sub>3</sub>, 3H), 2.18 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 192.4, 167.2, 147.8, 138.1, 135.9, 135.7, 135.5, 129.1, 129.1, 128.7, 128.4, 128.3, 127.0, 124.5, 123.4, 123.2, 119.2, 109.9, 52.0, 50.9, 28.6. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>25</sub>H<sub>22</sub>NO<sub>3</sub>: 384.1594; found: 384.1595.

**1-(4-(3-Acetyl-1-benzyl-1H-indol-4-yl)phenyl)ethan-1-one (9ah)**. Compound **9ah** was synthesized by following general procedure D



using 1-(1-benzyl-1*H*-indol-3-yl)ethan-1-one (**8a**, 50 mg, 0.2 mmol), AgOAc (98 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and 1-(4-iodophenyl)ethan-1-one (**2h**, 87 mg, 0.4 mmol) at 120 °C for 6 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9ah** (58 mg, 80%, a brown solid, mp: 92–93 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.05–7.97 (m, AA' part of AA'BB' system, CH, 2H), 7.85 (s, CH, 1H), 7.53–7.41 (m, BB' part of AA'BB' system, CH, 2H), 7.41–7.30 (m, CH, 5H), 7.24–7.14 (m, CH, 3H), 5.41 (s, CH<sub>2</sub>, 2H), 2.65 (s, CH<sub>3</sub>, 3H), 2.25 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 198.1, 192.5, 148.1, 138.2, 136.0, 135.8, 135.6, 135.3, 129.1(2C), 128.8, 128.3, 127.9, 127.0, 124.7, 123.5, 123.1, 118.9, 110.1, 50.9, 26.6. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>25</sub>H<sub>22</sub>NO<sub>2</sub>: 368.1645; found: 368.1666.

**1-(1-Benzyl-4-(4-(trifluoromethyl)phenyl)-1*H*-indol-3-yl)ethan-1-one (9ai).** Compound **9ai** was synthesized by following general procedure D using 1-(1-benzyl-1*H*-indol-3-yl)ethan-1-one (**8a**, 50 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and 1-iodo-4-(trifluoromethyl)benzene (**2i**, 59 μL, 0.4 mmol) at 100 °C for 5.5 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9ai** (60 mg, 77%, a white solid, mp: 169–170 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.80 (s, CH, 1H), 7.70–7.63 (m, AA' part of AA'BB' system, CH, 2H), 7.55–7.46 (m, BB' part of AA'BB' system, CH, 2H), 7.42–7.30 (m, CH, 5H), 7.25–7.15 (m, CH, 3H), 5.41 (s, CH<sub>2</sub>, 2H), 2.21 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 192.1, 146.7, 138.1, 135.7(2C), 135.6, 129.1, 128.9, 128.7 (q, *J* = 32.4 Hz), 128.3, 127.0, 124.7, 124.6 (q, *J* = 3.4 Hz), 124.3 (q, *J* = 272.4 Hz), 123.5, 123.2, 119.0, 110.0, 50.9, 28.5. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>24</sub>H<sub>19</sub>F<sub>3</sub>NO: 394.1413; found: 394.1413.

**1-(1-Benzyl-4-(3,4-dimethylphenyl)-1*H*-indol-3-yl)ethan-1-one (9ak).** Compound **9ak** was synthesized by following general procedure D using 1-(1-benzyl-1*H*-indol-3-yl)ethan-1-one (**8a**, 50 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and 4-iodo-1,2-dimethylbenzene (**2k**, 57 μL, 0.4 mmol) at 100 °C for 4 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9ak** (49 mg, 70%, a yellow gum); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.65 (s, CH, 1H), 7.34–7.27 (m, CH, 3H), 7.26–7.24 (m, CH, 1H), 7.23–7.19 (m, CH, 2H), 7.18–7.11 (m, CH, 5H), 5.33 (s, CH<sub>2</sub>, 2H), 2.28 (s, CH<sub>3</sub>, 3H), 2.26 (s, CH<sub>3</sub>, 3H), 1.92 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 194.7, 140.4, 137.9, 136.7, 136.3, 135.9, 135.3, 134.2, 129.8, 129.5, 129.1, 128.2, 127.1, 126.1, 123.9, 123.6, 123.3, 120.1, 109.0, 50.8, 29.4, 19.9, 19.7. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>25</sub>H<sub>24</sub>NO: 354.1852; found: 354.1852.

**1-(1-Benzyl-4-(3-nitrophenyl)-1*H*-indol-3-yl)ethan-1-one (9al).** Compound **9al** was synthesized by following general procedure D using 1-(1-benzyl-1*H*-indol-3-yl)ethan-1-one (**8a**, 50 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and 1-iodo-3-nitrobenzene (**2l**, 100 mg, 0.4 mmol) at 100 °C for 48 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate): **9al** (37 mg, 51%, a yellow solid, mp: 135–136 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.25–8.19 (m, CH, 2H), 7.85 (s, CH, 1H), 7.73–7.70 (m, CH, 1H), 7.56 (t, *J* = 7.9 Hz, CH, 1H), 7.41–7.33 (m, CH, 5H), 7.24–7.18 (m, CH, 3H), 5.44 (s, CH<sub>2</sub>, 2H), 2.34 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 191.5, 147.5, 144.7, 138.3, 136.6, 135.6, 134.8, 134.7, 129.3, 128.5, 128.3, 127.0, 125.0, 124.0, 123.8, 123.3, 121.7, 118.6, 110.4, 51.1, 28.3. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>23</sub>H<sub>19</sub>N<sub>2</sub>O<sub>3</sub>: 371.1390; found: 371.1389.

**1-(1-Benzyl-7-fluoro-4-phenyl-1*H*-indol-3-yl)ethan-1-one (9ba).** Compound **9ba** was synthesized by following general procedure D using 1-(1-benzyl-7-fluoro-1*H*-indol-3-yl)ethan-1-one (**8b**, 54 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and iodobenzene (**2a**, 45 μL, 0.4 mmol) at 90 °C for 6 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9ba** (57 mg, 83%, a dirty white solid, mp: 112–113 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.64 (s, CH, 1H), 7.44–7.31 (m, CH, 8H), 7.22 (d, *J* = 6.6 Hz, CH, 2H), 7.10–6.98 (m, CH, 2H), 5.54 (s, CH<sub>2</sub>, 2H), 1.97 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 194.1, 149.5 (d, *J* = 245.2 Hz), 142.2, 136.6, 135.5, 132.6 (d, *J* = 3.8

Hz), 129.0, 128.6, 128.20, 128.17, 127.1, 127.0, 126.9 (d, *J* = 5.0 Hz), 125.5 (d, *J* = 9.9 Hz), 124.2 (d, *J* = 6.7 Hz), 120.6, 109.2 (d, *J* = 18.1 Hz), 53.0 (d, *J* = 6.9 Hz), 29.2. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>23</sub>H<sub>19</sub>FNO: 344.1445; found: 344.1447.

**1-(1-Benzyl-7-bromo-4-phenyl-1*H*-indol-3-yl)ethan-1-one (9ca).** Compound **9ca** was synthesized by following general procedure D using 1-(1-benzyl-7-bromo-1*H*-indol-3-yl)ethan-1-one (**8c**, 66 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and iodobenzene (**2a**, 45 μL, 0.4 mmol) at 75 °C for 5 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9ca** (63 mg, 78%, a pale-brown solid, mp: 130–131 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.62 (s, CH, 1H), 7.50 (d, *J* = 7.9 Hz, A part of AB system, CH, 1H), 7.45–7.28 (m, CH, 8H), 7.08 (d, *J* = 6.9 Hz, CH, 2H), 7.04 (d, *J* = 7.9 Hz, B part of AB system, CH, 1H), 5.90 (s, CH<sub>2</sub>, 2H), 1.95 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 194.2, 142.0, 137.5, 136.9, 136.1, 134.1, 128.9 (2C), 128.4, 128.3, 127.9, 127.3, 126.5, 126.4, 124.9, 120.1, 103.2, 52.3, 29.3. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>23</sub>H<sub>19</sub>BrNO: 404.0645; found: 404.0644.

**1-(1-Methyl-4-phenyl-1*H*-indol-3-yl)ethan-1-one (9ea).** Compound **9ea** was synthesized by following general procedure D using 1-(1-methyl-1*H*-indol-3-yl)ethan-1-one (**8e**, 70 mg, 0.4 mmol) and iodobenzene (**2a**, 90 μL, 0.8 mmol) at 100 °C for 5 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9ea** (79 mg, 80%, a yellow gum); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.67 (s, CH, 1H), 7.45–7.32 (m, CH, 7H), 7.23 (d, *J* = 6.9 Hz, CH, 1H), 3.86 (s, CH<sub>3</sub>, 3H), 2.02 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 193.6, 143.1, 138.4, 136.7, 135.6, 128.7, 128.0, 126.9, 124.1, 123.3, 123.2, 119.2, 108.7, 33.6, 29.0. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>16</sub>NO: 250.1226; found: 250.1226.

**Methyl 4-(3-Acetyl-1-methyl-1*H*-indol-4-yl)benzoate (9eg).** Compound **9eg** was synthesized by following general procedure D using 1-(1-methyl-1*H*-indol-3-yl)ethan-1-one (**8e**, 70 mg, 0.2 mmol) and methyl 4-iodobenzoate (**2g**, 210 mg, 0.8 mmol) at 100 °C for 8 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate): **9eg** (100 mg, 82%, off-white solid, mp: 149–150 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.13–8.03 (m, AA' part of AA'BB' system, CH, 2H), 7.73 (s, CH, 1H), 7.49–7.42 (m, BB' part of AA'BB' system, CH, 2H), 7.40–7.32 (m, CH, 2H), 7.25–7.19 (m, CH, 1H), 3.92 (s, CH<sub>3</sub>, 3H), 3.89 (s, CH<sub>3</sub>, 3H), 2.18 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 192.0, 167.2, 147.9, 138.5, 136.3, 135.8, 129.1, 128.7, 128.3, 124.4, 123.3, 122.9, 118.5, 109.3, 52.0, 33.7, 28.5. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>19</sub>H<sub>18</sub>NO<sub>3</sub>: 308.1281; found: 308.1281.

**2-Phenyl-1*H*-indole (11a).**<sup>27</sup> Compound **11a** was synthesized by following general procedure D using 1-*H*-indole-3-carboxylic acid (**10a**, 65 mg, 0.4 mmol) and iodobenzene (**2a**, 90 μL, 0.8 mmol) at 120 °C for 3 h and purified by silica gel column chromatography (95:5 hexane/ethyl acetate): **11a** (66 mg, 85%, off-white solid, mp: 188–189 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.34 (bs, NH, 1H), 7.67 (t, *J* = 6.6 Hz, CH, 3H), 7.49–7.39 (m, CH, 3H), 7.34 (t, *J* = 7.3 Hz, CH, 1H), 7.22 (t, *J* = 7.4 Hz, CH, 1H), 7.15 (t, *J* = 7.5 Hz, CH, 1H), 6.85 (d, *J* = 1.3 Hz, CH, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 137.9, 136.8, 132.4, 129.3, 129.0, 127.7, 125.2, 122.4, 120.7, 120.3, 110.9, 100.0.

**2-(*p*-Tolyl)-1*H*-indole (11b).**<sup>27</sup> Compound **11b** was synthesized by following general procedure D using 1-*H*-indole-3-carboxylic acid (**10a**, 65 mg, 0.4 mmol) and 1-iodo-4-methylbenzene (**2b**, 175 mg, 0.8 mmol) at 120 °C for 3 h and purified by silica gel column chromatography (95:5 hexane/ethyl acetate): **11b** (62 mg, 75%, a white solid, mp: 214–215 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.30 (bs, NH, 1H), 7.63 (d, *J* = 7.7 Hz, CH, 1H), 7.60–7.54 (m, CH, 2H), 7.39 (d, *J* = 8.0 Hz, CH, 1H), 7.28–7.24 (m, CH, 2H), 7.22–7.17 (m, CH, 1H), 7.15–7.10 (m, CH, 1H), 6.80 (d, *J* = 1.5 Hz, CH, 1H), 2.40 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 138.1, 137.7, 136.7, 129.7, 129.6, 129.4, 125.1, 122.1, 120.5, 120.2, 110.8, 99.4, 21.2.

**2-(*m*-Tolyl)-1*H*-indole (11c).**<sup>28</sup> Compound **11c** was synthesized by following general procedure D using 1-*H*-indole-3-carboxylic acid (**10a**, 64 mg, 0.4 mmol) and 1-iodo-3-methylbenzene (**2c**, 102 μL, 0.8

mmol) at 120 °C for 3 h and purified by silica gel column chromatography (95:5 hexane/ethyl acetate): **11c** (65 mg, 78%, a white solid, mp: 128–129 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.32 (bs, NH, 1H), 7.65 (d, *J* = 7.7 Hz, CH, 1H), 7.54–7.46 (m, CH, 2H), 7.41 (d, *J* = 8.0 Hz, CH, 1H), 7.35 (t, *J* = 7.6 Hz, CH, 1H), 7.23–7.18 (m, CH, 1H), 7.18–7.12 (m, CH, 2H), 6.84 (d, *J* = 1.3 Hz, CH, 1H), 2.44 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 138.7, 138.1, 136.8, 132.3, 129.3, 129.0, 128.6, 125.9, 122.3, 122.3, 120.6, 120.2, 110.9, 99.9, 21.6.

**2-(4-(tert-Butyl)phenyl)-1H-indole (11d).**<sup>27</sup> Compound **11d** was synthesized by following general procedure D using 1-1H-indole-3-carboxylic acid (**10a**, 64 mg, 0.4 mmol) and 1-(tert-butyl)-4-iodobenzene (**2d**, 90 μL, 0.8 mmol) at 120 °C for 3 h and purified by silica gel column chromatography (95:5 hexane/ethyl acetate): **11d** (87 mg, 87%, a white solid, mp: 251–252 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.32 (bs, NH, 1H), 7.66–7.59 (m, CH, 3H), 7.48 (d, *J* = 8.4 Hz, CH, 2H), 7.40 (d, *J* = 7.9 Hz, CH, 1H), 7.22–7.16 (m, CH, 1H), 7.16–7.09 (m, CH, 1H), 6.81 (d, *J* = 1.5 Hz, CH, 1H), 1.37 (s, CH<sub>3</sub>, 9H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 150.9, 138.0, 136.7, 129.6, 129.4, 126.0, 124.9, 122.1, 120.5, 120.2, 110.8, 99.5, 34.7, 31.3.

**2-(4-Bromophenyl)-1H-indole (11e).**<sup>29</sup> Compound **11e** was synthesized by following general procedure D using 1-1H-indole-3-carboxylic acid (**10a**, 64 mg, 0.4 mmol) and 1-bromo-4-iodobenzene (**2e**, 227 mg, 0.8 mmol) at 120 °C for 3 h and purified by silica gel column chromatography (95:5 hexane/ethyl acetate): **11e** (90 mg, 84%, a white solid, mp: 209–210 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.29 (bs, CH, 1H), 7.63 (d, *J* = 7.9 Hz, CH, 1H), 7.59–7.55 (m, CH, 2H), 7.54–7.50 (m, CH, 2H), 7.40 (d, *J* = 8.1 Hz, CH, 1H), 7.24–7.19 (m, CH, 1H), 7.16–7.11 (m, CH, 1H), 6.83–6.82 (m, CH, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 136.9, 136.7, 132.2, 131.3, 129.2, 126.6, 122.8, 121.5, 120.8, 120.5, 111.0, 100.6.

**2-(3,4-Dimethylphenyl)-1H-indole (11k).**<sup>30</sup> Compound **11k** was synthesized by following general procedure D using 1-1H-indole-3-carboxylic acid (**10a**, 64 mg, 0.4 mmol) and 4-iodo-1,2-dimethylbenzene (**2k**, 114 μL, 0.8 mmol) at 120 °C for 3 h and purified by silica gel column chromatography (95:5 hexane/ethyl acetate): **11k** (74 mg, 83%, a white solid, mp: 142–143 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.29 (bs, NH, 1H), 7.66 (d, *J* = 7.4 Hz, CH, 1H), 7.47 (s, CH, 1H), 7.44–7.38 (m, CH, 2H), 7.25–7.12 (m, CH, 3H), 6.81 (s, CH, 1H), 2.36 (s, CH<sub>3</sub>, 3H), 2.33 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 138.2, 137.3, 136.7, 136.4, 130.3, 130.0, 129.4, 126.5, 122.6, 122.1, 120.5, 120.2, 110.8, 99.3, 20.0, 19.6.

**Gram-Scale Reaction and Synthetic Applications.** (a). *Gram-Scale Reaction of 3aa.* Compound **3aa** (1.05 g, 69%) was synthesized by following general procedure D using 1H-indole-3-carbaldehyde (**1a**) (1.0 g, 6.9 mmol), Pd(OAc)<sub>2</sub> (154 mg, 0.69 mmol, 10 mol %), AgOAc (2.3 g, 13.8 mmol, 2 equiv), and iodobenzene (**2a**) (1.54 mL, 13.8 mmol, 2 equiv) at 100 °C for 4 h and purified by silica gel column chromatography (80:20 hexane/ethyl acetate).

(b). *Gram-Scale Reaction of 5aa.* Compound **5aa** (960 mg, 65%) was synthesized by following general procedure D using 1-(1H-indol-3-yl)ethan-1-one (**4a**) (1.0 g, 6.3 mmol), Pd(OAc)<sub>2</sub> (141 mg, 0.63 mmol, 10 mol %), AgOAc (2.1 g, 0.8 mmol, 2 equiv), and iodobenzene (**2a**) (1.4 mL, 12.6 mmol, 2 equiv) at 120 °C for 10 h and purified by silica gel column chromatography (90:10 hexane/ethyl acetate).

(c). *N-Benzoylation Reaction of 5aa.* 1-(1-Benzyl-4-phenyl-1H-indol-2-yl)ethan-1-one (**12**). Compound **12** (80 mg, 82%) was prepared starting from **5aa** (71 mg, 0.3 mmol) according to the General Procedure C. Yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.61 (dd, *J* = 8.1, 1.1 Hz, CH, 2H), 7.45 (t, *J* = 7.6 Hz, CH, 2H), 7.42 (s, CH, 1H), 7.39–7.33 (m, CH, 1H), 7.33–7.27 (m, CH, 2H), 7.20–7.11 (m, CH, 4H), 7.04–6.97 (m, CH, 2H), 5.81 (s, CH<sub>2</sub>, 2H), 2.49 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 191.3, 140.5, 140.3, 138.3, 136.9, 134.7, 128.9, 128.8, 128.6, 127.6, 127.2, 126.6, 126.5, 124.6, 120.8, 112.5, 110.0, 48.4, 28.2. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>23</sub>H<sub>20</sub>NO: 326.1539; found: 326.1539.

(d). *Cross-Coupling of 2-Thienyl Boronic Acid and 5ca.*<sup>31</sup> 1-(4-Phenyl-7-(thiophen-2-yl)-1H-indol-2-yl)ethan-1-one (**5ca**, 63 mg, 0.2 mmol) and thiophen-2-ylboronic acid (31 mg, 0.24 mmol, 1.2 equiv) in DME/water (15 mL, 2:1 v/v, 0.013 M for **5ca**) was added Na<sub>2</sub>CO<sub>3</sub> (42 mg, 0.4 mmol, 2 equiv). After degassing, Pd(PPh<sub>3</sub>)<sub>4</sub> (12 mg, 10 μmol) was added and the mixture was boiled in a preheated oil bath at 100 °C for 18 h. Then, the mixture was cooled to room temperature and the solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 30 mL). The organic phase was combined, washed with water (3 × 20 mL), dried with Na<sub>2</sub>SO<sub>4</sub>, and concentrated under reduced pressure. The crude material was purified by column chromatography on silica gel (EtOAc/hexane, 1:9) to afford **13** as a red solid (41 mg, 85% yield, mp: 163–164 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.45 (bs, NH, 1H), 7.73–7.69 (m, CH, 2H), 7.59–7.53 (m, CH, 3H), 7.49–7.39 (m, CH, 4H), 7.27 (d, *J* = 7.6 Hz, CH, 1H), 7.23 (dd, *J* = 5.0, 3.6 Hz, CH, 1H), 2.59 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 190.2, 140.0, 139.9, 136.6, 136.0, 135.2, 128.82, 128.75, 128.2, 127.7, 126.8, 126.1, 125.5, 125.0, 121.1, 118.6, 109.8, 25.9. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>20</sub>H<sub>16</sub>NOS: 318.0947; found: 318.0947.

(e). *Construction of Functionalized Natural Product.* (1-Benzyl-4-(4-(tert-butyl)phenyl)-1H-indol-3-yl)(6-methoxy-9H-pyrido[3,4-b]indol-1-yl)methanone (**16**). Compounds **16** and **17** were prepared according to the method reported in the literature.<sup>32b</sup> 3-Acetylindole derivative **9ad** (57 mg, 0.15 mmol), 5-methoxytryptamine (**15**, 28 mg, 0.15 mmol, 1 equiv), I<sub>2</sub> (30 mg, 0.12 mmol, 0.8 equiv), hydrogen peroxide (30% aqueous solution, 1.5 equiv), and DMSO (2 mL, 2 M for **9ad**) were placed in a sealed tube (15 mL) with a magnetic stir bar. The resulting mixture was stirred in a preheated oil bath at 110 °C for 5 h, monitoring the reaction by TLC. Once the reaction was complete, the mixture was cooled to rt, diluted with water (50 mL), and then the reaction mixture was extracted with EtOAc (3 × 50 mL). The combined organic layers were washed with 10% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution, and then with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, and evaporated. The residue was purified by column chromatography on silica gel (80:20 hexane/ethyl acetate) to give **16** (63 mg, 75%, yellow solid, mp: 269–270 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 10.04 (bs, NH, 1H), 8.40–8.30 (m, CH, 2H), 7.90 (d, *J* = 5.0 Hz, CH, 1H), 7.50 (d, *J* = 2.4 Hz, CH, 1H), 7.36–7.27 (m, CH, 7H), 7.26–7.14 (m, CH, 5H), 7.10 (d, *J* = 8.3 Hz, CH, 2H), 5.42 (s, CH<sub>2</sub>, 2H), 3.89 (s, CH<sub>3</sub>, 3H), 1.07 (s, CH<sub>3</sub>, 9H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 189.9, 154.3, 148.9, 139.8, 138.1, 137.6, 137.3, 137.2, 136.7, 136.7, 136.1, 135.8, 130.8, 129.0, 128.1, 127.9, 127.2, 125.3, 124.4, 123.9, 123.3, 121.2, 118.7, 117.5, 116.4, 112.6, 109.1, 103.7, 56.1, 51.0, 34.2, 31.2. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>38</sub>H<sub>34</sub>N<sub>3</sub>O<sub>2</sub>: 564.2646; found: 564.2645.

(6-Methoxy-9H-pyrido[3,4-b]indol-1-yl)(4-phenyl-1H-indol-2-yl)-methanone (**17**). Compound **17** was synthesized starting from **5aa** (48 mg, 0.25 mmol) (prepared according to the literature procedure<sup>32b</sup>). The residue was purified by column chromatography on silica gel (80:20 hexane/ethyl acetate) to give **17** (86 mg, 83%, yellow solid, mp: >300 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 12.18 (bs, NH, 1H), 10.54 (bs, NH, 1H), 8.58 (d, *J* = 4.8 Hz, CH, 1H), 8.13 (d, *J* = 4.8 Hz, CH, 1H), 7.96 (s, CH, 1H), 7.77 (d, *J* = 7.2 Hz, CH, 2H), 7.62–7.41 (m, CH, 7H), 7.29–7.20 (m, CH, 2H), 3.93 (s, CH<sub>3</sub>, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 182.3, 154.7, 140.6, 138.4, 137.5, 137.4, 137.1, 136.8, 136.6, 136.0, 131.8, 128.9, 128.6, 127.4, 126.4, 125.6, 121.0, 120.3, 119.3, 118.6, 112.9, 111.7, 111.3, 103.8, 56.1. HRMS (ESI-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>27</sub>H<sub>20</sub>N<sub>3</sub>O<sub>3</sub>: 418.1550; found: 418.1552.

**Control Experiments.** a. *3,2-Carbonyl Migration from 6ca.* 1-(7-Bromo-4-phenyl-1H-indol-2-yl)ethan-1-one (**5ca**). Compound **6ca** (63 mg, 0.2 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and AgOAc (67 mg, 0.4 mmol, 2 equiv) were weighed in air and placed in a sealed tube (15 mL) with a magnetic stir bar. To the reaction mixture, HFIP/TFA (2 mL, 1:1, v/v) was added. The reaction mixture was then heated to 120 °C for 10 h under vigorous stirring. Upon completion, the reaction mixture was cooled to room temperature, the solvents were removed under reduced pressure, and



the resulting mixture was purified by a silica gel column chromatography column to give **5ca** (46 mg, 74%) using hexane/EtOAc (85:15 hexane/ethyl acetate).

**b. Reaction under Standard Conditions without 2a.** 2,2,2-Trifluoro-1-(1H-indol-3-yl)ethan-1-one (**18**).<sup>33</sup> Reaction of **10a** (or **10b**) under standard condition without **2a** gave the compound **18**. 2,2,2-trifluoro-1-(1H-indol-3-yl)ethan-1-one (**18**) (63 mg, 75% (or 59 mg, 70%), white solid, mp 153–154 °C); <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 12.72 (bs, NH, 1H), 8.51–8.47 (m, CH, 1H), 8.24–8.16 (m, CH, 1H), 7.62–7.56 (m, CH, 1H), 7.38–7.27 (m, CH, 2H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, DMSO-*d*<sub>6</sub>): δ 174.1 (q, *J* = 33.8 Hz), 137.6 (q, *J* = 4.8 Hz), 136.6, 125.7, 124.3, 123.4, 121.1, 116.9 (q, *J* = 291.7 Hz), 113.0, 108.8.

**c. Reaction under Standard Conditions of 19a and 2a.** Compound **11a** (27 mg, 70%) was synthesized by following general procedure D using 1H-indole (**19a**, 24 mg, 0.2 mmol), AgOAc (67 mg, 0.4 mmol), Pd(OAc)<sub>2</sub> (5 mg, 20 μmol, 10 mol %), and iodobenzene (**2a**, 45 μL, 0.4 mmol) at 120 °C for 3 h and purified by silica gel column chromatography (95:15 hexane/ethyl acetate).

**d. Reaction under Standard Conditions of 10c without 2a with 2a.** 2-Methyl-1H-indole (**19b**).<sup>46</sup> (Red solid; mp: 59–60 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.75 (bs, NH, 1H), 7.56 (d, *J* = 7.6 Hz, CH, 1H), 7.28 (d, *J* = 7.9 Hz, CH, 1H), 7.18–7.09 (m, CH, 2H), 6.25 (s, CH, 1H), 2.44 (s, CH, 3H). <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 136.1, 135.2, 129.1, 121.0, 119.7 (2C), 110.3, 100.4, 13.8.

**e. Reaction under Standard Conditions of 19b and 2a.** In the reaction of **19b** and **2a** under standard conditions, no products were observed.

## ■ ASSOCIATED CONTENT

### SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.joc.2c00716>.

NMR and HRMS spectra of compounds (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

Nurullah Saracoglu – Department of Chemistry, Faculty of Sciences, Atatürk University, Erzurum 25240, Turkey; [orcid.org/0000-0002-1504-7480](https://orcid.org/0000-0002-1504-7480); Email: [nsarac@atauni.edu.tr](mailto:nsarac@atauni.edu.tr)

### Authors

Yunus Taskesenligil – Department of Chemistry, Faculty of Sciences, Atatürk University, Erzurum 25240, Turkey; [orcid.org/0000-0002-5399-2520](https://orcid.org/0000-0002-5399-2520)

Murat Aslan – Department of Chemistry, Faculty of Sciences, Atatürk University, Erzurum 25240, Turkey; [orcid.org/0000-0001-9251-1462](https://orcid.org/0000-0001-9251-1462)

Tuba Cogurcu – Department of Chemistry, Faculty of Sciences, Atatürk University, Erzurum 25240, Turkey

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.joc.2c00716>

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

The authors thank Atatürk University, The Scientific Research Projects Coordination Unit, Turkey (FAD-2019-7025 and FAD-2018-6642), for financial support to research projects of Department of Chemistry. Y.T. thanks the Scientific and Technological Research Council of Turkey (TUBITAK—BIDEB 2211/C National Ph.D. Scholarship Program in the Priority Fields in Science and Technology) for a scholarship.

The authors thank Prof. Dr. Cavit Kazaz for his help in NOE experiments.

## ■ REFERENCES

- (1) (a) Park, Y.; Kim, Y.; Chang, S. Transition metal-catalyzed C–H amination: scope, mechanism, and applications. *Chem. Rev.* **2017**, *117*, 9247–9301. (b) Gensch, T.; Hopkinson, M. N.; Glorius, F.; Wencel-Delord, J. Mild metal-catalyzed C–H activation: examples and concepts. *Chem. Soc. Rev.* **2016**, *45*, 2900–2936.
- (2) Davies, H. M. L.; Morton, D. Recent advances in C–H functionalization. *J. Org. Chem.* **2016**, *81*, 343–350.
- (3) (a) Wan, Y.; Li, Y.; Yan, C.; Yan, M.; Tang, Z. Indole: A privileged scaffold for the design of anti-cancer agents. *Eur. J. Med. Chem.* **2019**, *183*, No. 111691. (b) Kochanowska-Karamyan, A. J.; Hamann, M. T. Marine indole alkaloids: potential new drug leads for the control of depression and anxiety. *Chem. Rev.* **2010**, *110*, 4489–4497. (c) Pegu, M.; Kazim, S.; Huang, P.; Lezama, L.; Ahmad, S. Tetra-indole core as a dual agent: a hole selective layer that passivates defects in perovskite solar cells. *J. Mater. Chem. C* **2021**, *9*, 7074–7082. (d) Miyasaka, M.; Fukushima, A.; Satoh, T.; Hirano, K.; Miura, M. Fluorescent Diarylindoles by Palladium-Catalyzed Direct and Decarboxylative Arylations of Carboxyindoles. *Chem. – Eur. J.* **2009**, *15*, 3674–3677.
- (4) (a) Lanke, V.; Prabhu, K. R. Highly regioselective C2-alkenylation of indoles using the *N*-benzoyl directing group: an efficient Ru-catalyzed coupling reaction. *Org. Lett.* **2013**, *15*, 2818–2821. (b) Li, B.; Ma, J.; Xie, W.; Song, H.; Xu, S.; Wang, B. Ruthenium-catalyzed regioselective C2 alkenylation of indoles and pyrroles via C–H bond functionalization. *J. Org. Chem.* **2013**, *78*, 9345–9353. (c) Bandini, M.; Eichholzer, A. Catalytic Functionalization of Indoles in a New Dimension. *Angew. Chem., Int. Ed.* **2009**, *48*, 9608–9644. (d) Bartoli, G.; Bencivenni, G.; Dalpozzo, R. Organocatalytic Strategies for the Asymmetric Functionalization of Indoles. *Chem. Soc. Rev.* **2010**, *39*, 4449–4465. (e) Sandtorv, A. H. Transition Metal-Catalyzed C–H Activation of Indoles. *Adv. Synth. Catal.* **2015**, *357*, 2403–2435. (f) Zhang, L. Q.; Yang, S.; Huang, X.; You, J.; Song, F. Aerobic Ru-catalyzed direct C2-olefination of *N*-heteroarenes with alkenes directed by a removable *N*-dimethylcarbamoyl group. *Chem. Commun.* **2013**, *49*, 8830–8832.
- (5) (a) Leitch, J. A.; Bhonoah, Y.; Frost, C. G. Beyond C2 and C3: Transition-Metal-Catalyzed C–H Functionalization of Indole. *ACS Catal.* **2017**, *7*, 5618–5627. (b) Yang, Y.; Shi, Z. Regioselective Direct Arylation of Indoles on the Benzenoid Moiety. *Chem. Commun.* **2018**, *54*, 1676–1685. (c) Wen, J.; Shi, Z. From C4 to C7: Innovative Strategies for Site-Selective Functionalization of Indole C–H Bonds. *Acc. Chem. Res.* **2021**, *54*, 1723–1736. (d) Kalepu, J.; Gandeepan, P.; Ackermann, L.; Pilarski, L. T. C4–H indole functionalisation: precedent and prospects. *Chem. Sci.* **2018**, *9*, 4203–4216. (e) Prabagar, B.; Yang, Y.; Shi, Z. Site-selective C–H functionalization to access the arene backbone of indoles and quinolines. *Chem. Soc. Rev.* **2021**, *50*, 11249–11269. (f) Lv, J.; Chen, X.; Xue, X. S.; Zhao, B.; Liang, Y.; Wang, M.; Jin, L.; Yuan, Y.; Han, Y.; Zhao, Y.; Lu, Y.; Zhao, J.; Sun, W. Y.; Houk, K. N.; Shi, Z. Metal-free directed sp<sup>2</sup>-C–H borylation. *Nature* **2019**, *575*, 336–340. (g) Wang, D.; Chen, X.; Wong, J. J.; Jin, L.; Li, M.; Zhao, Y.; Houk, K. N.; Shi, Z. Phosphorus(III)-assisted regioselective C–H silylation of heteroarenes. *Nat. Commun.* **2021**, *12*, No. 524. (h) Lanke, V.; Ramaiah Prabhu, K. Regioselective synthesis of 4-substituted indoles via C–H activation: a ruthenium catalyzed novel directing group strategy. *Org. Lett.* **2013**, *15*, 6262–6265. (i) Urbina, K.; Tresp, D.; Sipps, K.; Szostak, M. Recent Advances in Metal-Catalyzed Functionalization of Indoles. *Adv. Synth. Catal.* **2021**, *363*, 2723–2739. (j) Shah, T. A.; De, P. B.; Pradhan, S.; Punniyamurthy, T. Transition-metal-catalyzed site-selective C7-functionalization of indoles: advancement and future prospects. *Chem. Commun.* **2019**, *55*, 572–587.
- (6) (a) Qiu, X.; Wang, P.; Wang, D.; Wang, M.; Yuan, Y.; Shi, Z. P<sup>III</sup>-Chelation-Assisted Indole C7-Arylation, Olefination, Methylation, and Acylation with Carboxylic Acids/Anhydrides by Rhodium Catalysis. *Angew. Chem., Int. Ed.* **2019**, *58*, 1504–1508. (b) Han,



- X.; Yuan, Y.; Shi, Z. Rhodium-Catalyzed Selective C–H Trideuteromethylation of Indole at C7 Position Using Acetic- $d_6$  Anhydride. *J. Org. Chem.* **2019**, *84*, 12764–12772. (c) Borah, A. J.; Shi, Z. Rhodium-catalyzed, remote terminal hydroarylation of activated olefins through a long-range deconjugative isomerization. *J. Am. Chem. Soc.* **2018**, *140*, 6062–6066. (d) Shi, X.; Wang, Z.; Li, Y.; Li, X.; Li, X.; Shi, D. Palladium-Catalyzed Remote C–H Phosphonylation of Indoles at the C4 and C6 Positions by a Radical Approach. *Angew. Chem., Int. Ed.* **2021**, *60*, 13871–13876. (e) Borah, A. J.; Shi, Z. Palladium-catalyzed regioselective C–H fluoroalkylation of indoles at the C4-position. *Chem. Commun.* **2017**, *53*, 3945–3948. (f) Yang, Y.; Qiu, X.; Zhao, Y.; Mu, Y.; Shi, Z. Palladium-catalyzed C–H arylation of indoles at the C7 position. *J. Am. Chem. Soc.* **2016**, *138*, 495–498.
- (7) (a) Chen, S.; Zhang, M.; Su, R.; Chen, X.; Feng, B.; Yang, Y.; You, J. C2/C4 Regioselective heteroarylation of indoles by tuning C–H metalation modes. *ACS Catal.* **2019**, *9*, 6372–6379. (b) Chen, S.; Feng, B.; Zheng, X.; Yin, J.; Yang, S.; You, J. Iridium-catalyzed direct regioselective C4-amidation of indoles under mild conditions. *Org. Lett.* **2017**, *19*, 2502–2505.
- (8) (a) Yang, G.; Lindovska, P.; Zhu, D.; Kim, J.; Wang, P.; Tang, R. Y.; Movassaghi, M.; Yu, J. Q. Pd(II)-catalyzed *meta*-C–H olefination, arylation, and acetoxylation of indolines using a U-shaped template. *J. Am. Chem. Soc.* **2014**, *136*, 10807–10813. (b) Yang, G.; Zhu, D.; Wang, P.; Tang, R. Y.; Yu, J. Q. Remote C–H Activation of Various N-Heterocycles Using a Single Template. *Chem. – Eur. J.* **2018**, *24*, 3434–3438.
- (9) (a) Choi, I.; Messinis, A. M.; Ackermann, L. C7-Indole Amidations and Alkenylations by Ruthenium(II) Catalysis. *Angew. Chem., Int. Ed.* **2020**, *59*, 12534–12540. (b) Wang, W.; Wu, J.; Kuniyil, R.; Kopp, A.; Lima, R. N.; Ackermann, L. Peptide Late-Stage Diversifications by Rhodium-Catalyzed Tryptophan C7 Amidation. *Chem* **2020**, *6*, 3428–3439.
- (10) (a) Pradhan, S.; Mishra, M.; De, P. B.; Banerjee, S.; Punniyamurthy, T. Weak Coordination Enabled Switchable C4-Alkenylation and Alkylation of Indoles with Allyl Alcohols. *Org. Lett.* **2020**, *22*, 1720–1725. (b) Zhang, J.; Wu, M.; Fan, J.; Xu, Q.; Xie, M. Selective C–H acylation of indoles with  $\alpha$ -oxocarboxylic acids at the C4 position by palladium catalysis. *Chem. Commun.* **2019**, *55*, 8102–8105. (c) Thrimurtulu, N.; Dey, A.; Singh, A.; Pal, K.; Maiti, D.; Volla, C. M. R. Palladium Catalyzed Regioselective C4-Arylation and Olefination of Indoles and Azaindoles. *Adv. Synth. Catal.* **2019**, *361*, 1441–1446. (d) Sherikar, M. S.; Kapaniaiah, R.; Lanke, V.; Prabhu, K. R. Rhodium(III)-catalyzed C–H activation at the C4-position of indole: switchable hydroarylation and oxidative Heck-type reactions of maleimides. *Chem. Commun.* **2018**, *54*, 11200–11203.
- (11) (a) Banjare, S. K.; Nanda, T.; Pati, B. V.; Adhikari, G. K. D.; Dutta, J.; Ravikumar, P. C. Breaking the Trend: Insight into Unforeseen Reactivity of Alkynes in Cobalt-Catalyzed Weak Chelation-Assisted Regioselective C(4)–H Functionalization of 3-Pivaloyl Indole. *ACS Catal.* **2021**, *11*, 11579–11587. (b) Zhang, S. S.; Liu, Y. Z.; Zheng, Y. C.; Xie, H.; Chen, S. Y.; Song, J. L.; Shu, B. Rhodium(III)-Catalyzed Regioselective C–H Allylation and Prenylation of Indoles at C4-Position. *Adv. Synth. Catal.* **2022**, *364*, 64–70. (c) Kim, Y.; Park, Y.; Chang, S. Delineating physical organic parameters in site-selective C–H functionalization of indoles. *ACS Cent. Sci.* **2018**, *4*, 768–775.
- (12) Leitch, J. A.; McMullin, C. L.; Mahon, M. F.; Bhonoah, Y.; Frost, C. G. Remote C6-selective ruthenium-catalyzed C–H alkylation of indole derivatives via  $\sigma$ -activation. *ACS Catal.* **2017**, *7*, 2616–2623.
- (13) (a) Yang, Y.; Gao, P.; Zhao, Y.; Shi, Z. Regiocontrolled Direct C–H Arylation of Indoles at the C4 and C5 Positions. *Angew. Chem., Int. Ed.* **2017**, *56*, 3966–3971. (b) Yang, Y.; Li, R.; Zhao, Y.; Zhao, D.; Shi, Z. Cu-catalyzed direct C6-arylation of indoles. *J. Am. Chem. Soc.* **2016**, *138*, 8734–8737.
- (14) (a) Okada, T.; Sakai, A.; Hinoue, T.; Satoh, T.; Hayashi, Y.; Kawauchi, S.; Chandrabunaidu, K.; Miura, M. Rhodium(III)-catalyzed oxidative coupling of *N*-phenylindole-3-carboxylic acids with alkenes and alkynes via C4–H and C2–H/C2'–H bond cleavage. *J. Org. Chem.* **2018**, *83*, 5639–5649. (b) Banjare, S. K.; Nanda, T.; Ravikumar, P. C. Cobalt-catalyzed regioselective direct C4 alkenylation of 3-acetylindole with Michael acceptors using a weakly coordinating functional group. *Org. Lett.* **2019**, *21*, 8138–8143.
- (15) Lanke, V.; Prabhu, K. R. Iridium(III) catalyzed regioselective amidation of indoles at the C4-position using weak coordinating groups. *Chem. Commun.* **2017**, *53*, 5117–5120.
- (16) Pradhan, S.; De, P. B.; Punniyamurthy, T. Weak coordination-guided regioselective direct redox-neutral C4 allylation of indoles with Morita–Baylis–Hillman adducts. *Org. Lett.* **2019**, *21*, 9898–9903.
- (17) (a) Chen, X.; Zheng, G.; Li, Y.; Song, G.; Li, X. Rhodium-catalyzed site-selective coupling of indoles with diazo esters: C4-alkylation versus C2-annulation. *Org. Lett.* **2017**, *19*, 6184–6187. (b) Sherikar, M. S.; Devarajappa, R.; Prabhu, K. R. Weak Coordinating Carbonyl-Directed Rhodium(III)-Catalyzed C–H Activation at the C4-Position of Indole with Allyl Alcohols. *J. Org. Chem.* **2020**, *85*, 5516–5524.
- (18) (a) Liu, X.; Li, G.; Song, F.; You, J. Unexpected regioselective carbon–hydrogen bond activation/cyclization of indolyl aldehydes or ketones with alkynes to benzo-fused oxindoles. *Nat. Commun.* **2014**, *5*, No. 5030. (b) Bettadapur, K. R.; Kapaniaiah, R.; Lanke, V.; Prabhu, K. R. Weak Directing Group Steered Formal Oxidative [2+2+2]-Cyclization for Selective Benzannulation of Indoles. *J. Org. Chem.* **2018**, *83*, 1810–1818.
- (19) Li, J. F.; Zhao, R. F.; Zhou, F. Q.; She, M. Y.; Zhang, J.; Yin, B.; Zhang, S.-Y.; Li, J. L. Exploring the necessity of an acidic additive for Pd(II)-catalyzed exclusive C4-fluoroalkylation of 3-acetylindole: a detailed DFT study on the mechanism and regioselectivity. *Org. Chem. Front.* **2019**, *6*, 2607–2618.
- (20) Kuang, G.; Liu, D.; Chen, X.; Liu, G.; Fu, Y.; Peng, Y.; Li, H.; Zhou, Y. Transient Directing Group Strategy as a Unified Method for Site Selective Direct C4–H Halogenation of Indoles. *Org. Lett.* **2021**, *23*, 8402–8406.
- (21) Liu, X. H.; Park, H.; Hu, J. H.; Hu, Y.; Zhang, Q. L.; Wang, B. L.; Sun, B.; Yeung, K.-S.; Zhang, F.-L.; Yu, J. Q. Diverse *ortho*-C(sp<sup>2</sup>)–H functionalization of benzaldehydes using transient directing groups. *J. Am. Chem. Soc.* **2017**, *139*, 888–896.
- (22) Basak, S.; Paul, T.; Punniyamurthy, T. Palladium-Catalyzed 2-fold C–H Activation/C–C Coupling for C4-Arylation of Indoles Using Weak Chelation. *Org. Lett.* **2022**, *24*, 554–558.
- (23) (a) Nandi, D.; Jhou, Y. M.; Lee, J. Y.; Kuo, B. C.; Liu, C. Y.; Huang, P. W.; Lee, H. M. Pd(0)-catalyzed decarboxylative coupling and tandem C–H arylation/decarboxylation for the synthesis of heteroaromatic biaryls. *J. Org. Chem.* **2012**, *77*, 9384–9390. (b) Jose, A.; Vijaykumar, G.; Vardhanapu, P. K.; Mandal, S. K. Abnormal NHC supported palladacycles: Regioselective arylation of heteroarenes via decarboxylation. *J. Organomet. Chem.* **2018**, *865*, 51–57.
- (24) Arun, V.; Pilania, M.; Kumar, D. Access to 2-Arylindoles via Decarboxylative C–C Coupling in Aqueous Medium and to Heteroaryl Carboxylates under Base-Free Conditions using Diaryliodonium Salts. *Chem. Asian J.* **2016**, *11*, 3345–3349.
- (25) Cheng, Y.; Yu, S.; He, Y.; An, G.; Li, G.; Yang, Z. C4-arylation and domino C4-arylation/3,2-carbonyl migration of indoles by tuning Pd catalytic modes: Pd(I)–Pd(II) catalysis vs. Pd(II) catalysis. *Chem. Sci.* **2021**, *12*, 3216–3225.
- (26) Taskesenligil, Y.; Lafzi, F.; Kilic, H.; Saracoglu, N. Palladium-catalyzed regioselective C2-arylation of 5-aminoindole. *J. Heterocycl. Chem.* **2019**, *56*, 3289–3296.
- (27) Liu, C.; Ding, L.; Guo, G.; Liu, W.; Yang, F. L. Palladium-catalyzed direct arylation of indoles with arylsulfonyl hydrazides. *Org. Biomol. Chem.* **2016**, *14*, 2824–2827.
- (28) Modha, S. G.; Greaney, M. F. Atom-economical transformation of diaryliodonium salts: tandem C–H and N–H arylation of indoles. *J. Am. Chem. Soc.* **2015**, *137*, 1416–1419.
- (29) Deprez, N. R.; Kalyani, D.; Krause, A.; Sanford, M. S. Room temperature palladium-catalyzed 2-arylation of indoles. *J. Am. Chem. Soc.* **2006**, *128*, 4972–4973.

- (30) Bhunia, S. K.; Polley, A.; Natarajan, R.; Jana, R. Through-Space 1,4-Palladium Migration and 1,2-Aryl Shift: Direct Access to Dibenzo[*a,c*]carbazoles through a Triple C–H Functionalization Cascade. *Chem. – Eur. J.* **2015**, *21*, 16786–16791.
- (31) Aslan, M.; Taskesenligil, Y.; Piravadi, S.; Saracoglu, N. Functionalization at Nonperipheral Positions of Triazatruxene: Modular Construction of 1,6,11-Triarylated-Triazatruxenes for Potentially Organic Electronics and Optoelectronics. *J. Org. Chem.* **2022**, *87*, S037–S050.
- (32) (a) Szabó, T.; Hazai, V.; Volk, B.; Simig, G.; Milen, M. First total synthesis of the  $\beta$ -carboline alkaloids trigonostemine A, trigonostemine B and a new synthesis of pityriacitrin and hyrtiosulawesine. *Tetrahedron Lett.* **2019**, *60*, 1471–1475. (b) Zhu, Y. P.; Liu, M. C.; Cai, Q.; Jia, F. C.; Wu, A. X. A Cascade Coupling Strategy for One-Pot Total Synthesis of  $\beta$ -Carboline and Isoquinoline-Containing Natural Products and Derivatives. *Chem. – Eur. J.* **2013**, *19*, 10132–10137. (c) Liew, L. P.; Fleming, J. M.; Longeon, A.; Mouray, E.; Florent, I.; Bourguet-Kondracki, M. L.; Copp, B. R. Synthesis of 1-indolyl substituted  $\beta$ -carboline natural products and discovery of antimalarial and cytotoxic activities. *Tetrahedron* **2014**, *70*, 4910–4920.
- (33) Yao, S. J.; Ren, Z. H.; Wang, Y. Y.; Guan, Z. H. Friedel–Crafts fluoroacetylation of indoles with fluorinated acetic acids for the synthesis of fluoromethyl indol-3-yl ketones under catalyst- and additive-free conditions. *J. Org. Chem.* **2016**, *81*, 4226–4234.
- (34) Zhang, C.; Xu, D.; Wang, J.; Kang, C. Efficient synthesis and biological activity of novel Indole derivatives as VEGFR-2 tyrosine kinase inhibitors. *Russ. J. Gen. Chem.* **2017**, *87*, 3006–3016.
- (35) (a) Shan, G.; Yang, X.; Ma, L.; Rao, Y. Pd-Catalyzed C–H Oxygenation with TFA/TFAA: Expedient Access to Oxygen-Containing Heterocycles and Late-Stage Drug Modification. *Angew. Chem., Int. Ed.* **2012**, *51*, 13070–13074. (b) Vána, J.; Lang, J.; Soltesova, M.; Hanusek, J.; Ruzicka, A.; Sedlak, M.; Roithova, J. The role of trinuclear species in a palladium acetate/trifluoroacetic acid catalytic system. *Dalton Trans.* **2017**, *46*, 16269–16275.
- (36) Lane, B. S.; Sames, D. Direct C–H bond arylation: Selective palladium-catalyzed C2-arylation of *N*-substituted indoles. *Org. Lett.* **2004**, *6*, 2897–2900.
- (37) Coowar, D.; Bouissac, J.; Hanbali, M.; Paschaki, M.; Mohier, E.; Luu, B. Effects of indole fatty alcohols on the differentiation of neural stem cell derived neurospheres. *J. Med. Chem.* **2004**, *47*, 6270–6282.
- (38) Shi, X.; Xu, W.; Wang, R.; Zeng, X.; Qiu, H.; Wang, M. Ketone-directed cobalt(III)-catalyzed regioselective C2 amidation of indoles. *J. Org. Chem.* **2020**, *85*, 3911–3920.
- (39) Adams, S. E.; Parr, C.; Miller, D. J.; Allemann, R. K.; Hallett, M. B. Potent inhibition of  $\text{Ca}^{2+}$ -dependent activation of calpain-1 by novel mercaptoacrylates. *Med. Chem. Commun.* **2012**, *3*, S66–S70.
- (40) Yin, Z.; Wang, Z.; Wu, X. F. Selective nickel-catalyzed dehydrogenative–decarboxylative formylation of indoles with glyoxylic acid. *Org. Biomol. Chem.* **2018**, *16*, 3707–3710.
- (41) Betterley, N. M.; Kerdphon, S.; Chaturonrutsamee, S.; Kongsriprapan, S.; Surawatanawong, P.; Soorukram, D.; Kuhakarn, C.; et al. *et al.* Bi(OTf)<sub>3</sub> Enabled C–F Bond Cleavage in HFIP: Electrophilic Aromatic Formylation with Difluoro(phenylsulfanyl)-methane. *Asian J. Org. Chem.* **2018**, *7*, 1642–1647.
- (42) Abe, T.; Haruyama, T.; Yamada, K. C4 Pictet–Spengler Reactions for the Synthesis of Core Structures in Hyrtiazepine Alkaloids. *Synthesis* **2017**, *49*, 4141–4150.
- (43) Jones, B. T.; García-Cárceles, J.; Caiger, L.; Hazelden, I. R.; Lewis, R. J.; Langer, T.; Bower, J. F. Complex Polyheterocycles and the Stereochemical Reassignment of Pileamartine A via Aza-Heck Triggered Aryl C–H Functionalization Cascades. *J. Am. Chem. Soc.* **2021**, *143*, 15593–15598.
- (44) Giraud, F.; Alves, G.; Debiton, E.; Nauton, L.; Théry, V.; Durieu, E.; Ferandin, Y.; Lozach, O.; Meijer, L.; Anizon, F.; Pereira, E.; Moreau, P. Synthesis, protein kinase inhibitory potencies, and in vitro antiproliferative activities of meridianin derivatives. *J. Med. Chem.* **2011**, *54*, 4474–4489.
- (45) Ali, M. A.; Punniyamurthy, T. Domino ligand-free copper-catalyzed synthesis of polysubstituted indoles. *Synlett* **2011**, *5*, 623–626.
- (46) Maeda, K.; Matsubara, R.; Hayashi, M. Synthesis of substituted anilines from cyclohexanones using Pd/C–ethylene system and its application to indole Synthesis. *Org. Lett.* **2021**, *23*, 1530–1534.