

Synthesis of Hexamethyl-1,1'-spirobiindane-Based Chiral Spiro Cp Ligands and Their Application in Rhodium-Catalyzed Enantioselective Aryl C–H Addition to Nitroalkenes

Hui Yang, Ru Zhang, Su-Zhen Zhang, Qing Gu, and Shu-Li You*



Cite This: *ACS Catal.* 2023, 13, 8838–8844



Read Online

ACCESS |


Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: Spiro cyclopentadienyl rhodium (SCpRh) complexes are powerful catalysts for promoting asymmetric C–H functionalization reactions. However, the application of chiral SCp ligands is limited due to tedious synthetic procedures and expensive starting materials. Herein, we have developed a series of chiral spiro ligands (BCSCp) with 6–7 steps from commercially available and cheap Bisphenol C. Their corresponding rhodium complexes have been prepared and successfully applied in enantioselective aryl C–H addition to nitroalkenes, affording a series of C–H adducts in up to 88% yield with up to 98% ee.

KEYWORDS: asymmetric catalysis, C–H functionalization, C–H addition, chiral cyclopentadienyl, rhodium catalysis



Transition-metal-catalyzed asymmetric C–H functionalization has received considerable interest as an increasingly important strategy to access versatile chiral molecules.^{1–10} Notably, chiral cyclopentadienyl metal complexes have been developed as one of the most efficient catalysts to realize asymmetric C–H bond functionalization.^{11–20} Since the pioneering works by Cramer²¹ and Ward and Rovis,²² diverse chiral cyclopentadienyl ligands have been revealed,^{23–39} such as binaphthyl-derived Cp I,^{23–30} piperidine-fused Cp II,³¹ cyclopentane-fused Cp III,³² and bicyclo[2,2,2]octane-fused Cp IV,³³ as well as planar prochiral Cp ligands^{34–37} and others^{38,39} (Figure 1). Despite these advances, it remains essential to develop a concise synthetic approach for novel chiral ligands to meet the ever-growing demands of the rapid development of asymmetric C–H functionalization reactions.

1,1'-Spirobiindane has been recognized as a privileged chiral scaffold to provide an excellent platform for chiral ligand and catalyst diversification since the pioneering work of Zhou.^{40–46} In 2016, our group designed and synthesized a series of 1,1'-spirobiindane-derived Cp (SCp) ligands and successfully applied their Rh complexes in an asymmetric oxidative coupling reaction.⁴⁷ Since then, application of SCpRh in versatile catalytic asymmetric C–H functionalization reactions has been well demonstrated.^{48–60} Despite their successful applications in challenging C–H functionalization reactions, these SCp ligands are synthetically less practical due to the tedious synthesis and relatively expensive SPINOL starting material. Therefore, the design of novel SCp ligands with practical synthesis is of great significance and in high demand.

Transition-metal-catalyzed direct C–H addition to polar unsaturated bonds C=X (X = C, N, O) such as electron-deficient alkenes, aldehydes, and ketones has emerged as a




Figure 1. Previously and newly developed chiral cyclopentadienyl ligands.


Received: May 15, 2023

Revised: June 9, 2023

Published: June 20, 2023



Scheme 1. Synthesis of Hexamethyl-1,1'-spirobiindane Cp Ligands and Their Rhodium Complexes.




powerful approach for the formation of C–C bonds.^{61–68} The most significant feature of these reactions is that the crucial nucleophilic C–M species are generated catalytically *in situ* via C–H activation. In recent years, considerable studies on enantioselective C–H direct addition reactions have been successfully developed by Shibata,^{69–72} Cramer,^{30,73} Matsunaga,^{74–77} and others.^{78–83} Driven by our continuous interests in developing new chiral cyclopentadienyl ligands and asymmetric C–H functionalization,^{84–90} we synthesized a novel class of chiral spiro (BCSCp) ligands from Bisphenol C in 6–7 steps. In addition, these Rh^{III} complexes of BCSCps were found to be highly efficient for a challenging asymmetric C–H addition of aryl amides to nitroalkenes, affording a series of C–H adducts in excellent yields and enantioselectivities. Herein, we report the details of this study. Recently, the Wang group also reported the synthesis and application of this class of ligands during the submission of our manuscript.⁹¹

As shown in Scheme 1, chiral Cp ligands **5** could be conveniently synthesized from Bisphenol C. The preparation of enantiopure hexamethyl-1,1'-spirobiindane-6,6'-diol **1** (6,6'-HMSIOL) was carried out according to the known procedure from commercially available Bisphenol C.⁹² The Duff reaction of (S)-**1** with hexamethylenetetramine (HMTA) afforded dialdehyde (S)-**2** in 87% yield on a 10 g scale. O-Alkylation of (S)-**2** with MeI in the presence of potassium carbonate gave (S)-**3a** in an 87% yield. Then, dialdehyde (S)-**3a** was further transformed to the corresponding dichloride (S)-**4a** in 86% yield over two steps by NaBH₄ reduction, followed by chlorination with SOCl₂. Double alkylation of CpNa with (S)-**4a** gave substituted cyclopentadiene (S)-**5a** in 46% yield. The chiral CpRh^I complex (S)-Rh-**1** was smoothly prepared by reaction with [Rh(COD)-OAc]₂ in toluene/methanol at 70 °C under argon. At the same time, the corresponding CpRh^I complexes (S)-Rh-**2** to (S)-Rh-**4** were prepared by a similar strategy (see the Supporting

Information for details). Finally, upon oxidation of the CpRh^I complex with iodine, the CpRh^{III} complex (S)-Rh-**5** was obtained in 85% yield. The structure of (S)-Rh-**1** was confirmed unambiguously by X-ray crystallographic analysis.

Then, the utility of these newly developed BCSCpRh complexes in asymmetric C–H functionalization was explored. Recently, we reported a Rh-catalyzed enantioselective direct addition of aryl C–H bond to nitroalkenes,⁵³ and the highest enantioselectivity for the reaction of pyrrolidine benzamide **6a** with nitrostyrene **7a** could only reach 86% ee in the presence of the optimized catalyst Rh-**9**. Furthermore, the ee values of the products are generally lower than 90%. To our delight, with the newly developed BCSCpRh complex Rh-**1**, the reaction proceeded with excellent enantioselectivity to give **8aa**, albeit in a poor yield (Table 1, entry 1, 21% NMR yield, 92% ee). The existence of the 3,3,3',3'-tetramethyl groups might affect the dihedral angle of two indanyls, leading to a better chiral induction environment. When catalysts Rh-**2**, Rh-**3**, and Rh-**4** were employed, inferior results were given (see the Supporting Information for details). Gratifyingly, Rh-**5** led to a remarkable improvement in catalytic efficiency, giving **8aa** in 64% NMR yield with 92% ee (entry 2). In contrast, the utilization of Rh-**6** or Rh-**7** gave a lower efficiency and enantioselectivity (entries 3 and 4). In addition, the desired product **8aa** was obtained in 78% yield with only 23% ee when Rh-**8** was used, probably due to the decreased steric hindrance (entry 5). Therefore, Rh-**5** was identified as the optimized catalyst for further optimizations. Screening of additives indicated that NaOAc was beneficial, giving the product in 82% NMR yield with 94% ee (entries 6–8). As expected, both excellent enantioselectivity and good yield were maintained when the additive loading was reduced to 0.4 equiv (entry 9, 81% yield, 93% ee). Further reducing the amount of the additive to 0.2 equiv resulted in a measurable decrease in yield (entry 10, 72% yield, 93% ee). For comparison, Rh-**9** and

Table 1. Optimization of Reaction Conditions^a


Reaction scheme for Table 1: 6a + 7a with [Rh] (5 mol%), AgSbF₆ (40 mol%), additive (x equiv) in DCE at 80 °C for 24 h to form 8aa.

Chemical structures of Rh-1, Rh-5, Rh-6, Rh-7, Rh-8, Rh-9, and Rh-10 are shown.


Rh-5, R = OMe
Rh-6, R = OⁱPr
Rh-7, R = Ph
Rh-8, R = H

entry	[Rh]	additive (equiv)	yield (%) ^b	ee (%) ^c
1 ^d	Rh-1	BPO (0.05)	21	92
2	Rh-5		64	92
3	Rh-6		45	87
4	Rh-7		46	84
5	Rh-8		78	23
6	Rh-5	NaOAc (1.0)	82	94
7	Rh-5	KOAc (1.0)	30	92
8	Rh-5	Zn(OAc) ₂ (1.0)	54	94
9	Rh-5	NaOAc (0.4)	81	93
10	Rh-5	NaOAc (0.2)	72	93
11	Rh-9	NaOAc (0.4)	78	88
12	Rh-10	NaOAc (0.4)	56	84

^aReaction conditions unless specified otherwise: **6a** (0.15 mmol), **7a** (0.1 mmol), [Rh] (5 mol%), AgSbF₆ (40 mol%), and additive in DCE (0.5 mL) at 80 °C for 24 h under an argon atmosphere. ^bDetermined by ¹H NMR analysis of the crude reaction mixture using CH₂Br₂ as an internal standard. ^cThe ee values of **8aa** were determined by HPLC analysis on a chiral stationary phase. ^dAbsence of AgSbF₆.

Rh-10 were also tested under identical reaction conditions (entries 11 and 12). However, lower enantioselectivity was observed in both cases (88% and 84% ee, respectively). Finally, the optimized reaction conditions were obtained as the following: **6a** (1.5 equiv), **7a** (1.0 equiv), **Rh-5** (5 mol%), AgSbF₆ (40 mol%), and NaOAc (0.4 equiv) under argon in DCE at 80 °C (entry 9).

With the optimized conditions in hand, we first explored the reaction scope with a variety of aryl amides **6** with nitrostyrene **7a** (Table 2). Pyrrolidine benzamides bearing a series of *para* electron-donating and electron-withdrawing substituents all reacted with good efficiency with excellent enantioselectivity (**8aa–ea**, 51–82% yields, 93–94% ee). Arenes bearing a 3-methyl or 3-bromo group were also compatible (**8fa,ga**), and the C–H activation took place at the less hindered *ortho* C–H site in excellent enantioselectivity (90–95% ee). Heteroaromatic substrates such as furan-, thiophene-, and indole-derived amides afforded their corresponding adducts **8ha–ja** in reasonable yields with good to excellent enantioselectivity (42–64% yields, 82–92% ee). In addition to pyrrolidine benzamide, other amide directing groups such as piperidine benzamide and *N,N*-diethylbenzamide were also successful in this transformation

Table 2. Substrate Scope for Benzamides^a


Reaction scheme for Table 2: **6** + **7a** with **Rh-5** (5 mol%), AgSbF₆ (40 mol%), NaOAc (0.4 equiv) in DCE at 80 °C to form **8**.

1.5 equiv 0.2 mmol c = 0.2 mol/L

Chemical structures of **8aa** through **8ja** are shown.

8aa, R = H, 24 h, 75% yield, 93% ee
8ba, R = 4-OMe, 22 h, 82% yield, 93% ee
8ca, R = 4-Me, 24 h, 72% yield, 94% ee
8da, R = 4-Br, 30 h, 52% yield, 93% ee
8ea, R = 4-Cl, 30 h, 51% yield, 93% ee
8fa, R = 3-Me, 24 h, 63% yield, 95% ee
8ga, R = 3-Br, 30 h, 45% yield, 90% ee

8ha, X = O, 30 h, 42% yield, 87% ee
8ia, X = S, 22 h, 57% yield, 82% ee
8ka, 24 h, 58% yield, 96% ee


8la, 24 h, 51% yield, 87% ee
8ma, 24 h, 52% yield, 94% ee
8na, 32 h, 39% yield, 92% ee

^aReaction conditions: **6** (0.3 mmol), **7a** (0.2 mmol), **Rh-5** (5 mol%), AgSbF₆ (40 mol%), and additive in DCE (1.0 mL) at 80 °C under an argon atmosphere. Isolated yields of **8** are given. The ee values of **8** were determined by HPLC analysis on a chiral stationary phase.

(**8ka,la**, 51–58% yields, 87–96% ee). Notably, the *N*-methylbenzamide directing group was compatible with the reaction conditions, affording the product **8ma** in 52% yield with 94% ee. It is worth mentioning that an asymmetric reaction of such a substrate has not been realized previously. When the thiophene-derived amide **6n** was applied, product **8na** was obtained in 39% yield with 92% ee.

Next, the scope with respect to the nitroalkene was explored by reacting with benzamide **6b** as shown in Table 3. Aromatic rings bearing electron-donating or electron-withdrawing substituents were competent in this transformation, and products **8bb–bj** were obtained in moderate to good yields with excellent enantiocontrol (45–88% yields, 91–98% ee). When a 1-naphthyl-substituted nitroolefin was employed, product **8bk** was obtained in 86% yield with 95% ee. Furthermore, heteroaromatic substrates such as furan- and thiophene-derived nitroalkenes afforded products **8bl,bm** in acceptable yields with excellent enantioselectivity (70–74% yields, 90–93% ee). Notably, the reaction also worked well for alkyl nitroalkenes in moderate yields with good enantioselectivity (**8bn–bp**, 52–74% yields, 85–88% ee).

To demonstrate the practicality of this method, a 1.0 mmol scale reaction of **7a** was performed, and the desired product **8ba** was isolated in 79% yield with 93% ee (Scheme 2a). Moreover, 4-substituted dihydroisoquinolones **9ba,ia,bm** were obtained in good yields without erosion of enantioselectivity (Scheme 2b, 82–90% yields, 91–93% ee). Dihydroisoquinolone fragments are widely found in biologically active compounds and natural products.^{93,94} The access to the 4-substituted product isomer is

Table 3. Substrate Scope for Nitroolefins^a


^aReaction conditions: **6b** (0.3 mmol), **7** (0.2 mmol), **Rh-5** (5 mol%), **AgSbF₆** (40 mol%), and **NaOAc** (0.4 equiv) in **DCE** (1.0 mL) at 80 °C under an argon atmosphere. Isolated yields of **8** are given. The ee values of **8** were determined by HPLC analysis on a chiral stationary phase.

scarce by Rh(III)-catalyzed annulation of benzohydroxamic acid and styrene.^{95,96}

To shed light on the mechanism of this addition reaction, we carried out several experimental studies (Scheme 3). First, H/D exchange between **6a** and CD₃OD under the standard conditions in the presence or absence of nitrostyrene **7a** both gave significant deuteration at the *ortho* position of the benzene ring, suggesting the reversibility of the C–H cleavage. Two parallel reactions of **6a** and **6a-d₅** with **7a** were performed to determine the kinetic isotope effect, and $k_H/k_D = 1.0$ was observed at a low conversion under the standard conditions. Overall, these results suggest that C–H bond cleavage is not involved in the rate-determining step.


In conclusion, we have developed a series of novel chiral cyclopentadienyl (BCSCp) ligands based on a hexamethyl-1,1'-spirobiindane scaffold, which are synthesized efficiently from commercially available Bisphenol C. The corresponding rhodium complexes were successfully applied to the asymmetric C–H addition of aryl amides to nitroalkenes. Excellent enantioselectivity and good yields were obtained for a wide range of substrates, including those which previously failed. Products could be converted to chiral 4-substituted dihydroisoquinolones in high yields without loss of enantiopurity. The preliminary mechanistic studies support the idea that C–H

Scheme 2. (a) 1 mmol Scale Reaction and (b) Product Transformation




Scheme 3. Mechanistic Studies

H/D Exchange Experiments



KIE Experiments



bond cleavage is reversible and is not involved in the rate-determining step. Further applications of these BCSCpRh complexes in asymmetric catalysis are part of ongoing research in our laboratory.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acscatal.3c02199>.

Experimental procedures and compound characterization data (PDF)

X-ray crystallographic data for compound (S)-Rh-1 (CIF)

X-ray crystallographic data for compound 9ia (CIF)

AUTHOR INFORMATION

Corresponding Author

Shu-Li You – State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, People's Republic of China; orcid.org/0000-0003-4586-8359; Email: slyou@sioc.ac.cn

Authors

Hui Yang – State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, People's Republic of China

Ru Zhang – State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, People's Republic of China

Su-Zhen Zhang – State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, People's Republic of China

Qing Gu – State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, People's Republic of China; orcid.org/0000-0003-4963-2271

Complete contact information is available at:
<https://pubs.acs.org/10.1021/acscatal.3c02199>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank the National Key R&D Program of China (2021YFA1500100), the NSFC (21821002, 92256302, and 22071260), and Science and Technology Commission of Shanghai Municipality (21520780100) for generous financial support.

REFERENCES

- (1) Liu, C.-X.; Zhang, W.-W.; Yin, S.-Y.; Gu, Q.; You, S.-L. Synthesis of Atropisomers by Transition-Metal-Catalyzed Asymmetric C–H Functionalization Reactions. *J. Am. Chem. Soc.* **2021**, *143*, 14025–14040.
- (2) Li, Y.; Yuan, B.; Sun, Z.; Zhang, W. C–H bond functionalization reactions enabled by photobiocatalytic cascades. *Green Synth. Catal.* **2021**, *2*, 267–274.
- (3) Lam, N. Y. S.; Wu, K.; Yu, J.-Q. Advancing the Logic of Chemical Synthesis: C–H Activation as Strategic and Tactical Disconnections for C–C Bond Construction. *Angew. Chem., Int. Ed.* **2021**, *60*, 15767–15790.
- (4) Liu, C.-X.; Gu, Q.; You, S.-L. Asymmetric C–H Bond Functionalization of Ferrocenes: New Opportunities and Challenges. *Trends. Chem.* **2020**, *2*, 737–749.
- (5) Liao, G.; Zhang, T.; Lin, Z.-K.; Shi, B.-F. Transition Metal-Catalyzed Enantioselective C–H Functionalization via Chiral Transient Directing Group Strategies. *Angew. Chem., Int. Ed.* **2020**, *59*, 19773–19786.
- (6) Achar, T. K.; Maiti, S.; Jana, S.; Maiti, D. Transition Metal Catalyzed Enantioselective C(sp²)–H Bond Functionalization. *ACS Catal.* **2020**, *10*, 13748–13793.
- (7) Woźniak, Ł.; Cramer, N. Enantioselective C–H Bond Functionalizations by 3d Transition-Metal Catalysts. *Trends. Chem.* **2019**, *1*, 471–484.
- (8) Loup, J.; Dhawa, U.; Pesciaoli, F.; Wencel-Delord, J.; Ackermann, L. Enantioselective C–H Activation with Earth-Abundant 3d Transition Metals. *Angew. Chem., Int. Ed.* **2019**, *58*, 12803–12818.
- (9) Saint-Denis, T. G.; Zhu, R.-Y.; Chen, G.; Wu, Q.-F.; Yu, J.-Q. Enantioselective C(sp³)–H bond activation by chiral transition metal catalysts. *Science* **2018**, *359*, eaao4798.
- (10) Newton, C. G.; Wang, S.-G.; Oliveira, C. C.; Cramer, N. Catalytic Enantioselective Transformations Involving C–H Bond Cleavage by Transition-Metal Complexes. *Chem. Rev.* **2017**, *117*, 8908–8976.
- (11) Yang, H.; Gou, B.-B.; Dai, H.-T.; Gu, Q.; You, S.-L. Advances in the synthesis of chiral cyclopentadienyl ligands and metal complexes. *Sci. Sin. Chim.* **2023**, *53*, 359–374.
- (12) Davies, C.; Shaaban, S.; Waldmann, H. Asymmetric catalysis with chiral cyclopentadienyl complexes to access privileged scaffolds. *Trends. Chem.* **2022**, *4*, 318–330.
- (13) Yoshino, T.; Matsunaga, S. Chiral Carboxylic Acid Assisted Enantioselective C–H Activation with Achiral Cp[∗]M^{III} (M = Co, Rh, Ir) Catalysts. *ACS Catal.* **2021**, *11*, 6455–6466.
- (14) Wang, Q.; Liu, C.-X.; Gu, Q.; You, S.-L. Chiral Cp[∗]Rh complexes for C–H functionalization reactions. *Sci. Bull.* **2021**, *66*, 210–213.
- (15) Pan, C.; Yin, S.-Y.; Gu, Q.; You, S.-L. Cp[∗]M(III)-catalyzed enantioselective C–H functionalization through migratory insertion of metal–carbenes/nitrenes. *Org. Biomol. Chem.* **2021**, *19*, 7264–7275.
- (16) Mas-Roselló, J.; Herraiz, A. G.; Audic, B.; Laverny, A.; Cramer, N. Chiral Cyclopentadienyl Ligands: Design, Syntheses, and Applications in Asymmetric Catalysis. *Angew. Chem., Int. Ed.* **2021**, *60*, 13198–13224.
- (17) Yoshino, T.; Satake, S.; Matsunaga, S. Diverse Approaches for Enantioselective C–H Functionalization Reactions Using Group 9 Cp[∗]M^{III} Catalysts. *Chem. Eur. J.* **2020**, *26*, 7346–7357.
- (18) Shaaban, S.; Davies, C.; Waldmann, H. Applications of Chiral Cyclopentadienyl (Cp[∗]) Metal Complexes in Asymmetric Catalysis. *Eur. J. Org. Chem.* **2020**, *2020*, 6512–6524.
- (19) Newton, C. G.; Kossler, D.; Cramer, N. Asymmetric Catalysis Powered by Chiral Cyclopentadienyl Ligands. *J. Am. Chem. Soc.* **2016**, *138*, 3935–3941.
- (20) Ye, B.; Cramer, N. Chiral Cyclopentadienyls: Enabling Ligands for Asymmetric Rh(III)-Catalyzed C–H Functionalizations. *Acc. Chem. Res.* **2015**, *48*, 1308–1318.
- (21) Ye, B.; Cramer, N. Chiral Cyclopentadienyl Ligands as Stereocontrolling Element in Asymmetric C–H Functionalization. *Science* **2012**, *338*, 504–506.
- (22) Hyster, T. K.; Knörr, L.; Ward, T. R.; Rovis, T. Biotinylated Rh(III) Complexes in Engineered Streptavidin for Accelerated Asymmetric C–H Activation. *Science* **2012**, *338*, 500–503.
- (23) Pan, C.; Yin, S.-Y.; Wang, S.-B.; Gu, Q.; You, S.-L. Oxygen-Linked Cyclopentadienyl Rhodium(III) Complexes-Catalyzed Asymmetric C–H Arylation of Benzo[h]quinolines with 1-Diazonaphthoquinones. *Angew. Chem., Int. Ed.* **2021**, *60*, 15510–15516.
- (24) Wang, S.-G.; Cramer, N. Asymmetric Cp[∗]Rh(III)-Catalyzed Acrylic Acid C–H Functionalization with Allenes Provides Chiral γ -Lactones. *ACS Catal.* **2020**, *10*, 8231–8236.
- (25) Cui, W.-J.; Wu, Z.-J.; Gu, Q.; You, S.-L. Divergent Synthesis of Tunable Cyclopentadienyl Ligands and Their Application in Rh-Catalyzed Enantioselective Synthesis of Isoindolinone. *J. Am. Chem. Soc.* **2020**, *142*, 7379–7385.
- (26) Duchemin, C.; Smits, G.; Cramer, N. Rh^I, Ir^{III}, and Co^{III} Complexes with Atropchiral Biaryl Cyclopentadienyl Ligands: Syntheses, Structures, and Catalytic Activities. *Organometallics* **2019**, *38*, 3939–3947.
- (27) Sun, Y.; Cramer, N. Tailored trisubstituted chiral Cp[∗]Rh^{III} catalysts for kinetic resolutions of phosphinic amides. *Chem. Sci.* **2018**, *9*, 2981–2985.
- (28) Teng, H.-L.; Luo, Y.; Wang, B.; Zhang, L.; Nishiura, M.; Hou, Z. Synthesis of Chiral Aminocyclopropanes by Rare-Earth-Metal-

- Catalyzed Cyclopropene Hydroamination. *Angew. Chem., Int. Ed.* **2016**, *55*, 15406–15410.
- (29) Reddy Chidipudi, S.; Burns, D. J.; Khan, I.; Lam, H. W. Enantioselective Synthesis of Spiroindenes by Enol-Directed Rhodium-(III)-Catalyzed C–H Functionalization and Spiroannulation. *Angew. Chem., Int. Ed.* **2015**, *54*, 13975–13979.
- (30) Ye, B.; Cramer, N. A Tunable Class of Chiral Cp Ligands for Enantioselective Rhodium(III)-Catalyzed C–H Allylations of Benzamides. *J. Am. Chem. Soc.* **2013**, *135*, 636–639.
- (31) Jia, Z.-J.; Merten, C.; Gontla, R.; Daniliuc, C. G.; Antonchick, A. P.; Waldmann, H. General Enantioselective C–H Activation with Efficiently Tunable Cyclopentadienyl Ligands. *Angew. Chem., Int. Ed.* **2017**, *56*, 2429–2434.
- (32) Wang, S.-G.; Park, S. H.; Cramer, N. A Readily Accessible Class of Chiral Cp Ligands and their Application in Ru^{II}-Catalyzed Enantioselective Syntheses of Dihydrobenzoindoles. *Angew. Chem., Int. Ed.* **2018**, *57*, 5459–5462.
- (33) Li, G.; Yan, X.; Jiang, J.; Liang, H.; Zhou, C.; Wang, J. Chiral Bicyclo[2.2.2]octane-Fused CpRh Complexes: Synthesis and Potential Use in Asymmetric C–H Activation. *Angew. Chem., Int. Ed.* **2020**, *59*, 22436–22440.
- (34) Yan, X.; Jiang, J.; Wang, J. A Class of Readily Tunable Planar-Chiral Cyclopentadienyl Rhodium(III) Catalysts for Asymmetric C–H Activation. *Angew. Chem., Int. Ed.* **2022**, *61*, e202201522.
- (35) Kolos, A. V.; Nelyubina, Y. V.; Sundararaju, B.; Perekalin, D. S. Synthesis of Overloaded Cyclopentadienyl Rhodium(III) Complexes via Cyclotetramerization of tert-Butylacetylene. *Organometallics* **2021**, *40*, 3712–3719.
- (36) Farr, C. M. B.; Kazerouni, A. M.; Park, B.; Poff, C. D.; Won, J.; Sharp, K. R.; Baik, M.-H.; Blakey, S. B. Designing a Planar Chiral Rhodium Indenyl Catalyst for Regio- and Enantioselective Allylic C–H Amidation. *J. Am. Chem. Soc.* **2020**, *142*, 13996–14004.
- (37) Trifonova, E. A.; Ankudinov, N. M.; Mikhaylov, A. A.; Chusov, D. A.; Nelyubina, Y. V.; Perekalin, D. S. A Planar-Chiral Rhodium(III) Catalyst with a Sterically Demanding Cyclopentadienyl Ligand and Its Application in the Enantioselective Synthesis of Dihydroisquinolones. *Angew. Chem., Int. Ed.* **2018**, *57*, 7714–7718.
- (38) Liang, H.; Vasamsetty, L.; Li, T.; Jiang, J.; Pang, X.; Wang, J. A New Class of C₂-Symmetric Chiral Cyclopentadienyl Ligand Derived from Ferrocene Scaffold: Design, Synthesis and Application. *Chem. Eur. J.* **2020**, *26*, 14546–14550.
- (39) Pototskiy, R. A.; Kolos, A. V.; Nelyubina, Y. V.; Perekalin, D. S. Rhodium Catalysts with a Chiral Cyclopentadienyl Ligand Derived from Natural R-Myrtenal. *Eur. J. Org. Chem.* **2020**, *2020*, 6019–6025.
- (40) Xie, J.-H.; Zhou, Q.-L. Magical Chiral Spiro Ligands. *Acta Chim. Sinica* **2014**, *72*, 778–797.
- (41) Xie, J.-H.; Zhou, Q.-L. Chiral Diphosphine and Monodentate Phosphorus Ligands on a Spiro Scaffold for Transition-Metal-Catalyzed Asymmetric Reactions. *Acc. Chem. Res.* **2008**, *41*, 581–593.
- (42) Liu, B.; Zhu, S.-F.; Wang, L.-X.; Zhou, Q.-L. Preparation and application of bisoxazoline ligands with a chiral spirobiindane skeleton for asymmetric cyclopropanation and allylic oxidation. *Tetrahedron: Asymmetry* **2006**, *17*, 634–641.
- (43) Zhu, S.-F.; Yang, Y.; Wang, L.-X.; Liu, B.; Zhou, Q.-L. Synthesis and Application of Chiral Spiro Phospholane Ligand in Pd-Catalyzed Asymmetric Allylation of Aldehydes with Allylic Alcohols. *Org. Lett.* **2005**, *7*, 2333–2335.
- (44) Xie, J.-H.; Wang, L.-X.; Fu, Y.; Zhu, S.-F.; Fan, B.-M.; Duan, H.-F.; Zhou, Q.-L. Synthesis of Spiro Diphosphines and Their Application in Asymmetric Hydrogenation of Ketones. *J. Am. Chem. Soc.* **2003**, *125*, 4404–4405.
- (45) Hu, A.-G.; Fu, Y.; Xie, J.-H.; Zhou, H.; Wang, L.-X.; Zhou, Q.-L. Monodentate Chiral Spiro Phosphoramidites: Efficient Ligands for Rhodium-Catalyzed Enantioselective Hydrogenation of Enamides. *Angew. Chem., Int. Ed.* **2002**, *41*, 2348–2350.
- (46) Fu, Y.; Xie, J.-H.; Hu, A.-G.; Zhou, H.; Wang, L.-X.; Zhou, Q.-L. Novel monodentate spiro phosphorus ligands for rhodium-catalyzed hydrogenation reactions. *Chem. Commun.* **2002**, 480–481.
- (47) Zheng, J.; Cui, W.-J.; Zheng, C.; You, S.-L. Synthesis and Application of Chiral Spiro Cp Ligands in Rhodium-Catalyzed Asymmetric Oxidative Coupling of Biaryl Compounds with Alkenes. *J. Am. Chem. Soc.* **2016**, *138*, S242–S245.
- (48) Zou, Y.; Wang, P.; Kong, L.; Li, X. Rhodium-Catalyzed Atroposelective C–H Arylation of (Hetero)Arenes Using Carbene Precursors as Arylating Reagents. *Org. Lett.* **2022**, *24*, 3189–3193.
- (49) Zhang, W.-W.; Wang, Q.; Zhang, S.-Z.; Zheng, C.; You, S.-L. (SCp)Rhodium-Catalyzed Asymmetric Satoh–Miura Reaction for Building-up Axial Chirality: Counteranion-Directed Switching of Reaction Pathways. *Angew. Chem., Int. Ed.* **2023**, *62*, e202214460.
- (50) Zhang, W.-W.; Liu, C.-X.; Yang, P.; Zhang, S.-Z.; Gu, Q.; You, S.-L. Rhodium-Catalyzed Atroposelective C–H/C–H Cross-Coupling Reaction between 1-Aryl Isoquinoline Derivatives and Indolizines. *Org. Lett.* **2022**, *24*, 564–569.
- (51) Yin, S.-Y.; Pan, C.; Zhang, W.-W.; Liu, C.-X.; Zhao, F.; Gu, Q.; You, S.-L. SCpRh(III)-Catalyzed Enantioselective Synthesis of Atropisomers by C2-Arylation of Indoles with 1-Diazonaphthoquinones. *Org. Lett.* **2022**, *24*, 3620–3625.
- (52) Yang, C.; Shi, L.; Wang, F.; Su, Y.; Xia, J.-B.; Li, F. Rhodium-Catalyzed Asymmetric (3 + 2 + 2) Annulation via N–H/C–H Dual Activation and Internal Alkyne Insertion toward N-Fused 5/7 Bicycles. *ACS Catal.* **2022**, *12*, 14194–14208.
- (53) Wu, Z.-J.; Zhang, R.; Gu, Q.; You, S.-L. SCpRh(III)-Catalyzed Enantioselective Aryl C–H Addition to Nitroalkenes. *Asian J. Org. Chem.* **2021**, *10*, 1722–1725.
- (54) Wang, Q.; Zhang, W.-W.; Zheng, C.; Gu, Q.; You, S.-L. Enantioselective Synthesis of Azoniahelices by Rh-Catalyzed C–H Annulation with Alkynes. *J. Am. Chem. Soc.* **2021**, *143*, 114–120.
- (55) Wang, F.; Jing, J.; Zhao, Y.; Zhu, X.; Zhang, X.-P.; Zhao, L.; Hu, P.; Deng, W.-Q.; Li, X. Rhodium-Catalyzed C–H Activation-Based Construction of Axially and Centrally Chiral Indenes through Two Discrete Insertions. *Angew. Chem., Int. Ed.* **2021**, *60*, 16628–16633.
- (56) Yan, X.; Zhao, P.; Liang, H.; Xie, H.; Jiang, J.; Gou, S.; Wang, J. Rhodium(III)-Catalyzed Asymmetric C–H Activation of N-Methoxybenzamide with Quinone and Its Application in the Asymmetric Synthesis of a Dihydrolycoricidine Analogue. *Org. Lett.* **2020**, *22*, 3219–3223.
- (57) Wang, Q.; Zhang, W.-W.; Song, H.; Wang, J.; Zheng, C.; Gu, Q.; You, S.-L. Rhodium-Catalyzed Atroposelective Oxidative C–H/C–H Cross-Coupling Reaction of 1-Aryl Isoquinoline Derivatives with Electron-Rich Heteroarenes. *J. Am. Chem. Soc.* **2020**, *142*, 15678–15685.
- (58) Li, H.; Yan, X.; Zhang, J.; Guo, W.; Jiang, J.; Wang, J. Enantioselective Synthesis of C–N Axially Chiral N-Aryloxindoles by Asymmetric Rhodium-Catalyzed Dual C–H Activation. *Angew. Chem., Int. Ed.* **2019**, *58*, 6732–6736.
- (59) Li, T.; Zhou, C.; Yan, X.; Wang, J. Solvent-Dependent Asymmetric Synthesis of Alkynyl and Monofluoroalkenyl Isoindolinones by CpRh^{III}-Catalyzed C–H Activation. *Angew. Chem., Int. Ed.* **2018**, *57*, 4048–4052.
- (60) Zheng, J.; Wang, S.-B.; Zheng, C.; You, S.-L. Asymmetric Synthesis of Spiropyrazolones by Rhodium-Catalyzed C(sp²)-H Functionalization/Annulation Reactions. *Angew. Chem., Int. Ed.* **2017**, *56*, 4540–4544.
- (61) Hummel, J. R.; Boerth, J. A.; Ellman, J. A. Transition-Metal-Catalyzed C–H Bond Addition to Carbonyls, Imines, and Related Polarized π Bonds. *Chem. Rev.* **2017**, *117*, 9163–9227.
- (62) Dong, Z.; Ren, Z.; Thompson, S. J.; Xu, Y.; Dong, G. Transition-Metal-Catalyzed C–H Alkylation Using Alkenes. *Chem. Rev.* **2017**, *117*, 9333–9403.
- (63) Shi, X.-Y.; Han, W.-J.; Li, C.-J. Transition-Metal-Catalyzed Direct Addition of Aryl C–H Bonds to Unsaturated Electrophiles. *Chem. Rev.* **2016**, *16*, 1178–1190.
- (64) Yang, L.; Huang, H. Transition-Metal-Catalyzed Direct Addition of Unactivated C–H Bonds to Polar Unsaturated Bonds. *Chem. Rev.* **2015**, *115*, 3468–3517.
- (65) Li, C.-J. The Development of Catalytic Nucleophilic Additions of Terminal Alkynes in Water. *Acc. Chem. Res.* **2010**, *43*, 581–590.

- (66) Colby, D. A.; Bergman, R. G.; Ellman, J. A. Rhodium-Catalyzed C–C Bond Formation via Heteroatom-Directed C–H Bond Activation. *Chem. Rev.* **2010**, *110*, 624–655.
- (67) Trost, B. M.; Weiss, A. H. The Enantioselective Addition of Alkyne Nucleophiles to Carbonyl Groups. *Adv. Synth. Catal.* **2009**, *351*, 963–983.
- (68) Ritleng, V.; Sirlin, C.; Pfeffer, M. Ru-, Rh-, and Pd-Catalyzed C–C Bond Formation Involving C–H Activation and Addition on Unsaturated Substrates: Reactions and Mechanistic Aspects. *Chem. Rev.* **2002**, *102*, 1731–1770.
- (69) Shibata, T.; Kurita, H.; Onoda, S.; Kanyiva, K. S. Ir-Catalyzed Enantioselective Intra- and Intermolecular Formal C–H Conjugate Addition to β -Substituted α,β -Unsaturated Esters. *Asian J. Org. Chem.* **2018**, *7*, 1411–1418.
- (70) Shibata, T.; Michino, M.; Kurita, H.; Tahara, Y.-k.; Kanyiva, K. S. Enantioselective Formal C–H Conjugate Addition of Acetanilides to β -Substituted Acrylates by Chiral Iridium Catalysts. *Chem. Eur. J.* **2017**, *23*, 88–91.
- (71) Shibata, T.; Shizuno, T. Iridium-Catalyzed Enantioselective C–H Alkylation of Ferrocenes with Alkenes Using Chiral Diene Ligands. *Angew. Chem., Int. Ed.* **2014**, *53*, 5410–5413.
- (72) Tsuchikama, K.; Hashimoto, Y.-k.; Endo, K.; Shibata, T. Iridium-Catalyzed Selective Synthesis of 4-Substituted Benzofurans and Indoles via Directed Cyclodehydration. *Adv. Synth. Catal.* **2009**, *351*, 2850–2854.
- (73) Ye, B.; Cramer, N. Chiral Cyclopentadienyl Ligands Enable a Rhodium(III)-Catalyzed Enantioselective Access to Hydroxychromanes and Phthalides. *Synlett* **2015**, *26*, 1490–1495.
- (74) Kurihara, T.; Kojima, M.; Yoshino, T.; Matsunaga, S. Achiral Cp^{*}Rh(III)/Chiral Lewis Base Cooperative Catalysis for Enantioselective Cyclization via C–H Activation. *J. Am. Chem. Soc.* **2022**, *144*, 7058–7065.
- (75) Kurihara, T.; Kojima, M.; Yoshino, T.; Matsunaga, S. Cp^{*}Co(III)/Chiral Carboxylic Acid-Catalyzed Enantioselective 1,4-Addition Reactions of Indoles to Maleimides. *Asian J. Org. Chem.* **2020**, *9*, 368–371.
- (76) Huang, L.-T.; Fukagawa, S.; Kojima, M.; Yoshino, T.; Matsunaga, S. Rhodium(III)/Chiral Carboxylic Acid Catalyzed Enantioselective C(sp³)–H Alkylation of 8-Ethylquinolines with α,β -Unsaturated Carbonyl Compounds. *Org. Lett.* **2020**, *22*, 8256–8260.
- (77) Satake, S.; Kurihara, T.; Nishikawa, K.; Mochizuki, T.; Hatano, M.; Ishihara, K.; Yoshino, T.; Matsunaga, S. Pentamethylcyclopentadienyl rhodium(III)–chiral disulfonate hybrid catalysis for enantioselective C–H bond functionalization. *Nat. Catal.* **2018**, *1*, 585–591.
- (78) Wu, Z.-J.; Wu, Z.; Zhang, W.-W.; Gu, Q.; You, S.-L. Rh(III)-Catalyzed Enantioselective Intermolecular Aryl C–H Bond Addition to Aldehydes. *Chin. J. Chem.* **2022**, *40*, 2780–2784.
- (79) Wang, J.; Chen, H.; Kong, L.; Wang, F.; Lan, Y.; Li, X. Enantioselective and Diastereoselective C–H Alkylation of Benzamides: Synergized Axial and Central Chirality via a Single Stereodetermining Step. *ACS Catal.* **2021**, *11*, 9151–9158.
- (80) Potter, T. J.; Kamber, D. N.; Mercado, B. Q.; Ellman, J. A. Rh(III)-Catalyzed Aryl and Alkenyl C–H Bond Addition to Diverse Nitroalkenes. *ACS Catal.* **2017**, *7*, 150–153.
- (81) Shirai, T.; Yamamoto, Y. Scope and Mechanistic Studies of the Cationic Ir/Me-BIPAM-Catalyzed Asymmetric Intramolecular Direct Hydroarylation Reaction. *Organometallics* **2015**, *34*, 3459–3463.
- (82) Filloux, C. M.; Rovis, T. Rh(I)–Bisphosphine-Catalyzed Asymmetric, Intermolecular Hydroheteroarylation of α -Substituted Acrylate Derivatives. *J. Am. Chem. Soc.* **2015**, *137*, 508–517.
- (83) Shirai, T.; Ito, H.; Yamamoto, Y. Cationic Ir/Me-BIPAM-Catalyzed Asymmetric Intramolecular Direct Hydroarylation of α -Ketoamides. *Angew. Chem., Int. Ed.* **2014**, *53*, 2658–2661.
- (84) Wang, Q.; Nie, Y.-H.; Liu, C.-X.; Zhang, W.-W.; Wu, Z.-J.; Gu, Q.; Zheng, C.; You, S.-L. Rhodium(III)-Catalyzed Enantioselective C–H Activation/Annulation of Ferrocenecarboxamides with Internal Alkynes. *ACS Catal.* **2022**, *12*, 3083–3093.
- (85) Mi, R.; Chen, H.; Zhou, X.; Li, N.; Ji, D.; Wang, F.; Lan, Y.; Li, X. Rhodium-Catalyzed Atroposelective Access to Axially Chiral Olefins via C–H Bond Activation and Directing Group Migration. *Angew. Chem., Int. Ed.* **2022**, *61*, e202111860.
- (86) Huang, Y.-Q.; Wu, Z.-J.; Zhu, L.; Gu, Q.; Lu, X.; You, S.-L.; Mei, T.-S. Electrochemical Rhodium-Catalyzed Enantioselective C–H Annulation with Alkynes. *CCS Chem.* **2022**, *4*, 3181–3189.
- (87) Hu, P.; Liu, B.; Wang, F.; Mi, R.; Li, X.-X.; Li, X. A Stereodivergent–Convergent Chiral Induction Mode in Atroposelective Access to Biaryls via Rhodium-Catalyzed C–H Bond Activation. *ACS Catal.* **2022**, *12*, 13884–13896.
- (88) Hu, P.; Kong, L.; Wang, F.; Zhu, X.; Li, X. Twofold C–H Activation-Based Enantio- and Diastereoselective C–H Arylation Using Diarylacetylenes as Rare Arylating Reagents. *Angew. Chem., Int. Ed.* **2021**, *60*, 20424–20429.
- (89) Li, H.; Gontla, R.; Flegel, J.; Merten, C.; Ziegler, S.; Antonchick, A. P.; Waldmann, H. Enantioselective Formal C(sp³)–H Bond Activation in the Synthesis of Bioactive Spiropyrazolone Derivatives. *Angew. Chem., Int. Ed.* **2019**, *58*, 307–311.
- (90) Shan, G.; Flegel, J.; Li, H.; Merten, C.; Ziegler, S.; Antonchick, A. P.; Waldmann, H. C–H Bond Activation for the Synthesis of Heterocyclic Atropisomers Yields Hedgehog Pathway Inhibitors. *Angew. Chem., Int. Ed.* **2018**, *57*, 14250–14254.
- (91) Guo, W.; Pang, X.; Jiang, J.; Wang, J. Chiral 3,3,3',3'-Tetramethyl-1,1'-Spirobiindanyl Cyclopentadienyl (TMSCp) Ligands: Design, Synthesis, and Applications. *Org. Lett.* **2023**, *25*, 3823–3828.
- (92) Gu, H.; Han, Z.; Xie, H.; Lin, X. Iron-Catalyzed Enantioselective Si–H Bond Insertions. *Org. Lett.* **2018**, *20*, 6544–6549.
- (93) Zhang, L.; Wang, C.; Han, J.; Huang, Z.-B.; Zhao, Y. Palladium-Catalyzed Carbonylation of β -Arylethylamide Directed by Oxalyl Amide in the Presence of Carbon Monoxide. *J. Org. Chem.* **2016**, *81*, 5256–5262.
- (94) Murray, C. W.; Rees, D. C. Opportunity Knocks: Organic Chemistry for Fragment-Based Drug Discovery (FBDD). *Angew. Chem., Int. Ed.* **2016**, *55*, 488–492.
- (95) Hyster, T. K.; Dalton, D. M.; Rovis, T. Ligand design for Rh(III)-catalyzed C–H activation: an unsymmetrical cyclopentadienyl group enables a regioselective synthesis of dihydroisoquinolones. *Chem. Sci.* **2015**, *6*, 254–258.
- (96) Rakshit, S.; Grohmann, C.; Besset, T.; Glorius, F. Rh(III)-Catalyzed Directed C–H Olefination Using an Oxidizing Directing Group: Mild, Efficient, and Versatile. *J. Am. Chem. Soc.* **2011**, *133*, 2350–2353.