

Palladium-Catalyzed Cross-Coupling, Divergent Allene Generation, and Cycloadditions toward Cyclobuta[*b*]naphthalen-3(1*H*)-ones and 11*H*-Benzo[*b*]fluoren-11-onesShugao Zhu,* Dan Wang,[†] Shihan Liu,[†] Yufeng Ma,[†] Hong Wang, Yu Lan,* Ruopeng Bai, and Ruwei Shen*Cite This: *ACS Catal.* 2023, 13, 8402–8412


Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information



ABSTRACT: The Pd-catalyzed reaction of 1-(2-iodophenyl)-3-arylprop-2-yn-1-ones with propargyl sulfonamide produces cyclobuta[*b*]naphthalen-3(1*H*)-ones, while the reaction of 1-(2-iodophenyl)-3-arylprop-2-yn-1-ones with propargyl ethers under the similar palladium catalysis affords 11*H*-benzo[*b*]fluoren-11-ones as products. Combined experimental and theoretical studies on the reaction mechanism reveal that the former reaction proceeds through the Pd-catalyzed cross-coupling, propargyl-allenyl isomerization (1,3-*H* transfer) and [2 + 2] cycloaddition, whereas the latter involves a process of Pd-catalyzed cross-coupling, a propargylic Alder-ene-type reaction (1,8-*H* transfer) and 6*π*-electrocyclization leading to the formation of the formal [4 + 2] cycloadducts. The divergent formation of two types of allenic intermediates depending on propargylic substrates is confirmed and elucidated.

KEYWORDS: tandem reaction, propargyl-allenyl isomerization, Alder-ene reaction, allene, cycloaddition

INTRODUCTION

Allenes have attracted numerous attention in synthetic organic chemistry in recent years due to their rich properties and versatile reactivities.^{1–7} In particular, the relatively high-energy profile and unique geometry of allenes arising from the cumulated diene structural moieties render them valuable and versatile synthetic intermediates for the synthesis of complex carbo- and heterocyclic backbones via diversified cyclization reactions.^{8–17} Moreover, allenes are also frequently engaged as reactive intermediates in reaction design for rapid construction of cyclic frameworks that usually cannot be easily obtained from traditional methods.^{18–23} Based on a facile and base-promoted propargyl-allenyl isomerization reactions (CIRs, Scheme 1a), for example, Müller and we have developed a series of tandem transformations of the in situ generated vinyl-allene intermediates **I**, such as Diels-Alder reaction,^{24–28} Alder-ene reaction,²⁹ Claisen rearrangement,^{30–32} [3 + 2] azide-olefin cycloaddition,³³ and Schmitt cyclization,³⁴ to afford a

diversity of useful skeletons.^{35–39} As our further endeavor, we recently postulated (1) the possible formation of an aryl-tethered heteroatom-substituted allene-ynone intermediate **II** from the CIR of an *ortho*-iodophenyl-ynone and a propargylic substrate and questioned (2) the reactivity of such a functionalized allenic intermediate for cyclization reactions (initial design, Scheme 1b). It is worth noting that previous studies on cyclization reactions of relevant yne-allenes have proven extreme success in the synthesis of a variety of cyclic skeletons via cycloadditions,^{40–42} cycloisomerizations,⁴³ and other annulations⁴⁴ upon thermal or photochemical initiation


Received: February 23, 2023

Revised: April 24, 2023

Published: June 9, 2023




Scheme 1. Reaction Design Based on the Generation of the Heteroatom-Substituted Allene Intermediates

Table 1. Selected Results on Optimization for the Diverged Cyclization Reactions^{a,j}

entry	1a/2a	catalyst	base	solvent	4a, yield (%) ^b	entry	1a/3a	catalyst	base	solvent	5a, yield (%) ^b
1	1/1.2	Pd(PPh ₃) ₂ Cl ₂	DIPA	toluene	40	15	1.2/1	Pd(PPh ₃) ₂ Cl ₂	DIPA	toluene	50
2	1/1.2	Pd ₂ (dba) ₃	DIPA	toluene	39	16	1.2/1	Pd ₂ (dba) ₃	DIPA	toluene	45
3	1/1.2	Pd(PPh ₃) ₄	DIPA	toluene	58	17	1.2/1	Pd(PPh ₃) ₄	DIPA	toluene	60
4	1/1.2	Pd(OAc) ₂	DIPA	toluene	35	18	1.2/1	Pd(OAc) ₂	DIPA	toluene	48
5	1/1.2	Pd(PPh ₃) ₄	Et ₃ N	toluene	39	19	1.2/1	Pd(PPh ₃) ₄	Et ₃ N	toluene	20
6	1/1.2	Pd(PPh ₃) ₄	pyridine	toluene	0	20	1.2/1	Pd(PPh ₃) ₄	pyridine	toluene	0
7	1/1.2	Pd(PPh ₃) ₄	K ₂ CO ₃ ^c	toluene	0	21	1.2/1	Pd(PPh ₃) ₄	DIPA	THF	17
8	1.2/1	Pd(PPh ₃) ₄	DIPA	toluene	50	22	1.2/1	Pd(PPh ₃) ₄	DIPA	MeCN	37
9	1.5/1	Pd(PPh ₃) ₄	DIPA	toluene	53	23	1.2/1	Pd(PPh ₃) ₄	DIPA	DMF	14
10	2/1	Pd(PPh ₃) ₄	DIPA ^d	toluene	60	24	1.2/1	Pd(PPh ₃) ₄	DIPA	DCM	28
11	2/1	Pd(PPh₃)₄	DIPA^e	toluene	71	25	1/1	Pd(PPh ₃) ₄	DIPA	toluene	58
12	4/1	Pd(PPh ₃) ₄	DIPA	toluene	68	26	1/1.2	Pd(PPh₃)₄	DIPA	toluene	73
13	2/1	Pd(PPh ₃) ₄	DIPA ^f	toluene	65	27	1/1.2	Pd(PPh ₃) ₄	DIPA	toluene	68 ^h
14	2/1	Pd(PPh ₃) ₄	DIPA ^g	toluene	67	28	1/1.2	Pd(PPh ₃) ₄	DIPA	toluene	70 ⁱ

^aMixture of 1a (0.25–0.5 mmol), 2a or 3a (0.25–0.30 mmol), the Pd catalyst (5% mol), CuI (10% mol), toluene (2 mL), and the base was charged to a Schlenk tube under N₂ and heated at 60 °C. ^bIsolated yield. ^c3 equiv of K₂CO₃ was used. ^d1.5 mL. ^e0.1 mL. ^f0.5 mL. ^g1 mL. ^h80 °C. ⁱ100 °C. ^jThe boldface values for entries 11 and 26 highlight the best reaction conditions.

Table 2. Reaction Scope of the Coupling-Isomerization [2 + 2] Cycloaddition and Formal [4 + 2] Cycloaddition Sequence^a

^aReactions were performed on a 0.25 mmol scale. Isolated yields are given. For experimental details, see the SI.

and Lewis acid or transition metal catalysis.^{45–59} However, the reactivity on aryl-tethered ynone-allenes has been less explored, and their synthetic utilities remain to be exploited.⁶⁰


During the course of this project, we discovered that the cross-coupling intermediates from *ortho*-iodophenyl-ynones and propargylic substrates undergo divergent isomerizations to generate two types of allene species, i.e., ynone-allene **II–1** via propargyl-allenyl isomerization (involving 1,3-hydrogen transfer) and enone-allene **III** via propargylic Alder ene-type^{61–63} (involving 1,8-hydrogen transfer), depending on heteroatoms at the propargyl carbon (Scheme 1b). The two different types of allenic intermediates then fulfill formal [2 + 2] and [4 + 2] cycloadditions with high selectivity to provide polycyclic compounds of cyclobuta[*b*]naphthalen-3(1*H*)-one

and 11*H*-benzo[*b*]fluoren-11-one backbones, respectively (this finding, Scheme 1b). Here, we report the details on the development of these types of palladium-catalyzed tandem reactions involving divergent formation of allene intermediates with consecutive cyclizations, as well as a combined experimental and theoretical study to illustrate the reaction mechanism and the origin of the observed selectivity.

RESULTS AND DISCUSSION

Reaction Discovery and Optimization. We initially explored the reactions of 1-(2-iodophenyl)-3-phenylprop-2-yn-1-one (**1a**), with propargyl tosylamide **2a** and propargyl benzyl ether **3a** under the palladium catalysis in the presence of an excessive base, which was expected to facilitate the cross-

Scheme 2. Control Experiments




coupling reaction and promote the propargyl-allenyl isomerization. The preliminary experiments showed that **1a** coupled with **2a** in the presence of 5 mol % Pd(PPh₃)₂Cl₂, 10 mol % CuI and diisopropylamine (DIPA) to form the alkynylation intermediate, which was finally converted to a cyclobuta[*b*]-naphthalen-3(1*H*)-one product **4a** in 40% yield as the product (Table 1, entry 1). In contrast, the reaction of **1a** and **3a** under similar reaction conditions gave rise to a skeletally different product, i.e., 5-benzoyl-11*H*-benzo[*b*]fluoren-11-one (**5a**) in 50% yield (Table 1, entry 15). Encouraged by these intriguing results, we next performed a series of optimization experiments by varying reaction parameters including the palladium catalyst, base, solvent, reaction temperature, and molar ratios of the substrates to improve the product yields (for detailed results, see Tables S1 and S2 in the Supporting Information). It was found that the combination of a catalytic amount of Pd(PPh₃)₄ and CuI with an excess DIPA as the base in toluene was the best system for both reactions. The optimal conditions were obtained as follows: the reaction of **1a** and **2a** (2/1 molar ratio) with 5 mol % Pd(PPh₃)₄ and 10 mol % CuI in toluene/DIPA at 60 °C for 6 h gave **4a** in 71% yield (Table 1, entry 11); and the reaction of **1a** and **3a** (1/1.2 molar ratio) with 5

mol % Pd(PPh₃)₄ and 10 mol % CuI in toluene/DIPA at 60 °C for 3 h produced **5a** in 73% yield (Table 1, entry 26).

Reaction Scope. With the optimal conditions in hand, the scopes of the two reactions were investigated (Table 2). Thus, 1-(2-iodophenyl)-3-phenylprop-2-yn-1-ones bearing R¹ substituents as MeO, Cl, Br, and CF₃ were all competent substrates to yield the corresponding products (**4a–4i**). The incorporation of an electron-withdrawing group R¹ (CF₃) leads to a decreased product yield as indicated from the results of **4e**. In contrast, electronic nature of the R² group has little impact on the reaction, and the products **4f–4i** were all obtained in moderate to good yields (55–68%). Regarding propargyl tosylamides **2**, R³ substituents including *p*-Me, *p*-MeO, *p*-Cl, and *p*-Br were tolerated to furnish products **4j–4m** in moderate to good yields. The reaction of a naphthanenyl-derived substrate was also applicable, albeit producing **4n** in a low yield. In addition, the performance of propargyl benzenesulfonamide and propargyl *o*-nitrobenzenesulfonamide were checked, and the reactions gave the products **4o** and **4p** in 75% and 45% yields, respectively. Noted that the reaction of 1-(2-iodocyclohex-1-en-1-yl)-3-phenylprop-2-yn-1-one failed

Scheme 3. Deuterium-Labeled Experiments



to generate the expected product **4q**. The structure of **4a** was further characterized by an X-ray diffraction study.


The reactions of various 1-(2-iodophenyl)-3-phenylprop-2-yn-1-ones **1** with propargylic benzyl ethers **2** were also evaluated (Table 2, right column). 1-(2-Iodophenyl)-3-phenylprop-2-yn-1-ones **1** bearing R^1 substituents as Me, MeO, Cl, and even the Br atom all reacted smoothly with **3a**, to afford the corresponding products **5a–5e** in 38–73% yields. Varied R^2 substituents like Et, Cl, MeO, CN, and CF_3 were allowed to give the expected products **5f–5j**, **5r** in 57–78% yields. Several variants on R^3 like *p*-Cl, *p*-MeO, and *o*-MeO groups in propargyl benzyl ethers **3** were examined, and the reactions gave the expected products (**5k–5m**, **5o–5t**) in good to high yields. It is notable that the incorporation of the electron-donating *p*-MeO group as R^3 gave the products high yields (**5l** and **5t**). However, the reaction of a 2-thiophenyl substrate with **1a** gave the product **5n** in a low yield (37%). Similarly, 1-(2-iodocyclohex-1-en-1-yl)-3-phenylprop-2-yn-1-one was not a suitable substrate to give the desired product **5u**. The structure of **5h** was further determined by an X-ray single crystal diffraction analysis unambiguously.

Mechanism Study. The distinguished chemoselectivity of the above reactions clearly suggests remarkably different reactivities between the oxygen- and nitrogen-substituted alkyne-ynone intermediates engaged in the reactions. Thus, a series of control experiments were conducted to probe the reaction mechanism. As shown in Scheme 2, the Pd/Cu-catalyzed reaction of **1a** and **2a** performed at room temperature for 1.5 h afforded the cross-coupling product **6a** in 90% yield. Treatment of **6a** with DIPA in toluene for 6 h afforded **4a** in 76–82% yield, while the reaction in the absence of DIPA did not give **4a** (Scheme 2, eqs 1–3). In addition, it was observed that the presence of radical scavengers, i.e., 2,2,6,6-tetramethyl-1-piperidinyloxy, butylated hydroxytoluene, and 1,1-diphenylethane, led to a significant loss of product yield of **4a** (Scheme 2, eq 4).


On the other hand, the cross-coupling intermediate **6b** from **1a** and **3a** was also isolated. Heating of **6b** with toluene affords **5a** in 75% yield (Scheme 2, eq 4). We also checked the reactions in the presence of the stronger base, i.e., DBU. It was found that a catalytic amount of DBU promoted the transformation of **6b** into the cyclobuta[*b*]naphthalen-3(1H)-one **5a'** in 35% yield (Scheme 2, eq 6).⁶⁴ These results imply that the acidity of propargylic C–H in the alkyne-ynone intermediate (**6a/6b**) may play a critical role in the selectivity of the reaction. Moreover, it was found that the presence of radical scavengers did not shut down the transformation, suggesting that a free radical pathway is unlikely for the reaction (Scheme 2, eq 7). In addition, the detection of PhCH_2OH by GC/MS analysis of the reaction mixture indicated the involvement of a hydrolysis step for the formation of the final product **5a**.

The labeling studies using deuterated substrates were performed to further gain insights into the mechanism (Scheme 3). The reaction of **1a** and **2a–d** gave **4a–d1** in 68% yield, in which only ca. 53% content of D was observed at C(8) (Scheme 3, eq 1). The significant loss of D content supports the notion that H/D exchange takes place in the DIPA-assisted propargyl-allenyl isomerization step (Scheme 5, vide infra). Conducting the reaction using the aprotic Et_3N as the base afforded **4a–d2** in which 84% D was incorporated, further supporting this assumption (Scheme 3, eq 2). On the other hand, the reaction of **1a** and **3a–d** under the optimal conditions produced **5a–d** in 65% yield. ^1H NMR analysis of this product showed that about 78% content of the D atom was incorporated into the C(10) position, almost consistent with the D content of **3a–d** at the propargylic carbon (Scheme 3, eq 3). This result led us to postulate that a direct 1,8-H transfer of the cross-coupling intermediate **6b** might take place during the reaction to form a different allenic intermediate (like species **15–Z** in Scheme 5, vide infra) to fulfill the sequence.

Scheme 4. Intramolecular Trapping of the Enone-Allene Intermediate



Scheme 5. Proposed Mechanisms for the Reactions



To verify the above hypothesis, we performed the reactions of **1a** with allyl-substituted propargyl ethers **7a** and **7b** to

capture the possible intermediates. Gratifyingly, both reactions proceeded smoothly under the optimal conditions and gave




Figure 1. (a) Free energy profile for the formal [2 + 2] or [4 + 2] cycloaddition of 1,6-diyne **6a** with DIPA. The energy values are in kcal mol⁻¹ and represent the relative free energies calculated at the M06/6-311+G(d,p)-SDD/SMD(toluene)//M06/6-31G(d)-SDD/SMD(toluene) level of theory in toluene solvent. (b) Free energy profile for the formal [2 + 2] or [4 + 2] cycloaddition of 1,6-diyne **6b** with DIPA. The energy values are in kcal mol⁻¹ and represent the relative free energies calculated at the M06/6-311+G(d,p)-SDD/SMD(toluene)//M06/6-31G(d)-SDD/SMD(toluene) level of theory in toluene solvent.

rise to the fused pentacyclic and tetracyclic products **8a** and **8b** with excellent diastereoselectivity in 54 and 80% yield, respectively (Scheme 4). The structure of **8a** was unambiguously confirmed by an X-ray diffraction study. The product **8b** was fully characterized by NMR spectra including COSY,

NOESY, HSQC, and HMBC. It was interestingly observed that H_(ph) atoms of the indanone moiety in compound **8b** show NMR signals at a relatively high field probably because of the shielding effect of the remote phenyl ring incorporated at the 2,3-dihydrofuran moiety. The formation of these products

thus can be rationalized by a process of intramolecular Diels–Alder reaction of the enone-allene intermediate **10/12**, generated from a propargylic Alder-ene-type reaction of **9/11**.

At this stage, the plausible mechanism of the reaction was proposed (Scheme 5). The Pd/Cu-catalyzed Sonogashira coupling reaction first generates the coupling intermediate. In the case of propargyl tosylamide **2a** as the coupling partner, the cross-coupling intermediate **6a** undergoes DIPA-assisted propargyl-allenyl isomerization to form an ynone-allene intermediate **14**. The isomerization should proceed through an ion-pair intermediacy **13**, as indicated by the deuterium-labeled experiments (Scheme 5, eqs 1 and 2). Subsequent intramolecular [2 + 2] cycloaddition of **14** furnishes the product **4a**. When propargyl benzyl ether **3a** was employed as the coupling partner, the resulted intermediate **6b** may undergo a propargylic Alder-ene-type reaction to give a cyclized enone-allene intermediate **15-Z**. This step may be influenced by the sterically hindering effect of a substitute *ortho* to the carbonyl group in substrate **1**, which is indicated by a low yield of **5e** (Table 2). Subsequent 6 π -electrocyclization forms a fused tetracyclic intermediate **16**. Successive hydrolysis of the vinyl ether motif and aromatization finally form **5a**.

To validate the mechanism of the cycloaddition step to reveal the substituent-controlled chemoselectivity, we conducted density functional theory (DFT) calculations at the M06/6-311 + G(d,p)-SDD/SMD(toluene)//M06/6-31G(d)-SDD/SMD(toluene) level of theory. As shown in Figure 1, 1,6-diynes **6a** and **6b**, in situ generated from Sonogashira coupling, were chosen as the starting species in our theoretical study. In this type of diyne, the acidity of the propargylic hydrogen plays an important role in chemoselectivity. Either base-assisted 1,3-hydrogen transfer or Alder-ene-type hydrogen transfer could occur to achieve different transformations, which would depend on the neighboring substituent group. As shown in Figure 1a, **6a** was chosen as relative zero in calculated free energy profiles. The [2 + 2] cycloaddition could start from a DIPA-assisted intramolecular 1,3-hydrogen transfer. The DIPA-promoted deprotonation of **6a** at the propargyl position could occur to generate an allenyl anion species **13** via transition state **TS-1**. The calculated activation free energy is 25.4 kcal/mol for this process, which would be the rate-determining step. Subsequent re-protonation of the allenyl carbon in **13** could achieve 1,3-hydrogen transfer to generate the ynone-allene intermediate **14**. The following [2 + 2] cycloaddition could take place through a stepwise pathway.⁶⁵ The formation of the first C–C bond could occur to generate the diradical intermediate **17** via a close-shell transition state **TS-3** with a free energy barrier of 20.7 kcal/mol. The radical–radical coupling takes place via an open-shell singlet transition state **TS-4** to yield the product **4a** in a highly exergonic step. The competitive intramolecular Alder-ene-type pathway was also considered. As shown in Figure 1a (left part), the Alder-ene-type hydrogen transfer would occur via transition state **TS-5** with a free energy barrier of 26.7 kcal/mol to provide the enone-allene intermediate **18-Z** for further transformations. The calculated activation free energy of this step is 1.3 kcal/mol higher than that of DIPA-assisted 1,3-hydrogen transfer. Therefore, a stepwise process of 1,3-hydrogen transfer followed by [2 + 2] cycloaddition is preferred. On the other hand, DFT calculation revealed that the deprotonation of 1,6-diyne **6b** needs to traverse a free energy barrier of 28.4 kcal/mol via proton transfer transition state **TS-11** (Figure 1b). The free energy barrier of this step is 3.0 kcal/mol higher than the

corresponding step using propargyl sulfonamido substituted 1,6-diyne **6a**. It means the deprotonation of **6b** is difficult compared with **6a**. Based on this observation, we considered the alternative intramolecular Alder-ene-type reaction of **6b**. DFT calculation clearly revealed that the free energy barrier of Alder-ene-type reaction via transition state **TS-6** is 25.8 kcal/mol, which is 2.6 kcal/mol lower than that of deprotonation via transition state **TS-11**. Therefore, Alder-ene-type reaction would take precedence over deprotonation, which leads to different chemoselectivity. The Alder-ene reaction gives an enone-allene intermediate **15-Z** with exergonic energy of 38.9 kcal/mol leading to an irreversible process. Subsequently, the copper-promoted *cis-trans* isomerization occurs, eventually resulting in the generation of **15-E**. This isomerization process is endothermic by 1.3 kcal mol^{−1}, with an overall activation free energy of 19.2 kcal mol^{−1}. The subsequent pericyclic 6 π -electrocyclic reaction takes place via transition state **TS-8**, which reversibly provides a dearomatized benzofluorenone intermediate **16**. Then, hydrolysis of vinyl ether moiety results in a ketone intermediate **19**, which could undergo a dehydrogenation to achieve aromatization and yield the product **5a** (see Figure S6 in the Supporting Information). DFT calculation clearly revealed that the chemoselectivity is determined by the acidity of propargyl hydrogen in the 1,6-diyne molecule. We calculated the acidity of intermediate **6a** and **6b** using B3LYP/6-311++G(2df,2p)/IEFPCM(DMSO)//B3LYP/6-31G(d) level of theory.^{66–70} We found that the pK_a of **6a** is 9.8, which is 0.9 lower than that of **6b**, which could be attributed to the substituent effect. Therefore, **5a** would be the major product when **6b** is used as the reactant. Our DFT calculations are in complete agreement with experimental observation.

CONCLUSIONS

In conclusion, we have reported two sets of palladium-catalyzed tandem reactions involving cross-coupling, allene formation, and formal [2 + 2] or [4 + 2] cycloaddition, thereby providing a rapid synthesis of polycyclic compounds bearing cyclobuta[*b*]naphthalen-3(1H)-one and 5-benzoyl-11H-benzo[*b*]fluoren-11-one skeletons. The method has advantages such as the use of readily available starting materials, operational simplicity, catalytic conditions, and high selectivity. Mechanism studies indicated that the reactions involved the generation of two different types of allenic intermediates, i.e., ynone-allenes via propargyl-allenyl isomerization (1,3-hydrogen transfer) and enone-allenes via the propargylic ene-type reaction (1,8-hydrogen transfer), from the corresponding cross-coupling intermediates. Detailed computational calculations were performed to elucidate the reaction mechanism and provide insights on the origin of the selectivity of the reactions. Ongoing efforts are currently directed to further expanding the scope and applications of the methodology.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acscatal.3c00857>.

Detailed experimental procedures, compound characterization data, X-ray data, computational details, and copies of NMR spectra for new compounds (PDF)

Crystallographic data for **4a** (cif)

Crystallographic data for **5a** (cif)

Crystallographic data for 8a (cif)

AUTHOR INFORMATION

Corresponding Authors

Shugao Zhu – School of Chemistry and Chemical Engineering, Nanchang University, Nanchang 330031, China; orcid.org/0000-0001-8814-4161; Email: shugaozhu@ncu.edu.cn

Yu Lan – School of Chemistry and Chemical Engineering, Chongqing Key Laboratory of Theoretical and Computational Chemistry, Chongqing University, Chongqing 400030, China; College of Chemistry and Institute of Green Catalysis, Zhengzhou University, Zhengzhou 450001, China; orcid.org/0000-0002-2328-0020; Email: lanyu@cqu.edu.cn

Ruwei Shen – State Key Laboratory of Materials-Oriented Chemical Engineering, College of Chemical Engineering, Nanjing Tech University, Nanjing 211800, China; orcid.org/0000-0002-8834-3256; Email: shenrw@njtech.edu.cn

Authors

Dan Wang – School of Chemistry and Chemical Engineering, Nanchang University, Nanchang 330031, China

Shihan Liu – School of Chemistry and Chemical Engineering, Chongqing Key Laboratory of Theoretical and Computational Chemistry, Chongqing University, Chongqing 400030, China

Yufeng Ma – School of Chemistry and Chemical Engineering, Nanchang University, Nanchang 330031, China

Hong Wang – School of Chemistry and Chemical Engineering, Nanchang University, Nanchang 330031, China

Ruopeng Bai – School of Chemistry and Chemical Engineering, Chongqing Key Laboratory of Theoretical and Computational Chemistry, Chongqing University, Chongqing 400030, China; orcid.org/0000-0002-1097-8526

Complete contact information is available at: <https://pubs.acs.org/10.1021/acscatal.3c00857>

Author Contributions

[†]D.W., S.L., and Y.M. contributed equally to this work.

Author Contributions

S.Z. and R.S. contributed to the conception and design of the experiments. D.W., Y.M., and H.W. performed the experiments and analyzed the data. Y.L. and R.B. directed the calculation section, and S.L. completed the DFT calculation. S.Z. directed the whole project. The manuscript was written through the contributions of all authors.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (22261033, 22271034, and 22271149) and the start-up Fund of Nanchang University. We acknowledge Beijing PARATERA Tech CO, for providing HPC resources that have contributed to the research results. Reported within this paper. URL <http://cloud.paratera.com>.

REFERENCES

- (1) Sydnes, L. K. Allenes from cyclopropanes and their use in organic synthesis recent developments. *Chem. Rev.* **2003**, *103*, 1133–1150.
- (2) Krause, N.; Hashmi, A. S. *Modern allene chemistry*; Wiley-VCH, 2004; p 1143.
- (3) Hoffmann-Röder, A.; Krause, N. Synthesis and properties of allenic natural products and pharmaceuticals. *Angew. Chem., Int. Ed.* **2004**, *43*, 1196–1216.
- (4) Ma, S. Some typical advances in the synthetic applications of allenenes. *Chem. Rev.* **2005**, *105*, 2829–2872.
- (5) Yu, S.; Ma, S. Allenes in catalytic asymmetric synthesis and natural product syntheses. *Angew. Chem., Int. Ed.* **2012**, *51*, 3074–3112.
- (6) Rivera-Fuentes, P.; Diederich, F. Allenes in molecular materials. *Angew. Chem., Int. Ed.* **2012**, *51*, 2818–2828.
- (7) Neff, R. K.; Frantz, D. E. Recent advances in the catalytic syntheses of allenenes: a critical assessment. *ACS Catal.* **2014**, *4*, 519–528.
- (8) Bates, R. W.; Satcharoen, V. Nucleophilic transition metal based cyclization of allenenes. *Chem. Soc. Rev.* **2002**, *31*, 12–21.
- (9) Ma, S. Electrophilic addition and cyclization reactions of allenenes. *Acc. Chem. Res.* **2009**, *42*, 1679–1688.
- (10) Alcaide, B.; Almendros, P. Novel cyclization reactions of aminoallenenes. *Adv. Synth. Catal.* **2011**, *353*, 2561–2576.
- (11) Krause, N.; Winter, C. Gold-catalyzed nucleophilic cyclization of functionalized allenenes: a powerful access to carbo- and heterocycles. *Chem. Rev.* **2011**, *111*, 1994–2009.
- (12) Allen, A. D.; Tidwell, T. T. Ketenes and other cumulenes as reactive intermediates. *Chem. Rev.* **2013**, *113*, 7287–7342.
- (13) Alcaide, B.; Almendros, P. Gold-catalyzed cyclization reactions of allenol and alkynol derivatives. *Acc. Chem. Res.* **2014**, *47*, 939–952.
- (14) Tius, M. A. Allene ether Nazarov cyclization. *Chem. Soc. Rev.* **2014**, *43*, 2979–3002.
- (15) Ye, J.; Ma, S. Palladium-catalyzed cyclization reactions of allenenes in the presence of unsaturated carbon–carbon bonds. *Acc. Chem. Res.* **2014**, *47*, 989–1000.
- (16) Alcaide, B.; Almendros, P.; Aragoncillo, C. Cyclization reactions of bis (allenenes) for the synthesis of polycarbo (hetero) cycles. *Chem. Soc. Rev.* **2014**, *43*, 3106–3135.
- (17) Ye, Z.; Adhikari, S.; Xia, Y.; Dai, M. Expedient syntheses of N-heterocycles via intermolecular amphoteric diamination of allenenes. *Nat. Commun.* **2018**, *9*, 721.
- (18) Qian, D.; Zhang, J. Gold-catalyzed cyclopropanation reactions using a carbenoid precursor toolbox. *Chem. Soc. Rev.* **2015**, *44*, 677–698.
- (19) Lemiere, G.; Gandon, V.; Cariou, K.; Fukuyama, T.; Dhiman, A.-L.; Fensterbank, L.; Malacria, M. Tandem gold (I)-catalyzed cyclization/electrophilic cyclopropanation of vinyl allenenes. *Org. Lett.* **2007**, *9*, 2207–2209.
- (20) Tejedor, D.; Méndez-Abt, G.; Cotos, L.; García-Tellado, F. Propargyl Claisen rearrangement: allene synthesis and beyond. *Chem. Soc. Rev.* **2013**, *42*, 458–471.
- (21) Wang, Y.; Burton, D. J. Site-specific preparation of 2-carboalkoxy-4-substituted naphthalenes and 9-alkylphenanthrenes and evidence for an allene intermediate in the novel base-catalyzed cyclization of 2-alkynylbiphenyls. *Org. Lett.* **2006**, *8*, 5295–5298.
- (22) Barber, J. S.; Yamano, M. M.; Ramirez, M.; Darzi, E. R.; Knapp, R. R.; Liu, F.; Houk, K.; Garg, N. K. Diels–Alder cycloadditions of strained azacyclic allenenes. *Nat. Chem.* **2018**, *10*, 953–960.
- (23) Yamano, M. M.; Kelleghan, A. V.; Shao, Q.; Giroud, M.; Simmons, B. J.; Li, B.; Chen, S.; Houk, K.; Garg, N. K. Intercepting fleeting cyclic allenenes with asymmetric nickel catalysis. *Nature* **2020**, *586*, 242–247.
- (24) Shen, R.; Huang, X. Pd-catalyzed sequential reactions via allene intermediate for the synthesis of polycyclic frameworks containing 2, 3-dihydrofuran units. *Org. Lett.* **2008**, *10*, 3283–3286.
- (25) Shen, R.; Huang, X.; Chen, L. A Facile and Efficient Synthesis of Dihydroisobenzofuran Derivatives via Tandem Palladium-Cata-

- lyzed Coupling, Propargyl-Allenyl Rearrangement, [4+ 2] Cycloaddition and Aromatization Reaction. *Adv. Synth. Catal.* **2008**, 350, 2865–2870.
- (26) Zhu, S.; Cao, J.; Wu, L.; Huang, X. Synthesis of Polycyclic Isoindoline Derivatives via Tandem Pd-Catalyzed Coupling, Propargyl-Allenyl Isomerization, [4+ 2] Cycloaddition and Aromatization Reaction. *J. Org. Chem.* **2012**, 77, 10409–10415.
- (27) D'Souza, D. M.; Rominger, F.; Mueller, T. J. A domino sequence consisting of insertion, coupling, isomerization, and diels-alder steps yields highly fluorescent spirocycles. *Angew. Chem., Int. Ed.* **2005**, 44, 153–158.
- (28) D'Souza, D. M.; Kiel, A.; Herten, D. P.; Rominger, F.; Mueller, T. J. Synthesis, structure and emission properties of spirocyclic benzofuranones and dihydroindolones: a domino insertion-coupling-isomerization-diels-alder approach to rigid fluorophores. *Chem. – Eur. J.* **2008**, 14, 529–547.
- (29) Shen, R.; Zhu, S.; Huang, X. Unexpected Pd-Catalyzed Coupling, Propargyl-Allenyl Isomerization and Alder-Ene Reaction: Facile Synthesis of Some Not Readily Available 2, 3-Dihydrofuran Derivatives. *J. Org. Chem.* **2009**, 74, 4118–4123.
- (30) Zhu, S.; Wu, L.; Huang, X. Palladium-catalyzed sequential reaction via Sonogashira coupling, isomerization, Claisen rearrangement and [4+ 2] cycloaddition sequence for the rapid synthesis of tricyclo [3.2.1.0^{2,7}] oct-3-ene derivatives. *RSC Adv.* **2012**, 2, 132–134.
- (31) D'Souza, D. M.; Rominger, F.; Müller, T. J. Coupling-isomerization-Claisen sequences-mechanistic dichotomies in hetero domino reactions. *Chem. Commun.* **2006**, 4096–4098.
- (32) D'Souza, D. M.; Liao, W.-W.; Rominger, F.; Müller, T. J. Dichotomies in microwave-assisted propargyl-isomerization-Claisen domino sequences dependent on base strengths. *Org. Biomol. Chem.* **2008**, 6, 532–539.
- (33) Huang, X.; Zhu, S.; Shen, R. Palladium-Catalyzed Sequential Reactions via Allene Intermediates for the Rapid Synthesis of Fused Polycyclic Pyrrole Derivatives. *Adv. Synth. Catal.* **2009**, 351, 3118–3122.
- (34) Shen, R.; Chen, L.; Huang, X. Facile Synthesis of Polycyclic Fluorene Derivatives via a Palladium-Catalyzed Coupling, Propargyl-Allenyl Isomerization and Schmitt Cyclization Sequence. *Adv. Synth. Catal.* **2009**, 351, 2833–2838.
- (35) Müller, T. J. J.; Ansorge, M.; Aktah, D. An Unexpected Coupling-Isomerization Sequence as an Entry to Novel Three-Component-Pyrazoline Syntheses. *Angew. Chem., Int. Ed.* **2000**, 39, 1253–1256.
- (36) Cao, J.; Huang, X. Facile Synthesis of Tetrahydro-1 H-isoindolones via a Sequential Three-Component Copper-Catalyzed Coupling/Propargyl-Allenyl Isomerization/[4+ 2] Cyclization Reaction. *Org. Lett.* **2010**, 12, 5048–5051.
- (37) Mueller, T. J. Synthesis of carbo-and heterocycles via coupling-isomerization reactions. *Synthesis* **2012**, 44, 159–174.
- (38) Shen, R.; Chen, K.; Deng, Q.; Yang, J.; Zhang, L. Highly stereoselective generation of complex oxy-bicyclic scaffolds via an atom-economic Pd (II)-catalyzed hydroalkynylation, isomerization and Diels-Alder cycloaddition sequence. *Org. Lett.* **2014**, 16, 1208–1211.
- (39) Ghazvini, H. J.; Armaghan, M.; Janiak, C.; Balalaie, S.; Müller, T. J. Coupling-Isomerization-Cycloisomerization Reaction (CICIR) – An Unexpected and Efficient Domino Approach to Luminescent 2-(Hydroxymethylene) indenones. *Eur. J. Org. Chem.* **2019**, 2019, 7058–7062.
- (40) Hori, H.; Arai, S.; Nishida, A. A 2-Benzothiazolylphenyl Group Accelerates the Intramolecular [2+ 2] Cycloaddition of Allene-Ynes. *Asian J. Org. Chem.* **2014**, 3, 41–43.
- (41) Brummond, K. M.; Osbourn, J. M. Complete transfer of chirality in an intramolecular, thermal [2+ 2] cycloaddition of allenes to form non-racemic spirooxindoles. *Beilstein J. Org. Chem.* **2011**, 7, 601–605.
- (42) Ni, Q.; Song, X.; Png, C. W.; Zhang, Y.; Zhao, Y. Access to substituted cyclobutenes by tandem [3, 3]-sigmatropic rearrangement/[2+ 2] cycloaddition of dipropargylphosphonates under Ag/Co relay catalysis. *Chem. Sci.* **2020**, 11, 12329–12335.
- (43) Matsuda, T.; Kadowaki, S.; Goya, T.; Murakami, M. A direct entry to bicyclic cyclobutenes via platinum-catalyzed cycloisomerization of allenes. *Synlett* **2006**, 2006, 0575–0578.
- (44) Mukai, C.; Inagaki, F.; Yoshida, T.; Kitagaki, S. Rh (I)-catalyzed ring-closing reaction of allenes: selective construction of cycloheptene, bicyclo [5.3. 0] decadienone, and bicyclo [5.2. 0] nonene frameworks. *Tetrahedron Lett.* **2004**, 45, 4117–4121.
- (45) Lee-Ruff, E.; Mladenova, G. Enantioselectively pure cyclobutane derivatives and their use in organic synthesis. *Chem. Rev.* **2003**, 103, 1449–1484.
- (46) Xu, H.; Zhang, W.; Shu, D.; Werness, J. B.; Tang, W. Synthesis of Cyclobutenes by Highly Selective Transition-Metal-Catalyzed Ring Expansion of Cyclopropanes. *Angew. Chem., Int. Ed.* **2008**, 47, 8933–8936.
- (47) Mohamed, R. K.; Peterson, P. W.; Alabugin, I. V. Concerted reactions that produce diradicals and zwitterions: electronic, steric, conformational, and kinetic control of cycloaromatization processes. *Chem. Rev.* **2013**, 113, 7089–7129.
- (48) Noucti, N. N.; Alexanian, E. J. Stereoselective Nickel-Catalyzed [2+ 2] Cycloadditions of Ene-Allenes. *Angew. Chem., Int. Ed. Engl.* **2015**, 127, 5447–5450.
- (49) Xu, Y.; Conner, M. L.; Brown, M. K. Cyclobutane and cyclobutene synthesis: Catalytic enantioselective [2+ 2] cycloadditions. *Angew. Chem., Int. Ed.* **2015**, 54, 11918–11928.
- (50) Kumar, R.; Tamai, E.; Ohnishi, A.; Nishimura, A.; Hoshimoto, Y.; Ohashi, M.; Ogoshi, S. Nickel-catalyzed enantioselective synthesis of cyclobutenes via [2+ 2] cycloaddition of α , β -unsaturated carbonyls with 1, 3-enynes. *Synthesis* **2016**, 48, 2789–2794.
- (51) Misale, A.; Niyomchon, S.; Maulide, N. Cyclobutenes: at a crossroad between diastereoselective syntheses of dienes and unique palladium-catalyzed asymmetric allylic substitutions. *Acc. Chem. Res.* **2016**, 49, 2444–2458.
- (52) Antonsen, S.; Østby, R. B.; Stenström, Y. Naturally occurring cyclobutenes: their biological significance and synthesis. *Stud. Nat. Prod. Chem.* **2018**, 57, 1–40.
- (53) Wen, K.-G.; Peng, Y.-Y.; Zeng, X.-P. Advances in the catalytic asymmetric synthesis of quaternary carbon containing cyclobutenes. *Org. Chem. Front.* **2020**, 7, 2576–2597.
- (54) Biswas, S.; Parsutkar, M. M.; Jing, S. M.; Pagar, V. V.; Herbort, J. H.; RajanBabu, T. A New Paradigm in Enantioselective Cobalt Catalysis: Cationic Cobalt (I) Catalysts for Heterodimerization, Cycloaddition, and Hydrofunctionalization Reactions of Olefins. *Acc. Chem. Res.* **2021**, 54, 4545–4564.
- (55) Zhu, M.; Huang, X.-L.; Xu, H.; Zhang, X.; Zheng, C.; You, S.-L. Visible-light-mediated synthesis of cyclobutene-fused indolizidines and related structural analogs. *CCS Chem.* **2021**, 3, 652–664.
- (56) Wang, J. Y.; Li, C. L.; Xu, T.; Li, M. F.; Hao, W. J.; Tu, S. J.; Wang, J.; Li, G.; Yu, Z. X.; Jiang, B. Catalytic Enantioselective Construction of 6-4 Ring-Junction All-Carbon Stereocenters and Mechanistic Insights. *Chin. J. Chem.* **2022**, 40, 1767–1776.
- (57) Shen, Q.; Hammond, G. B. Regiospecific Synthesis of Bicyclo- and Heterobicyclo-gem-difluorocyclobutenes Using Functionalized Fluoroallenes and a Novel Mo-Catalyzed Intramolecular [2+ 2] Cycloaddition Reaction. *J. Am. Chem. Soc.* **2002**, 124, 6534–6535.
- (58) Cao, J.; Huang, X.; Wu, L. Facile Synthesis of Selenium/Sulfur-Substituted 3-Oxa-bicyclo [4. 2. 0] octa-1 (8), 5-diene and Tetrahydro-1 H-isochromene via Sequential Three-Component Conjugate Addition/Condensation/Elimination/[2+ 2] or [4+ 2] Cyclization Reactions. *J. Org. Chem.* **2011**, 76, 1440–1443.
- (59) Jiang, X.; Ma, S. Intramolecular [2+ 2]-cycloaddition of propargylic 2, 3-allenoates for the efficient synthesis of 3-oxabicyclo [4.2. 0] octa-1 (8), 5-dien-4-ones: a dramatic substituent effect. *Tetrahedron* **2007**, 63, 7589–7595.
- (60) Wang, J. Y.; Hao, W. J.; Tu, S. J.; Jiang, B. Engaging Yne-Allenenes in Cycloaddition Reactions: Recent Developments. *Chin. J. Chem.* **2022**, 40, 1224–1242.

(61) Sakai, T.; Danheiser, R. L. Cyano Diels–Alder and cyano ene reactions. Applications in a formal [2+ 2+ 2] cycloaddition strategy for the synthesis of pyridines. *J. Am. Chem. Soc.* **2010**, *132*, 13203–13205.

(62) Robinson, J. M.; Sakai, T.; Okano, K.; Kitawaki, T.; Danheiser, R. L. Formal [2+ 2+ 2] cycloaddition strategy based on an intramolecular propargylic ene reaction/Diels–Alder cycloaddition cascade. *J. Am. Chem. Soc.* **2010**, *132*, 11039–11041.

(63) Sasaki, M.; Hamzik, P. J.; Ikemoto, H.; Bartko, S. G.; Danheiser, R. L. Formal Bimolecular [2+ 2+ 2] Cycloaddition Strategy for the Synthesis of Pyridines: Intramolecular Propargylic Ene Reaction/Aza Diels–Alder Reaction Cascades. *Org. Lett.* **2018**, *20*, 6244–6249.

(64) The reaction of **1a** and **3a** under the Pd/Cu catalysis using 1.1 equiv of DBU afforded **5a** in 29% yield. However, the reaction with 3 equiv of DBU generated a complex mixture. For these results, also see Section 8.2 in Supporting Information.

(65) Siebert, M. R.; Osbourn, J. M.; Brummond, K. M.; Tantillo, D. J. Differentiating mechanistic possibilities for the thermal, intramolecular [2 + 2] cycloaddition of allene–ynes. *J. Am. Chem. Soc.* **2010**, *132*, 11952–11966.

(66) Miertuš, S.; Scrocco, E.; Tomasi, J. Electrostatic interaction of a solute with a continuum. A direct utilization of AB initio molecular potentials for the prevision of solvent effects. *Chem. Phys.* **1981**, *55*, 117–129.

(67) Scalmani, G.; Frisch, M. J. Continuous surface charge polarizable continuum models of solvation I. General formalism. *J. Chem. Phys.* **2010**, *132*, No. 114110.

(68) Chengteh, L.; Weitao, Y.; Robert, G. P. Development of the Colle–Salvetti correlation-energy formula into a functional of the electron density. *Phys. Rev. B* **1988**, *37*, 785–789.

(69) Becke, A. D. Density-functional thermochemistry. I. The effect of the exchange-only gradient correction. *J. Chem. Phys.* **1992**, *96*, 2155–2160.

(70) Shen, K.; Fu, Y.; Li, J.-N.; Liu, L.; Guo, Q.-X. What are the pK_a values of C–H bonds in aromatic heterocyclic compounds in DMSO? *Tetrahedron* **2007**, *63*, 1568–1576.

Recommended by ACS

Diverting the 5-*exo*-Trig Oxypalladation to Formally 6-*endo*-Trig Fluorocycloetherification Product through 1,2-O/Pd(IV) Dyotropic Rearrangement

Jing Gong, Jieping Zhu, *et al.*

JULY 18, 2023

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Pd-Catalyzed Domino Narasaka–Heck/C–H Activation/Amination Reactions: Synthesis of Bis-heterocyclic Spirocycles

Jonathan Bajohr, Mark Lautens, *et al.*

JULY 12, 2023

ORGANIC LETTERS

READ 

Asymmetric Dearomatization of Indoles with Azodicarboxylates via Cascade Electrophilic Amination/Aza-Prins Cyclization/Phenonium-like Rearrangement

Xian-Yun Huang, Shu-Li You, *et al.*

MAY 19, 2023

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Regio-, Site-, and Stereoselective Three-Component Aminofluorination of 1,3-Dienes via Cooperative Silver Salt and Copper Catalysis

Yang Li, Junkai Fu, *et al.*

FEBRUARY 02, 2023

ACS CATALYSIS

READ 

Get More Suggestions >