

Mn-Mediated Direct Regioselective C–H Trifluoromethylation of Imidazopyridines and Quinoxalines

Mahdieh Esi Firuz, Saideh Rajai-Daryasarei, Frank Rominger, Abbas Biglari, and Saeed Balalaie*

Cite This: <https://doi.org/10.1021/acs.joc.3c00621>

Read Online

ACCESS |




Metrics & More



Article Recommendations



Supporting Information



ABSTRACT: A simple and highly efficient strategy has been developed for direct C–H trifluoromethylation at C-3 of imidazopyridines and C-8 of quinoxalines with readily available Langlois reagent through $\text{KMnO}_4/\text{AcOH}$ system. This protocol showed broad substrate scope and afforded moderate-to-excellent yields of both products. It is the first report that the functionalization of quinoxalines occurred regioselectively at the C-8 position of quinoxalines. Mechanistic studies revealed that reaction proceeded via radical pathway.

INTRODUCTION

Trifluoromethylated compounds have received a significant amount of attention in medicinal chemistry due to the incorporation of a CF_3 group into organic molecules that can significantly enhance their physical, chemical, and biological properties such as membrane permeability, favorable protein–ligand interactions, lipophilicity, bioavailability, and metabolic stability in comparison with their nonfluorinated analogs (Figure 1).¹ Over the past decades, various protocols for the preparation of trifluoromethylated organic compounds utilizing numerous trifluoromethylating agents including CF_3I , TMSCF_3 , $\text{CF}_3\text{SO}_2\text{Na}$, $\text{CF}_3\text{SO}_2\text{Cl}$, Baran's reagent, Umemoto's reagent, Togni's reagent, etc., have been developed.² In particular, sodium trifluoromethanesulfonate (NaSO_2CF_3) was reported by Langlois et al. in 1980 and represents one of the cheapest and most readily available reagents for trifluoromethylation of organic compounds through radical reaction.³ Among the reported strategies, the direct trifluoromethylation of organic compounds via C–H functionalization reaction is of high interest, due to its high atom economy and no require to prefuctionalization.⁴

On the other hand, imidazopyridines and quinoxalines are recognized as important class of nitrogen-containing heterocycles and found in a wide variety of natural products and pharmaceuticals and materials science. These scaffolds exhibit varied biological activities, such as antitumor, anti-inflammatory, antiparasitic, antipyretic, cytotoxic, antiviral, antiulcer,





Figure 1. The trifluoromethyl-containing pharmaceuticals.

Received: March 22, 2023

Scheme 1. Strategies to Regioselective Trifluoromethylation of Nitrogen-Containing Heterocycles



antifungal, anti-rhinoviral and anticancer.⁵ Therefore, synthesis and functionalization of these scaffolds have been paid much attention from the synthetic community.⁶ Various methods have been developed for the C–H bond functionalization of the C-3 position of the imidazopyridine structures using different substituents such as aryl, alkyl, halogen, thiol, and trifluoromethyl.⁷ Among these reported methods, direct trifluoromethylation of imidazo[1,2-*a*]pyridines via trifluoromethyl radicals have received considerable attention. In 2015, Hajra and co-workers were the first group to report the successful synthesis of 3-trifluoromethylimidazo[1,2-*a*]pyridines using Langlois' reagent (NaSO_2CF_3) as CF_3 radical source in the presence of TBHP and AgNO_3 at room temperature (Scheme 1a).⁸ The direct C3-trifluoromethylation of imidazo[1,2-*a*]pyridine derivatives with Ruppert-Prakash reagent (TMSCF_3) in the presence of $\text{PhI}(\text{OAc})_2$ and CsF in CH_3CN was described by Wu et al. in 2019 (Scheme 1b).⁹ Very recently, Deng and co-workers disclosed a visible-light-promoted C3-trifluoromethylation of imidazopyridine derivatives using tetramethylguanidine trifluoromethyl iodide complex ($\text{TMG}\cdot\text{CF}_3\text{I}$) as CF_3 radical source in the presence of DBU as base (Scheme 1c).¹⁰ Despite such advances, there still remains need for novel and simple synthetic strategies that are facile and effective. In addition, the strategy C–H functionalization of quinoxaline derivatives have remained largely unexplored, and this area of research is still in its infancy. To the best of our knowledge, no examples involving trifluoromethylation of quinoxaline derivatives have been reported to date. Therefore, the development of new synthetic methods to the functionalization these compounds remains a high-value achievement. Inspired by these studies and our continuing interest in the radical reactions, herein we report regioselective C3-trifluoromethylation of imidazo[1,2-*a*]pyridines and C8-trifluoromethylation of quinoxalines using Langlois' reagent (NaSO_2CF_3) as CF_3 radical source in the presence of Mn^{III} species, which is generated in situ by the $\text{KMnO}_4/\text{AcOH}$ system.

RESULTS AND DISCUSSION

To verify the feasibility of our hypothesis, reaction of 2-phenylimidazo[1,2-*a*]pyridine **1a** with Langlois' reagent (NaSO_2CF_3) **2** was selected as the model substrates to optimize the reaction conditions (Table 1). Initially, the reaction of **1a** with $\text{CF}_3\text{SO}_2\text{Na}$ **2** was studied in the presence


Table 1. Optimization of Reaction Conditions^a

entry	solvent	temp (°C)	yield %
1	AcOH	80	45
2	AcOH/DMSO (1:1)	80	68
3	AcOH/PhCl (1:1)	80	35
4	AcOH/PhCH ₃ (1:1)	80	20
5	AcOH/DCE (1:1)	80	43
6	AcOH/acetone (1:1)	80	10
7	AcOH/DMF (1:1)	80	20
8	AcOH/CH ₃ CN (1:1)	80	55
9	AcOH/EtOAc (1:1)	80	61
10	AcOH/DMSO (1:1)	100	72
11	AcOH/DMSO (1:2)	100	65
12	AcOH/DMSO (1:4)	100	52
13 ^b	AcOH/DMSO (1:1)	100	59
14 ^c	AcOH/DMSO (1:1)	100	47
15 ^d	AcOH/DMSO (1:1)	100	72

^aReaction conditions: **1a** (0.2 mmol), **2** (0.6 mmol), solvent (1.0 mL), KMnO_4 (3.0 equiv.), at 100 °C for 2 h. ^b KMnO_4 (4.0 equiv.).

^cReaction time 4 h. ^d N_2 atmosphere.

3.0 equiv of KMnO_4 as oxidant in AcOH as solvent, at the temperature of 80 °C for 2h. To our delight, desired product **3a** was obtained in 45% yield (entry 1). Encouraged by this initial result, we examined the effect of various common solvents. Screening of several solvent systems including (DMSO, PhCl, PhCH₃, DCE, DMF, CH₃CN, EtOAc, and acetone)/AcOH revealed that AcOH/DMSO gives the best result for this reaction (entries 2–9). When the reaction temperature was raised to 100 °C, the desired product **3a** was formed in 72% yields (entry 10). In addition, the effect of different AcOH/DMSO ratios on the reaction yield was tested (entries 11 and 12), unfortunately, no results were obtained on the improvement of the product yield. Increasing the usage of KMnO_4 to 4.0 equiv led to form product **3a** in 59% yield (entry 13). No further improvement of the yield was observed with the increasing reaction time to 4 h (entry 14). As a note, when the model reaction was conducted under N_2 atmosphere, the yield of the desired product was not changed (entry 15). The above optimization results suggested that the maximum

Table 2. Substrate Scope of the Trifluoromethylation of Imidazo[1,2-*a*]pyridines and Quinoxalines^a

^aReaction conditions: **1** (0.2 mmol), **2/4** (0.6 mmol), solvent (1.0 mL), KMnO_4 (3.0 equiv), at 100°C for 2 h.


yield of the desired product **3a** (72%) can be obtained under condition of using 3.0 equiv of KMnO_4 as the oxidant in AcOH/DMSO as solvent at 100°C for 2 h.

With the optimized reaction conditions in hand, the generality of this method using Langlois' reagent (NaSO_2CF_3) **2** and variously substituted imidazo[1,2-*a*]pyridines **1** was investigated and the results are summarized in Table 2. Imidazo[1,2-*a*]pyridines with electron-donating (OMe), halogen (F, Cl, Br), as well as some other electron-withdrawing (CN, NO_2) groups on the benzene ring participated in the reaction well to deliver the desired products **3a–3g** in 72–87% yields. Subsequently, imidazopyridines bearing substituents on the pyridine rings were also explored. 6-Methyl-2-arylimidazo[1,2-*a*]pyridine **2h** and **2i** react with trifluoromethylating agent to afford the desired products **3h** and **3i** in 80 and 83% yields, respectively. When Br and Cl were substituted on the C-6 or C-7 position of imidazo[1,2-*a*]pyridines, the reaction performed well and gave the corresponding products **3j–3q**, in 82–90% yields. In addition, we investigated the trifluorome-


thylation of 1-methyl-1H-imidazole and 2-arylbenzo[*d*]-imidazo[2,1-*b*]thiazoles under standard reaction conditions. Fortunately, trifluoromethylation of benzo[*d*]imidazo[2,1-*b*]thiazoles were led to form desired products **3a'** and **3b'** in 83 and 84% yields, respectively. But, 1-methyl-1H-imidazole was not compatible with this transformation, and the corresponding products **3c'** were not detected after 6 h. It is noteworthy to mention, to show the diversity of the reaction, we also employed quinoxalines **4** as a coupling partner and studied it to the same reaction conditions. Surprisingly, we observed desired product **5a** in 48% yield. Hence, to increase the product yield, we explored different solvent and temperature. A better yield was obtained in AcOH as solvent (please see the optimization table included in the Supporting Information). As depicted in Table 2, a variety of quinoxalines bearing electron-donating, -withdrawing and neutral groups on the phenyl ring were tolerated, allowing the construction of C8-trifluoromethylated quinoxalines **5a–5i** in moderate yields. The structure of **5d** also was unambiguously confirmed by X-ray analysis, the

Scheme 2. Mechanistic Study and Proposed Reaction Mechanism


(A) Gram scale synthesis



(B) Control experiments



(C) Reaction mechanism



ellipsoid probability level is 50%. Additionally, 6,7-dimethyl-2-arylquinoxalines performed smoothly to deliver the desired product **5g–5i** in 48–55% yields. To demonstrate the practicability of this approach, we next carried out the gram-scale synthesis of **3a** and **5a** under the standard conditions and the desired products **3a** and **5a** were obtained in 68% (357 mg) and 42% (230 mg) yields, respectively (Scheme 2A).

To gain insight to the reaction mechanism, a radical scavenger 2,2,6,6-tetramethylpiperidine-1-oxyl (TEMPO) was added to the reaction mixture in varying quantities, and yields of the desired products **3a** and **5a** were sharply decreased (Scheme 2B). These results suggested that this reaction might involve a radical pathway. On the basis of experimental results above and previous reports,^{3,11} a plausible reaction mechanism

is depicted in Scheme 2C. First, Mn^{III} species are generated from the reaction KMnO_4 and AcOH under the thermal condition, which then reacts with the Langlois' reagent (NaSO_2CF_3) **2** to deliver trifluoromethyl radical A. Subsequently, the trifluoromethyl radical is added selectively to C3 of imidazo[1,2-a]pyridine/C8 of quinoxalines to give radical intermediate B/D. Then, a single-electron transfer (SET) from intermediate B/D to Mn^{III} leads to release the cation intermediate C/E and simultaneously Mn^{III} reduces into Mn^{II} . Probably, both the intermediates B/D and C/E can be stabilized through conjugation with adjacent aryl group. Finally, deprotonation of intermediate C/E affords the desired product **3/5**.

In summary, we have developed an operationally simple and efficient approach for regioselective trifluoromethylation of imidazopyridines and quinoxalines using Langlois's reagent as CF_3 radical source by $\text{KMnO}_4/\text{AcOH}$ system. The Mn^{III} species is generated in situ from $\text{KMnO}_4/\text{RCOOH}$ system for the formation of trifluoromethyl radicals. A variety of trifluoromethylated imidazo[1,2-*a*]pyridines and quinoxalines with high functional group tolerance were obtained in moderate-to-excellent yields. To the best of our knowledge, this is the first report for the direct C8-trifluoromethylation of quinoxalines. The attractive features of this protocol are the use of inexpensive and readily available materials, short time, and regioselectivity. Further investigations on the detailed mechanism and applications of this method on the other heteroaromatic compounds are currently underway in our laboratory.

EXPERIMENTAL SECTION

General Information. All chemicals and solvents used in this study were purchased from chemical suppliers (Merck, Sigma-Aldrich, Fluka, and Alpha-Aesar), and used without further purification. The reaction process was monitored by thin-layer chromatography (TLC) using aluminum-coated plates of silica gel (MERCK, 60F254), via detection under UV fluorescence (wavelength of 254 nm). Chromatographic separations of products were carried out by flash column chromatography on MACHEREY-NAGEL silica gel 60 (230–400 mesh). Melting points (m.p.) were determined with an Electrothermal 9100 digital melting point apparatus and are uncorrected. Nuclear Magnetic Resonance (NMR) spectra (^1H and ^{13}C NMR) were recorded on a Bruker AVANCE 300, 400 and 500 MHz (DRX) spectrometer. Chemical shifts (δ) are quoted in ppm and the coupling constants (*J*) in Hz. Multiplets were indicated by the following abbreviations: s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet, complex pattern), dd (doublet of doublet), ddd (doublet of doublet of doublet), td (triplet of doublet), tt (triplet of triplet), and bs (broad singlet). High-resolution mass spectrum (HRMS) (ESI-TOF) was recorded using Waters LCT Premier XE mass spectrometer. X-ray crystal data were measured on a Bruker APEX-II Quazar area detector.

General Experimental Procedures. General Procedure for Preparation of Imidazo[1,2-*a*]pyridines 2a–b'.¹² A mixture of 2-aminopyridine (5.0 mmol), α -bromo ketone (5.5 mmol), and NaHCO_3 (6.0 mmol) in EtOH (10.0 mL) was heated at 70 °C in an oil bath for 1–2 h. The completion of reaction as was indicated by TLC monitoring, then the solvent was removed under the reduced pressure. The product was extracted with ethyl acetate (2 \times 10.0 mL). The combined organic phase was dried over Na_2SO_4 . The solvent was removed under the reduced pressure. The solid product was recrystallized by ethanol affording desired imidazo[1,2-*a*]pyridines.

General Procedure for Trifluoromethylation of 2-Arylimidazo[1,2-*a*]pyridine 3a–b'. First, the solution of potassium permanganate (0.6 mmol, 0.094 g, 3.0 equiv)/acetic acid (0.5 mL) was heated at 100 °C in an oil bath for 15 min. When the reaction mixture completely changed color from purple to brown, time was given for the mixture to reach ambient temperature. Then, dimethyl sulfoxide (0.5 mL) was added to the reaction vessel and 2-arylimidazo[1,2-*a*]pyridine 1 (0.2 mmol, 0.038 g, 1.0 equiv) and sodium trifluoromethanesulfinate 2 (0.6 mmol, 0.096 g, 3.0 equiv) were added to it. Then, the reaction mixture was heated at 100 °C in an oil bath for 2 h. After completion of the reaction as was indicated by TLC monitoring, the reaction mixture was cooled to ambient temperature. A saturated solution of NaHCO_3 (3.0 mL) was added and the product was extracted with ethyl acetate (3 \times 4.0 mL). The combined organic phase was dried over Na_2SO_4 . The solvent was removed under the reduced pressure. The residue was purified by column chromatography using *n*-hexane/EtOAc (8:1) as eluent to afford pure products 3a–b'.

General Procedure for the Preparation of Compound 3a in a Scale of 2.0 mmol. The solution of potassium permanganate (6.0

mmol, 0.948 g, 3.0 equiv)/acetic acid (2.0 mL) was heated at 100 °C in an oil bath for 15 min. When the reaction mixture completely changed color from purple to brown, time was given for the mixture to reach ambient temperature. Then, dimethyl sulfoxide (2.0 mL) was added to the reaction vessel and 2-phenylimidazo[1,2-*a*]pyridine 1 (2.0 mmol, 0.388 g, 1.0 equiv) and sodium trifluoromethanesulfinate 2 (6.0 mmol, 0.936 g, 3.0 equiv) were added to it. Then, the reaction mixture was heated at 100 °C in an oil bath for 2 h. After completion of the reaction as was indicated by TLC monitoring, the reaction mixture was cooled to ambient temperature. A saturated solution of NaHCO_3 (9.0 mL) was added and the product was extracted with ethyl acetate (3 \times 6.0 mL). The combined organic phase was dried over Na_2SO_4 . The solvent was removed under the reduced pressure. The residue was purified by column chromatography using *n*-hexane/EtOAc (8:1) as eluent to afford pure product 3a in 68% (357 mg) yield.

General Procedure for Preparation of Quinoxalines 4a–i.¹³ *o*-Phenylenediamine (5.0 mmol) was added to the solution of arylglyoxal (5.0 mmol) in acetic acid as a solvent (10.0 mL) and the reaction mixture was heated at 80 °C in an oil bath for 1 h and the progress of the reaction was monitored by thin TLC. The reaction mixture was cooled to ambient temperature. A saturated solution of NaHCO_3 (40.0 mL) was added and the product was extracted with ethyl acetate (3 \times 20.0 mL). The combined organic phase was dried over Na_2SO_4 . The solvent was removed under the reduced pressure. The solid product was recrystallized by ethyl acetate affording desired quinoxalines 4a–i.

General Procedure for Trifluoromethylation of 2-Arylquinoxaline 5a–i. First, the solution of potassium permanganate (0.6 mmol, 0.094, 3.0 equiv)/acetic acid (0.5 mL) was heated at 100 °C in an oil bath for 15 min. When the reaction mixture completely changed color from purple to brown, time was given for the mixture to reach ambient temperature. Then, 2-arylquinoxaline 4 (0.2 mmol, 0.041, 1.0 equiv) and sodium trifluoromethanesulfinate 2 (0.6 mmol, 0.096, 3.0 equiv) were added to the reaction vessel and the reaction mixture was heated at 100 °C in an oil bath for 2 h. After completion of the reaction as was indicated by TLC monitoring, the reaction mixture was cooled to ambient temperature. A saturated solution of NaHCO_3 (3.0 mL) was added and the product was extracted with ethyl acetate (3 \times 4.0 mL). The combined organic phase was dried over Na_2SO_4 . The solvent was removed under the reduced pressure. The residue was purified by column chromatography using *n*-hexane/EtOAc (8:1) as eluent to afford pure products 5a–i.

General Procedure for the Preparation of Compound 5a in a Scale of 2 mmol. The solution of permanganate (6.0 mmol, 0.948 g, 3.0 equiv)/acetic acid (2.0 mL) was heated at 100 °C in an oil bath for 15 min. When the reaction mixture completely changed color from purple to brown, time was given for the mixture to reach ambient temperature. Then, 2-phenylquinoxaline 4 (2.0 mmol, 0.412 g, 1.0 equiv) and sodium trifluoromethanesulfinate 2 (6.0 mmol, 0.936 g, 3.0 equiv) were added to the reaction vessel and the reaction mixture was heated at 100 °C in an oil bath for 2 h. After completion of the reaction as was indicated by TLC monitoring, the reaction mixture was cooled to ambient temperature. A saturated solution of NaHCO_3 (9.0 mL) was added and the product was extracted with ethyl acetate (3 \times 5.0 mL). The combined organic phase was dried over Na_2SO_4 . The solvent was removed under the reduced pressure. The residue was purified by column chromatography using *n*-hexane/EtOAc (8:1) as eluent to afford pure product 5a in 42% (230 mg) yield.

6-Methyl-2-phenylimidazo[1,2-*a*]pyridine (2h). Yield: 958 mg, 92%; yellow solid; mp 164–167 °C [lit.^{12d} mp 176–177 °C]. ^1H NMR (400 MHz, CDCl_3): δ 7.94 (dd, *J* = 8.4, 1.4 Hz, 2H), 7.80 (br s, 1H), 7.70 (s, 1H), 7.59 (d, *J* = 9.0 Hz, 1H), 7.43 (t, *J* = 7.6 Hz, 2H), 7.32 (tt, *J* = 7.3, 1.5 Hz, 1H), 6.99 (dd, *J* = 9.3, 1.7 Hz, 1H), 2.26 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 145.1, 144.6, 133.7, 128.7, 128.1, 127.9, 125.9, 123.4, 122.2, 116.6, 108.0, 18.1. HRMS (ESI-TOF) *m/z*: [*M* + *H*]⁺ calcd for $\text{C}_{14}\text{H}_{13}\text{N}_2$, 209.1079; found, 209.1075.

7-Bromo-2-(*p*-tolyl)imidazo[1,2-*a*]pyridine (2n). Yield: 1364 mg, 95%; yellow solid; mp 177–180 °C [lit.^{12e} mp 280 °C]. ^1H NMR

(400 MHz, CDCl_3): δ 8.04 (dd, J = 7.1, 1.0 Hz, 1H), 7.89 (d, J = 2.0 Hz, 1H), 7.86 (s, 1H), 7.82 (d, J = 8.2 Hz, 2H), 7.25 (d, J = 7.9 Hz, 2H), 6.93 (dd, J = 7.1, 1.9 Hz, 1H), 2.41 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 146.3, 145.5, 138.4, 130.0, 129.6, 126.0, 125.8, 119.4, 118.5, 116.5, 108.0, 21.4. HRMS (ESI-TOF): $[M + H]^+$ calcd for $\text{C}_{14}\text{H}_{12}\text{BrN}_2$, 287.0184; found, 287.0176.

2-Phenyl-3-(trifluoromethyl)imidazo[1,2-*a*]pyridine (3a). Yield: 38 mg, 72%; yellow oil; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (300 MHz, CDCl_3): δ 8.32 (d, J = 7.0 Hz, 1H), 7.69–7.76 (m, 3H), 7.43–7.51 (m, 3H), 7.39 (t, J = 7.9 Hz, 1H), 6.99 (t, J = 6.8 Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3): δ 148.0 (q, J = 1.3 Hz), 146.1, 132.9, 129.6 (q, J = 1.1 Hz), 128.9, 128.2, 127.0, 125.5 (q, J = 3.8 Hz), 121.9 (q, J = 265.6 Hz), 118.1, 113.9, 109.8 (q, J = 39.2 Hz). HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{14}\text{H}_{10}\text{F}_3\text{N}_2$, 263.0796; found, 263.0790.

2-(4-Methoxyphenyl)-3-(trifluoromethyl)imidazo[1,2-*a*]pyridine (3b). Yield: 50 mg, 86%; light yellow solid; mp 78–80 °C [lit.⁹ mp 81–83 °C]; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (400 MHz, CDCl_3): δ 8.33 (d, J = 6.8 Hz, 1H), 7.78 (d, J = 9.1 Hz, 1H), 7.68 (d, J = 8.3 Hz, 2H), 7.42 (dt, J = 8.0, 0.8 Hz, 1H), 7.02 (d, J = 8.8 Hz, 2H), 7.01 (t, J = 7.3 Hz, 1H), 3.90 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 160.4, 147.6, 145.8, 130.9, 127.2, 125.5 (q, J = 3.1 Hz), 124.8, 122.0 (q, J = 265.6 Hz), 117.8, 114.0, 113.7, 109.1 (q, J = 40.5 Hz), 55.4. ^{19}F NMR (376 MHz, CDCl_3): δ -57.6. HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{15}\text{H}_{12}\text{F}_3\text{N}_2\text{O}$, 293.0902; found, 293.0895.

2-(4-Fluorophenyl)-3-(trifluoromethyl)imidazo[1,2-*a*]pyridine (3c). Yield: 47 mg, 84%; light yellow solid; mp 78–80 °C [lit.⁹ mp 55–57 °C]; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (300 MHz, CDCl_3): δ 8.32 (d, J = 7.0 Hz, 1H), 7.66–7.75 (m, 3H), 7.37–7.43 (m, 1H), 7.13–7.19 (m, 2H), 7.00 (td, J = 7.5, 1.3 Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3): δ 163.3 (d, J = 247.1 Hz), 147.0 (q, J = 1.8 Hz), 146.1, 131.4 (dq, J = 8.7, 1.3 Hz), 129.0 (d, J = 3.3 Hz), 127.1, 125.5 (q, J = 3.8 Hz), 121.8 (q, J = 265.6 Hz), 118.1, 115.2 (d, J = 21.3 Hz), 114.0, 109.5 (q, J = 39.2 Hz). HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{14}\text{H}_9\text{F}_4\text{N}_2$, 281.0702; found, 281.0694.

2-(4-Bromophenyl)-3-(trifluoromethyl)imidazo[1,2-*a*]pyridine (3d). Yield: 55 mg, 81%; light yellow solid; mp 76–78 °C [lit.⁹ mp 94–96 °C]; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (300 MHz, CDCl_3): δ 8.30 (d, J = 7.0 Hz, 1H), 7.72 (d, J = 8.7 Hz, 1H), 7.55–7.61 (m, 4H), 7.39 (t, J = 8.0 Hz, 1H), 6.99 (t, J = 7.0 Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3): δ 146.8 (q, J = 1.8 Hz), 146.1, 131.8, 131.5, 131.1 (q, J = 1.3 Hz), 127.2, 125.5 (q, J = 3.1 Hz), 123.5, 121.8 (q, J = 265.6 Hz), 118.2, 114.2, 109.6 (q, J = 39.5 Hz). HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{14}\text{H}_9\text{BrF}_3\text{N}_2$, 340.9901; found, 340.9896.

2-(4-Chlorophenyl)-3-(trifluoromethyl)imidazo[1,2-*a*]pyridine (3e). Yield: 48 mg, 81%; light yellow solid; mp 93–95 °C [lit.⁹ mp 84–86 °C]; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (300 MHz, CDCl_3): δ 8.31 (d, J = 7.0 Hz, 1H), 7.73 (d, J = 9.1 Hz, 1H), 7.64 (d, J = 8.3 Hz, 2H), 7.44 (d, J = 8.3 Hz, 2H), 7.40 (d, J = 8.4 Hz, 1H), 7.02 (t, J = 6.9 Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3): δ 146.8, 146.1, 135.2, 131.3, 130.9, 128.5, 127.2, 125.5 (q, J = 4.0 Hz), 121.8 (q, J = 265.9 Hz), 118.1, 114.1, 109.6 (q, J = 39.7 Hz). HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{14}\text{H}_9\text{ClF}_3\text{N}_2$, 297.0406; found, 297.0405.

4-(3-(Trifluoromethyl)imidazo[1,2-*a*]pyridin-2-yl)benzonitrile (3f). Yield: 50 mg, 87%; light yellow solid; mp 163–165 °C [lit.⁹ mp 147–148 °C]; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (300 MHz, CDCl_3): δ 8.33 (d, J = 7.0 Hz, 1H), 7.82 (d, J = 8.3 Hz, 2H), 7.75 (d, J = 8.3 Hz, 2H), 7.73 (d, J = 7.8 Hz, 1H), 7.44 (t, J = 7.9 Hz, 1H), 7.06 (t, J = 6.9 Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3): δ 146.3, 145.8 (q, J = 1.6 Hz), 137.5, 132.1, 130.4, 127.5, 125.6 (q, J = 3.2 Hz), 121.6 (q, J = 266.2 Hz), 118.6, 118.4, 114.7, 112.7, 110.2 (q, J = 39.8 Hz). HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{15}\text{H}_8\text{F}_3\text{N}_3$, 288.0749; found, 288.0742.

2-(4-Nitrophenyl)-3-(trifluoromethyl)imidazo[1,2-*a*]pyridine (3g). Yield: 50 mg, 81%; light yellow solid; mp 145–147 °C [lit.⁹ mp 139–140 °C]; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (300 MHz, CDCl_3): δ 8.34 (d, J = 6.8 Hz, 1H), 8.32 (d, J = 8.9 Hz, 2H),

7.89 (d, J = 8.9 Hz, 2H), 7.77 (d, J = 9.1 Hz, 1H), 7.43–7.49 (m, 1H), 7.07 (dt, J = 7.0, 1.5 Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3): δ 148.1, 146.3, 145.4 (q, J = 2.2 Hz), 139.2, 130.7 (q, J = 1.5 Hz), 127.7, 125.6 (q, J = 3.8 Hz), 123.4, 121.5 (q, J = 268.1 Hz), 118.4, 114.7, 110.5 (q, J = 39.5 Hz). HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{14}\text{H}_9\text{F}_3\text{N}_3\text{O}_2$, 308.0647; found, 308.0642.

6-Methyl-2-phenyl-3-(trifluoromethyl)imidazo[1,2-*a*]pyridine (3h). Yield: 44 mg, 80%; light yellow solid; mp 73–75 °C [lit.¹⁴ mp 97–98 °C]; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (300 MHz, CDCl_3): δ 8.09 (s, 1H), 7.69 (d, J = 7.6 Hz, 2H), 7.64 (d, J = 9.3 Hz, 1H), 7.44–7.50 (m, 3H), 7.24 (d, J = 9.1 Hz, 1H), 2.41 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3): δ 147.7 (q, J = 1.5 Hz), 145.1, 133.0, 130.1, 129.6 (q, J = 1.2 Hz), 128.8, 128.1, 123.8, 123.1 (q, J = 3.4 Hz), 121.9 (q, J = 265.7 Hz), 117.4, 109.2 (q, J = 38.6 Hz), 18.4. HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{15}\text{H}_{12}\text{F}_3\text{N}_2$, 277.0953; found, 277.0947.

2-(4-Bromophenyl)-6-methyl-3-(trifluoromethyl)imidazo[1,2-*a*]pyridine (3i). Yield: 59 mg, 83%; white solid; mp 130–132 °C; [silica gel (8:1 hexane/EtOAc)]. ^1H NMR (300 MHz, CDCl_3): δ 8.08 (s, 1H), 7.62 (d, J = 8.3 Hz, 2H), 7.53–7.58 (m, 3H), 7.25 (d, J = 9.1 Hz, 1H), 2.41 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3): δ 146.5 (q, J = 1.8 Hz), 145.2, 132.0, 131.8, 131.4 (q, J = 2.2 Hz), 131.2, 130.3, 124.1, 123.3, 121.8 (q, J = 265.9 Hz), 117.4, 109.5 (q, J = 38.0 Hz), 18.4. HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{15}\text{H}_{11}\text{BrF}_3\text{N}_2$, 355.0058; found, 355.0051.

6-Bromo-2-phenyl-3-(trifluoromethyl)imidazo[1,2-*a*]pyridine (3j). Yield: 59 mg, 86%; white solid; mp 110–112 °C [lit.¹⁴ mp 104–106 °C]; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (300 MHz, CDCl_3): δ 8.45 (s, 1H), 7.67–7.71 (m, 2H), 7.64 (d, J = 9.5 Hz, 1H), 7.44–7.50 (m, 4H). $^{13}\text{C}\{^1\text{H}\}$ NMR (75 MHz, CDCl_3): δ 148.6 (q, J = 1.9 Hz), 144.5, 132.3, 130.5, 129.6 (q, J = 1.6 Hz), 129.4, 128.2, 125.6 (q, J = 4.2 Hz), 121.6 (q, J = 266.2 Hz), 118.7, 109.9 (q, J = 39.9 Hz), 108.8. HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{14}\text{H}_9\text{BrF}_3\text{N}_2$, 340.9901; found, 340.9893.

6-Bromo-2-(4-chlorophenyl)-3-(trifluoromethyl)imidazo[1,2-*a*]pyridine (3k). Yield: 67 mg, 89%; white solid; mp 150–152 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (400 MHz, CDCl_3): δ 8.46 (s, 1H), 7.64 (d, J = 9.2 Hz, 3H), 7.45–7.51 (m, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 147.4 (q, J = 1.5 Hz), 144.6, 135.5, 130.8 (q, J = 1.3 Hz), 129.1, 128.6, 127.5, 125.6 (q, J = 3.9 Hz), 122.8 (q, J = 266.5 Hz), 118.6, 110.0 (q, J = 39.4 Hz), 109.0. HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{14}\text{H}_8\text{BrClF}_3\text{N}_2$, 374.9511; found, 374.9501.

6-Bromo-2-(4-bromophenyl)-3-(trifluoromethyl)imidazo[1,2-*a*]pyridine (3l). Yield: 76 mg, 90%; white solid; mp 159–161 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (400 MHz, CDCl_3): δ 8.46 (s, 1H), 7.61–7.66 (m, 3H), 7.57 (d, J = 8.6 Hz, 2H), 7.49 (dd, J = 9.5, 2.0 Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 147.3 (q, J = 1.3 Hz), 144.6, 131.6, 131.3, 131.1 (q, J = 1.3 Hz), 130.9, 125.6 (q, J = 3.7 Hz), 123.8, 121.5 (q, J = 266.5 Hz), 118.7, 109.9 (q, J = 39.4 Hz), 109.0. HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{14}\text{H}_8\text{Br}_2\text{F}_3\text{N}_2$, 418.9006; found, 418.9007.

4-(6-Bromo-3-(trifluoromethyl)imidazo[1,2-*a*]pyridin-2-yl)benzonitrile (3m). Yield: 64 mg, 87%; white solid; mp 179–172 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (400 MHz, CDCl_3): δ 8.48 (s, 1H), 7.83 (d, J = 8.6 Hz, 2H), 7.78 (d, J = 8.4 Hz, 2H), 7.67 (d, J = 9.6 Hz, 1H), 7.53 (dd, J = 9.5, 1.8 Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 146.3 (q, J = 1.4 Hz), 144.7, 136.9, 132.1, 131.3, 130.2 (q, J = 1.2 Hz), 125.6 (q, J = 3.6 Hz), 121.6 (q, J = 266.2 Hz), 118.8, 118.5, 113.0, 110.4 (q, J = 40.2 Hz), 109.5. HRMS (ESI-TOF) m/z : $[M + H]^+$ calcd for $\text{C}_{15}\text{H}_8\text{BrF}_3\text{N}_3$, 365.9854; found, 365.9846.

7-Bromo-2-(*p*-tolyl)-3-(trifluoromethyl)imidazo[1,2-*a*]pyridine (3n). Yield: 58 mg, 82%; white solid; mp 118–120 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (400 MHz, CDCl_3): δ 8.19 (d, J = 7.1 Hz, 1H), 7.96–7.98 (d, J = 1.0 Hz, 1H), 7.61 (d, J = 8.1 Hz, 2H), 7.31 (d, J = 8.0 Hz, 2H), 7.14 (dd, J = 7.3 Hz, 1.9 Hz, 1H), 2.45 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 148.4, 145.9, 139.5, 129.5, 129.1, 129.0, 125.7 (q, J = 3.2 Hz), 121.6 (q, J = 263.3 Hz), 121.5, 120.1, 118.1, 109.7 (q, J = 39.2 Hz), 21.4. ^{19}F NMR (376 MHz,

CDCl₃): δ -57.4. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₅H₁₁BrF₃N₂, 355.0058; found, 355.0053.

7-Bromo-2-(4-chlorophenyl)-3-(trifluoromethyl)imidazo[1,2-a]-pyridine (30). Yield: 64 mg, 85%; white solid; mp 158–160 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ¹H NMR (400 MHz, CDCl₃): δ 8.19 (d, J = 7.4 Hz, 1H), 7.97 (d, J = 1.0 Hz, 1H), 7.65 (d, J = 8.0 Hz, 2H), 7.47 (d, J = 8.5 Hz, 2H), 7.17 (dd, J = 7.4 Hz, 1.9 Hz, 1H). ¹³C{¹H} NMR (100 MHz, CDCl₃): δ 147.1, 146.0, 135.7, 130.9, 130.4, 128.7, 125.7 (q, J = 3.1 Hz), 121.5 (q, J = 265.4 Hz), 121.9, 120.2, 118.4, 110.0 (q, J = 39.6 Hz). ¹⁹F NMR (376 MHz, CDCl₃): δ -50.5. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₄H₈BrClF₃N₂, 374.9511; found, 374.9509.

2-(4-Bromophenyl)-6-chloro-3-(trifluoromethyl)imidazo[1,2-a]-pyridine (3p). Yield: 65 mg, 86%; white solid; mp 141–142 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ¹H NMR (500 MHz, CDCl₃): δ 8.34 (s, 1H), 7.68 (d, J = 9.5 Hz, 1H), 7.60 (d, J = 8.5 Hz, 2H), 7.55 (d, J = 8.3 Hz, 2H), 7.38 (dd, J = 9.5, 1.5 Hz, 1H). ¹³C{¹H} NMR (125 MHz, CDCl₃): δ 147.4 (q, J = 2.8 Hz), 144.4, 132.2, 131.53, 131.52, 131.0 (q, J = 1.9 Hz), 128.8, 123.6 (q, J = 270.7 Hz), 123.5 (q, J = 3.8 Hz), 122.7, 118.3, 110.1 (q, J = 44.1 Hz). ¹⁹F NMR (470 MHz, CDCl₃): δ -57.7. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₄H₈BrClF₃N₂, 374.9511; found, 374.9505.

6-Chloro-2-(4-chlorophenyl)-3-(trifluoromethyl)imidazo[1,2-a]-pyridine (3q). Yield: 54 mg, 82%; white solid; mp 90–92 °C [lit.¹⁴ mp 83–85 °C]; [silica gel (8:1 *n*-hexane/EtOAc)]. ¹H NMR (500 MHz, CDCl₃): δ 8.34 (s, 1H), 7.69 (d, J = 9.5 Hz, 1H), 7.62 (d, J = 8.3 Hz, 2H), 7.44 (d, J = 8.3 Hz, 2H), 7.38 (d, J = 9.4 Hz, 1H). ¹³C{¹H} NMR (125 MHz, CDCl₃): δ 147.4 (q, J = 2.4 Hz), 144.4, 135.5, 130.8 (q, J = 1.5 Hz), 130.7, 128.8, 128.6, 123.6 (q, J = 266.0 Hz), 123.4 (q, J = 3.9 Hz), 122.6, 118.3, 110.2 (q, J = 40.9 Hz). ¹⁹F NMR (470 MHz, CDCl₃): δ -57.7. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₄H₈Cl₂F₃N₂, 331.0017; found, 331.0011.

2-(4-Chlorophenyl)-3-(trifluoromethyl)benzo[d]imidazo[2,1-b]-thiazole (3a'). Yield: 59 mg, 83%; white solid; mp 174–176 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ¹H NMR (500 MHz, CDCl₃): δ 7.91 (d, J = 8.4 Hz, 1H), 7.74 (d, J = 8.0 Hz, 1H), 7.58 (d, J = 8.4 Hz, 2H), 7.51 (t, J = 8.0 Hz, 1H), 7.40–7.44 (m, 3H). ¹³C{¹H} NMR (125 MHz, CDCl₃): δ 151.0, 148.7 (q, J = 2.5 Hz), 135.1, 132.0, 131.2, 130.8 (q, J = 1.5 Hz), 130.1, 128.4, 126.9, 125.7, 124.3, 121.3 (q, J = 266.3 Hz), 114.8 (q, J = 4.6 Hz), 112.6 (q, J = 40.8 Hz). ¹⁹F NMR (470 MHz, CDCl₃): δ -55.2. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₆H₉ClF₃N₂S, 353.0127; found, 353.0129.

2-(4-Bromophenyl)-3-(trifluoromethyl)benzo[d]imidazo[2,1-b]-thiazole (3b'). Yield: 67 mg, 84%; white solid; mp 191–193 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ¹H NMR (500 MHz, CDCl₃): δ 7.92 (d, J = 8.4 Hz, 1H), 7.75 (d, J = 8.0 Hz, 1H), 7.58 (d, J = 8.4 Hz, 2H), 7.51–7.54 (m, 3H), 7.43 (t, J = 7.6 Hz, 1H). ¹³C{¹H} NMR (125 MHz, CDCl₃): δ 151.0, 148.7 (q, J = 1.2 Hz), 132.0, 131.6, 131.4, 131.1, 130.1 (q, J = 2.7 Hz), 126.9, 125.8, 124.4, 123.4, 121.3 (q, J = 263.5 Hz), 114.8, 112.6 (q, J = 38.0 Hz). ¹⁹F NMR (470 MHz, CDCl₃): δ -55.1. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₆H₈BrF₃N₂S, 396.9622; found, 396.9619.

2-(4-Chlorophenyl)quinoxaline (4b). Yield: 1119 mg, 93%; yellow solid; mp 130–132 °C [lit.^{13d} mp 135–137 °C]. ¹H NMR (400 MHz, CDCl₃): δ 9.38 (s, 1H), 8.25–8.31 (m, 2H), 8.22 (d, J = 8.5 Hz, 2H), 7.84–7.93 (m, 2H), 7.60 (d, J = 8.5 Hz, 2H). ¹³C{¹H} NMR (100 MHz, CDCl₃): δ 150.8, 142.4, 141.9, 140.4, 137.0, 134.7, 131.0, 130.9, 130.4, 129.6, 128.9, 128.5. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₄H₁₀ClN₂, 241.0533; found, 241.0529.

2-(4-Fluorophenyl)quinoxaline (4c). Yield: 1017 mg, 91%; yellow solid; mp 119–120 °C [lit.^{13d} mp 119–121 °C]. ¹H NMR (400 MHz, CDCl₃): δ 9.33 (s, 1H), 8.25 (d, J = 9.0 Hz, 1H), 8.24 (d, J = 9.0 Hz, 1H), 8.14–8.19 (m, 2H), 7.76–7.86 (m, 2H), 7.29 (t, J = 8.7 Hz, 2H). ¹³C{¹H} NMR (100 MHz, CDCl₃): δ 164.3 (d, J = 249.1 Hz), 150.8, 142.9, 142.3, 141.4, 132.9, 130.5, 129.7, 129.6, 129.5 (d, J = 8.7 Hz), 129.1, 116.3 (d, J = 21.7 Hz). HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₄H₁₀FN₂, 225.0828; found, 225.0826.

2-(4-(Trifluoromethyl)phenyl)quinoxaline (4d). Yield: 1289 mg, 94%; yellow solid; mp 147–148 °C [lit.^{13d} mp 143–144 °C]. ¹H NMR (400 MHz, CDCl₃): δ 9.37 (s, 1H), 8.34 (d, J = 8.1 Hz, 2H),

8.18 (t, J = 8.0 Hz, 2H), 7.79–7.86 (m, 4H). ¹³C{¹H} NMR (100 MHz, CDCl₃): δ 150.3, 142.7, 142.3, 141.7, 140.0, 132.0 (q, J = 32.5 Hz), 130.8, 130.4, 129.8, 129.1, 127.8, 126.1, 124.0 (q, J = 270.8 Hz). HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₅H₁₀F₃N₂, 275.0796; found, 275.0798.

6,7-Dimethyl-2-phenylquinoxaline (4g). Yield: 1043 mg, 89%; yellow solid; mp 125–127 °C [lit.^{13d} mp 130–131 °C]. ¹H NMR (400 MHz, CDCl₃): δ 9.25 (s, 1H), 8.18–8.22 (m, 2H), 7.96 (s, 1H), 7.91 (s, 1H), 7.53–7.62 (m, 3H), 2.55 (s, 6H). ¹³C{¹H} NMR (100 MHz, CDCl₃): δ 151.1, 141.5, 141.4, 141.3, 140.9, 139.4, 136.7, 130.2, 129.2, 128.6, 127.4, 20.5, 20.4. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₆H₁₅N₂, 235.1235; found, 235.1226.

6,7-Dimethyl-2-(4-(trifluoromethyl)phenyl)quinoxaline (4h). Yield: 1390 mg, 92%; yellow solid; mp 120–123 °C [lit.^{13e} mp 109–111 °C]. ¹H NMR (400 MHz, CDCl₃): δ 9.24 (s, 1H), 8.30 (d, J = 8.1 Hz, 2H), 7.91 (d, J = 5.0 Hz, 2H), 7.81 (d, J = 8.2 Hz, 2H), 2.54 (s, 6H). ¹³C{¹H} NMR (100 MHz, CDCl₃): δ 149.4, 141.6, 141.5, 141.3, 141.2, 140.1, 131.6 (q, J = 32.5 Hz), 128.7, 127.7, 127.6, 126.0, 123.9 (q, J = 269.7 Hz), 20.5, 20.4. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₇H₁₄F₃N₂, 303.1109; found, 303.1107.

6,7-Dimethyl-2-(4-nitrophenyl)quinoxaline (4i). Yield: 1271 mg, 91%; yellow solid; mp 135–137 °C [lit.^{13f} mp 197–200 °C]. ¹H NMR (400 MHz, CDCl₃): δ 9.30 (s, 1H), 8.33 (d, J = 8.0 Hz, 2H), 8.00–8.03 (m, 2H), 7.85 (d, J = 8.2 Hz, 2H), 2.57 (s, 6H). ¹³C{¹H} NMR (100 MHz, CDCl₃): δ 149.6, 142.0, 141.9, 141.5, 140.7, 140.0, 139.4, 128.7, 127.7, 127.3, 126.1, 20.52, 20.51. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₆H₁₄N₃O₂, 280.1086; found, 280.1087.

2-Phenyl-8-(trifluoromethyl)quinoxaline (5a). Yield: 31 mg, 56%; white solid; mp 120–122 °C; [silica gel (8:1 hexane/EtOAc)]. ¹H NMR (300 MHz, CDCl₃): δ 9.46 (s, 1H), 8.29–8.35 (m, 3H), 8.14 (d, J = 7.4 Hz, 1H), 7.79 (t, J = 8.0 Hz, 1H), 7.55–7.63 (m, 3H). ¹³C{¹H} NMR (75 MHz, CDCl₃): δ 151.6, 143.6, 141.4, 139.3, 135.9, 133.5, 130.8 (q, J = 4.2 Hz), 129.3, 128.8, 128.5, 127.8 (q, J = 29.9 Hz), 127.7, 123.7 (q, J = 266.6 Hz). HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₅H₁₀F₃N₂, 275.0796; found, 275.0790.

2-(4-Chlorophenyl)-8-(trifluoromethyl)quinoxaline (5b). Yield: 33 mg, 53%; yellow solid; mp 150–153 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ¹H NMR (300 MHz, CDCl₃): δ 9.43 (s, 1H), 8.31 (d, J = 8.5 Hz, 1H), 8.27 (d, J = 8.6 Hz, 2H), 8.15 (d, J = 7.2 Hz, 1H), 7.81 (t, J = 7.9 Hz, 1H), 7.57 (d, J = 8.7 Hz, 2H). ¹³C{¹H} NMR (100 MHz, CDCl₃): δ 150.5, 143.2, 141.5, 139.1, 137.3, 134.3, 133.5, 129.5, 128.9, 128.7 (q, J = 3.9 Hz), 128.2, 128.0 (q, J = 29.9 Hz), 123.6 (q, J = 263.0 Hz). ¹⁹F NMR (376 MHz, CDCl₃): δ -59.6. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₅H₉ClF₃N₂, 309.0406; found, 309.0401.

2-(4-Fluorophenyl)-8-(trifluoromethyl)quinoxaline (5c). Yield: 29 mg, 50%; yellow solid; mp 181–183 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ¹H NMR (400 MHz, CDCl₃): δ 9.45 (s, 1H), 8.30–8.38 (m, 3H), 8.17 (d, J = 7.3 Hz, 1H), 7.82 (t, J = 7.8 Hz, 1H), 7.30 (dd, J = 7.6, 5.0 Hz, 2H). ¹³C{¹H} NMR (100 MHz, CDCl₃): δ 164.7 (d, J = 250.2 Hz), 150.6, 143.2, 141.2, 133.4, 132.1 (d, J = 1.8 Hz), 129.8 (d, J = 8.6 Hz), 128.7 (q, J = 5.2 Hz), 128.2, 128.2 (q, J = 29.2 Hz), 128.1, 122.3 (q, J = 271.0 Hz), 116.4 (d, J = 21.7 Hz). ¹⁹F NMR (376 MHz, CDCl₃): δ -59.6, δ -109.2. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₅H₉F₄N₂, 293.0702; found, 293.0692.

8-(Trifluoromethyl)-2-(4-(trifluoromethyl)phenyl)quinoxaline (5d). Yield: 36 mg, 53%; white solid; mp 159–161 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ¹H NMR (400 MHz, CDCl₃): δ 9.48 (s, 1H), 8.43 (d, J = 8.1 Hz, 2H), 8.35 (d, J = 8.4 Hz, 1H), 8.19 (d, J = 7.2 Hz, 1H), 7.84–7.88 (m, 3H). ¹³C{¹H} NMR (100 MHz, CDCl₃): δ 150.1, 143.4, 141.8, 139.2, 139.1, 133.6, 132.5 (q, J = 32.5 Hz), 128.9 (q, J = 5.4 Hz), 128.7, 128.3 (q, J = 30.1 Hz), 128.0, 126.2 (q, J = 3.5 Hz), 123.9 (q, J = 271.8 Hz), 122.2 (q, J = 272.4 Hz). ¹⁹F NMR (376 MHz, CDCl₃): δ -59.7, -62.9. HRMS (ESI-TOF) m/z : [M + H]⁺ calcd for C₁₆H₈F₆N₂, 343.0670; found, 343.0661. Single crystal was obtained using EtOAc/*n*-hexane (5:1).

2-(4-Nitrophenyl)-8-(trifluoromethyl)quinoxaline (5e). Yield: 34 mg, 53%; yellow solid; mp 218–220 °C; [silica gel (8:1 hexane/EtOAc)]. ¹H NMR (400 MHz, CDCl₃): δ 9.54 (s, 1H), 8.52 (d, J = 9.2 Hz, 2H), 8.47 (d, J = 9.0 Hz, 2H), 8.39 (dd, J = 8.5, 0.8 Hz, 1H),

8.23 (d, $J = 7.3$ Hz, 1H), 7.91 (t, $J = 7.8$ Hz, 1H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 149.2, 143.3, 141.9, 141.6, 139.0, 133.6, 129.3, 129.1 (q, $J = 4.7$ Hz), 128.6, 128.5 (q, $J = 30.0$ Hz), 128.4, 124.4, 123.5 (q, $J = 272.5$ Hz). ^{19}F NMR (376 MHz, CDCl_3): δ -59.7. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{15}\text{H}_9\text{F}_3\text{N}_3\text{O}_2$, 320.0647; found, 320.0645.

4-(8-(Trifluoromethyl)quinoxalin-2-yl)benzonitrile (5f). Yield: 31 mg, 52%; white solid; mp 250–252 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (400 MHz, CDCl_3): δ 9.51 (s, 1H), 8.46 (td, $J = 8.4$, 1.7 Hz, 2H), 8.38 (dd, $J = 8.4$, 0.6 Hz, 1H), 8.22 (d, $J = 7.2$ Hz, 1H), 7.92 (d, $J = 7.8$ Hz, 2H), 7.90 (t, $J = 8.5$ Hz, 1H). ^{19}F NMR (376 MHz, CDCl_3): δ -59.7. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{16}\text{H}_9\text{F}_3\text{N}_3$, 300.0749; found, 300.0748.

6,7-Dimethyl-2-phenyl-8-(trifluoromethyl)quinoxaline (5g). Yield: 33 mg, 55%; white solid; mp 149–151 °C; [silica gel (8:1 hexane/EtOAc)]. ^1H NMR (400 MHz, CDCl_3): δ 9.34 (s, 1H), 8.29–8.33 (m, 2H), 8.06 (s, 1H), 7.54–7.62 (m, 3H), 2.68 (q, $J = 2.9$ Hz, 3H), 2.60 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 150.2, 142.2, 141.5, 140.4, 139.2 (q, $J = 4.3$ Hz), 136.4, 132.1, 130.5, 129.4, 129.2, 127.5, 125.5 (q, $J = 27.6$ Hz), 125.2 (q, $J = 276.0$ Hz), 21.8, 17.5 (q, $J = 4.7$ Hz). ^{19}F NMR (376 MHz, CDCl_3): δ -50.7. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{14}\text{F}_3\text{N}_2$, 303.1109; found, 303.1104.

6,7-Dimethyl-8-(trifluoromethyl)-2-(4-(trifluoromethyl)phenyl)-quinoxaline (5h). Yield: 39 mg, 53%; white solid; mp 142–145 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (400 MHz, CDCl_3): δ 9.38 (s, 1H), 8.43 (d, $J = 8.1$ Hz, 2H), 8.12 (s, 1H), 7.86 (d, $J = 8.1$ Hz, 2H), 2.71 (q, $J = 3.0$ Hz, 3H), 2.64 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (125 MHz, CDCl_3): δ 148.6, 142.0, 141.9, 141.2, 139.7, 139.6, 132.2, 131.9, 127.7, 126.0 (q, $J = 3.6$ Hz), 125.8 (q, $J = 32.4$ Hz), 125.6 (q, $J = 28.2$ Hz), 124.0 (q, $J = 276.0$ Hz), 123.9 (q, $J = 271.0$ Hz), 21.8 (q, $J = 2.8$ Hz), 17.4 (q, $J = 4.5$ Hz). ^{19}F NMR (376 MHz, CDCl_3): δ -50.8, -62.7. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{18}\text{H}_{13}\text{F}_6\text{N}_2$, 371.0983; found, 371.0977.

6,7-Dimethyl-2-(4-nitrophenyl)-8-(trifluoromethyl)quinoxaline (5i). Yield: 33 mg, 48%; yellow solid; mp 219–221 °C; [silica gel (8:1 *n*-hexane/EtOAc)]. ^1H NMR (400 MHz, CDCl_3): δ 9.42 (s, 1H), 8.50 (d, $J = 9.1$ Hz, 2H), 8.45 (d, $J = 9.1$ Hz, 2H), 8.13 (s, 1H), 2.27 (q, $J = 2.9$ Hz, 3H), 2.65 (s, 3H). $^{13}\text{C}\{^1\text{H}\}$ NMR (125 MHz, CDCl_3): δ 148.9, 142.2, 141.9, 141.8, 140.0, 139.9, 132.2, 128.3 (q, $J = 31.0$ Hz), 128.2, 124.9 (q, $J = 3.1$ Hz), 124.3, 123.3 (q, $J = 267.5$ Hz), 120.5, 21.9, 17.5 (q, $J = 3.7$ Hz). ^{19}F NMR (376 MHz, CDCl_3): δ -50.8. HRMS (ESI-TOF) m/z : $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{13}\text{F}_3\text{N}_3\text{O}_2$, 348.0960; found, 348.0945.

■ ASSOCIATED CONTENT

Data Availability Statement

The data underlying this study are available in the published article and its [Supporting Information](#).

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.joc.3c00621>.

^1H , ^{19}F , and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra, and HRMS for compounds **3a–q**, **3a'–b'**, **5a–i** as well as X-ray crystallographic data for compound **5d** (PDF)

Accession Codes

CCDC 2250178 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Author

Saeed Balalaie – Peptide Chemistry Research Institute, K. N. Toosi University of Technology, Tehran 19697-64499, Iran; orcid.org/0000-0002-5764-0442; Email: balalaie@kntu.ac.ir

Authors

Mahdieh Esi Firuz – Peptide Chemistry Research Institute, K. N. Toosi University of Technology, Tehran 19697-64499, Iran

Saideh Rajai-Daryasarei – Peptide Chemistry Research Institute, K. N. Toosi University of Technology, Tehran 19697-64499, Iran; orcid.org/0000-0003-0418-2211

Frank Rominger – Organisch-Chemisches Institut der Universität Heidelberg, Heidelberg 69120, Germany

Abbas Biglari – Department of Chemistry, Institute for Advanced Studies in Basic Sciences (IASBS), Zanjan 45137-66731, Iran

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.joc.3c00621>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Saeed Balalaie thanks Alexander von Humboldt for the research fellowship. Dr. Rajai-Daryasarei gratefully acknowledge the financial support of Basic Sciences Research Fund (no. BSRF-chem-399-08). We thank Iran National Science Foundation (INSF, grant no. 99029802) for the financial support.

■ REFERENCES

- (1) (a) Wang, J.; Sánchez-Roselló, M.; Aceña, J. L.; Del Pozo, C.; Sorochinsky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. Fluorine in pharmaceutical industry: fluorine-containing drugs introduced to the market in the last decade (2001–2011). *Chem. Rev.* **2014**, *114*, 2432–2506. (b) Dong, X.; Wang, W.; Li, H.; Xu, Q.; Ye, L.; Li, X.; Zhao, Z.; Li, X. Stereoselective synthesis of 2-trifluoromethylated and 2-difluoromethylated dihydrofurans via organocatalytic cascade Michael/alkylation reaction. *Org. Chem. Front.* **2021**, *8*, 3260–3267. (c) Sindhe, H.; Chaudhary, B.; Chowdhury, N.; Kamble, A.; Kumar, V.; Lad, A.; Sharma, S. Recent advances in the transition-metal catalyzed directed C–H functionalization with fluorinated building blocks. *Org. Chem. Front.* **2022**, *9*, 1742–1775. (d) Liu, T.; Shen, Q. Progress in copper-mediated formation of trifluoromethylated arenes. *Eur. J. Org. Chem.* **2012**, *2012*, 6679–6687. (e) Wiehn, M. S.; Vinogradova, E. V.; Togni, A. Electrophilic trifluoromethylation of arenes and N-heteroarenes using hypervalent iodine reagents. *J. Fluorine Chem.* **2010**, *131*, 951–957. (f) Lundgren, R. J.; Stradiotto, M. Übergangsmetallkatalysierte trifluormethylierung von arylhalogeniden. *Angew. Chem.* **2010**, *122*, 9510–9512. (g) Shimizu, R.; Egami, H.; Hamashima, Y.; Sodeoka, M. Copper-catalyzed trifluoromethylation of allylsilanes. *Angew. Chem., Int. Ed.* **2012**, *51*, 4577–4580. (h) Jana, S.; Ashokan, A.; Kumar, S.; Verma, A.; Kumar, S. Copper-catalyzed trifluoromethylation of alkenes: synthesis of trifluoromethylated benzoxazines. *Org. Biomol. Chem.* **2015**, *13*, 8411–8415. (i) Jana, S.; Verma, A.; Kadu, R.; Kumar, S. Visible-light-induced oxidant and metal-free dehydrogenative cascade trifluoromethylation and oxidation of 1, 6-enynes with water. *Chem. Sci.* **2017**, *8*, 6633–6644. (2) (a) Zhang, Z.; Zhang, L.; Cao, Y.; Li, F.; Bai, G.; Liu, G.; Yang, Y.; Mo, F. Mn-mediated electrochemical trifluoromethylation/C(sp²)–H functionalization cascade for the synthesis of azahetero-

- cycles. *Org. Lett.* **2019**, *21*, 762–766. (b) Wang, H.; Zhang, J.; Shi, J.; Li, F.; Zhang, S.; Xu, K. Organic photoredox-catalyzed synthesis of δ -fluoromethylated alcohols and amines via 1, 5-hydrogen-transfer radical relay. *Org. Lett.* **2019**, *21*, 5116–5120. (c) Fujiwara, Y.; Dixon, J. A.; O'Hara, F.; Funder, E. D.; Dixon, D. D.; Rodriguez, R. A.; Baxter, R. D.; Herle, B.; Sach, N.; Collins, M. R.; et al. Practical and innate carbon–hydrogen functionalization of heterocycles. *Nature* **2012**, *492*, 95–99. (d) Sun, B.; Huang, P.; Yan, Z.; Shi, X.; Tang, X.; Yang, J.; Jin, C. Self-catalyzed phototandem perfluoroalkylation/cyclization of unactivated alkenes: synthesis of perfluoroalkyl-substituted quinazolinones. *Org. Lett.* **2021**, *23*, 1026–1031. (e) Studer, A. A “renaissance” in radical trifluoromethylation. *Angew. Chem., Int. Ed.* **2012**, *51*, 8950–8958. (f) Barata-Vallejo, S.; Lantaño, B.; Postigo, A. Recent advances in trifluoromethylation reactions with electrophilic trifluoromethylating reagents. *Chem. - Eur. J.* **2014**, *20*, 16806–16829. (g) Eisenreich, F.; Palmans, A. R. Direct C–H trifluoromethylation of (hetero) arenes in water enabled by organic photoredox-active amphiphilic nanoparticles. *Eurasian J. Chem.* **2022**, *28*, No. e202201322.
- (3) (a) Tordeux, M.; Langlois, B.; Wakselman, C. Reactions of bromotrifluoromethane and related halides. 8. Condensations with dithionite and hydroxymethanesulfinate salts. *J. Org. Chem.* **1989**, *54*, 2452–2453. (b) Langlois, B. R.; Laurent, E.; Roidot, N. Trifluoromethylation of aromatic compounds with sodium trifluoromethanesulfinate under oxidative conditions. *Tetrahedron Lett.* **1991**, *32*, 7525–7528. (c) Shen, J.; Li, L.; Xu, J.; Shen, C.; Zhang, P. Recent advances on the application of Langlois' reagent in olefin difunctionalization. *Org. Biomol. Chem.* **2023**, *21*, 2046–2058. (d) Shen, J.; Xu, J.; He, L.; Liang, C.; Li, W. Application of Langlois' reagent (NaSO_2CF_3) in C–H functionalisation. *Chin. Chem. Lett.* **2022**, *33*, 1227–1235. (e) Zhang, C. Application of Langlois' reagent in trifluoromethylation reactions. *Adv. Synth. Catal.* **2014**, *356*, 2895–2906. (f) Wang, J.; Sun, B.; Zhang, L.; Xu, T.; Xie, Y.; Jin, C. Visible-light-induced trifluoromethylation of quinoxalin-2(1H)-ones under photocatalyst-free conditions. *Asian J. Org. Chem.* **2019**, *8*, 1942–1946.
- (4) (a) Liu, H.; Gu, Z.; Jiang, X. Direct trifluoromethylation of the C–H bond. *Adv. Synth. Catal.* **2013**, *355*, 617–626. (b) Wang, L.; Zhang, Y.; Li, F.; Hao, X.; Zhang, H. Y.; Zhao, J. Direct C–H trifluoromethylation of quinoxalin-2 (1H)-ones under transition-metal-free conditions. *Adv. Synth. Catal.* **2018**, *360*, 3969–3977. (c) Yang, X.; Sun, R.; Li, S.; Zheng, X.; Yuan, M.; Xu, B.; Jiang, W.; Chen, H.; Fu, H.; Li, R. Regioselective Direct C–H Trifluoromethylation of Pyridine. *Org. Lett.* **2020**, *22*, 7108–7112. (d) Mi, X.; Kong, Y.; Yang, H.; Zhang, J.; Pi, C.; Cui, X. Visible-light-promoted metal-free C–H trifluoromethylation of imidazopyridines. *Eur. J. Org. Chem.* **2020**, *2020*, 1019–1022. (e) Kumar, S.; Rathore, V.; Verma, A.; Prasad, C. D.; Kumar, A.; Yadav, A.; Jana, S.; Sattar, M.; Kumar, S.; Kumar, S. KOTBu-mediated aerobic transition-metal-free regioselective β -arylation of indoles: synthesis of β -(2-/4-nitroaryl)-indoles. *Org. Lett.* **2015**, *17*, 82–85. (f) Jha, R. K.; Upadhyay, A.; Kanika; Jain, S.; KA, N.; Kumar, S. Light-Driven Carbon–Carbon Coupling of α -sp³–CH of Aliphatic Alcohols with sp²–CH Bond of 1,4-Naphthoquinones. *Org. Lett.* **2022**, *24*, 7605–7610. (g) Hou, H.; Wang, C.; Cheng, X.; Chen, H.; Sun, W.; Zheng, Z.; Ke, F. Visible light-induced direct and highly selective C–H functionalization of quinoxalin-2 (1H)-one without orientating group. *Catal. Sci. Technol.* **2023**, *13*, 305–309. (h) Li, Y.; Song, G.-T.; Tang, D.-Y.; Xu, Z.-G.; Chen, Z.-Z. Acid-promoted direct C–H carbamoylation at the C-3 position of quinoxalin-2 (1H)-ones with isocyanide in water. *ACS Omega* **2023**, *8*, 1577–1587. (i) Sattar, M.; Rathore, V.; Prasad, C. D.; Kumar, S. Transition-metal-free chemoselective oxidative C–C coupling of the sp³ C–H bond of oxindoles with arenes and addition to alkene: synthesis of 3-aryl oxindoles, and benzofuro- and indoloindoles. *Chem.-Asian J.* **2017**, *12*, 734–743.
- (5) (a) Jaso, A.; Zarranz, B.; Aldana, I.; Monge, A. Synthesis of new quinoxaline-2-carboxylate 1, 4-dioxide derivatives as anti-mycobacterium tuberculosis agents. *J. Med. Chem.* **2005**, *48*, 2019–2025. (b) Sarges, R.; Howard, H. R.; Browne, R. G.; Lebel, L. A.; Seymour, P. A.; Koe, B. K. 4-Amino [1, 2, 4] triazolo [4, 3-a] quinoxalines. A novel class of potent adenosine receptor antagonists and potential rapid-onset antidepressants. *J. Med. Chem.* **1990**, *33*, 2240–2254. (c) Loriga, M.; Piras, S.; Sanna, P.; Paglietti, G. Quinoxaline chemistry. Part 7. 2-[aminobenzoates]- and 2-[aminobenzoylglutamate]-quinoxalines as classical antifolate agents. Synthesis and evaluation of in vitro anticancer, anti-HIV and antifungal activity. *Farmaco* **1997**, *52*, 157–166. (d) Rodrigues, F. A.; Bomfim, I. d. S.; Cavalcanti, B. C.; Pessoa, C. d. Ó.; Wardell, J. L.; Wardell, S. M.; Pinheiro, A. C.; Kaiser, C. R.; Nogueira, T. C.; Low, J. N.; et al. Design, synthesis and biological evaluation of (E)-2-(2-arylhydrazinyl) quinoxalines, a promising and potent new class of anticancer agents. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 934–939. (e) Corona, P.; Carta, A.; Loriga, M.; Vitale, G.; Paglietti, G. Synthesis and in vitro antitumor activity of new quinoxaline derivatives. *Eur. J. Med. Chem.* **2009**, *44*, 1579–1591. (f) Rival, Y.; Grassy, G.; Taudou, A.; Ecalte, R. Antifungal activity in vitro of some imidazo [1,2-a] pyrimidine derivatives. *Eur. J. Med. Chem.* **1991**, *26*, 13–18. (g) Samanta, S.; Mondal, S.; Ghosh, D.; Hajra, A. Rhodium-catalyzed directed C–H amidation of imidazoheterocycles with dioxazolones. *Org. Lett.* **2019**, *21*, 4905–4909. (h) Shukla, N. M.; Salunke, D. B.; Yoo, E.; Mutz, C. A.; Balakrishna, R.; David, S. A. Antibacterial activities of Groebke–Blackburn–Bienaymé-derived imidazo [1, 2-a] pyridin-3-amines. *Bioorg. Med. Chem.* **2012**, *20*, 5850–5863. (i) Wen, J.; Niu, C.; Yan, K.; Cheng, X.; Gong, R.; Li, M.; Guo, Y.; Yang, J.; Wang, H. Electrochemical-induced regioselective C-3 thiomethylation of imidazopyridines via a three-component cross-coupling strategy. *Green Chem.* **2020**, *22*, 1129–1133.
- (6) (a) Maikhuri, V. K.; Prasad, A. K.; Jha, A.; Srivastava, S. Recent advances in the transition metal catalyzed synthesis of quinoxalines: a review. *New J. Chem.* **2021**, *45*, 13214–13246. (b) Yashwantrao, G.; Saha, S. Recent advances in the synthesis and reactivity of quinoxaline. *Org. Chem. Front.* **2021**, *8*, 2820–2862. (c) Zhang, X.; Chen, J.; Khan, R.; Shen, G.; He, Z.; Zhou, Y.; Fan, B. Rhodium-catalyzed transfer hydrogenation of quinoxalines with water as a hydrogen source. *Org. Biomol. Chem.* **2019**, *17*, 10142–10147. (d) Ghosh, D.; Ghosh, S.; Hajra, A. Electrochemical functionalization of imidazopyridine and indazole: An overview. *Adv. Synth. Catal.* **2021**, *363*, 5047–5071. (e) Bagdi, A. K.; Santra, S.; Monir, K.; Hajra, A. Synthesis of imidazo [1, 2-a] pyridines: a decade update. *Chem. Commun.* **2015**, *51*, 1555–1575.
- (7) (a) Gernet, A.; Sevrain, N.; Volle, J.-N.; Ayad, T.; Pirat, J.-L.; Virieux, D. Diversity-oriented synthesis toward aryl- and phosphoryl-functionalized imidazo [1,2-a] pyridines. *J. Org. Chem.* **2020**, *85*, 14730–14743. (b) Okai, H.; Tanimoto, K.; Ohkado, R.; Iida, H. Multicomponent synthesis of imidazo [1, 2-a] pyridines: Aerobic oxidative formation of C–N and C–S bonds by flavin–iodine-coupled organocatalysis. *Org. Lett.* **2020**, *22*, 8002–8006. (c) Chaudhary, N. R.; Kapdi, A. R. HFIP promoted thio (hetero) arylation of imidazoheterocycles under metal- and base-free conditions. *Chem. Commun.* **2021**, *57*, 8202–8205. (d) Nipate, D. S.; Jaspal, S.; Shinde, V. N.; Rangan, K.; Kumar, A. TEMPO-mediated cross-dehydrogenative coupling of indoles and imidazo [1, 2-a] pyridines with fluorinated alcohols. *Org. Lett.* **2021**, *23*, 1373–1377. (e) Neto, J. S. S.; Balaguez, R. A.; Franco, M. S.; de Sa Machado, V. C.; Saba, S.; Rafique, J.; Galetto, F. Z.; Braga, A. L. Trihaloisocyanuric acids in ethanol: an eco-friendly system for the regioselective halogenation of imidazo-heteroarenes. *Green Chem.* **2020**, *22*, 3410–3415. (f) Chen, X.; Ding, L.; Li, L.; Li, J.; Zou, D.; Wu, Y.; Wu, Y. Transition metal-free direct CH trifluoromethylation of (hetero) arenes with Togni's reagent. *Tetrahedron Lett.* **2020**, *61*, 151538.
- (8) Monir, K.; Bagdi, A. K.; Ghosh, M.; Hajra, A. Regioselective oxidative trifluoromethylation of imidazoheterocycles via C (sp²)–H bond functionalization. *J. Org. Chem.* **2015**, *80*, 1332–1337.
- (9) Han, S.; Gao, X.; Wu, Q.; Li, J.; Zou, D.; Wu, Y.; Wu, Y. Transition-metal-free direct trifluoromethylation and perfluoroalkylation of imidazopyridines under mild conditions. *Adv. Synth. Catal.* **2019**, *361*, 1559–1563.

(10) Li, M.; Li, G.; Dai, C.; Zhou, W.; Zhan, W.; Gao, M.; Rong, Y.; Tan, Z.; Deng, W. Visible-light-promoted direct C3-trifluoromethylation and perfluoroalkylation of imidazopyridines. *Org. Biomol. Chem.* **2021**, *19*, 8301–8306.

(11) (a) Yuan, J.-W.; Yang, L.-R.; Yin, Q.-Y.; Mao, P.; Qu, L.-B. $\text{KMnO}_4/\text{AcOH}$ -mediated C3-selective direct arylation of coumarins with arylboronic acids. *RSC Adv.* **2016**, *6*, 35936–35944. (b) Demir, A. S.; Findik, H. Potassium permanganate/carboxylic acid/organic solvent: a powerful reagent for enone oxidation and aryl coupling reactions. *Tetrahedron* **2008**, *64*, 6196–6201. (c) Cao, X.-H.; Pan, X.; Zhou, P.-J.; Zou, J.-P.; Asekun, O. T. Manganese (iii)-mediated direct C sp²–H radical trifluoromethylation of coumarins with sodium trifluoromethanesulfonate. *Chem. Commun.* **2014**, *50*, 3359–3362.

(12) (a) Zhang, H.; Wei, Q.; Wei, S.; Qu, J.; Wang, B. Highly efficient and practical thiocyanation of imidazopyridines using an N-chlorosuccinimide/NaSCN combination. *Eur. J. Org. Chem.* **2016**, *2016*, 3373–3379. (b) Godugu, K.; Nallagonda, C. G. R. Solvent and catalyst-free synthesis of imidazo [1, 2-a] pyridines by grindstone chemistry. *J. Heterocycl. Chem.* **2021**, *58*, 250–259. (c) Xue, C.; Han, J.; Zhao, M.; Wang, L. Rapid construction of fused heteropolycyclic aromatics via palladium-catalyzed domino arylations of imidazopyridine derivatives. *Org. Lett.* **2019**, *21*, 4402–4406. (d) Wang, Y.; Li, S.; Wang, X.; Yao, Y.; Feng, L.; Ma, C. A multi pathway coupled domino strategy: I_2/TBHP -promoted synthesis of imidazopyridines and thiazoles via sp³, sp² and sp C–H functionalization. *RSC Adv.* **2022**, *12*, 5919–5927. (e) Quattrini, L.; Gelardi, E. L. M.; Coviello, V.; Sartini, S.; Ferraris, D. M.; Mori, M.; Nakano, I.; Garavaglia, S.; La Motta, C. Imidazo [1, 2-a] pyridine derivatives as aldehyde dehydrogenase inhibitors: Novel chemotypes to target glioblastoma stem cells. *J. Med. Chem.* **2020**, *63*, 4603–4616.

(13) (a) Meshram, H.; Santosh Kumar, G.; Ramesh, P.; Chennakesava Reddy, B. A mild and convenient synthesis of quinoxalines via cyclization–oxidation process using DABCO as catalyst. *Tetrahedron Lett.* **2010**, *51*, 2580–2585. (b) Zhang, R.; Qin, Y.; Zhang, L.; Luo, S. Oxidative synthesis of benzimidazoles, quinoxalines, and benzoxazoles from primary amines by ortho-quinone catalysis. *Org. Lett.* **2017**, *19*, 5629–5632. (c) Nguyen, L. A.; Nguyen, T. T. T.; Ngo, Q. A.; Nguyen, T. B. Sulfur-catalyzed oxidative condensation of aryl alkyl ketones with o-phenylenediamines: Access to quinoxalines. *Adv. Synth. Catal.* **2022**, *364*, 2748–2752. (d) Yang, H. R.; Hu, Z. Y.; Li, X. C.; Wu, L.; Guo, X. X. Cobalt-catalyzed effective access to quinoxalines with insights in annulation of terminal alkynes and o-phenylenediamines. *Org. Lett.* **2022**, *24*, 8392–8396. (e) Wang, X.; Wang, B.; Miao, J.; Chen, Y.; Zhu, G.; Zhang, J.; Wang, J.; Wang, H.; Tang, L.; Wang, C. Direct C-2 arylation of quinoxaline with arylhydrazine salts as arylation reagents. *J. Heterocycl. Chem.* **2022**, *59*, 1272–1279. (f) Huang, T.; Zhang, Q.; Chen, J.; Gao, W.; Ding, J.; Wu, H. Synthesis of quinoxalines catalysed by cetyltrimethyl ammonium bromide (CTAB) in aqueous media. *J. Chem. Res.* **2009**, *2009*, 761–765.

(14) Wei, T.; Wang, K.; Yu, Z.; Hou, J.; Xie, Y. Electrochemically mediated trifluoromethylation of 2H-indazole derivatives using $\text{CF}_3\text{SO}_2\text{Na}$. *Tetrahedron Lett.* **2021**, *86*, 153313.