

Chemoenzymatic Dynamic Kinetic Resolution of Secondary Alcohols and Amines Employing Copper-Based Photocatalysis

Clémentine Minozzi, Nicolas Dowe, Noémie Beaucage, and Shawn K. Collins*

Cite This: *ACS Catal.* 2023, 13, 8347–8353

Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: A tandem photo- and biocatalytic dynamic kinetic resolution of secondary alcohols and amines was achieved employing a commercially available lipase, *Candida antarctica* lipase B (CALB), and a heteroleptic copper complex. The process is an example of using copper-based photocatalysis to promote hydrogen atom transfer (HAT) processes employing thiyl radicals. Screening of 48 complexes and 5 disulfides identified Cu(dtbbpy)(DPEPhos)BF₄ and (Ph₃SiS)₂ as an optimal catalyst system to promote the HAT process. The catalytic system represents a general system for radical-mediated racemization applicable to both aliphatic and aromatic alcohols and amines (18 examples, 60 → 97% yield, 76 → 99% ee).

KEYWORDS: copper, photocatalysis, dynamic kinetic resolution, secondary alcohols, biocatalysis



1. INTRODUCTION

Photocatalysis¹ and biocatalysis² are both regarded as green tools for molecular synthesis that proceed under relatively mild reaction conditions. It is not surprising that the merger of the two fields has garnered significant interest. Where photocatalysis can be exploited to promote redox transformations or generate reactive intermediates, biocatalysis is utilized to perform asymmetric synthesis with often exquisite levels of selectivity. Photo-biocatalysis can involve enzymes that require light for catalytic activity and take advantage of the chiral environment to induce asymmetry.³ Such processes can be used to address long-standing challenges in molecular synthesis, such as enantioselective C_{sp}³–C_{sp}³ couplings.⁴ Photo-biocatalysis can also utilize both individual enzymes and photocatalysts in linear cascades. Such systems often manipulate photocatalysis to engage challenging redox transformations, in tandem with a stereoselective enzymatic step, often, but not exclusively, with co-factors.⁵ The coupling of biocatalysis and transition metal catalysis also provides powerful tools for molecular synthesis. The Bäckvall dynamic kinetic resolution of alcohols is a potent chemoenzymatic strategy for preparing enantioenriched secondary alcohols and amines from racemic precursors.⁶ The process typically employs lipases and proteases as enantioselective acylating agents and transition metal complexes as catalysts to promote racemization of the secondary alcohol or amine. Different catalysts can promote racemization either through sequential oxidation/reduction⁷ or via reversible carbocation formation.⁸

In 1998, Roberts studied an alternative mechanism for racemization via radical chain mechanisms employing thiyl radicals (Figure 1). Alkanethiols substituted by electron-withdrawing groups could racemize (*R*)-tetrahydrofurfuryl


acetate,⁹ and subsequent reports demonstrated that thiyl radicals could also epimerize 1,2-diols.¹⁰ In 2006, Bertrand et al. showed that the same strategy could be employed for the racemization of non-activated aliphatic amines using thio-glycolate/AIBN to thermally generate thiyl radicals.¹¹ Exploiting knowledge of important S–H and C–H bond dissociation energies (BDEs) for designing such processes, a dynamic kinetic resolution (DKR) process was later developed for aliphatic amines.¹²

Recently, photocatalysis has emerged as a mild and efficient route to hydrogen atom transfer chemistry via thiyl radicals. Wendlandt et al. reported that Ir-based photocatalysts under blue light irradiation promote selective, catalytic isomerization of *cis*-1,2-diols to *trans*-diequatorial-1,2-diols employing triphenylsilanethiol (Ph₃SiSH) as a co-catalyst.¹³ The power of the strategy was particularly effective for selective “editing” of stereocenters in complex carbohydrates.¹⁴ Other photocatalytic systems employing decatungstate complexes¹⁵ and thermally promoted processes¹⁶ have also been reported for selective isomerizations via hydrogen atom transfer (HAT) and many other thiyl-mediated HAT processes continue to be reported.¹⁷ Not surprisingly, the first photocatalytic and enzymatic DKR for unactivated aliphatic amines was reported in 2018 by Zhou et al.¹⁸ Using an Ir-based catalyst and *n*-OctSH, a variety of primary aliphatic amines were converted to


Received: April 14, 2023

Revised: May 25, 2023

Published: June 8, 2023



Past Work: Thiyl Radical-Mediated Hydrogen Atom Transfer (HAT)



This Work:




Figure 1. A timeline of thiyl-radical-promoted HAT/isomerization toward photocatalytic chemoenzymatic dynamic kinetic resolution.




Figure 2. Optimization of a heteroleptic copper complex for racemization of secondary alcohols.

their corresponding amides in high yields and enantioselectivities. Despite the ability of photocatalysis to promote mild racemization of stereocenters, it is surprising that its compatibility with biocatalysis in the context of DKR has not been further developed. While radical-based approaches have been studied in the past two decades for aliphatic amines, extensions to aliphatic alcohols and more widely used aromatic substrates are noticeably absent, and investigation of alternative photocatalysts is still in its infancy. Our group has demonstrated that heteroleptic copper complexes^{19,20} of the type Cu(NN)(PP)X can be fine-tuned for various applications in photocatalysis,^{21,22} and it is probable that they could be designed for HAT-type processes. Herein, we report on the design of a heteroleptic copper-based complex for a photocatalytic and chemoenzymatic DKR of secondary alcohols and amines with broad scope.

2. RESULTS AND DISCUSSION

2.1. Optimization of the Catalyst System. To optimize a heteroleptic copper complex for HAT, a series of different diimine (NN) and bisphosphine (PP) ligands were selected for evaluation (Figure 2).

Four bisphosphines were selected: two bisphosphines with wide bite angles, DPEPhos and XantPhos, and two with smaller bite angles, SEGPhos and BINAP. Twelve diimine ligands were selected that had varied electronics, steric effects, and coordination geometry, leading to a total of 48 different catalyst structures. As a starting point, (Ph₃SiS)₂ was selected as a disulfide, given previous reports of the parent silane as an effective HAT promoter.¹⁴ DABCO was chosen as an electron donor and acetonitrile as the solvent, and the reaction mixtures were irradiated under blue LEDs.²³ Efficiency was judged by the degree of racemization of secondary alcohol **1** over 24 h (Figure 2). In examining general trends, heteroleptic






Figure 3. Optimization of the disulfide for the photocatalytic racemization of secondary alcohols.


a Proposed Mechanism



b Photophysical Data (All Potentials vs. SCE)



c Optimization of the DKR



d Stern-Volmer Plots




Figure 4. Proposed mechanism: (a) possible catalytic cycles, (b) catalyst photophysical data, (c) optimization of the resulting DKR of secondary alcohols, and (d) Stern–Volmer plots for the racemization reaction.

Table 1. Photochemical DKR of Secondary Alcohols and Amines Employing the Cu(dtbbpy)(DPEPhos)BF₄ Catalyst^a

$\text{R}^1\text{CH}(\text{X})\text{CH}(\text{R}^2)\text{R}^3$		$\text{H}_{n-1}\text{X}\text{R}^3$	
Cu(dtbbpy)(DPEPhos)BF ₄ (3 mol %) SS1 (30 mol %) DABCO (2 equiv) acyl donor (3 equiv) CALB (20 mg/mmol), MeCN Blue LEDs, 24 h			
1	2	3	4
76 % 99 %ee	90 % 99 %ee	76 % 99 %ee	89 % 99 %ee
5	6	7	8
77 % 99 %ee			
9	10	11	12
60 % 99 %ee	83 % 99 %ee	72 % 76 %ee	80 % 99 %ee
13	14	15	16
60 % 99 %ee	76 % 99 %ee	75 % 96 %ee	81 % 99 %ee
17	18	19	20
82 % 99 %ee			
21	22	23	24
84 % 99 %ee	97 % 99 %ee	84 % 99 %ee	

^aYields following chromatography.

complexes derived from SEGPhos tend to show good levels of racemization of **1** over 24 h (\sim <60% ee of **1**). Previous reports have suggested that bisphosphines with small bite angles tend to afford complexes that are good excited-state electron acceptors.²¹ Among the diimines, four ligands (dtbbpy (4,4'-di-*tert*-butyl-2,2'-dipyridyl), batho (4,7-diphenyl-1,10-phenanthroline), bathocup (2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline), dq (2,2'-biquinoline)) displayed generally good levels of racemization.

Several of the ligands have also been used in other heteroleptic copper complexes for single-electron transfer chemistry. Surprisingly, only two complexes afforded secondary alcohol **1** in less than 20% ee after 24 h, with the complex Cu(dtbbpy)(DPEPhos)BF₄ being the best (10% ee of **1**). The large-bite-angle DPEPhos is well-established to reinforce the tetrahedral geometry of the ground and excited states, enhancing the emissive properties of the complex. While in most cases, the absorption coefficient is lower compared to those of analogous homoleptic diimine complexes, excited-state reduction potentials remain similar, while oxidation potentials are much improved. The sum effect results in complexes that are both efficient reducing and oxidizing agents in their excited states.^{19e} Subsequently, a survey of different bases to replace DABCO was carried out (see inset in Figure

2); however, none was as effective as DABCO. Among the organic and inorganic bases selected, the inorganic bases were largely unproductive, while others that could serve as good electron donors (quinuclidine, TMP, NPh₃) afforded some racemization of alcohol **1** (<50% ee). Note that when the racemization of secondary alcohol **1** with disulfide **SS1** was conducted in the dark, no racemization was observed either at room temperature (79% recovered **1**, 99% ee) or at 40 °C (70% **1**, 99% ee). Similarly, no racemization was observed in the absence of the copper catalyst (83% **1**, 99% ee).²⁴ When disulfide **SS1** was replaced by its corresponding thiol (Ph₃SiSH), the racemization process shut down.²⁵ Next, the effect of the nature of the disulfide was investigated. Five disulfides were selected, namely, the original (Ph₃SiS)₂ **SS1**; two with heteroaromatic substituents (**SS2** and **SS3**); one with aromatic substituents, **SS4** (BDE = 70 kcal/mol); and one with aliphatic substituents, **SS5** (BDE = 86 kcal/mol) (Figure 3).²⁶ The racemization of aliphatic secondary alcohol **1** was investigated at various time points over 24 h to gauge its reactivity (Figure 3, left). The heteroaromatic-substituted disulfides did not produce almost any racemization over a 24 h period. The racemization of alcohol **1** plateaued at approximately 50% ee with aromatic disulfide **SS3**. Encouraging results were obtained with the dicycysteine derivative **SS5**,




arriving at 33% ee of **1** after 24 h. Despite the encouraging reactivity, **SS1** remained the most active. Both **SS1** and **SS5** were then investigated in the racemization of benzylic alcohol **2** (Figure 3, right). Gratifyingly, both **SS5** and **SS1** again produced good levels of racemization of the benzylic alcohol, with **SS1** affording the lowest ee after 24 h (15% ee of **2**).

The racemization of the secondary alcohols is proposed to occur via HAT with a thiyl radical. There are several possibilities for generating thiyl radicals (Figure 4a). Upon irradiation with blue LEDs, Cu(dtbbpy)(DPEPhos)BF₄ is promoted to its excited state. A report from Glorius et al. suggested that disulfides can undergo transformation to thiyl radicals via an energy-transfer mechanism.²⁷ Stern–Volmer plots display no quenching with disulfide **SS1**, or a secondary alcohol, but rather with the DABCO additive. One possibility is that following reductive quenching, subsequent SET from the reduced ground state of Cu(dtbbpy)(DPEPhos)BF₄ to **SS1** generates the thiyl radical (catalytic cycle A, Figure 4a). While some disulfides are reduced at low potentials (~1.6 to –2.8 V),²⁸ the cyclic voltammograms of **SS1** showed a much lower reduction potential ($E_{1/2}$ red = –0.88 V vs SCE). If such reduction were to occur, the thiyl radical could undergo productive reversible HAT to racemize the secondary alcohol. TEMPO trapping of a radical intermediate for a secondary amine substrate was observed via ESI–MS analysis. As no racemization is observed when the corresponding thiol is used in place of the disulfide, this suggests that some mechanism is operative to generate thiyl radicals from the disulfide. Reduction of the disulfide would also result in the formation of the thiolate anion. Such thiolates are good single-electron reducing agents and could also quench the excited state of the copper complex in an alternate catalytic cycle (catalytic cycle B, Figure 4a). In doing so, another equivalent of the thiyl radical would be generated for HAT with the secondary alcohol. With conditions in hand for photochemical racemization, a DKR process was investigated using racemic secondary alcohol **2**, solid-supported *Candida antarctica* lipase B (CALB), and isopropenyl acetate as the acyl donor (Figure 4c). Experimentation revealed that **2** could be converted to (R)-**3** at 73% yield and 99% ee after 24 h. Heating the reaction to 40 °C to promote enzyme activity had no significant effect on the outcome of the DKR. Repeating the DKR in either PhMe or THF resulted in a significant drop in the yield of (R)-**3** (27–48% yield).

2.2. Evaluation of Scope. Following identification of the Cu(dtbbpy)(DPEPhos)BF₄/(Ph₃SiS)₂ (**SS1**) catalytic system, it was evaluated in the biocatalytic/photocatalytic DKR of various alcohols and amines (Table 1). As model secondary alcohol **2** underwent DKR to afford acetate (R)-**3** at 76% isolated yield (99% ee), further evaluation of the tolerance to aromatic substituents was probed. Nitro-substituted acetate **4** was isolated at 90% yield (99% ee), demonstrating that the nitro substituent did not interfere with the photochemistry. Furthermore, the presence of *p*-fluoride (**5**), *p*-methoxy (**6**), *p*-methylcarboxy (**7**), or *o*-chloride (**8**) all afforded the corresponding acetates in good yields and enantioselectivities (60 → 89%, 99% ee). Unactivated aliphatic alcohols were equally reactive: acetate **9** was isolated at 99% ee, while indole-based acetate **10** was isolated at 76% ee.²⁹ Secondary alcohols bearing substituents other than Me that are accepted by CALB were also well tolerated (**11** → **13**). In particular, the successful DKR to afford alkenyl-substituted **12** and alkynyl-substituted **13** are noteworthy. During DKR, the α -substituted

radical intermediates could undergo delocalization with the alkene or alkyne to afford rearranged products. However, acceptable yields and ee's were obtained for all substrates. Secondary alcohols with other aromatics, such as 1-benzosuberyl, 2-thiophenyl, 1-naphthyl, or 2-naphthyl, were also all well tolerated in the biocatalytic/photocatalytic protocol (**14** → **17**, 75 → 82%, 96–99% ee). Finally, secondary amines were also subjected to DKR (Table 1, entries 16–18). Amide **18** was isolated at 84% yield and 99% ee. It should be noted that in previous attempts at racemization using thermal conditions (PhSH, AIBN, Δ),³⁰ low yields of the racemized amine (30–50%) were obtained as competing oxidation and imine formation were observed. Unactivated amines afforded amides **19** and **20** at high yields and enantioselectivities as well and were used to demonstrate the DKR on a larger scale. Amine **19a** was subjected to the optimized conditions for a reaction time of 3 days (Scheme 1) and afforded amide **19** at 70% (99% ee).

Scheme 1. Gram-Scale Synthesis via DKR



In summary, a tandem photo- and biocatalytic DKR of secondary alcohols and amines was achieved employing a commercially available lipase and a heteroleptic copper complex. The racemization was made possible by a Cu(NN)-(PP)X-type complex, an example of using copper-based photocatalysis to promote HAT processes employing common thiyl radicals. A screening approach identified Cu(dtbbpy)-(DPEPhos)BF₄ from among 48 complexes, and (Ph₃SiS)₂ from among five disulfides, as an optimal catalyst system to promote the HAT process. In contrast to previous radical-mediated racemization methods (thermal and photochemical), the catalytic system described herein is the first general system applicable to aliphatic and aromatic alcohols and amines (18 examples, 60 → 97% yield, 76 → 99% ee). The above exemplifies the potential utility to exploit photocatalysis using base metal complexes in modern HAT processes and synthesis mediated by thiyl radicals.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acscatal.3c01713>.

Experimental procedures and characterization data, development of racemization, tabular data from screening, substrate synthesis, DKR procedures and experimental data, absorbance and emission data, excited-state lifetime data, Stern–Volmer experiments, electrochemical data, TEMPO experiments, NMR data for new compounds (PDF)

AUTHOR INFORMATION

Corresponding Author

Shawn K. Collins – Centre for Green Chemistry and Catalysis, Département de Chimie, Université de Montréal, Montréal, QC H2V 0B3, Canada; orcid.org/0000-0001-9206-5538; Email: shawn.collins@umontreal.ca

Authors

Clémentine Minozzi – Centre for Green Chemistry and Catalysis, Département de Chimie, Université de Montréal, Montréal, QC H2V 0B3, Canada

Nicolas Dowe – Centre for Green Chemistry and Catalysis, Département de Chimie, Université de Montréal, Montréal, QC H2V 0B3, Canada

Noémie Beaucage – Centre for Green Chemistry and Catalysis, Département de Chimie, Université de Montréal, Montréal, QC H2V 0B3, Canada

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acscatal.3c01713>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors acknowledge the Natural Sciences and Engineering Research Council of Canada (NSERC, Discovery 1043344 and RGPIN-2020-05006) and the Fonds de recherche Nature et technologie via the Centre in Green Chemistry and Catalysis (FRQNT-2020-RS4-265155-CCVC) for generous funding. The authors thank Ms. Alizée Debais (Rocheffort group, UdeM) for help obtaining CV data, D. Chartrand for help obtaining excited-state lifetime data, and Dr. Calvine Lai (Lebel group, UdeM) for access to HPLC equipment.

REFERENCES

- (1) (a) Shaw, M. H.; Twilton, J.; MacMillan, D. W. C. Photoredox Catalysis in Organic Chemistry. *J. Org. Chem.* **2016**, *81*, 6898–6926. (b) Twilton, J.; Le, C.; Zhang, P.; Shaw, M. H.; Evans, R. W.; MacMillan, D. W. C. The merger of transition metal and photocatalysis. *Nat. Rev. Chem.* **2017**, *1*, No. 0052. (c) Kärkäs, M. D.; Porco, J. A.; Stephenson, C. R. J. Photochemical Approaches to Complex Chemotypes: Applications in Natural Product Synthesis. *Chem. Rev.* **2016**, *116*, 9683–9747. (d) Romero, N. A.; Nicewicz, D. A. Organic Photoredox Catalysis. *Chem. Rev.* **2016**, *116*, 10075–10166. (e) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis. *Chem. Rev.* **2013**, *113*, 5322–5363. (f) Skubi, K. L.; Blum, T. R.; Yoon, T. P. Dual Catalysis Strategies in Photochemical Synthesis. *Chem. Rev.* **2016**, *116*, 10035–10074. (g) Teegardin, K.; Day, J. I.; Chan, J.; Weaver, J. Advances in Photocatalysis: A Microreview of Visible Light Mediated Ruthenium and Iridium Catalyzed Organic Transformations. *Org. Process Res. Dev.* **2016**, *20*, 1156–1163. (h) Marzo, L.; Pagire, S. K.; Reiser, O.; König, B. Visible-Light Photocatalysis: Does It Make a Difference in Organic Synthesis? *Angew. Chem., Int. Ed.* **2018**, *57*, 10034–10072. (i) Visible light photocatalysis in organic chemistry; Eds. Stephenson, C. R. J.; Yoon, T. P.; MacMillan, D. W. C.; Wiley-VCH Verlag: 2017.
- (2) (a) Sheldon, R. A.; Brady, D. Green Chemistry, Biocatalysis, and the Chemical Industry of the Future. *ChemSusChem* **2022**, *15*, No. e202102628. (b) Lubberink, M.; Schnepel, C.; Citolter, J.; Derrington, S. R.; Finnigan, W.; Hayes, M. A.; Turner, N. J.; Flitsch, S. L. Biocatalytic monoacylation of sym. diamines and its application to the synthesis of pharmaceutically relevant amides. *ACS Catal.* **2020**, *10*, 10005–10009. (c) Sheldon, R. A.; Brady, D. Broadening the scope of biocatalysis in sustainable organic synthesis. *ChemSusChem* **2019**, *12*, 2859–2881. (d) Sheldon, R. A.; Woodley, J. M. Role of Biocatalysis in Sustainable Chemistry. *Chem. Rev.* **2018**, *118*, 801–838.
- (3) (a) Meyer, L.-E.; Eser, B. E.; Kara, S. Coupling light with biocatalysis for sustainable synthesis—very recent developments and future perspectives. *Curr. Opin. Green Sustainable Chem.* **2021**, *31*, No. 100496. (b) Harrison, W.; Huang, X.; Zhao, H. Photobiocatalysis for Abiological Transformations. *Acc. Chem. Res.* **2022**, *55*, 1087–1096. (c) Peng, Y.; Chen, Z.; Xu, J.; Wu, Q. Recent Advances in Photobiocatalysis for Selective Organic Synthesis. *Org. Process Res. Dev.* **2022**, *26*, 1900–1913. (d) Schmermund, L.; Jurkaš, V.; Özgen, F. F.; Barone, G. D.; Büchsenstschütz, H. C.; Winkler, C. K.; Schmidt, S.; Kourist, R.; Kroutil, W. Photo-Biocatalysis: Biotransformations in the Presence of Light. *ACS Catal.* **2019**, *9*, 4115–4144. (e) DeHovitz, J. S.; Hyster, T. K. Photoinduced Dynamic Radical Processes for Isomerizations, Deracemizations, and Dynamic Kinetic Resolutions. *ACS Catal.* **2022**, *12*, 8911–8924. (f) Bierbaumer, S.; Schmermund, L.; List, A.; Winkler, C. K.; Glueck, S. M.; Kroutil, W. Synthesis of Enantiopure Sulfoxides by Concurrent Photocatalytic Oxidation and Biocatalytic Reduction. *Angew. Chem., Int. Ed.* **2022**, *134*, No. e202117103. (g) DeHovitz, J. S.; Loh, Y. Y.; Kautzky, J. A.; Nagao, K.; Meichan, A. J.; Yamauchi, M.; MacMillan, D. W. C.; Hyster, T. K. Static to inducibly dynamic stereocontrol: The convergent use of racemic b-substituted ketones. *Science* **2020**, *369*, 1113–1118. (h) Guo, X.; Okamoto, Y.; Schreier, M. R.; Ward, T. R.; Wenger, O. S. Enantioselective synthesis of amines by combining photoredox and enzymatic catalysis in a cyclic reaction network. *Chem. Sci.* **2018**, *9*, 5052–5056.
- (4) (a) Fu, H.; Cao, J.; Qiao, T.; Qi, Y.; Charnock, S. J.; Garfinkle, S.; Hyster, T. K. An asymmetric sp³–sp³ cross-electrophile coupling using 'ene'-reductases. *Nature* **2022**, *610*, 302–307. (b) Emmanuel, M. A.; Greenberg, N. R.; Oblinsky, D. G.; Hyster, T. K. Accessing non-natural reactivity by irradiating nicotinamide-dependent enzymes with light. *Nature* **2016**, *540*, 414–417.
- (5) (a) Ge, R.; Zhang, P.; Dong, X.; Li, Y.; Sun, Z.; Zeng, Y.; Chen, B.; Zhang, W. Photobiocatalytic Decarboxylation for the Synthesis of Fatty Epoxides from Renewable Fatty Acids. *ChemSusChem* **2022**, *15*, No. e202201275. (b) Peng, Y.; Li, D.; Fan, J.; Xu, W.; Xu, J.; Yu, H.; Lin, X.; Wu, Q. Enantiocomplementary C–H Bond Hydroxylation Combining Photo-Catalysis and Whole-Cell Biocatalysis in a One-Pot Cascade Process. *Eur. J. Org. Chem.* **2020**, *7*, 821–825. (c) Sosa, V.; Melkie, M.; Sulca, C.; Li, J.; Tang, L.; Li, T. J.; Faris, J.; Foley, B.; Banh, T.; Kato, M.; Cheruzel, L. Selective light-driven chemo-enzymatic trifluoromethylation/hydroxylation of substituted arenes. *ACS Catal.* **2018**, *8*, 2225–2229. (d) Litman, Z. C.; Wang, Y.; Zhao, H.; Hartwig, J. F. Cooperative asymmetric reactions combining photocatalysis and enzymatic catalysis. *Nature* **2018**, *560*, 355–359.
- (6) Verho, O.; Bäckvall, J.-E. Chemoenzymatic Dynamic Kinetic Resolution: A Powerful Tool for the Preparation of Enantiomerically Pure Alcohols and Amines. *J. Am. Chem. Soc.* **2015**, *137*, 3996–4009.
- (7) (a) Conley, B. L.; Pennington-Boggio, M. K.; Boz, E.; Williams, T. J. Discovery, Applications, and Catalytic Mechanisms of Shvo's Catalyst. *Chem. Rev.* **2010**, *110*, 2294–2312. (b) Shvo, Y.; Czarkie, D.; Rahamim, Y.; Chodosh, D. F. A new group of ruthenium complexes: structure and catalysis. *J. Am. Chem. Soc.* **1986**, *108*, 7400–7402.
- (8) (a) Akai, S.; Tanimoto, K.; Kanao, Y.; Egi, M.; Yamamoto, T.; Kita, Y. A Dynamic Kinetic Resolution of Allyl Alcohols by the Combined Use of Lipases and [VO(OSiPh₃)₃]. *Angew. Chem., Int.*

- Ed. **2006**, 45, 2592–2595. (b) Akai, S.; Hanada, R.; Fujiwara, N.; Kita, Y.; Egi, M. One-Pot Synthesis of Optically Active Allyl Esters via Lipase–Vanadium Combo Catalysis. *Org. Lett.* **2010**, 12, 4900–4903.
- (9) Cai, Y. Radical-chain racemisation of tetrahydrofurfuryl acetate under conditions of polarity-reversal catalysis: possible implications for the radical-induced strand cleavage of DNA. *Chem. Commun.* **1998**, 1145–1146.
- (10) Dang, H.-S.; Roberts, B. P. Strategy for contra-thermodynamic radical-chain epimerisation of 1,2-diols using polarity-reversal catalysis. *Tetrahedron Lett.* **2000**, 41, 8595–8599.
- (11) Escoubet, S.; Gastaldi, S.; Vanthuyne, N.; Gil, G.; Siri, D.; Bertrand, M. P. Thiyl Radical Mediated Racemization of Non-activated Aliphatic Amines. *J. Org. Chem.* **2006**, 71, 7288–7292.
- (12) (a) Gastaldi, S.; Escoubet, S.; Vanthuyne, N.; Gil, G.; Bertrand, M. P. Dynamic Kinetic Resolution of Amines Involving Biocatalysis and in Situ Free Radical Mediated Racemization. *Org. Lett.* **2007**, 9, 837–839. (b) Coniglio, A.; Galli, C.; Gentili, P.; Vadalà, R. Hydrogen atom abstraction from C–H bonds of benzylamides by the aminoxyl radical BTNO: A kinetic study. *Org. Biomol. Chem.* **2009**, 7, 155–160.
- (13) Zhang, Y.-A.; Gu, X.; Wendlandt, A. E. A Change from Kinetic to Thermodynamic Control Enables trans Selective Stereochemical Editing of Vicinal Diols. *J. Am. Chem. Soc.* **2022**, 144, 599–605.
- (14) (a) Zhang, Y.-A.; Palani, V.; Seim, A. E.; Wang, Y.; Wang, K. J.; Wendlandt, A. E. Stereochemical editing logic powered by the epimerization of unactivated tertiary stereocenters. *Science* **2022**, 378, 383–390. (b) Carder, H. M.; Wang, Y.; Wendlandt, A. E. Selective Axial-to-Equatorial Epimerization of Carbohydrates. *J. Am. Chem. Soc.* **2022**, 144, 11870–11877. (c) Wang, Y.; Carder, H. M.; Wendlandt, A. E. Synthesis of rare sugar isomers through site-selective epimerization. *Nature* **2020**, 578, 403–408.
- (15) Oswood, C. J.; MacMillan, D. W. C. Selective Isomerization via Transient Thermodynamic Control: Dynamic Epimerization of trans to cis Diols. *J. Am. Chem. Soc.* **2022**, 144, 93–98.
- (16) Wang, Y.; Hu, X.; Morales-Rivera, C. A.; Li, G.-X.; Huang, X.; He, G.; Liu, P.; Chen, G. Epimerization of Tertiary Carbon Centers via Reversible Radical Cleavage of Unactivated C(sp³)–H Bonds. *J. Am. Chem. Soc.* **2018**, 140, 9678–9684.
- (17) (a) Loh, Y. Y.; Nagao, K.; Hoover, A. J.; Hesk, D.; Rivera, N. R.; Colletti, S. L.; Davies, I. W.; MacMillan, D. W. C. Photoredox-catalyzed deuteration and tritiation of pharmaceutical compounds. *Science* **2017**, 358, 1182–1187. (b) Dang, H.-S.; Franchi, P.; Roberts, B. P. Radical-chain deoxygenation of tertiary alcohols, protected as their methoxymethyl (MOM) ethers, using thiols as polarity-reversal catalysts. *Chem. Commun.* **2000**, 499–500. (c) Panferova, L. I.; Zubkov, M. O.; Kokorekin, V. A.; Levin, V. V.; Dilman, A. D. Using the Thiyl Radical for Aliphatic Hydrogen-Atom Transfer: Thiolation of Unactivated C–H Bonds. *Angew. Chem., Int. Ed.* **2021**, 60, 2849–2854.
- (18) Yang, Q.; Zhao, F.; Zhang, N.; Liu, M.; Hu, H.; Zhang, J.; Zhou, S. Mild dynamic kinetic resolution of amines by coupled visible-light photoredox and enzyme catalysis. *Chem. Commun.* **2018**, 54, 14065–14068.
- (19) (a) Lévêque, C.; Levernier, E.; Corcé, V.; Fensterbank, L.; Malacria, M.; Ollivier, C. Photoredox Catalysis, an Opportunity for Sustainable Radical Chemistry In *Advanced Green Chemistry, Part 2: From Catalysis to Chemistry Frontiers*; 2019, 6, 49–121, DOI: 10.1142/9789811210587_0003. (b) König, B. *Science of Synthesis: Photocatalysis in Organic Synthesis*; Thieme: 2019, pp. 271–296. (c) Hossain, A.; Bhattacharyya, A.; Reiser, O. Copper's rapid ascent in visible light photoredox catalysis. *Science* **2019**, 364, No. eaav9713. (d) Hockin, B. M.; Li, C.; Robertson, N.; Zysman-Colman, E. Photoredox catalysts based on earth-abundant metal complexes. *Catal. Sci. Technol.* **2019**, 9, 889–915. (e) Beaudelot, J.; Oger, S.; Peruško, S.; Phan, T.-A.; Teunens, T.; Moucheron, C.; Evano, G. Photoactive Copper Complexes: Properties and Applications. *Chem. Rev.* **2022**, 122, 16365–16609.
- (20) (a) Li, C.; Dickson, R.; Rockstroh, N.; Rabeah, J.; Cordes, D. B.; Slawin, A. M. Z.; Hünemörder, P.; Spannenberg, A.; Bühl, M.; Mejía, E.; Zysman-Colman, E.; Kamer, P. C. J. Ligand electronic fine-tuning and its repercussion on the photocatalytic activity and mechanistic pathways of the copper-photocatalysed aza-Henry reaction. *Catal. Sci. Technol.* **2020**, 10, 7745–7756. (b) Gracia, L.-L.; Luci, L.; Bruschi, C.; Sambri, L.; Weis, P.; Fuhr, O.; Bizzarri, C. *Chem. – Eur. J.* **2020**, 26, 9929–9937. (c) Hunter, C. J.; Boyd, M. J.; May, G. D.; Fimognari, R. Visible-Light-Mediated N-Desulfonylation of N-Heterocycles Using a Heteroleptic Copper(I) Complex as a Photocatalyst. *J. Org. Chem.* **2020**, 85, 8732–8739. (d) Matsuo, K.; Yamaguchi, E.; Itoh, A. Atom-Transfer Radical Addition Photocatalysis Using a Heteroleptic Copper Complex. *Asian J. Org. Chem.* **2018**, 7, 2435–2438. (e) Mejía, E.; Luo, S.-P.; Karnahl, M.; Friedrich, A.; Tschierlei, S.; Surkus, A.-E.; Junge, H.; Gladiali, S.; Lochbrunner, S.; Beller, M. A Noble-Metal-Free System for Photocatalytic Hydrogen Production from Water. *Chem. – Eur. J.* **2013**, 19, 15972–15978. (f) Engl, S.; Reiser, O. Copper Makes the Difference: Visible Light-Mediated Atom Transfer Radical Addition Reactions of Iodoform with Olefins. *ACS Catal.* **2020**, 10, 9899–9906. (g) Hossain, A.; Engl, S.; Lutscher, E.; Reiser, O. Visible-Light-Mediated Regioselective Chlorosulfonylation of Alkenes and Alkynes: Introducing the Cu(II) Complex [Cu(dap)Cl₂] to Photocatalyzed ATRA Reactions. *ACS Catal.* **2019**, 9, 1103–1109. (h) Hossain, A.; Vidyasagar, A.; Eichinger, C.; Lankes, C.; Phan, J.; Rehbein, J.; Reiser, O. Visible-Light-Accelerated Copper(II)-Catalyzed Regio- and Chemoselective Oxo-Azidation of Vinyl Arenes. *Angew. Chem., Int. Ed.* **2018**, 57, 8288–8292. (i) Alkan-Zambada, M.; Hu, X. Cu Photoredox Catalysts Supported by a 4,6-Disubstituted 2,2'-Bipyridine Ligand: Application in Chlorotrifluoromethylation of Alkenes. *Organometallics* **2018**, 37, 3928–3935.
- (21) Minozzi, C.; Caron, A.; Grenier-Petel, J.-C.; Santandrea, J.; Collins, S. K. Heteroleptic Copper(I)-Based Complexes for Photocatalysis: Combinatorial Assembly, Discovery, and Optimization. *Angew. Chem., Int. Ed.* **2018**, 57, 5477–5481.
- (22) Caron, A.; Morin, É.; Collins, S. K. Bifunctional Copper-Based Photocatalyst for Reductive Pinacol-Type Couplings. *ACS Catal.* **2019**, 9, 9458–9464.
- (23) All reactions irradiated with blue LEDs were cooled by an external fan that maintained temperatures below 30 °C.
- (24) Some other common photocatalysts were also evaluated. See the Supporting Information for details.
- (25) When the racemization of **1** was conducted under optimized conditions with Cu(bathocup)(XantPhos)BF₄ (2 mol %), DABCO (10 mol %) and (Ph₃SiS)₂ (30 mol %), MeCN, blue LEDs, 18 h, a 37% ee of **1** was observed. When (Ph₃SiS)₂ was exchanged for Ph₃SiSH (30 mol %), the ee of recovered **1** was 99%.
- (26) Dénès, F.; Pichowicz, M.; Povie, G.; Renaud, P. Thiyl Radicals in Organic Synthesis. *Chem. Rev.* **2014**, 114, 2587–2693.
- (27) Teders, M.; Henkel, C.; Anhäuser, L.; Strieth-Kalthoff, F.; Gómez-Suárez, A.; Kleinmans, R.; Kahnt, A.; Rentmeister, A.; Guld, D.; Glorius, F. The energy-transfer-enabled biocompatible disulfide–ene reaction. *Nat. Chem.* **2018**, 10, 981–988.
- (28) (a) Qin, Y.; Zhu, Q.; Sun, R.; Ganley, J. M.; Knowles, R. R.; Nocera, D. G. Mechanistic Investigation and Optimization of Photoredox Anti-Markovnikov Hydroamination. *J. Am. Chem. Soc.* **2021**, 143, 10232–10242. (b) Christensen, T. B.; Daasbjerg, K.; Valentin-Hansen, P.; Pedersen, E. B.; Rissanen, K.; Shi, W.; Styring, S.; Tommos, C.; Warncke, K.; Wood, B. R. Investigation of the Direct and Indirect Reduction Processes of Some Disulfides by Electrochemical Means. *Acta Chem. Scand.* **1997**, 51, 307–317.
- (29) The lower ee is not a result of the DKR but a limitation of CALB. The kinetic resolution of the corresponding alcohol occurs at only 65% ee. See the Supporting Information for details.
- (30) Escoubet, S.; Gastaldi, S.; Vanthuyne, N.; Gil, G.; Siri, D.; Bertrand, M. P. Thiyl Radical Mediated Racemization of Benzylic Amines. *Eur. J. Org. Chem.* **2006**, 3242–3250.