

Characterization of the Biosynthetic Gene Cluster of Rebeccamycin from *Lechevalieria aerocolonigenes* ATCC 39243

Hiroyasu ONAKA,[†] Shin-ichi TANIGUCHI, Yasuhiro IGARASHI, and Tamotsu FURUMAI

Biotechnology Research Center, Toyama Prefectural University, Kosugi, Toyama 939-0398, Japan

Received July 29, 2002; Accepted August 19, 2002

The biosynthetic gene cluster for rebeccamycin, an indolocarbazole antibiotic, from *Lechevalieria aerocolonigenes* ATCC 39243 has 11 ORFs. To clarify their functions, mutants with *rebG*, *rebD*, *rebC*, *rebP*, *rebM*, *rebR*, *rebH*, *rebT*, or *orfD2* disrupted were constructed, and the gene products were examined. *rebP* disruptants produced 11,11'-dichlorochromopyrrolic acid, found to be a biosynthetic intermediate by a bioconversion experiment. Other genes encoded *N*-glycosyltransferase (*rebG*), monooxygenase (*rebC*), methyltransferase (*rebM*), a transcriptional activator (*rebR*), and halogenase (*rebH*). *rebT* disruptants produced rebeccamycin as much as the wild strain, so *rebT* was probably not involved in rebeccamycin production. Biosynthetic genes of staurosporine, an another indolocarbazole antibiotic, were cloned from *Streptomyces* sp. TP-A0274. *staO*, *staD*, and *staP* were similar to *rebO*, *rebD*, and *rebP*, respectively, all of which are responsible for indolocarbazole biosynthesis. But a *rebC* homolog, encoding a putative enzyme oxidizing the C-7 site of pyrrole rings, was not found in the staurosporine biosynthetic gene cluster. These results suggest that indolocarbazole is constructed by oxidative decarboxylation of chromopyrrolic acid (11,11'-dichlorochromopyrrolic acid in rebeccamycin) generated from two molecules of tryptophan by coupling and that the oxidation state at the C-7 position depends on the additional

enzyme(s) encoded by the biosynthetic genes.

Key words: rebeccamycin; indolocarbazole; biosynthetic pathway; *Lechevalieria aerocolonigenes* ATCC 39243; chromopyrrolic acid

Rebeccamycin (Fig. 1) is an indolocarbazole alkaloid produced by *Lechevalieria aerocolonigenes* ATCC 39243.* It inhibits DNA topoisomerase I with an IC₅₀ of 1.75 μ M *in vitro* and inhibits the growth of human lung adenocarcinoma cells with a GI₅₀ of 6.0 μ g/ml *in vivo*.¹⁾ Indolocarbazole compounds from natural sources have attracted attention because of they are inhibitors of protein kinases or DNA topoisomerases. Staurosporine (Fig. 1), the strongest inhibitor of phospholipid-Ca²⁺-dependent protein kinase (protein kinase C), with an IC₅₀ of 2.7 nM, is frequently used to study cellular processes.²⁾ Protein kinases and DNA topoisomerases are important in cell proliferation, so specific inhibitors are promising candidates for antitumor drugs. In fact, UCN-01,³⁾ BMY-27557,⁴⁾ and J-109384⁵⁾ are indolocarbazoles now in clinical trials for cancer chemotherapy.

Most indolocarbazole compounds are isolated from actinomycetes: rebeccamycin is from *Lechevalieria aerocolonigenes* ATCC 39243, staurosporine




Fig. 1. Chemical Structures of Indolocarbazole Compounds in This Study, and of Violacein.

[†] To whom correspondence should be addressed. Tel: +81-766-56-7500 ext. 522; Fax: +81-766-56-2498; E-mail: onaka@pu-toyama.ac.jp

Abbreviations: CCA, 11,11'-dichlorochromopyrrolic acid; ORF, open reading frame

* The genus name for strain ATCC 39243 was changed from *Saccharothrix* to *Lechevalieria* in 2001.²⁷⁾

is from *L. aerocolonigenes* subsp. *staurosporeus* and *Streptomyces* sp. strain TP-A0274, K252a is from *Nocardioopsis* sp. strain K252⁶⁾ and J-104303 is from *Streptomyces mobaraensis* BA 13793.⁷⁾ Biosynthetic studies done with isotope-labeled precursors indicated that the indolocarbazole structure is derived from two molecules of tryptophan, probably *via* indolepyruvic acid as an intermediate.⁸⁻¹⁰⁾ Ohuchi *et al.* reported cloning of the *ngt* gene encoding an *N*-glycosyltransferase from *L. aerocolonigenes* ATCC 39243.⁵⁾ The Ngt protein converted an indolocarbazole, J-104303, to its *N*-glucoside, indicating that *ngt* is responsible for *N*-glycosylation in rebeccamycin biosynthesis. Very recently, in a first report about indolocarbazole antibiotics, Sanchez *et al.* cloned the rebeccamycin biosynthetic genes, using an internal fragment of *ngt* as a probe.¹¹⁾ Nucleotide sequence analysis of a 25.6-kb fragment of the *L. aerocolonigenes* ATCC 39243 genome found 18 open reading frames (ORFs). On the basis of sequence analysis and database searches, they proposed four genes for indolocarbazole biosynthesis (*rebO*, *rebD*, *rebC*, and *rebP*), two genes for halogenation (*rebH*, and *rebF*), glycosylation (*rebG*, the name was changed from *ngt*), and sugar methylation (*rebM*), as well as a regulatory gene (*rebR*) and two resistance and secretion genes (*rebU* and *rebT*). Heterologous gene expression of the set *rebO*, *rebD*, *rebC*, and *rebP* for indolocarbazole biosynthesis resulted in production of deschlororebeccamycin aglycone. We have been independently investigating the biosynthetic genes for rebeccamycin,¹²⁾ and discovered its biosynthetic pathway with the identification of several key biosynthetic intermediates. We here characterize the biosynthetic gene cluster for rebeccamycin using a newly developed host-vector system for the genus *Lechevalieria* and gene-disruption experiments. The structures of the accumulated products of each *reb* disruptant were identified and an overall biosynthetic pathway of rebeccamycin is proposed.

Materials and Methods

Bacterial strains, plasmids, and growth condition. *L. aerocolonigenes* ATCC 39243 was obtained from the American Type Culture Collection and used as the source of DNA in the construction of the genomic libraries. *Streptomyces* sp. strain TP-A0274 was newly isolated from soil collected in Kosugi-machi, Toyama, Japan. *Escherichia coli* DH-5 α served as the host for plasmid subcloning in pUC19 and pK18mob.¹³⁾ *E. coli* XL1-Blue MR was used for cloning of the pK402¹⁴⁾ and the pTOYAMAcos cosmid (Onaka, in preparation). *E. coli* S17-1 was used for transconjugation.¹⁵⁾ Growth conditions and manipulations of *E. coli* were as described by Sambrook and Russell.¹⁶⁾ The production medium for *L. aero-*

colonigenes was medium G134.¹⁾ The seed culture for *L. aerocolonigenes* was in V-22 medium.¹⁷⁾ Bennett's glucose agar,¹⁷⁾ nutrient broth agar, and mannitol soya flour agar¹⁸⁾ were used for conjugational transformation.

General recombinant DNA techniques. Restriction enzymes, T4 DNA ligase, and *Taq* polymerase were purchased from New England Biolabs. DNA fragments were labelled with [α -³²P]-dCTP (6000 Ci mmol⁻¹; Amersham) and the BcaBEST labelling kit (Takara Shuzo Co., Ltd.). PCR was carried out with the PTC-200 DNA Engine (MJ Research, MA, USA). DNA manipulations were as described by Sambrook and Russell.¹⁶⁾

Cloning of the rebeccamycin biosynthetic genes. The two PCR primers, 5'-TCGGAATTCATGGGGCACGAGTGCTG-3' (NGT1) and 5'-AGGAAGCTTGAACGGGCGACGAACCT-3' (NGT2), were synthesized on the basis of the *ngt* DNA sequence recorded as accession No. AB023953 in GenBank. Underlining letters indicate *Eco*RI and *Hind*III restriction enzyme sites for cloning to vectors. A 731-bp DNA fragment was amplified by PCR with each of the PCR primers and *L. aerocolonigenes* chromosomal DNA. The amplified products were radiolabelled by the random priming method and put through colony hybridization. For the construction of a genomic library, chromosomal DNA of *L. aerocolonigenes* was prepared as described by Kieser *et al.*¹⁸⁾ and partially digested with *Sau*3AI, and DNA fragments of > 30 kb were purified by gel electrophoresis. The purified DNA fragments were ligated with pKU402 and packaged into λ phage. A hybridized cosmid, pREBC5, was digested with *Bam*HI and the hybridizing 8-kb fragment was subcloned into pUC19 to give pREB1. 3.4-kb *Bam*HI, 2.9-kb *Bam*HI, 2.7-kb *Sph*I, 2-kb *Pst*I, and 2.3-kb *Xho*I fragments, which were upstream from of the 8-kb fragment, also were cloned into pUC19 to give pREB5, pREB7, pREB9, pREB10, and pREB11, (see Fig. 2(A)).

Construction of plasmids for gene inactivation. (i) *pDrebG*. pDrebG was constructed by the cloning of a 776-bp *Pma*CI-*Sph*I fragment prepared from pREB1, which contained an internal region of the *rebG* gene, into the *Sma*I-*Sph*I sites of pK18mob. (ii) *pDrebD*, *pDrebP*, *pDrebM*, *pDrebH*, *pDrebT*, and *pDorfD2*. A 543-bp fragment was amplified by PCR with primers designed on the basis of the internal region of the *rebD* and pREB1 as the template. The fragment was inserted into pK18mob to give pDrebD. Five plasmids, pDrebP, pDrebM, pDrebH, pDrebT, and pDorfD2, which contained internal regions of *rebP*, *rebM*, *rebH*, *rebT*, and *orfD2* as shown in Fig. 2(A), also were constructed by the




Fig. 2. Restriction and Gene Organization Map of Cloned DNA Fragments from *L. aerocolonigenes* (A) and *Streptomyces* sp. Strain TP-A0274 (B).

Filled black bars indicate cloning regions of the plasmids used for single crossover disruption. Open bars indicate deleted regions of the in-frame deletion. B: *Bam*HI, E: *Eco*RI, N: *Nco*I, P: *Pst*I, S: *Sph*I, X: *Xho*I.

same method as that for pDrebD with pREB5 (pDrebP, pDrebM), pREB1 (pDorfd2), pREB10 (pDrebH), or pREB11 (pDrebT) as the templates for PCR. The primer sequences are as follows: *rebD*-N; 5'-GAAGAATTCGTGATGCTGCAGTACCTGTA-3', *rebD*-C; 5'-CGAAAGCTTCAGGAACAGGTG-GTGCTCGCC-3', *rebP*-N; 5'-CCGAAGCTTCAC-ACGGCGCTGAGATCGTTG-3', *rebP*-C; 5'-GTC GAATTCGCGCCAGCGAGTGACCATCTGCAC-3', *rebM*-N; 5'-ACGAAGCTTTCGGCTTGCGG-AACATCGGAC-3', *rebM*-C; 5'-CGGGAATTCG-ACGTCGGCTGCGGGATCGGC-3', *rebH*-N; 5'-AGCGAATTCGCTGTCGCACTACTGGTTCG-3', *rebH*-C; 5'-GTCAAGCTTCTCCTGCATGC-CGTCCGCCAG-3', *rebT*-N; 5'-CGGGAATTCCTGTTCGGTGCCGCCTCCCT-3', *rebT*-C; 5'-CCGAAGCTTTCTCCGCGAGTCCCGCGACGG-3', *orfD2*-N; 5'-CACGAATTCGAGGAGGTTT-CGCGAGAAGGTGGG-3', *orfD2*-C; 5'-GAAAAG-CTTGAAGCCGTCGGGATAGCGGTGGTG-3', in which *Eco*RI or *Hind*III sites (underlining letters indicated) were created for facilitation of the cloning. (iii) *pDrebC* and *pDrebR*. For in-frame deletion of the *rebC*, a 3.0-kb *Sph*I fragment was prepared from pREB1 and cloned into the *Sph*I site of pUC19 to generate pREB1A. The plasmid was digested with *Nco*I and the resulting *rebC*-truncated large fragment was purified and self-ligated to give pREB1B. A 2.9-kb *Bam*HI-*Eco*RI fragment was prepared from

pREB5 and subcloned into the *Bam*HI-*Eco*RI site of pREB1B to construct pREB1C. A 5-kb *Eco*RI-*Hind*III fragment was prepared from pREB1C and cloned into the *Eco*RI-*Hind*III site of pK18mob to construct pDrebC. For in-frame deletion of *rebR*, a 2.2-kb *Sph*I fragment was prepared from pREB7 and cloned into the *Sph*I site of pK18mob to generate pREB7A. A 2.3-kb *Pst*I fragment (one of the *Pst*I sites is on a multiple cloning site of pUC19) was prepared from pREB5 and subcloned into the *Pst*I site of pREB7A to construct pDrebR. Both of the deletion regions are shown in Fig. 2(A).

Gene disruption and transformation in Lechevalieria aerocolonigenes. For disruption of the chromosomal genes, we used the method of insertional inactivation via a single crossover with the non-replicating *E. coli* plasmid, pK18mob, derivatives (pDrebG, pDrebD, pDrebP, pDrebM, pDrebH, pDrebT, and pDorfd2) or via a double crossover with pDrebC and pDrebR. *rebG* and *rebC* double disruptants were constructed from the *rebC*-truncated mutant strain to use the method via a single crossover with pDrebG. The method of transconjugation is modified from the method for *Streptomyces* species as described by Mazodier *et al.*¹⁵ After transconjugation, 3 ml of 0.5% nutrient broth agar (Difco) with 200 µg/ml kanamycin was overlaid on a manitol soya flour agar plate for transformant selection

and the selected colonies were transferred to a Bennett's glucose agar plate containing 5 $\mu\text{g}/\text{ml}$ nalidixic acid and 20 $\mu\text{g}/\text{ml}$ kanamycin. The chromosomal DNAs were prepared from the candidate transformants and the restriction map of the chromosomal DNA was checked by Southern hybridization. For disruption of the chromosomal genes with pDrebC and pDrebR, first crossover events were done with the same method as for single crossover disruption. The pDrebC- or pDrebR-integrated strains were cultivated in a 500-ml K-1 flask containing 100 ml of V-22 medium with no antibiotics. After two rounds of cultivation, we selected colonies that were kanamycin-sensitive to obtain in-frame deletion mutants. The genomic structure of the strains obtained was checked by Southern hybridization.

HPLC of the products from gene-disrupted mutants. HPLC was done on a HP1090 system with a diode array detector (Hewlett Packard) and a C18 Rainin Microsorb column (3 μm , 100 \times 4.6 mm, i.d.; Rainin Instrument Co., MA, USA). Acetonitrile-0.15% KH_2PO_4 was the elution buffer. The temperature was 40°C, the flow rate was 1.2 ml/min, the solvent was acetonitrile-0.15% KH_2PO_4 (pH 3.5), and detection was at 254 nm. (the gradient diagram is in Fig. 4(A)) Each of the disrupted mutants was used to inoculate a 500-ml K-1 flask containing 100 ml of V-22 medium. After incubation at 30°C for 2 days on a rotary shaker at 200 rpm, 5-ml portions of the seed culture were transferred into 500-ml K-1 flasks containing 100 ml of medium G134. Fermentation was at 30°C for 11 days on the same rotary shaker. The culture broth was centrifuged to collect a mycelial cake, which was extracted with 100 ml of acetone. After the acetone was evaporated off, the resultant solution was extracted with 100 ml of ethyl acetate three times, concentrated under reduced pressure, and dissolved in Me_2SO for HPLC. The supernatant of the culture broth also was checked for production of rebeccamycin or its related compounds.

Purification of rebeccamycin intermediates. To purify the rebeccamycin aglycon (5), *L. aerocolonigenes rebG::km^r* was used to inoculate 500-ml K-1 flasks containing 100 ml of V-22 medium. After cultivation at 30°C for 2 days on a rotary shaker, 5-ml portions of the seed culture were transferred into fifty 500-ml K-1 flasks, each containing 100 ml of medium G134. The culture broth (5 liters) was centrifuged at 5000 $\times g$ for 10 min. The supernatant was discarded and the mycelial cake was extracted with 5 liters of acetone. After the evaporation, the resultant aqueous solution was extracted with an equal volume of ethyl acetate two times and concentrated under reduced pressure to give 1.2 g of crude extract. The residue was dissolved in 2.5 ml of methanol and put on an ODS-AM reverse-phase silica gel column

(4.6 i.d. \times 20 cm; YMC) equilibrated with 20% aqueous methanol. The column was eluted with 50% aqueous methanol and then 100% methanol. The rebeccamycin aglycon eluted in the 100% methanol fraction. The fractions containing the rebeccamycin aglycon were pooled, and evaporated under reduced pressure to give pure rebeccamycin aglycon (61.2 mg). For purification of 4 (11,11'-dichlorochromopyrrolic acid), 10 (4'-O-demethylrebeccamycin), 8 (7-deoxo-7-hydroxyrebeccamycin aglycon), 9 (7-deoxorebeccamycin aglycon), and 11 (1,11-dechloro-4'-O-demethylrebeccamycin), strain *rebD::km^r* (4), strain *rebM::km^r* (10), strain $\Delta\text{rebC} + \text{rebG}::\text{km}^r$ (8, 9), or strain *rebH::km^r* (11) was used and the purification method was the same as for rebeccamycin aglycon except that the materials eluted in the 50% methanol fraction from the ODS-AM column. Further HPLC was done of the products from strain $\Delta\text{rebC} + \text{rebG}::\text{km}^r$ (8, 9) and strain *rebH::km^r* (11) to obtain purity high enough for NMR. HPLC purification was done on a Shimadzu HPLC system (SPD-M10A) with a diode array detector (LC10-AT, Shimadzu) and an XTerra RP18 column (19 i.d. \times 300 mm; Waters). Partially purified samples containing the intermediates were dissolved in methanol and put on an HPLC column. The temperature was 30°C, flow rate 15 ml/min, solvent 50% acetonitrile-0.15% KH_2PO_4 (pH 3.5). Each intermediate was fractionated on the basis of the information from the diode array detector, with detection at 254 nm.

Analysis of metabolites. ^1H and ^{13}C NMR spectra were measured at 400 and 100 MHz in $\text{Me}_2\text{SO}-d_6$ on a JNM-LA400 apparatus (JEOL). LC-MS spectra were obtained on an API165 machine (Applied Biosystems). UV-visible spectra were taken on a HP1090 system. NMR data for compounds 4, 5, 8, 9, 10, and 11 are shown in Tables 1 to 3. MS data are as follows. Compound 4: ESI-MS $[\text{M} + \text{H}]^+$: m/z = 454.2, $[\text{M} - \text{H}]^-$: m/z 452.2. Compound 5: ESI-MS $[\text{M} - \text{H}]^-$: m/z 392.0. Compound 8: ESI-MS $[\text{M} + \text{H}]^+$: m/z 396.2. Compound 9: ESI-MS $[\text{M} + \text{H}]^+$: m/z 380.2. Compound 10: ESI-MS $[\text{M} + \text{H}]^+$: m/z 556.0, $[\text{M} - \text{H}]^-$: m/z 554.0. Compound 11: ESI-MS $[\text{M} + \text{H}]^+$: m/z 488.6, $[\text{M} - \text{H}]^-$: m/z 486.0.

Bioconversion experiments. Two hundred μg of a putative purified intermediate was added to a tube containing 10 ml of medium G134 and already cultured *L. aerocolonigenes rebH::km^r* was used to inoculate the same tube. After cultivation at 30°C for 5 days and 7 days on a rotary shaker, 5-ml portions were extracted. HPLC conditions and sample preparation were as described in the section on HPLC of the products from gene-disrupted mutants. The rebeccamycin production was identified with the reten-

tion time by HPLC and from the UV-visible spectrum.

Cloning and sequencing of the staurosporine biosynthetic genes. The *staD* gene fragment was amplified from *Streptomyces* sp. strain TP-A0274 genomic DNA with the primers rebBN and rebBC, designed on the basis of the *rebD* and *vioB* sequence, and was used as a colony hybridization probe. The sequences of primers were (rebBN) 5'-GCCAA-GCTTATCTGGCAGATGTGCGACCCG-3' and (rebBC) 5'-CAGGAATTCGATCATCTCCTCGC-GGGCGA-3', in which *Eco*RI or *Hind*III sites (underlining letters indicated) were created for facilitation of the cloning. PCR was done at 98°C for 20 sec, 60°C for 30 sec, and 72°C for 1 min, for 30 cycles. A 426-bp amplified fragment was radiolabelled by the random priming method and used for hybridization. For the construction of a genomic library, chromosomal DNA of *Streptomyces* sp. strain TP-A0274 was partially digested with *Sau*3AI, and DNA fragments of > 30 kb were purified by gel electrophoresis. The resulting DNA fragments were ligated with *Bam*HI-digested pTOYAMAcos, which is an actinomycetes-*E. coli* shuttle cosmid vector, and packaged into λ phage. Five hybridized cosmids were isolated. A selected cosmid was designated pTYMCsta and digested with *Bam*HI, and the hybridizing 5.6-kb fragment was subcloned into pUC19 to give pSTA1. Two-kilobase and 2.3-kb *Bam*HI fragments, which were in the upstream and downstream regions of the 5.6-kb fragment, respectively, also were cloned into pUC19 to give pSTA2 and pSTA3, as shown in Fig. 2(B). These fragments were sequenced with an ABI PRISM cycle sequencing kit and analyzed on a ABI PRISM 310 automated sequencer (Applied Biosystems). The templates for DNA sequencing were prepared with the Locus Pocus transposon insertion system kit (Novagen). The DNA and protein sequence similarity searches were done with the BLAST server at the National Center for Biotechnology Information (Bethesda, MD).

Nucleotide sequence accession number. The nucleotide sequence for the rebeccamycin biosynthetic genes has been deposited in the DDBJ database under DDBJ Accession Number AB071405 and that for the staurosporine biosynthetic genes has been deposited under DDBJ Accession Number AB071406.

Results and Discussion

Construction of disrupted mutants for biosynthesis

The conventional transformation protocol involving protoplasting and regeneration of mycelia usually

used for *Streptomyces* strains could not be used for *L. aerocolonigenes* ATCC 39243. We therefore developed a protocol for conjugational transfer from *E. coli* to strain ATCC 39243 by modifying the *Streptomyces* method.¹⁵ A plasmid, pK18mob, carrying the ColE1 replicon and an origin essential for conjugal transfer was used. Chromosomal DNAs were prepared from transformants that grew well on the selection plates, and chromosomal recombination events was monitored by Southern hybridization (Figs. 3(C) and 3(D)). Except for *rebC* and *rebR* disruptants, insertional inactivation via a single crossover was confirmed (Fig. 3(A)). *rebC* and *rebR* were disrupted by in-frame deletion (Fig. 3(B)) because insertional inactivation of *rebC* or *rebR* probably would have a polar effect on the transcription of genes downstream from *rebC* or *rebR* (Fig. 2(A)). Double disruptants of *rebC* and *rebG* were constructed from an in-frame *rebC* deletion mutant by a single crossover in the *rebG* region. Finally, three mutants were independently isolated and further analyzed for each of the following: *rebG*, *rebD*, *rebC*, *rebC*+*rebG*, *rebP*, *rebR*, *rebH*, *rebT*, and *orfD2* disruptants. Two mutants were isolated from *rebM* disruptants and analyzed.

Characterization of rebeccamycin biosynthetic genes

The *rebG*, *rebD*, *rebC*, *rebP*, *rebM*, *rebR*, and *rebH* genes were confirmed to constitute the rebeccamycin biosynthetic gene cluster by gene disruption experiments. Although the cloning and expression of *rebG* was previously described,⁵ it was not proved there that *rebG* is involved in rebeccamycin biosynthesis. We constructed a *rebG* mutant by gene disruption and found that it produced the rebeccamycin aglycon but not rebeccamycin (Fig. 4(B)). The *rebG* product therefore was the *N*-glycosyltransferase responsible for rebeccamycin biosynthesis.

rebD encoded a protein of 1,013 amino acid residue. A BLAST search indicated 36.6% identity over the entire sequence of RebD to VioB, which is involved in violacein (Fig. 1) biosynthesis in *Chromobacterium violaceum* and catalyzes a condensation reaction of 5-hydroxytryptophan and indolepyruvic acid to generate the bisindole intermediate for violacein.¹⁹ The *rebD*-disrupted mutant did not produce compounds related to indolocarbazoles (Fig. 4(C)); in the same way, the *vioB* disruptant does not produce any violacein related metabolites.¹⁹ *rebO*, *rebD*, *rebC*, and *rebP* seemed to be translationally coupled, suggesting that at least these four genes might be transcribed polycistronically. Strain *rebD::km^r* might have polar effects on the expression of *rebC* and *rebP*, but we could ignore such effects in this experiment because *rebC* and *rebP* products were involved in later biosynthetic steps than the *rebD* product, as described in the following paragraph




Fig. 3. Inactivation of the *reb* Cluster Genes by Gene Disruption Via Homologous Recombination.

(A) Schematic presentation of *rebG* gene disruption via single crossover. The 0.78-kb *PmaCI-SphI* DNA fragment used as the probe is indicated as a thin black bar. X, *XhoI*; *km'*, kanamycin resistance gene (*aphII*). *rebD*, *rebP*, *rebC + rebG*, *rebM*, *rebH*, *rebT*, and *orfD2* disruptants were prepared by a similar procedure. (B) Schematic presentation of *rebC* gene disruption with an in-frame deletion. N, *NcoI*. (C) Southern blotting of *L. aerocolonigenes* wild type (W) and a disruptant of *rebG*, *rebD*, *rebP*, *rebC + rebG*, *rebM*, *rebH*, *rebT*, or *orfD2*. The chromosomal DNAs were digested with *XhoI* (*rebG* and *rebH*), *SalI* (*rebC + rebG* and *rebD*), *BamHI* (*orfD2*, *rebP*, *rebT*, and *rebM*). The 0.78-kb internal *rebG* fragment from *pDrebG*, 0.54-kb *rebD* fragment from *pDrebD*, 0.61-kb *rebP* fragment from *pDrebP*, 0.42-kb *rebM* fragment from *pDrebM*, 0.42-kb *rebH* fragment from *pDrebH*, 0.92-kb *rebT* fragment from *pDrebT*, and 0.57-kb *orfD2* fragment from *pDorfD2* were used as ^{32}P -labelled probes. (D) Southern blotting of a disruptant of *rebC* or *rebR*. The chromosomal DNAs were digested with *SacI* (*rebC*) and *PvuII* (*rebR*). The 0.54-kb *rebD* fragment from *pDrebD* and 2.2-kb *SphI* fragment from *pREB7* were used as *rebC* and *rebR* probes, respectively. Lanes 1 and 2 indicate *pDrebC*- and *pDrebR*-integrated strains.

about results of the analysis of *rebP* and *rebC* disruptants. Considering the structural similarities of violacein and rebeccamycin, both of which are derived from two molecules of tryptophan, this observation led us to propose that *RebD* is involved in the construction of the bisindole structure by catalyzing a coupling reaction between 7-chlorotryptophan and 7-chloroindole-3-pyruvic acid.

The *rebP* gene product is quite similar to a cytochrome-P-450-like enzyme. The major product that the *rebP* disruptant accumulated was 11,11'-dichlorochromopyrrolic acid (CCA, 4) (Fig. 4(D)). The structure of CCA was identified from NMR and MS spectra. A ^{13}C NMR spectrum showed 11 sp^2 carbons, suggesting that CCA was a symmetrical molec-

ule and that all of the carbons in tryptophan were retained in CCA. ^1H - ^{13}C long-range couplings observed in the HMBC spectrum confirmed the chloroindole moiety (Fig. 5(A)). The presence of a pyrrole ring was not directly confirmed by HMBC, but the structure of CCA was deduced to be the chlorinated analog of chromopyrrolic acid²⁰ by the LC-MS, which gave a molecular ion $[\text{M} + \text{H}]^+$ at m/z 454.2, and by comparison of NMR results with those reported for chromopyrrolic acid. Chromopyrrolic acid [3,4-di-(3-indolyl)-pyrrole-2,5-dicarboxylic acid] is a secondary metabolite found in *C. violaceum*, a producer of violacein.²⁰ The close similarity of *vioB* and *rebD* suggested that CCA was biosynthesized directly by *RebD*. Conversion of CCA to indolocarpa-




Fig. 4. HPLC Chromatograph and UV-Vis Spectrum of Products of Wild Type and Gene Disruptant Mutants.

HPLC conditions and sample preparation were described in the text. The elution was done with a linear gradient as indicated on the right-hand scale in A.

zole through oxidative decarboxylation is generally thought to be catalyzed by RebP. In a check of our proposal, a bioconversion experiment was done (Fig. 6). CCA was added to the culture broth of the *rebH* disruptant and the products were analyzed by HPLC after 5 and 7 days. The peak of CCA (9.9 min) disappeared and the production of

rebecamycin (13.3 min) was detected after 7 days (Fig. 6(B)). Because the *rebH* disruptant was unable to produce chlorinated indolocarbazoles, including rebecamycin, we concluded that CCA was converted to rebecamycin as a biosynthetic intermediate.

The *rebC* gene product was similar to a flavin binding site containing monooxygenase. The *rebC*-trun-

Table 1. ^1H NMR Data for Compounds **5**, **8**, **9**, **10**, and **11**

Position	Compound				
	5	8	9	10	11
1	—	—	—	—	7.68(1H, d, 8.1)
2	7.50(2H, d, 7.6)	7.56(1H, d, 8.1)	7.54(1H, d, 7.6)	7.66(1H, d, 7.8)	7.55(1H, t, 8.3)
3	7.36(2H, t, 8.0)	7.28(1H, t, 7.8)	7.26(1H, t, 7.8)	7.36(1H, t, 7.8)	7.35(1H, t, 7.6)
4	8.85(2H, d, 8.1)	9.09(1H, d, 7.8)	9.16(1H, d, 8.0)	8.97(1H, d, 8.1)	9.15(1H, d, 8.1)
6-NH	11.09(1H, s)	8.87(1H, s)	8.61(1H, s)	11.29(1H, s)	11.11(1H, s)
7	—	6.41(1H, d, 10.0)	4.96(2H, s)	—	—
7-OH	—	6.52(1H, d, 10.2)	—	—	—
8	—	8.32(1H, d, 8.0)	8.02(1H, d, 7.8)	9.18(1H, dd, 1.2, 7.8)	9.15(1H, d, 8.1)
9	—	7.31(1H, t, 7.8)	7.32(1H, t, 7.8)	7.37(1H, t, 7.8)	7.35(1H, t, 7.6)
10	—	7.58(1H, d, 8.1)	7.58(1H, d, 7.8)	7.63(1H, d, 7.6)	7.57(1H, t, 8.8)
11	—	—	—	—	7.97(1H, d, 8.8)
12-NH	11.86(2H, br. s)	11.77(1H, s)	11.76(1H, s)	—	—
13-NH	—	11.63(1H, s)	11.58(1H, s)	10.67(1H, s)	11.66(1H, s)
1'	—	—	—	6.90(1H, d, 9.2)	6.27(1H, d, 9.0)
2'	—	—	—	3.75(1H, dt, 2.7, 7.8)	3.54(1H, m)
3'	—	—	—	3.36(1H, br. t)	3.56(1H, m)
4'	—	—	—	3.68(1H, t, 9.3)	3.98(1H, m)
5'	—	—	—	3.58, 3.75(1H, m)	3.97(1H, m)
6'	—	—	—	3.97(2H, m)	3.82, 4.08(2H, m)
2'-OH	—	—	—	4.92(1H, d, 5.8)	4.92(1H, d, 5.2)
3'-OH	—	—	—	5.22(1H, d, 4.9)	5.15(1H, d, 4.6)
4'-OH	—	—	—	5.39(1H, d, 5.4)	5.41(1H, d, 4.2)
6'-OH	—	—	—	5.12(1H, br. t)	6.01(1H, t, 4.0)

The $\text{Me}_2\text{SO}_4-d_6$ signal (2.49 ppm) was used as a reference. Integral multiplicities, and coupling constants (Hz) are in parentheses.

Table 2. ^{13}C NMR Data for Compounds **5**, **8**, **9**, **10**, and **11**

Position	Compound				
	5	8	9	10	11
1	115.65	115.27	115.21	116.02	112.12
2	126.05	124.68*	124.46*	126.92	126.94
3	121.39	120.34	120.16	122.08	120.36
4	123.24	124.05	124.24	123.53	124.44
4a	123.15	124.26	124.36	123.17	121.43
4b	115.74	115.52	116.04	117.46	116.92
4c	120.84	119.16	119.88	120.59	121.09*
5	170.81	170.35	171.84	170.31	170.99**
7	—	78.49	45.25	170.50	171.06**
7a	—	136.21	133.79	122.62	119.41
7b	—	115.52	114.75	119.37	118.42
7c	—	123.88	124.09	125.13	121.05*
8	—	122.04	120.33	124.10	124.44
9	—	120.93	121.14	122.57	120.61
10	—	124.70*	124.52*	130.00	126.76
11	—	115.63	115.70	116.24	111.80
11a	—	136.21	135.94	137.72	142.12
12a	—	128.02	127.72	129.65	128.32
12b	—	126.15	125.35	129.77	129.69
13a	136.96	136.10	135.80	137.00	140.77
1'	—	—	—	84.57	84.44
2'	—	—	—	71.98	73.04
3'	—	—	—	77.73	76.59
4'	—	—	—	69.92	67.51
5'	—	—	—	82.07	78.55
6'	—	—	—	60.55	58.27

The $\text{Me}_2\text{SO}_4-d_6$ signal (39.5 ppm) was used as a reference. * and **, inter-changeable.

Table 3. ^1H NMR and ^{13}C NMR Data for CCA

11,11'-Dichlorochromopyrrolic acid		
Position	^{13}C	^1H
1-NH		11.67(1H, s)
2	123.24	
3	123.50	
4-NH		11.13(2H, d, 1.9)
5	126.25	7.14(2H, d, 1.7)
6	109.37	
7	129.64	
8	118.43	7.06(2H, d, 8.1)
9	119.23	6.76(2H, t, 7.6)
10	119.89	7.00(2H, d, 7.6)
11	115.48	
12	132.22	
13	161.47	
13-COOH		12.39(2H, br. s)

The $\text{Me}_2\text{SO}_4-d_6$ signals (2.49 ppm for ^1H ; 39.5 ppm for ^{13}C) were used as references. Integral, multiplicities, and coupling constants (Hz) are in parentheses.

cated mutant produced several indolocarbazole products (Fig. 4(E)). The UV-visible spectrum showed that products **6** and **7** contained the indolocarbazole chromophore. This result indicates that indolocarbazole could be constructed without the involvement of the *rebC* product. The compounds of **6** and **7** were found to be 7-deoxo-7-hydroxyrebeccamycin and its 4'-O-demethyl derivative, on the basis of the following observations: in




Fig. 5. HMBC Correlations with Metabolites from Gene-disrupted Mutants.




Fig. 6. HPLC of Products of a *rebH*-Disruptant Grown in Medium with CCA.

The whole broth of 5-days (A) and 7-days (B) cultures were prepared as HPLC samples. The peaks of 9.9 and 13 min correspond to CCA and rebeccamycin, respectively.

LC-MS, the peaks at 7.7 (6) and 9 (7) min showed the molecular ion $[M+H]^+$ at m/z 572 and 558, two units higher than that of rebeccamycin (m/z 570) and 4'-O-demethylrebeccamycin (m/z 556), respectively; NMR results showed that each peak consisted of a mixture of 12-*N*- and 13-*N*-glycosides not separable on HPLC (NMR results not shown). Taken together, with the function of RebC assumed from the sequence similarity, the structures of 6 and 7 were deduced to be 7-deoxy-7-hydroxyrebeccamycin and

its 4'-O-demethyl derivative, although it is not clear whether the glycosylation site was at the 12- or 13-nitrogen. The *rebG* disruption experiment showed that *N*-glycosylation happened after indolocarbazole formation. We concluded that these two glycosylated products are shunt products. We then constructed *rebG* and *rebC* double disruptants to eliminate the glycosylated products, and analyzed the fermented products. Three peaks were detected in HPLC of the culture broth (Fig. 4(F), peaks 5, 8, and 9), and they were identified as rebeccamycin aglycon (5), 7-deoxy-7-hydroxyrebeccamycin aglycon (8), and 7-deoxorebeccamycin aglycon (9), respectively, by NMR. NMR assignments and HMBC results for 5, 8, and 9 are summarized in Tables 1 and 2 and in Figs. 5(B)–5(D). NMR results for these compounds showed similarity to reported values.^{21,22} Bioconversion experiments with 8 or 9 as an added sample were done to confirm which was the real intermediate, but neither compound was converted to rebeccamycin. Presumably, in the *rebC* disruptants, the other oxidase enzymes recognize the real intermediate of *rebC* disruptants as a substrate and produce 8 and 9. The RebC products were not confirmed in these disruption experiments, although, considering that rebeccamycin aglycon was identified as one of the products, we suppose that RebP took part in the conversion from CCA to rebeccamycin aglycone and that RebC controlled the oxidation state at the C-7 site.

The *rebM* product was similar to methyltransferases from various organisms. The *rebM* disruptant accumulated compound 10 (Fig. 4(G)),

which was identified as 4'-*O*-demethylrebeccamycin²³⁾ by NMR (Tables 1 and 2 and Fig. 5(E)) and LC-MS, which gave a $[M+H]^+$ ion at m/z 556.0. Therefore, *rebM* encoded the methyltransferase responsible for methylation at the 4-hydroxy group of the glucose moiety. Additionally, 4'-*O*-demethylrebeccamycin was converted to rebeccamycin by an *rebH* disruptant (not shown), which confirmed that the methylation was the last step of the biosynthesis.

The *rebH* gene product was similar to the tryptophan halogenase from *Pseudomonas chlororaphis*, which enzyme catalyzes the chlorination reaction from L-tryptophan to 7-chloro-L-tryptophan in pyrrolnitrin biosynthesis.²⁴⁾ The *rebH* disruptant produced an indolocarbazole compound, **11**, corresponding to the peak at 7.9 min on HPLC (Fig. 4(H)). The mass spectrum showed $[M+H]^+$ at m/z 487 and NMR confirmed compound **11** to be 1,11-dechloro-4'-*O*-demethylrebeccamycin. The NMR results for **11** were in accordance with those reported earlier.¹¹⁾ We concluded that RebH is responsible for chlorination of the indole moiety. Taking into account the similarity to the tryptophan halogenase from *P. chlororaphis*, chlorination probably takes place with tryptophan at the beginning of the biosynthetic cascade. The production of 1,11-dechloro-4'-*O*-demethylrebeccamycin might be allowed by substrate recognition by RebO, RebD, RebC, RebP, and RebG.

Regulatory and other genes

A database search found that RebR was similar to an *luxR* family transcriptional regulator. In our experiment, the *rebR*-truncated mutant did not produce compounds related to rebeccamycin (Fig. 4(I)). Sanchez *et al.* reported that *rebR* is essential for production in the heterologous expression of indolocarbazole biosynthesis genes.¹¹⁾ These findings indicate that RebR is a pathway-specific activator for rebeccamycin production.

rebT was at the end of the rebeccamycin biosynthetic gene cluster and was similar to transmembrane efflux proteins, which are candidates for rebeccamycin resistance or secretion. Disruption of *rebT* did not decrease the production level: the wild type produced rebeccamycin at the concentration of 52.5 ± 7 mg/l in 9 days of culture, and the *rebT* disruptants produced it at the concentration of 52.6 ± 1 mg/l. Sanchez *et al.* reported that *rebT* introduced into *Streptomyces albus* confers resistance to rebeccamycin.¹¹⁾ RebT might protect cells from the influx of an excess of rebeccamycin. We concluded that *rebT* is not involved in the rebeccamycin production in *L. aerocolonigenes* ATCC 39243.

orfD2 was immediately downstream of *rebG* with the reverse orientation on the chromosomal DNA. The *orfD2*-disrupted mutants produced as much rebeccamycin as the wild type strain did (Fig. 4(J)),

indicating that *orfD2* was not related to rebeccamycin biosynthesis and that the *rebG* gene was at the end of the rebeccamycin biosynthetic cluster.

Similarities and differences in indolocarbazole biosynthetic genes between rebeccamycin and staurosporine

The staurosporine-producing strain *Streptomyces* sp. strain TP-A0274 was newly isolated from a soil sample collected in Toyama, Japan, and the staurosporine biosynthetic gene cluster was cloned on the basis of information from the *rebD* nucleotide sequence. The genomic library of strain TP-A0274 was constructed and the biosynthesis gene was screened for with colony hybridization. The selected cosmid, pTYMCsta, was confirmed to include the entire staurosporine biosynthetic genes by heterologous expression in *Streptomyces lividans* TK23.²⁵⁾ The gene organization map of *rebD* homolog loci is shown in Fig. 2(B). A comparison between the amino acid sequences encoded by *staG* and *rebG* from *L. aerocolonigenes* found 55.8% identity, suggesting that *staG* encoded an *N*-glycosyltransferase. Upstream from *staG*, there were three ORFs, *staO*, *staD*, and *staP*, and the deduced amino acid sequences were similar to *rebO* (53.2% identity, 80.3% similarity), *rebD* (55.2% identity, 77.4% similarity), and *rebP* (51.8% identity, 79.8% similarity), respectively, suggesting that these ORFs encoded indolocarbazole biosynthesis genes. There was no gene similar to *rebC* in the staurosporine biosynthetic genes in strain TP-A0274. The differences in the structures of staurosporine and rebeccamycin aglycon were the presence or absence of a carbonyl group at C-7. *rebC*-disrupted mutants produced 7-deoxo-rebeccamycin, so RebC seemed to catalyze the oxidation at the C-7-position to the carbonyl group.

Proposed overall biosynthetic pathway of rebeccamycin

The most striking finding in this study was the identification of CCA as a biosynthetic intermediate: decarboxylation occurs after the coupling of two tryptophan-molecules derivative. On the basis of our results, the biosynthetic pathway for rebeccamycin shown in Fig. 7 is proposed. The initial step is the chlorination of tryptophan by the putative halogenase RebH. The produced 7-chlorotryptophan (**13**) is oxidatively deaminated by RebO to give 7-chloroindole-3-pyruvic acid (**14**). The *rebO* gene product has 25.3% identity in its 473 amino acid sequence to the L-amino acid oxidase from *Crotalus adamanteus* (eastern diamondback rattlesnake),²⁶⁾ which is a component of snake poison and catalyzes an oxidation reaction of L-tryptophan to indole-3-pyruvic acid. Therefore RebO may be responsible for the deamination of 7-chlorotryptophan. 7-Chloroindole-3-pyruvic acid (**14**) is then used in the coupling reaction




Fig. 7. Proposed Overall Biosynthetic Pathway of Rebeccamycin.

with 7-chlorotryptophan catalyzed by RebD to afford CCA (**4**). This dicarboxylic bisindole compound **4** is transformed to the rebeccamycin aglycon (**5**) via a putative oxidative reaction catalyzed by RebP, and oxidation at the 7-position to the carbonyl by RebC follows. Finally, the rebeccamycin aglycon is glycosylated with D-glucose by RebG (**10**), and the methylation of the hydroxyl group of the glucose moiety by RebM ends the biosynthesis of rebeccamycin.

Compared with the rebeccamycin biosynthetic pathway, staurosporine biosynthesis in strain TP-A0274 is proposed to be as follows. Tryptophan would be oxidatively deaminated by StaO to 3-indole pyruvic acid, and it would be used for the coupling reaction with tryptophan catalyzed by StaD to produce chromopyrrolic acid. As the last step of the aglycon biosynthesis, StaP would convert chromopyrrolic acid to staurosporine aglycon.

Acknowledgments

We thank Dr. H. Ikeda at Kitasato University for providing the plasmids and *E. coli* strains, and Dr. T. Daiiri at Toyama Prefectural University for valuable comments. We also thank Y. Ikeda at Toyama Prefectural University for obtaining the MS spectra and Nikka Fats & Oils Co., Ltd., for the gift of linseed meal in medium G134. This work was supported in part by a Grant-in-Aid for Scientific Research to H.O. from the Japan Society for the Promotion of Science.

References

- 1) Bush, J. A., Long, B. H., Catino, J. J., Bradner, W. T., and Tomita, K., Production and biological activity of rebeccamycin, a novel antitumor agent. *J. Antibiot.*, **40**, 668–678 (1987).
- 2) Omura, S., Sasaki, Y., Iwai, Y., and Takeshima, H., Staurosporine, a potentially important gift from a microorganism. *J. Antibiot.*, **48**, 535–548 (1995).
- 3) Akinaga, S., Sugiyama, K., and Akiyama, T., UCN-01 (7-hydroxystaurosporine) and other indolocarbazole compounds: a new generation of anticancer agents for the new century? *Anticancer Drug Des.*, **15**, 43–52 (2000).
- 4) Kaneko, T., Wong, H., Utzig, J., Schurig, J., and Doyle, T., Water-soluble derivatives of rebeccamycin. *J. Antibiot.*, **43**, 125–127 (1990).
- 5) Ohuchi, T., Ikeda-Araki, A., Watanabe-Sakamoto, A., Kojiri, K., Nagashima, M., Okanishi, M., and Suda, H., Cloning and expression of a gene encoding *N*-glycosyltransferase (*ngt*) from *Saccarothrix aerocolonigenes* ATCC 39243. *J. Antibiot.*, **53**, 393–403 (2000).
- 6) Kase, H., Iwahashi, K., and Matsuda, Y., K-252a, a potent inhibitor of protein kinase C from microbial origin. *J. Antibiot.*, **39**, 1059–1065 (1986).
- 7) Kojiri, K., Kondo, H., Yoshinari, T., Arakawa, H., Nakajima, S., Satoh, F., Kawamura, K., Okura, A., Suda, H., and Okanishi, M., A new antitumor substance BE-13793C, produced by a streptomycete: taxonomy, fermentation, isolation, structure determination and biological activity. *J. Antibiot.*, **44**, 723–728 (1991).
- 8) Pearce, C. J., Doyle, T. W., Forenza, S., Lam, K. S., and Schroeder, D. R., The biosynthetic origins of rebeccamycin. *J. Nat. Prod.*, **51**, 937–940 (1988).
- 9) Yang, S. W., Lin, L. J., Cordell, G. A., Wang, P., and Corley, D. G., *O*- and *N*-methylation in the biosynthesis of staurosporine. *J. Nat. Prod.*, **62**,

- 1551–1553 (1999).
- 10) Lam, K. S., Forenza, S., Doyle, T. W., and Pearce, C. J., Identification of indolepyruvic acid as an intermediate of rebeccamycin biosynthesis. *J. Ind. Microbiol.*, **6**, 291–294 (1990).
 - 11) Sanchez, C., Butovich, A. I., Brana, F. A., Rohr, J., Mendez, C., and Salas, A. J., The biosynthetic gene cluster for the antitumor rebeccamycin: characterization and generation of indolocarbazole derivatives. *Chem. Biol.*, **9**, 519–531 (2002).
 - 12) Onaka, H., Taniguchi, S., Igarashi, Y., and Furumai, T., The analysis of rebeccamycin biosynthetic pathway mediated by the biosynthetic gene disruption experiments. *Nippon Nougakagaku Kaishi Rinzi Zokan* (in Japanese), **75**, 359 (2001).
 - 13) Schafer, A., Tauch, A., Jager, W., Kalinowski, J., Thierbach, G., and Puhler, A., Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene*, **145**, 69–73 (1994).
 - 14) Pang, C. H., Shiiyama, M., Ikeda, H., Tanaka, H., and Omura, S., Cosmid vector for cloning and analysis of *Streptomyces* DNA. *Actinomycetol.*, **8**, 21–25 (1994).
 - 15) Mazodier, P., Petter, R., and Thompson, C., Inter-generic conjugation between *Escherichia coli* and *Streptomyces* species. *J. Bacteriol.*, **171**, 3583–3585 (1989).
 - 16) Sambrook, J., and Russell, D. W., Molecular cloning, a laboratory manual, third edition, Cold Spring Harbor Laboratory Press, Cold spring harbor, NY (2001).
 - 17) Onaka, H., Tabata, H., Igarashi, Y., Sato, Y., and Furumai, T., Goadsporin, a chemical substance which promotes secondary metabolism and morphogenesis in streptomycetes. I. Purification and characterization. *J. Antibiot.*, **54**, 1036–1044 (2001).
 - 18) Kieser, T., Bibb, M. J., Buttner, M. J., Chater, K. F., and Hopwood, D. A., Practical *Streptomyces* genetics, John Innes Foundation, Norwich, UK (2000).
 - 19) August, P. R., Grossman, T. H., Minor, C., Draper, M. P., MacNeil, I. A., Pemberton, J. M., Call, K. M., Holt, D., and Osburne, M. S., Sequence analysis and functional characterization of the violacein biosynthetic pathway from *Chromobacterium violaceum*. *J. Mol. Microbiol. Biotechnol.*, **2**, 513–519 (2000).
 - 20) Hoshino, T., Kojima, Y., Hayashi, T., Uchiyama, T., and Kaneko, K., A new metabolite of tryptophan, chromopyrrolic acid, produced by *Chromobacterium violaceum*. *Biosci. Biotechnol. Biochem.*, **57**, 775–781 (1993).
 - 21) Fabre, S., Prudhomme, M., and Rapp, M., Preparation of synthons for the synthesis of protein kinase C inhibitors from rebeccamycin. *Bioorg. Med. Chem.*, **2**, 449–452 (1992).
 - 22) Fabre, S., Prudhomme, M., Sancelme, M., and Rapp, M., Indolocarbazole protein kinase C inhibitors from rebeccamycin. *Bioorg. Med. Chem.*, **2**, 73–77 (1994).
 - 23) Lam, K. S., McDonald, L. A., Mattei, J., Forenza, S., and Matson, J. A., U.S. Patent 830969 (1994).
 - 24) Hammer, P. E., Burd, W., Hill, D. S., Ligon, J. M., and van Pee, K., Conservation of the pyrrolnitrin biosynthetic gene cluster among six pyrrolnitrin-producing strains. *FEMS Microbiol. Lett.*, **180**, 39–44 (1999).
 - 25) Onaka, H., Taniguchi, S., Igarashi, Y., and Furumai, T., Cloning of the staurosporine biosynthetic gene cluster from *Streptomyces* sp. TP-A0274 and its heterologous expression in *Streptomyces lividans*. *J. Antibiot.*, **55**, 1063–1071 (2002).
 - 26) Raibekas, A. A., and Massey, V., Primary structure of the snake venom L-amino acid oxidase shows high homology with the mouse B cell interleukin 4-induced Fig. 1 protein. *Biochem. Biophys. Res. Commun.*, **248**, 476–478 (1998).
 - 27) Labeda, D. P., Hatano, K., Kroppenstedt, R. M., and Tamura, T., Revival of the genus *Lentzea* and proposal for *Lechevalieria* gen. nov. *Int. J. Syst. Evol. Microbiol.*, **51**, 1045–1050 (2001).