

Reverse Clustering Impact: Geometry-Driven Minimal Parameters for Clustering via Morse Theory

Reinaldo Elias de Souza Junior
Faculdade de Medicina, Universidade Federal de Goiás, Brazil
resj3336@gmail.com

June 2026

Abstract

Classical clustering methods lack a principled mechanism for coupling *local* geometric variation with *global* multi-scale structure [17, 29, 8]. We introduce *Reverse Clustering Impact* (RCI), a parameter-minimal framework that requires only a single operational scale $r > 0$, with all remaining structural quantities determined automatically by small-ball asymptotics [14, 10, 3], nearest-neighbour statistics [1, 7], and doubling geometry [16, 9].

At each scale, RCI constructs the farthest-point cover of the dataset [13], generating a multi-scale profile $M(r)$ that quantifies effective volume growth with respect to the underlying metric. Using classical expansions for geodesic balls and empirical k NN radii, we obtain the approximation

$$\Delta^2 M_c(k) \approx -C_d S(x_c) r_k^2,$$

exhibiting a curvature-sensitive signature: negative on regions of positive curvature and positive on regions of negative curvature.

Globally, the farthest-point rule induces a discrete Morse scaffold in the sense of Forman [11, 12], whose critical transitions align with geometric jumps in $M(r)$. We prove that $M(r)$ is quantitatively comparable to the classical box-counting function $N(r)$ [9, 25], with constants depending only on the doubling dimension.

To evaluate geometric fidelity, we introduce the *Morse Erasure Index* (MEI), a parameter-free benchmark measuring how much variation of the Morse density field persists across cluster boundaries. MEI provides an objective comparison between clustering algorithms based on geometric consistency rather than hyperparameter tuning.

RCI thus achieves complete geometric clustering: metrically faithful at small scales, topologically coherent at large scales, and statistically controlled throughout—all from a single operational parameter. A complete, production-grade reference implementation is publicly available in a companion GitHub repository, ensuring full reproducibility of all experiments.

1 Introduction

Clustering methods widely used in practice—including KMeans, DBSCAN, spectral clustering, and Gaussian mixture models—lack a principled mechanism for coupling *local* geometric information with *global* multi-scale structure [17, 29, 8]. Their behavior depends sensitively on heuristic hyperparameters, often with weak geometric meaning. This work introduces **Reverse Clustering Impact (RCI)**, a parameter-minimal geometric framework in which the user specifies only a single analysis scale $r > 0$; all remaining quantities arise automatically from the metric structure itself: the radial base, local windows, cells, invariants, and admissible strata are generated from points and distances, while analytic stability estimates enter only as quantitative control in regular regimes.

Reproducibility. A complete, production-grade reference implementation of RCI is publicly available in a companion GitHub repository.¹ All experiments, spectral embeddings, MEI evaluations, and benchmark pipelines can be reproduced exactly using the provided code and fixed random seeds. The repository includes full documentation, a verified test suite, and scripts for regenerating every figure in this paper.

¹<https://github.com/Regis3336/rci-clustering>

Multi-scale profile via farthest-point covers. At each scale r , RCI computes the farthest-point cover [13], selecting successive points that maximize distance from the previously chosen centers. The number of selected centers, denoted

$$M(r) := |\text{FP}(r)|,$$

defines the *multi-scale profile*, a coarse geometric surrogate for the classical box-counting function $N(r)$ [9, 25]. Using only the doubling dimension, we establish the two-sided comparability inequalities

$$C_1 M(\lambda r) \leq N(r) \leq C_2 M(\lambda' r),$$

showing that $M(r)$ captures global metric complexity at substantially lower computational cost.

Local analytic layer: small-ball geometry, k NN radii, and curvature. The local layer combines classical small-ball asymptotics for geodesic volumes [14, 10, 3] with the statistical behavior of k NN radii [1, 7]. Concentration inequalities for shell averages [20, 2, 18] give stability of the centerwise growth profile $M_c(k)$. The discrete second difference satisfies the approximation

$$\Delta^2 M_c(k) \approx -C_d S(x_c) r_k^2,$$

where $S(x_c)$ is the scalar curvature at the center c . This curvature signature provides a local geometric probe governing the onset of multi-scale transitions and explains the contrasting behaviors on positively curved (sphere) and negatively curved (saddle) domains.

Global Morse structure. Section 4 shows that the farthest-point rule induces a discrete Morse scaffold in the sense of Forman [11, 12]. This scaffold mirrors classical Morse theory [21, 22, 19]: its critical transitions coincide precisely with the geometric jumps detected in the multi-scale profile $M(r)$ and the curvature signature $\Delta^2 M_c(k)$, providing a coherent global interpretation of cluster formation across scales.

Geometric fidelity via MEI. To compare clustering methods on intrinsic geometric grounds, we introduce the *Morse Erasure Index* (MEI), a parameter-free benchmark that quantifies how much variation of the Morse density field persists across cluster boundaries. The construction is compatible with stability theorems in topological data analysis [6, 4, 5] and with consistency results for geometric complexes on metric spaces [15, 23]. MEI thus measures geometric fidelity rather than tuning sensitivity.

Structural certification by admissible restriction. Beyond analytic stability, RCI admits a structural certification principle. Starting only from the finite metric data (X, d) , the algorithm generates a radial operational base, local invariant data, restriction maps, and walls of type change. On any admissible sieve avoiding these walls, the empirical RCI invariant restricts to a certified section:

$$\mathcal{S}_t \cap \mathcal{W}_n = \emptyset \implies \widehat{\mathcal{I}}_n|_{\mathcal{S}_t} \cong \mathcal{I}|_{\mathcal{S}_t}.$$

This shows that robustness is not only a probabilistic conclusion obtained after the fact, but also a structural mode of existence of the RCI object on the admissible metric base.

Empirical behavior. Section 6 shows that RCI achieves the highest MEI on all positively curved and mixed-geometry datasets. The sole exception is the saddle surface, where negative curvature yields $\Delta^2 M_c(k) > 0$, producing fragmentation patterns naturally aligned with density-threshold methods such as DBSCAN.

Contributions. This work provides:

1. a unified geometric framework relating farthest-point covers, curvature signatures, and discrete Morse structure;
2. quantitative comparability between $M(r)$ and classical box-counting;
3. a parameter-minimal clustering method requiring only one user choice;
4. a parameter-free geometric benchmark (MEI) grounded in TDA stability;
5. a structural robustness theorem showing that RCI-generated invariants become certified sections after restriction to admissible sieves avoiding walls of type change.

Structure of the appendices. Appendix 14 develops the analytic foundations of RCI: small-ball asymptotics, stability of k NN radii, and the bounded-difference estimates for shell averages. Appendix 13 constructs the topological layer, formalizing the scale poset, the spectral cover, the Čech nerve complexes, and the induced type- A persistence module. Appendix 11 presents the computational implementation, including the *Laplacian on demand* built from local k NN stencils, its spectral embedding, the full RCI pipeline, and the empirical validation across geometric benchmarks. Appendix 15 gives a complementary structural proof of robustness: starting from the minimal metric base generated by RCI, it defines walls of type change and proves that restriction to an admissible sieve yields a certified RCI section. Together, these appendices provide the analytic, topological, and algorithmic foundations of RCI.

RCI thus unifies analytic, topological, and fractal structure into a single geometric mechanism, providing metrically faithful, topologically coherent, and statistically controlled clustering across scales.

2 Methodological Note: AI-Assisted Mathematical Development

This work was developed through an unusual methodological approach that merits explicit documentation. The author is trained as a physician rather than as a professional research mathematician; for this reason, the development process deliberately incorporated adversarial formalization, repeated reconstruction, and computational validation as safeguards against informal reasoning. The author also employed a systematic AI-assisted verification protocol to ensure mathematical rigor throughout the formalization process.

2.1 Adversarial Reconstruction Dynamics

The theoretical development was systematic in its adversarial structure, but did not follow a fixed or predetermined verification pipeline in the procedural sense. Instead, it emerged through a highly iterative and adversarial dialogue between the author and multiple independent AI reasoning systems. Beyond that scaffold, the process was exploratory rather than scripted: ideas were constructed, dismantled, and reconstructed repeatedly until conceptual and mathematical coherence was achieved.

At different stages of the project, the workflow oscillated between the following informal modes:

- the author formulated candidate statements and geometric mechanisms based on intuition and empirical behavior observed in the algorithm;
- AI systems attempted formalizations, which were frequently incomplete or incorrect on first pass;
- the author challenged these attempts, requested alternative formulations, or rejected lines of reasoning when they contradicted intuition or computational evidence;
- independent systems were consulted in parallel, acting as external critics and exposing hidden assumptions, gaps, or inconsistencies;
- proofs were rewritten—sometimes several times—until the interaction between intuition, computation, and formal argument stabilized.

Role of AI systems. Large language models participated as active collaborators within the development process, occasionally proposing directions, alternative formalisms, or structural reorganizations that were not explicitly requested. These contributions, however, never operated autonomously: every suggestion, critique, or deviation was subjected to the author’s independent evaluation. The workflow functioned as a continuous dialectic in which the AI systems introduced variations and pressures, while the author determined—based on geometric intuition, conceptual coherence, and computational evidence—whether such proposals were consistent, misleading, incomplete, or worth pursuing further. The mathematical framework presented here therefore emerges from a genuinely collaborative interaction: AI systems contributed high-bandwidth reformulation and adversarial critique, but the author retained full epistemic authority, deciding when to accept, reject, reshape, or reverse the directions generated during the dialogue.

The process was not linear. It proceeded through cycles of construction and destruction, with entire sections of theory being rebuilt when contradictions or conceptual tensions surfaced. No formal count was recorded, but the author estimates that the overall development involved *several thousand* such adversarial rounds.

Certain components required particularly intensive cycles of refinement:

- The **Curvature Law** (Theorem 3.5) was repeatedly reformulated to achieve precise discretization and expectation bounds;
- The **bi-Lipschitz embedding theorem** (Theorem 3.3) went through multiple complete reconstructions to clarify failure modes and scale transitions;
- The **fractal completeness theorem** (Theorem 4.19) required several rebuilds to ensure the covering argument and dimension estimate were both tight and constructive;
- The **sheaf structure** (Appendix 13) emerged from extended debate on how to formulate separateness, gluing, and functorial coherence on the scale poset;
- The **stability theorem** (Theorem 3.4) required careful negotiation of bottleneck distances, interleavings, and morphisms between persistence modules.

Throughout the project, understanding was prioritized over acceptance. Every proof was rewritten in the author’s own words, reconstructed from first principles, and checked against computational intuition. AI systems accelerated formalization and supplied adversarial pressure, but responsibility for all mathematical content remains entirely with the author.

2.2 Computational Validation

Due to the author’s background outside pure mathematics, computational validation played a central role in validating the theoretical framework:

- (i) **Empirical observation preceded formalization.** The core geometric phenomena—curvature-driven boundary detection, scale-dependent cluster evolution, and topological transitions—were observed computationally across diverse synthetic datasets before any formal theorems were written. These observations guided which properties merited rigorous proof.
- (ii) **Implementation correctness was extensively tested.** The implementation includes comprehensive test coverage for edge conditions (degenerate geometries, extreme densities, pathological inputs), adversarial scenarios (non-isometric warps, outlier injection, density bias), and numerical stability (floating-point arithmetic, matrix conditioning, convergence behavior).
- (iii) **Benchmark results are fully reproducible.** All experimental results reported in Section 6 are generated by public scripts with fixed random seeds. Complete hyperparameter logs, dataset generation code, and evaluation pipelines are provided in the repository.

Scope of computational validation. The author prioritized *end-to-end empirical validation*—confirming that the complete framework produces the predicted adaptive-cover, boundary-detection, and topological behavior across the tested regimes—over *statement-by-statement numerical verification* of individual lemmas and bounds. This reflects standard practice in theoretical work: mathematical proofs establish correctness via logical argument, while computational experiments validate that the overall theory has predictive power.

The torture suite (Appendix 11) and benchmark results confirm that RCI behaves as the theory predicts across diverse geometric scenarios and adversarial transformations. Individual technical lemmas (e.g., Lemma 4.3, bounds in Appendix 14) are verified through mathematical reasoning rather than targeted numerical experiments for each statement.

The implementation and all benchmark scripts are publicly available, enabling independent verification of any theoretical claim by interested readers.

2.3 Limitations and Scope

This methodology has inherent limitations:

No substitute for peer review. While the adversarial LLM protocol provides internal consistency checks, it cannot replace scrutiny by expert human mathematicians. The convergence of multiple AI instances to the same conclusion increases confidence but does not constitute formal verification in the proof-assistant sense (e.g., Coq, Lean, Isabelle).

Proofs remain human-readable and standard. All proofs in this work use classical techniques (ε - δ arguments, compactness, continuity, measure-theoretic bounds, algebraic manipulations). No proof relies on brute-force computation or on numerical evidence as a substitute for argument. When probabilistic or compactness arguments are used, they are stated explicitly and remain within standard mathematical practice. Every argument can be checked by a trained mathematician with pencil and paper.

Possibility of subtle errors. Despite extensive adversarial review and computational validation, the possibility of errors remains. The author welcomes corrections and has established a public issue tracker for the repository where errors can be reported and addressed.

Author responsibility. The author takes full responsibility for any errors. AI systems served as tools for formalization, critique, and verification, not autonomous theorem-provers. Every claim was accepted only after the author reconciled geometric intuition, computational evidence, and formal argument. Where AI assistance suggested directions, the author evaluated and accepted or rejected them based on conceptual coherence and empirical evidence.

2.4 Broader Context

This work demonstrates that rigorous mathematical research can be conducted by individuals outside traditional academic mathematics, provided:

- (i) geometric intuition is grounded in computational experimentation;
- (ii) formalization is subjected to systematic adversarial scrutiny;
- (iii) the overall framework is validated empirically;
- (iv) limitations are acknowledged transparently;
- (v) the work is made fully public for community review.

The author views this methodology not as a replacement for traditional mathematical training but as an *expansion* of what is possible: enabling practitioners with domain expertise and strong computational intuition to formalize their insights with rigor, while maintaining intellectual honesty about the collaborative role of AI tools.

The iterative, adversarial nature of the development process mirrors, in some respects, the traditional peer review and seminar culture of mathematics—but compressed in time and augmented by AI capability to rapidly test formal statements, identify gaps, and suggest refinements. Its ultimate value—and its place relative to traditional mathematical practice—remains for the community to assess.

2.5 Acknowledgment

Mathematical formalization was developed with substantial assistance from large language models. The adversarial reconstruction process assisted the formal development, but all final decisions on mathematical content and full responsibility for correctness remain solely with the author. The author is grateful for the role these tools played in making a high level of formal rigor attainable outside traditional academic training, and equally grateful for any critical feedback from readers who identify gaps, improve arguments, or point out subtle errors that may have escaped the development process.

3 RCI: Formal Setup, Well-Posedness, and Robustness

This section introduces the **RCI algorithm** in its geometric form. The mathematical core of RCI is metric: once a finite metric space (X, d) is specified, the radial orderings, local density estimators, centerwise profiles, discrete curvature signals, and boundary indices are computed from distances. The reference implementation used in this work instantiates this metric through a finite-dimensional Euclidean representation $Z = \{z_i\}_{i=1}^N$. This representation may be either the original data coordinates or the spectral embedding produced by the matrix-on-demand Laplacian. Thus the execution is chart-based, while the RCI mechanism itself is the metric procedure of radial ordering, profile construction, and boundary detection. A spectral implementation provides a computationally efficient Euclidean surrogate; its fidelity is justified by the local bi-Lipschitz guarantees proved in Appendix 14. We proceed as follows.

Section 3.1 defines RCI on a general metric space. Section 3.2 describes the spectral surrogate. Section 3.3 establishes well-posedness of the boundary index. Section 3.4 records structural invariances. Section 3.5 states geometric and statistical guarantees, with proofs deferred to Appendix 14.

3.1 RCI on a General Metric Space

Let (X, d) be a finite metric space equipped with its counting measure, with $|X| = N < \infty$. This is the intrinsic metric form of RCI: after a metric has been fixed, the algorithm depends only on distance comparisons, local radii, density estimates, radial profiles, and finite differences. In the reference implementation, this metric space is realized by an operational Euclidean representation $Z = \{z_i\}_{i=1}^N \subset \mathbb{R}^q$, with

$$d(z_i, z_j) = \|z_i - z_j\|.$$

The representation Z may be the original coordinate data or the spectral embedding. The definitions below are written metrically, while the implementation realizes them through this Euclidean metric.

Local density estimator. For each $x_i \in X$, let $\mathcal{S}_\rho(i)$ be the m_ρ nearest neighbours of x_i in $X \setminus \{x_i\}$, and let $r_\rho(i)$ be the radius of the smallest closed d -ball containing $\mathcal{S}_\rho(i)$. We define the m_ρ -nearest-neighbour density estimator as:

$$\hat{\rho}(x_i) = \frac{m_\rho}{NV_m r_\rho(i)^m},$$

where $V_m := \pi^{m/2}/\Gamma(1 + m/2)$ is the volume of the unit ball in $\mathbb{R}^m$, and m denotes the intrinsic metric dimension estimated as in Appendix 14. This surrogate is standard for doubling spaces and captures the local scaling of the measure. Set

$$f_X(x_i) = \log \hat{\rho}(x_i).$$

Algorithmic centers and local profiles. The notation $M_c(k)$ denotes the local RCI profile computed after a centre c has been selected. In the analysis below, one may fix c in order to study the local profile. In the reference implementation, however, centres are not arbitrary: RCI runs on $Z = \{z_i\}_{i=1}^N \subset \mathbb{R}^q$, and the seeding rule—first centre via the empirical barycentre of Z , later centres via the residual surface in **Global RCI** below—produces centres c_j attached to each local profile. Thus every profile used by the algorithm has the form $M_{c_j}(k)$, where c_j is given by the implemented dynamics.

RCI profile. Fix a centre $c \in X$. Order the points of $X \setminus \{c\}$ by distance from c :

$$d(x_{(1)}, c) = r_1 \leq r_2 \leq \dots \leq r_{N-1},$$

and define

$$B_k(c) := \{c, x_{(1)}, \dots, x_{(k)}\}.$$

Assuming d separates points (no duplicated sites), $r_k > 0$ for $k \geq 1$. For a contrast function ϕ and a penalty $\alpha \geq 0$, for $1 \leq k \leq N - 1$ set

$$A_c(k) = \frac{1}{k+1} \sum_{x \in B_k(c)} \phi(\hat{\rho}(x)), \quad M_c(k) = A_c(k) - \alpha \log r_k.$$

The theoretical guarantees in Appendix 14 are stated for the logarithmic contrast $\phi(t) = \log t$. Other implementation-level contrasts may be used, but the curvature and stability results below refer to the logarithmic regime unless explicitly stated otherwise. The penalty term removes the trivial growth component from metric expansion, isolating the density–curvature interaction. Let $N_c := N - 1$ denote the number of admissible radii r_k in the local profile. Discrete curvature is encoded by the centred finite differences:

$$\begin{aligned} \Delta M_c(k) &= M_c(k+1) - M_c(k), & 1 \leq k \leq N_c - 1, \\ \Delta^2 M_c(k) &= M_c(k+1) - 2M_c(k) + M_c(k-1), & 2 \leq k \leq N_c - 1. \end{aligned}$$

Boundary index. After a warm-up of $m_0 \geq 3$,

$$\kappa^* = \min\{m_0 \leq k \leq N_c - 1 : \Delta^2 M_c(k) < 0 \text{ and } \Delta^2 M_c(k-1) \geq 0\},$$

with fallback to the smallest maximizer of $\Delta M_c(k)$ over $k \in \{m_0, \dots, N_c - 1\}$ if no such k exists. This ensures deterministic output and avoids pathological oscillatory behavior. The reference implementation uses a numerically robust version of this criterion: it smooths the second-difference signal, selects the strongest negative concavity region, and refines the boundary inside a local window using the first-difference profile. This implementation-level stabilization preserves the same Morse interpretation while improving finite-sample robustness. The index κ^* is the fundamental geometric observable of RCI, marking the first concavity change of the profile and acting as a discrete Morse boundary detector.

Global RCI. In the reference implementation, RCI is run on an operational Euclidean representation $Z = \{z_i\}_{i=1}^N$, either the original data representation or the spectral chart. The first centre is chosen as the point farthest from the empirical barycentre:

$$\bar{z} := \frac{1}{N} \sum_{i=1}^N z_i, \quad c_1 \in \arg \max_{z_i \in Z} \|z_i - \bar{z}\|,$$

with deterministic tie-breaking. For a selected centre c_j , let r_j^* be the radius associated with its boundary index κ_j^* . After J centres, define the covered region and absorbed data points by

$$U_J = \bigcup_{j=1}^J B(c_j, r_j^*), \quad A_J := X \cap U_J.$$

The implementation updates the residual surface

$$s_J(x) := \min_{1 \leq j \leq J} \max\{0, d(x, c_j) - r_j^*\}.$$

The next centre is selected by

$$c_{J+1} \in \arg \max_{x \in X \setminus A_J} s_J(x).$$

Equivalently, the next centre is the uncovered point with maximal residual distance outside the metric balls already constructed. The procedure stops when $A_J = X$, or when the uncovered set falls below the prescribed stopping tolerance.

Remark: fully metric initialization. The barycentric initialization above describes the reference implementation, where RCI is executed on an operational Euclidean representation. A fully coordinate-free initialization is also possible. For example, one may select the first centre by metric remoteness,

$$\text{rem}(x) := \frac{1}{|X| - 1} \sum_{y \in X \setminus \{x\}} d(x, y), \quad c_1 \in \arg \max_{x \in X} \text{rem}(x).$$

This variant requires no origin, barycentre, or ambient coordinates. It is not the initialization used in the reference implementation, but it shows that the RCI seeding mechanism admits a purely metric formulation.

3.2 Spectral Implementation of RCI: The Matrix-On-Demand Laplacian

The spectral construction used in RCI is not an embedding-dependent redefinition of the algorithm, but a computational surrogate that accelerates neighbourhood queries while preserving the intrinsic metric structure. The key object is a *matrix-on-demand Laplacian*: a Laplacian operator whose entries are never formed as a global matrix, but are generated locally and dynamically from the m_L -nearest-neighbour stencils. This preserves the geometric locality of RCI while enabling efficient, scalable spectral computations.

Local stencils. For each point $x_i \in X$, the neighbourhood oracle returns a stencil $\mathcal{S}_L(i)$ consisting of the m_L nearest neighbours of x_i , together with the squared distances $d(x_i, x_j)^2$ for all $j \in \mathcal{S}_L(i)$. No global matrix is built; all quantities are instantiated only when needed.

Matrix-on-demand affinities. For each point x_i , let $\mathcal{S}_L(i)$ denote its m_L nearest neighbours and define the local scale parameter

$$\sigma_i = d(x_i, x_{(m_L)}).$$

The density-adaptive affinity kernel is given by

$$K_{ij} = \exp\left(-\frac{d(x_i, x_j)^2}{\sigma_i \sigma_j}\right), \quad j \in \mathcal{S}_L(i) \text{ or } i \in \mathcal{S}_L(j).$$

Symmetrisation is applied lazily: the pair (i, j) contributes only when requested by the eigensolver or by a local consistency check. The degree entries are evaluated on demand by local summation,

$$D_{ii} = \sum_j K_{ij}.$$

Matrix-on-demand Laplacian. The normalized Laplacian acts functionally as

$$Lv = v - D^{-1/2} K D^{-1/2} v,$$

with every product involving K or D computed without ever forming global matrices. Local stencils provide the entries of K as needed, while the diagonal D is assembled dynamically from its local contributions. This matrix-free representation is compatible with Krylov, Lanczos, and power-iteration methods, and scales efficiently to large datasets while preserving the geometric locality intrinsic to RCI.

Spectral chart. The diffusion map

$$\Phi_q : X \rightarrow \mathbb{R}^q$$

is obtained from the first non-trivial eigenvectors of L , computed via matrix-on-demand operator calls. Appendix 14 proves that this chart is locally bi-Lipschitz: for all sufficiently close $x, y \in X$,

$$(1 - \varepsilon_N) d(x, y) \leq \|\Phi_q(x) - \Phi_q(y)\| \leq (1 + \varepsilon_N) d(x, y),$$

with $\varepsilon_N \rightarrow 0$ as $N \rightarrow \infty$. Thus the spectral embedding provides a computational Euclidean atlas that faithfully represents the intrinsic geometry at the scales relevant to RCI.

Metric core versus spectral implementation. The formal RCI mechanism is metric: radial orderings, profiles, finite differences, and boundary indices are computed from the metric supplied to the algorithm. In the spectral implementation, this metric is instantiated by Euclidean distances in the spectral representation $\Phi_q(X)$. The local bi-Lipschitz estimates of Appendix 14 justify using these spectral distances as faithful local surrogates for the intrinsic geometry at the scales relevant to RCI.

Conclusion. The matrix-on-demand Laplacian preserves the locality of RCI while providing a scalable numerical mechanism for spectral queries and implementing this controlled surrogate viewpoint.

3.3 Boundary Index and Well-Posedness

Theorem 3.1 (Well-Posedness per Center). *Fix deterministic tie-breaking in all arg max and nearest-neighbour selections. Assume $N_c - 1 \geq m_0$. For any finite dataset, any selected centre c , and parameters (ϕ, α, m_0) with $m_0 \geq 3$, the boundary index $\kappa^*(c)$ is uniquely defined, where N_c denotes the number of admissible radii r_k in the local profile (here $N_c = N - 1$), and satisfies*

$$m_0 \leq \kappa^*(c) \leq N_c - 1.$$

Proof. Assume $N_c - 1 \geq m_0$ (equivalently $N \geq m_0 + 2$ since $N_c = N - 1$), so the candidate index sets $\{m_0, \dots, N_c - 1\}$ in the definition of κ^* are nonempty. The profile $M_c(k)$ is defined for $k = 1, \dots, N_c = N - 1$ (Section 3.1). The centred second difference $\Delta^2 M_c(k)$ is thus defined only for $2 \leq k \leq N_c - 1$.

The set of indices satisfying the concavity-change condition

$$\{m_0 \leq k \leq N_c - 1 : \Delta^2 M_c(k) < 0 \text{ and } \Delta^2 M_c(k - 1) \geq 0\}$$

is finite (empty or not), so its minimum is well-defined whenever it is nonempty.

If the set is empty, the fallback rule selects the smallest maximizer of $\Delta M_c(k)$ over $k \in \{m_0, \dots, N_c - 1\}$. Under the deterministic tie-breaking assumed in Theorem 3.1, that maximizer is uniquely specified, hence so is $\kappa^*(c)$.

In either case, $\kappa^*(c)$ satisfies $m_0 \leq \kappa^*(c) \leq N_c - 1$. $\square$

3.4 Structural Invariances

The RCI construction depends on the metric d through: (i) nearest-neighbour sets, and (ii) the radii entering the density estimator and the penalty term. We record the precise invariance under positive rescaling of the metric.

Proposition 3.2 (Scaling Invariance). *Assume deterministic tie-breaking in all arg max and nearest-neighbour operations. Let $d_s(x, y) = s d(x, y)$ with $s > 0$. Then, for every center c and all k ,*

$$M_c^{(s)}(k) = M_c(k) - C_s,$$

where C_s is a constant independent of k . In particular,

$$\Delta M_c^{(s)}(k) = \Delta M_c(k), \quad \Delta^2 M_c^{(s)}(k) = \Delta^2 M_c(k),$$

and the boundary index κ^* and all cluster labels are unchanged.

Proof. (Log contrast.) The derivation below assumes $\phi = \log$ on $\hat{\rho}$, i.e., $\phi(\hat{\rho}(x_i)) = f_X(x_i) = \log \hat{\rho}(x_i)$.

Write all quantities with a superscript (s) when computed with respect to d_s . For the density estimator, we have

$$r_\rho^{(s)}(i) = s r_\rho(i), \quad \hat{\rho}^{(s)}(x_i) = \frac{m_\rho}{NV_m (r_\rho^{(s)}(i))^m} = \frac{m_\rho}{NV_m (s r_\rho(i))^m} = s^{-m} \hat{\rho}(x_i).$$

Taking logs,

$$f_X^{(s)}(x_i) = \log \hat{\rho}^{(s)}(x_i) = \log \hat{\rho}(x_i) - m \log s = f_X(x_i) - m \log s.$$

Therefore

$$A_c^{(s)}(k) = \frac{1}{k+1} \sum_{x \in B_k(c)} f_X^{(s)}(x) = \frac{1}{k+1} \sum_{x \in B_k(c)} (f_X(x) - m \log s) = A_c(k) - m \log s.$$

For the radii, $r_k^{(s)} = s r_k$ and hence

$$\log r_k^{(s)} = \log(s r_k) = \log r_k + \log s.$$

Thus the profile transforms as

$$M_c^{(s)}(k) = A_c^{(s)}(k) - \alpha \log r_k^{(s)} = (A_c(k) - m \log s) - \alpha(\log r_k + \log s) = M_c(k) - (m + \alpha) \log s.$$

The shift $C_s := (m + \alpha) \log s$ is independent of k , so first and second differences are preserved:

$$\Delta M_c^{(s)}(k) = M_c^{(s)}(k+1) - M_c^{(s)}(k) = (M_c(k+1) - C_s) - (M_c(k) - C_s) = \Delta M_c(k),$$

and similarly $\Delta^2 M_c^{(s)}(k) = \Delta^2 M_c(k)$.

The barycentric first-centre rule used in the reference implementation is also invariant under positive rescaling in the operational chart. Indeed, if $z_i^{(s)} = s z_i$, then

$$\bar{z}^{(s)} = s \bar{z}, \quad \|z_i^{(s)} - \bar{z}^{(s)}\| = s \|z_i - \bar{z}\|,$$

so the ordering of distances from the empirical barycentre is unchanged. Since the definition of κ^* depends only on the sign pattern of $\Delta^2 M_c(k)$ (and, in the fallback, on maximizers of $\Delta M_c(k)$), the boundary index is unchanged under rescaling whenever tie-breaking remains fixed.

The residual surface satisfies

$$s_J^{(s)}(x) = \min_{1 \leq j \leq J} \max\{0, s d(x, c_j) - s r_j^*\} = s s_J(x),$$

so its maximizers are unchanged under positive rescaling.

Because positive rescaling preserves all distance comparisons, and because tie-breaking is fixed, the same centres, boundary indices, and labels are obtained. $\square$

Remark (no general affine shift). A general affine transformation $d'(x, y) = ad(x, y) + b$ with $b > 0$ does not act as a simple rescaling on the radii entering the density estimator; in general $r'_\rho(i)$ is not equal to a constant multiple of $r_\rho(i)$. Therefore we do *not* claim full affine invariance for the profile $M_c(k)$, and restrict ourselves to the rigorously valid scaling invariance above.

3.5 Geometric Fidelity and Statistical Robustness

We summarize the main analytic guarantees for RCI. All proofs are given in Appendix 14.

Local Spectral Bi-Lipschitz Regularity.

Theorem 3.3 (Appendix 14, Theorem D.1). *Let $X \subset M$ be sampled from a compact C^3 Riemannian manifold (M, g) . With probability $1 - o(1)$ as $N \rightarrow \infty$, there exists $\varepsilon_N \rightarrow 0$ such that for all sufficiently close $x, y \in X$,*

$$(1 - \varepsilon_N) d_g(x, y) \leq \|\Phi_q(x) - \Phi_q(y)\| \leq (1 + \varepsilon_N) d_g(x, y).$$

Stability of the Boundary Index.

Theorem 3.4 (Appendix 14, Theorem D.2). *If the expected profile $M_c^\circ(k)$ admits a unique first concavity change at k° with a non-degenerate curvature gap, and if k° grows at least polylogarithmically in N , then*

$$\Pr(\kappa^\star = k^\circ) \geq 1 - N^{-C}, \quad C > 0.$$

Curvature law for RCI.

Theorem 3.5 (Curvature Law; Appendix 14, Theorem D.3). *Let the data be sampled from an m -dimensional Riemannian manifold (M, g) with scalar curvature S_g . When the volumetric penalty is set to $\alpha = m$, the leading r^m -term in the small-ball expansion is cancelled exactly. This cancellation renders the RCI profile $M_c(k)$ dimensionally neutral, with its second difference governed purely by curvature effects.*

For sufficiently large k , we obtain the approximation

$$\mathbb{E}[\Delta^2 M_c(k)] \approx -C_m S_g(c) \Delta(r_k^2),$$

where $C_m > 0$ depends only on m , and $\Delta(r_k^2) := r_{k+1}^2 - r_k^2$ measures the radial acceleration of the k -NN growth sequence. In particular, $\Delta^2 M_c(k)$ is typically negative on positively curved regions and positive on negatively curved regions.

Crucially, although the expression above is derived under smooth Riemannian assumptions, the validity of the RCI mechanism does not depend on global smoothness. The only structural requirement is local metric regularity: whenever the metric space behaves approximately Euclidean at sufficiently small scales—as is the case in Alexandrov spaces, metric-measure spaces, high-dimensional embeddings, and even effectively infinite-dimensional Hilbert-like settings—the cancellation of the volumetric term still holds and the Morse curvature signature remains well-defined. Thus, the curvature law extends far beyond the smooth manifold setting.

Summary. The results in this section show that the **RCI algorithm** is:

- metric at its core: once the operational metric is fixed, all radial profiles, curvature signals, and boundary indices depend only on distance comparisons and local radii;
- well-posed at every center, with a uniquely defined boundary index;
- invariant under positive rescaling of the metric (Proposition 3.2);
- dimensionally neutral, with the volumetric term cancelled exactly when $\alpha = m$, allowing RCI to operate uniformly in high-dimensional or even infinite-dimensional regimes;
- statistically stable, reliably recovering population-level geometric transitions;
- and geometrically faithful, with curvature information preserved and spectral surrogates controlled by local bi-Lipschitz estimates.

The guarantees apply to the metric RCI mechanism and transfer to spectral implementations whenever the stated local bi-Lipschitz assumptions hold. Thus densities, profiles, and curvature signatures are defined from the metric supplied to RCI; in the spectral implementation, this metric is provided by the spectral representation and controlled by the estimates above. The RCI boundary detector therefore retains its validity in metrically regular settings, independent of global smoothness or global manifold structure.

4 Metric-Measure and Morse Foundations of RCI

Structure of the section. Although RCI is algorithmically simple, its geometric structure is deep. This section establishes the metric-measure principles that justify the formal setup given in Section 3, providing the analytic and geometric backbone on which the algorithm rests.

The multiscale consistency of these constructions — sheaf-theoretic, topological, and categorical — is formalized in Appendix 13, while their quantitative stability and robustness are proved in Appendix 14. Readers primarily interested in empirical behavior may skip directly to Section 6, returning here when the underlying guarantees become relevant.

This section develops the continuum foundations of RCI, organized into three interacting layers:

1. the local volumetric behavior of a Borel probability measure ν on a compact metric space (X, d) ;
2. the geometric encoding of mass density via the *population* k -NN radii;
3. the global multiscale complexity captured by the covering profile $M(r)$.

We show that the scale-dependent complexity profile $M(r)$ is fractal complete, and that its structural transitions are reflected directly in the discrete curvature signatures $\Delta^2 M_c(k)$ used algorithmically by RCI. The computational alignment of these continuous quantities with their finite-sample estimators is validated through the full homology and nerve suite of Appendix 13.

Throughout, (X, d) denotes a compact metric space and ν a Borel probability measure on X .

Analytic centers versus algorithmic centers. Throughout the local analysis, expressions such as $M_c(k)$, $r_k(c)$, or $\Delta^2 M_c(k)$ are studied after fixing a centre c . This should not be read as an arbitrary initialization rule for the RCI algorithm. Analytically, fixing c localizes the estimates and allows one to prove uniform statements over possible centres. Algorithmically, however, RCI selects centres through a residual farthest-point procedure: the first centre is chosen as an extremal point of the dataset according to the global farthest-point criterion, and subsequent centres are chosen among the points not yet absorbed by previous clusters, maximizing their distance from the already clustered region. Thus every local profile $M_c(k)$ used by the algorithm is attached to a centre $c = c_j$ selected by the RCI residual seeding dynamics.

4.1 Small-Ball Regularity and Population k -NN Radii

In Section 3, the *empirical* k -NN radius $\hat{r}_k(x)$ was defined as a finite-sample proxy for local mass concentration. To analyze its statistical meaning and geometric content, we now introduce the corresponding *population* radius $r_k(x)$, defined with respect to the measure ν . The analytic properties established in this subsection provide the continuum foundation for the discrete RCI profile defined in Section 3. Their discrete-to-continuum consistency is demonstrated computationally in Appendix 11, while the full benchmarking pipeline used to validate these claims across synthetic manifolds is detailed in Appendix 12 for complete reproducibility.

Assumption 1 (Small-ball regularity). *There exist an integer $m \geq 1$, a constant $\omega_m > 0$, and a function*

$$R : X \times (0, r_0] \rightarrow (0, \infty)$$

such that:

1. For each $x \in X$ and $r \in (0, r_0]$,
- $$\nu(B_r(x)) = \omega_m r^m R(x, r).$$

2. For each $x \in X$, the map $r \mapsto R(x, r)$ is continuous on $(0, r_0]$ and admits a continuous extension to $r = 0$, denoted by

$$R_0(x) := \lim_{r \downarrow 0} R(x, r) \in (0, \infty).$$

3. The function $R_0 : X \rightarrow (0, \infty)$ is continuous and satisfies

$$0 < R_{\min} \leq R_0(x) \leq R_{\max} < \infty \quad \text{for all } x \in X.$$

The integer m serves as the effective local dimension, and R_0 as an effective density.

Definition 4.1 (Population k -NN radius). Fix $N \in \mathbb{N}$ and $k \in \{1, \dots, N\}$. For $x \in X$, the *population k -NN radius* $r_k(x)$ is defined as the unique number $r_k(x) \in (0, r_0]$ such that

$$\nu(B_{r_k(x)}(x)) = \frac{k}{N}.$$

Remark 4.2 (Radial inversion of scale). The population radius $r_k(x)$ should be understood as the radius revealed by radial expansion, not as an externally prescribed scale. In the classical small-ball viewpoint one fixes a radius r and studies the mass $\nu(B_r(x))$. In RCI the direction is reversed: the mass level k/N is fixed, the ball around x expands, and the radius $r_k(x)$ is the value at which

$$\nu(B_{r_k(x)}(x)) = \frac{k}{N}$$

is reached.

Thus the algorithm does not require prior knowledge of a privileged local scale. It generates the radial profile and lets the relevant scale appear through the expansion. The small-ball formula

$$\nu(B_r(x)) = \omega_m r^m R(x, r)$$

is used only to interpret the asymptotic behaviour of these revealed radii in the regime $k/N \rightarrow 0$, where $r_k(x) \rightarrow 0$. In that regime, the population radii enter the local mass geometry and yield the scaling law used below. They are not imposed as an additional stopping rule.

Lemma 4.3 (Existence and scaling of $r_k(x)$). *Let Assumption 1 hold. Then $r_k(x)$ is unique and satisfies the uniform scaling bounds:*

$$c_1 \left(\frac{k}{N} \right)^{1/m} \leq r_k(x) \leq c_2 \left(\frac{k}{N} \right)^{1/m},$$

where $0 < c_1 \leq c_2 < \infty$ are independent of x, k, N , and $r_k(x) \rightarrow 0$ uniformly in x as $k/N \rightarrow 0$.

The uniform scaling bounds in this lemma are crucial for ensuring that the RCI algorithm behaves consistently across different regions of the data space, regardless of variations in sampling density.

Proof. Under Assumption 1, for each fixed $x \in X$, the small-ball expansion gives

$$\nu(B_r(x)) = \omega_m r^m R(x, r),$$

where $R(x, r)$ is continuous in r and uniformly bounded above and below on $(0, r_0]$:

$$0 < R_{\min} \leq R(x, r) \leq R_{\max} < \infty.$$

Additionally, $\lim_{r \rightarrow 0} R(x, r) = R_0(x)$.

Step 1: Existence and uniqueness. For each fixed x , the map $r \mapsto \nu(B_r(x))$ is continuous and strictly increasing, with value 0 at $r = 0$ and at least 1 at $r = r_0$. Since $k/N \in (0, 1)$, the intermediate value theorem guarantees a unique solution $r_k(x) \in (0, r_0]$ to $\nu(B_{r_k(x)}(x)) = \frac{k}{N}$.

Step 2: Upper and lower bounds. Using the small-ball expansion and the uniform bounds on $R(x, r)$,

$$\omega_m r^m R_{\min} \leq \nu(B_r(x)) \leq \omega_m r^m R_{\max}.$$

Setting $\nu(B_r(x)) = k/N$ and solving for r yields

$$\left(\frac{k}{N\omega_m R_{\max}}\right)^{1/m} \leq r_k(x) \leq \left(\frac{k}{N\omega_m R_{\min}}\right)^{1/m}.$$

Thus, the constants c_1 and c_2 are given by

$$c_1 = (\omega_m R_{\max})^{-1/m}, \quad c_2 = (\omega_m R_{\min})^{-1/m},$$

which depend only on Assumption 1.

Step 3: Uniform vanishing as $k/N \rightarrow 0$. The above estimates directly imply

$$r_k(x) \leq c_2 \left(\frac{k}{N}\right)^{1/m},$$

and the right-hand side tends to 0 as $k/N \rightarrow 0$. Since the constants are uniform in x , the convergence is uniform in x .

Conclusion. The function $r_k(x)$ is uniquely defined, satisfies the uniform two-sided bounds, and tends to zero uniformly as $k/N \rightarrow 0$. $\square$

4.2 Uniform C^0 -Conjugacy Between Radius and Mass Potentials

The population radii $r_k(x)$ encode local density information. We formalize this by relating $r_k(x)$ to the mass profile $R_0(x)$ via logarithmic potentials.

Definition 4.4 (Radius and mass potentials). For each k and $x \in X$ set

$$u_k(x) := -\log r_k(x), \tag{1}$$

and define the mass profile

$$v(x) := \log R_0(x), \quad x \in X. \tag{2}$$

This definition sets the stage for Theorem 4.4 by translating the multiplicative relationship in Assumption 1 into an additive one, making it easier to compare the radius and mass potentials.

Theorem 4.5 (Uniform C^0 conjugacy). *Let Assumption 1 hold. For each k with k/N sufficiently small there exist a constant $C_k \in \mathbb{R}$ and a function $\eta_k : X \rightarrow \mathbb{R}$ such that*

$$v(x) = mu_k(x) + C_k + \eta_k(x), \quad x \in X, \tag{3}$$

with $C_k = \log(k/N) - \log \omega_m$, and $\|\eta_k\|_\infty \rightarrow 0$ as $k/N \rightarrow 0$. In particular, the family $\{u_k\}_k$ is uniformly C^0 -conjugate to the mass potential v .

This theorem provides a crucial link between the theoretical small-ball behavior and the practical k-NN radii. It shows that the k-NN radii effectively capture the underlying mass distribution, up to a well-controlled error term.

Proof. The definition of $r_k(x)$ gives

$$\nu(B_{r_k(x)}(x)) = \omega_m r_k(x)^m R(x, r_k(x)) = \frac{k}{N}. \tag{4}$$

Taking logarithms

$$\log \omega_m + m \log r_k(x) + \log R(x, r_k(x)) = \log \left(\frac{k}{N}\right). \tag{5}$$

Rearranging

$$\log R(x, r_k(x)) = \log \left(\frac{k}{N}\right) - \log \omega_m - m \log r_k(x). \tag{6}$$

Recall the potentials

$$u_k(x) = -\log r_k(x), \quad v(x) = \log R_0(x). \tag{7}$$

Define

$$C_k := \log(k/N) - \log \omega_m. \quad (8)$$

Then

$$\log R(x, r_k(x)) = C_k + mu_k(x). \quad (9)$$

Now write

$$v(x) = \log R_0(x) = \log R(x, r_k(x)) + [\log R_0(x) - \log R(x, r_k(x))]. \quad (10)$$

Define the error term

$$\eta_k(x) := \log R_0(x) - \log R(x, r_k(x)). \quad (11)$$

By Assumption 1, $R(x, r) \rightarrow R_0(x)$ uniformly in x as $r \rightarrow 0$. Since Lemma 4.3 shows $r_k(x) \rightarrow 0$ uniformly when $k/N \rightarrow 0$, it follows that

$$\sup_{x \in X} |\eta_k(x)| \rightarrow 0 \quad \text{as } k/N \rightarrow 0. \quad (12)$$

Substituting into (3):

$$v(x) = mu_k(x) + C_k + \eta_k(x). \quad (13)$$

This is the claimed C^0 -conjugacy. $\square$

4.3 Stability of Discrete Boundary Signals

The RCI boundary index κ^* (defined in Section 3) is determined by the sign change of the discrete second difference $\Delta^2 M_c(k)$, which acts as a curvature-sensitive boundary signal. Since the profile $M_c(k)$ is estimated from empirical data and therefore subject to sampling noise, it is essential to verify that the derived boundary index remains stable under perturbations.

The next lemma establishes this robustness: whenever the population profile admits a non-degenerate transition in curvature, the empirical estimator recovers the correct boundary index with high probability.

Definition 4.6 (Discrete differences and critical points). For a function $H : \{1, \dots, K\} \rightarrow \mathbb{R}$, we define:

$$\Delta^1 H(k) := H(k+1) - H(k),$$

$$\Delta^2 H(k) := H(k+1) - 2H(k) + H(k-1).$$

k_c is a *non-degenerate discrete critical point* of H if $|\Delta^2 H(k_c)| \geq \gamma$ for some non-degeneracy margin $\gamma > 0$.

Lemma 4.7 (Discrete Morse stability under uniform perturbations). *Let $F, G : \{1, \dots, K\} \rightarrow \mathbb{R}$ be discrete functions satisfying $\|F - G\|_\infty \leq \varepsilon$. Assume that every critical index k_c of F is non-degenerate in the sense that $|\Delta^2 F(k_c)| \geq \gamma$ for some $\gamma > 0$. If $\varepsilon < \gamma/8$, then:*

1. *For each non-degenerate critical point k_c of F , $\Delta^2 G(k_c)$ has the same sign as $\Delta^2 F(k_c)$.*
2. *If moreover $|\Delta^2 F(k)| \geq \gamma$ for all critical k and $|\Delta^2 F(k)| \leq \gamma/2$ for all non-critical k , then F and G have exactly the same set of critical indices and the same Morse types.*

Proof. For each $k \in \{2, \dots, K-1\}$ we have

$$\Delta^2 F(k) - \Delta^2 G(k) = (F(k+1) - G(k+1)) - 2(F(k) - G(k)) + (F(k-1) - G(k-1)),$$

so

$$|\Delta^2 F(k) - \Delta^2 G(k)| \leq 4\varepsilon. \quad (*)$$

(1) Preservation of signs. Let k_c be a non-degenerate critical point of F , so $|\Delta^2 F(k_c)| \geq \gamma$. Using $(*)$ and $\varepsilon < \gamma/8$,

$$|\Delta^2 G(k_c) - \Delta^2 F(k_c)| < \gamma/2.$$

Thus $\Delta^2 G(k_c)$ cannot cross zero, hence has the same sign as $\Delta^2 F(k_c)$. This proves (1).

(2) Preservation of critical indices. We argue by contradiction in both directions.

(a) *A critical index of F cannot become non-critical in G .* Suppose k_c is critical for F ($|\Delta^2 F(k_c)| \geq \gamma$) but non-critical for G ($|\Delta^2 G(k_c)| \leq \gamma/2$). Then by the reverse triangle inequality,

$$|\Delta^2 F(k_c) - \Delta^2 G(k_c)| \geq \gamma - \gamma/2 = \gamma/2.$$

Together with (*), we obtain

$$4\varepsilon \geq \gamma/2,$$

i.e. $\varepsilon \geq \gamma/8$, contradicting the hypothesis.

Thus every critical index of F remains critical for G .

(b) *A non-critical index of F cannot become critical in G .* Let k be non-critical for F , so $|\Delta^2 F(k)| \leq \gamma/2$. Suppose k becomes critical for G , so $|\Delta^2 G(k)| \geq \gamma$. Then

$$|\Delta^2 G(k) - \Delta^2 F(k)| \geq \gamma - \gamma/2 = \gamma/2,$$

but from (*) we must have

$$|\Delta^2 G(k) - \Delta^2 F(k)| \leq 4\varepsilon < \gamma/2,$$

contradiction.

Thus no new critical index can appear.

Having shown that the critical sets coincide and that signs are preserved, the Morse types also match. $\square$

Upper bound: $M(r) \leq C_2 N(r)$.

4.4 Controlled Geometry of RCI Covers in Doubling Spaces

We assume X is a doubling space, a common condition met by many datasets (including data sampled from low-dimensional manifolds).

Assumption 2 (Doubling metric space). *The space (X, d) is doubling: there exists $C_D \geq 1$ such that for every $x \in X$ and $r > 0$, the ball $B_r(x)$ can be covered by at most C_D balls of radius $r/2$.*

Definition 4.8 (Residual farthest-point seeding and r -nets). Let $S \subset X$ be the finite dataset. The RCI seeding procedure selects centres recursively. The first centre c_1 is chosen as an extremal point of S according to the global farthest-point criterion. After clusters $C_1, \dots, C_j$ have been formed, let

$$A_j := C_1 \cup \dots \cup C_j$$

denote the set of points already absorbed by the algorithm. The next centre is selected by

$$c_{j+1} \in \arg \max_{x \in S \setminus A_j} d(x, A_j), \quad d(x, A_j) := \inf_{a \in A_j} d(x, a).$$

The procedure stops when no unabsorbed point remains, or, at fixed scale r , when the maximal residual distance is at most r . In the fixed-scale idealization, this produces an r -net satisfying r -density and r -separation up to the residual absorption rule.

Proposition 4.9 (Residual farthest-point seeding produces controlled nets). *Under Assumption 2, the fixed-scale residual farthest-point procedure of Definition 4.8 terminates after finitely many steps and produces a controlled r -net of X , in the sense that the selected centres are r -separated and the resulting absorbed regions cover X at scale r .*

Definition 4.10 (RCI-type ball cover). Given an r -net $\{c_j\}_{j=1}^M$, we define the ball cover

$$\mathcal{C}(r) := \{B_{2r}(c_j) : 1 \leq j \leq M\}.$$

We denote by $M(r) := |\mathcal{C}(r)|$ the number of balls (or centers) in the cover. $M(r)$ is the ****RCI Complexity Profile****.

Lemma 4.11 (Bounded overlap). *Under Assumption 2, there exists a constant $Q = Q(C_D)$ such that any point $x \in X$ belongs to at most Q balls of $\mathcal{C}(r)$, i.e., the overlap is uniformly bounded.*

Proof. Fix $x \in X$ and consider all balls of the cover $\mathcal{C}(r)$ that contain x , say

$$x \in B_{2r}(c_{j_1}), B_{2r}(c_{j_2}), \dots, B_{2r}(c_{j_L}).$$

We show that L is uniformly bounded by a constant depending only on the doubling constant C_D .

Step 1: All corresponding centers lie in a small ball. If $x \in B_{2r}(c_{j_\ell})$, then $d(c_{j_\ell}, x) \leq 2r$, hence

$$c_{j_\ell} \in B_{2r}(x) \quad \text{for all } \ell.$$

Thus all the centers $\{c_{j_\ell}\}$ lie inside the ball $B_{2r}(x)$.

Step 2: Balls of radius $r/2$ around these centers are disjoint. Because $\{c_j\}$ is an r -net, it is r -separated:

$$d(c_i, c_j) > r \quad \text{for } i \neq j.$$

Hence the balls $B_{r/2}(c_{j_\ell})$ are pairwise disjoint.

Step 3: All these disjoint balls fit inside a slightly larger ball. For any ℓ ,

$$B_{r/2}(c_{j_\ell}) \subset B_{2r+r/2}(x) = B_{5r/2}(x).$$

Thus we have L disjoint balls of radius $r/2$ inside $B_{5r/2}(x)$.

Step 4: Use the doubling condition. By Assumption 2, a ball of radius $5r/2$ can be covered by at most

$$N := C_D^{\lceil \log_2(5) \rceil}$$

balls of radius $r/2$.

Since the L balls $\{B_{r/2}(c_{j_\ell})\}$ are disjoint and each must lie inside one of these N covering balls, we obtain

$$L \leq N.$$

Conclusion. Setting $Q := N$ gives a bound depending only on the doubling constant C_D , and therefore any point $x \in X$ belongs to at most Q balls of $\mathcal{C}(r)$. $\square$

The controlled overlap ensures that a finite number of balls are guaranteed to cover X .

Remark. The controlled overlap established in Lemma 4.11 ensures that the induced cover $\mathcal{C}(r)$ satisfies fundamental hypotheses for topological inference, allowing us to use the Nerve Theorem (Section 4.5).

4.5 Topological Transitions via the Nerve Complex

We now connect the metric cover to the topology of X using the Nerve Theorem. This allows tracking topological changes (component mergers) as discrete Morse transitions.

Assumption 3 (Local contractibility of ball intersections). *There exists $r_* > 0$ such that, for every $r \in (0, r_*]$ and for any finite family of indices J , the intersection*

$$\bigcap_{j \in J} B_{2r}(c_j)$$

is either empty or contractible.

Ball intersections are contractible, making the space topologically simple enough that it has the same basic shape.

Definition 4.12 (Nerve complex). The *nerve* $\mathcal{N}(r)$ of the cover $\mathcal{C}(r)$ is the abstract simplicial complex where a subset J forms a simplex if and only if the corresponding intersection of balls is non-empty.

Lemma 4.13 (Nerve equivalence). *Under Assumptions 2 and 3, for all $r \in (0, r_*]$, the complex $\mathcal{N}(r)$ and the union of balls*

$$U(r) := \bigcup_{j=1}^{M(r)} B_{2r}(c_j)$$

have the same homotopy type. In particular, $\beta_i(\mathcal{N}(r)) = \beta_i(U(r))$ for all $i \geq 0$.

The nerve and its union are topologically equivalent, so all the important properties of our cover are actually captured by the abstract simplicial complex.

Definition 4.14 (Component-count profile). Define the component-count profile

$$B(r) := \beta_0(U(r)) = \beta_0(\mathcal{N}(r)),$$

where β_0 is the number of connected components.

Lemma 4.15 (Nerve-theoretic Morse transitions). *The values of r at which the component-count profile $B(r)$ decreases correspond exactly to the r values where:*

1. *two distinct connected components of $U(r^-)$ intersect for the first time at r , causing a merger;*
2. *or, equivalently, two components of the nerve $\mathcal{N}(r^-)$ become connected by the addition of an edge (or higher simplex).*

Each drop in $B(r)$ is interpreted as a discrete Morse transition in β_0 .

The number of components decreasing indicates some high scale structural transition.

4.6 Fractal Completeness of RCI

We show that the RCI Complexity Profile $M(r)$ is closely related to the Minkowski dimension of X .

Definition 4.16 (Covering number). For $r > 0$, let $N(r)$ be the smallest number of balls of radius r required to cover X .

Lemma 4.17 (Comparison between $M(r)$ and $N(r)$). *Under Assumption 2, there exist constants $C_1, C_2 > 0$ and $r_1 > 0$ such that, for all $r \in (0, r_1]$,*

$$N(2r) \leq M(r) \leq C_2 N(r). \quad (14)$$

In particular, $M(r)$ and $N(r)$ are equivalent on a logarithmic scale as $r \downarrow 0$.

This lemma shows that counting the centres of balls in our construction actually gives us a reasonable approximation of the number needed to cover the space, and helps connect $M(r)$ to dimension. The RCI is capturing the space efficiently, neither being too redundant in coverage or leaving holes.

Proof. Let $\{c_j\}_{j=1}^{M(r)}$ be the controlled r -net produced by the residual farthest-point procedure of Definition 4.8, and let

$$\mathcal{C}(r) = \{B_{2r}(c_j) : 1 \leq j \leq M(r)\}$$

be the corresponding RCI cover.

Lower bound: $N(2r) \leq M(r)$. Each ball in $\mathcal{C}(r)$ has radius $2r$, so $\mathcal{C}(r)$ is a $2r$ -cover of X . Since $N(2r)$ is the *minimal* number of such balls,

$$N(2r) \leq M(r).$$

Doubling refinement. If the minimal cover uses balls of radius r , the RCI cover uses balls of radius $2r$. By the doubling property, replacing radius r by $2r$ affects the covering number by at most a factor depending only on C_D :

$$N(r) \leq C_D^{\lceil \log_2(2) \rceil} N(2r) = C_D N(2r).$$

Upper bound: $M(r) \leq C_2 N(r)$. Let $\{B_r(x_i)\}_{i=1}^{N(r)}$ be a minimal r -cover of X . For each i , define the index set

$$J_i := \{j : c_j \in B_r(x_i)\}.$$

Since the balls $\{B_r(x_i)\}$ cover X , the sets $\{J_i\}$ cover all centers, and therefore

$$M(r) \leq \sum_{i=1}^{N(r)} |J_i|.$$

Fix one such ball $B_r(x_i)$. All centers c_j with $j \in J_i$ lie inside $B_r(x_i)$, and since the set $\{c_j\}$ is r -separated, the balls $\{B_{r/2}(c_j)\}_{j \in J_i}$ are pairwise disjoint. Moreover,

$$B_{r/2}(c_j) \subset B_{r+r/2}(x_i) = B_{3r/2}(x_i).$$

Thus we obtain $|J_i|$ disjoint balls of radius $r/2$ contained in $B_{3r/2}(x_i)$. By the doubling property, the ball $B_{3r/2}(x_i)$ can be covered by at most $C_D^{\lceil \log_2 3 \rceil}$ balls of radius $r/2$, so

$$|J_i| \leq C_D^{\lceil \log_2 3 \rceil}.$$

Summing over all i ,

$$M(r) \leq \sum_{i=1}^{N(r)} |J_i| \leq C_D^{\lceil \log_2 3 \rceil} N(r).$$

Setting $C_2 := C_D^{\lceil \log_2 3 \rceil}$ gives the desired upper bound.

Conclusion. Combining the inequalities above, we obtain

$$N(2r) \leq M(r) \leq C_2 N(r),$$

which completes the proof. $\square$

Definition 4.18 (Minkowski dimension). The *Minkowski dimension* of X (if it exists) is

$$\dim_M(X) := \lim_{r \downarrow 0} \frac{\log N(r)}{-\log r}.$$

Theorem 4.19 (Fractal completeness of $M(r)$). *Under the hypotheses above, suppose that $\dim_M(X)$ exists. Then*

$$\lim_{r \downarrow 0} \frac{\log M(r)}{-\log r} = \dim_M(X).$$

In particular, the complexity profile $M(r)$ is fractal complete: its growth exponent exactly recovers the Minkowski dimension.

This is the key fractal property that connects the RCI to the fundamental definition of dimension; $M(r)$ contains full information to reflect the underlying geometry.

Proof. By Lemma 4.17, there exist constants $C_1, C_2 > 0$ and $r_1 > 0$ such that for every $r \in (0, r_1]$,

$$N(2r) \leq M(r) \leq C_2 N(r).$$

Taking logarithms,

$$\log N(2r) \leq \log M(r) \leq \log C_2 + \log N(r).$$

Divide by $-\log r$ (positive as $r \downarrow 0$):

$$\frac{\log N(2r)}{-\log r} \leq \frac{\log M(r)}{-\log r} \leq \frac{\log C_2}{-\log r} + \frac{\log N(r)}{-\log r}.$$

As $r \downarrow 0$, the term $\frac{\log C_2}{-\log r}$ tends to 0. Moreover,

$$\frac{\log N(2r)}{-\log r} = \frac{\log N(2r)}{-\log(2r)} \cdot \frac{-\log(2r)}{-\log r} \longrightarrow \dim_M(X),$$

since $\frac{-\log(2r)}{-\log r} \rightarrow 1$ and the Minkowski dimension is invariant under the scale change $r \mapsto 2r$. Similarly,

$$\frac{\log N(r)}{-\log r} \longrightarrow \dim_M(X).$$

By the squeeze theorem,

$$\frac{\log M(r)}{-\log r} \longrightarrow \dim_M(X).$$

□

Connection to the local profile $M_c(k)$. The global profile $M(r)$ (the number of centers at scale r) aggregates the density structure of X across all locations. When the geometry exhibits a transition—for example, a change in local dimension or curvature—the growth exponent of $M(r)$ changes accordingly. The next subsection explains how such global transitions manifest as local signals through the discrete second differences $\Delta^2 M_c(k)$.

4.7 Local–Global Scale Correspondence

The preceding results establish two complementary structural layers of RCI: (i) a *local* layer, governed by the behaviour of the population k -NN radii and the radius potential $u_k(x)$ (Theorem 4.5); and (ii) a *global* layer, governed by the multiscale growth of the farthest-point complexity profile $M(r)$, which is fractal complete (Theorem 4.19). This subsection describes how these two layers interact at the level of scale. The correspondence stated below is consistent with all structural properties established in earlier sections, but its full regularity theory requires additional analytical control (e.g., differentiability of $M(r)$), and is therefore presented as a heuristic statement.

Proposition 4.20 (Local–Global Scale Correspondence (Heuristic)). *Let $M(r)$ denote the global complexity profile at scale r , and let $M_c(k)$ denote the local RCI profile centred at c , computed at the population radius $r_k(c)$. Under Assumptions 1 and 2, and in the asymptotic regime $k/N \rightarrow 0$, the following heuristic relations hold:*

$$M_c(k) \approx M(r_k(c)), \quad \Delta^2 M_c(k) \approx M''(r_k(c)) (r_{k+1} - r_k)^2.$$

Scale correspondence mechanism . The r -net structure of the RCI cover $\mathcal{C}(r)$ implies that $M(r)$ measures, up to uniformly bounded overlap, the number of r -separated regions needed to cover X . For any analytically fixed centre c —and in particular for each algorithmic centre $c = c_j$ selected by the residual RCI seeding dynamics—the population radius $r_k(c)$ satisfies $\nu(B_{r_k(c)}(c)) = k/N$, so $M_c(k)$ counts the number of sample points within a ball of radius $r_k(c)$. Since both quantities capture volumetric complexity at comparable scales, one expects $M_c(k) \approx M(r_k(c))$, up to $O(1)$ fluctuations.

Assuming that $M(r)$ varies smoothly at the relevant scale, a discrete second-difference expansion yields

$$\Delta^2 M_c(k) \approx M''(r_k(c)) (\Delta r)^2, \quad \Delta r = r_{k+1} - r_k,$$

with higher-order contributions suppressed under the scaling $r_k(c) \lesssim (k/N)^{1/m}$. This identifies $\Delta^2 M_c(k)$ as a discrete curvature probe for $M(r)$ at the matched scale $r = r_k(c)$.

Corollary 4.21 (Detection of Growth-Exponent Transitions (Heuristic)). *Let*

$$D(r) := -\frac{d \log M(r)}{d \log r}$$

denote the local growth exponent of the global profile. If $D(r)$ undergoes a discontinuity, sharp inflection, or regime change at $r = r^$, then for any centre c such that $r_{k^*}(c) \approx r^*$, the discrete curvature statistic $\Delta^2 M_c(k^*)$ exhibits a corresponding peak, sign change, or non-degenerate extremum. Thus $\Delta^2 M_c(k)$ serves as a detector of structural transitions in $M(r)$.*

Relation to the curvature law. The discussion above identifies the intermediate-scale mechanism linking the *local* geometric behavior—captured by the curvature law for $\Delta^2 M_c(k)$ in smooth Riemannian regions (Theorem 3.5)—with the *global* fractal completeness of the complexity profile $M(r)$ (Theorem 4.19). In essence, variations in local curvature influence the discrete second differences of the local profiles $M_c(k)$, while aggregated changes across the space are reflected in the growth exponent of $M(r)$.

Connection to Appendix C. The heuristic correspondences developed in this subsection rely on the functorial scale structure and geometric compatibility formalized in Appendix C, where the scale sheaf, the spectral Čech nerve, and the induced type- A persistence module provide the regularity framework that justifies matching the local profile $M_c(k)$ with the global complexity profile $M(r_k(c))$.

4.8 Summary

The results above assemble a unified metric–measure and Morse–fractal foundation for RCI:

- Small-ball regularity yields a well-defined local dimension m together with a mass profile $R_0(x)$, and the radius potential $u_k(x)$ is uniformly C^0 -conjugate to the mass potential $v(x)$.
- The one-dimensional discrete Morse structure in k is stable under small uniform perturbations (Lemma 4.7).
- In doubling spaces, the residual farthest-point seeding used by RCI produces geometrically controlled covers $\mathcal{C}(r)$ whose complexity profile is $M(r)$.
- Under contractibility assumptions, the nerve $\mathcal{N}(r)$ captures the topology of $U(r)$, and drops in $B(r) = \beta_0(U(r))$ correspond to Morse-type transitions in the nerve.
- The global complexity profile $M(r)$ is fractal complete and recovers the Minkowski dimension (Theorem 4.19).
- The local profile $M_c(k)$ and the global profile $M(r)$ are asymptotically matched through the scale correspondence $M_c(k) \approx M(r_k(c))$. The structural conditions supporting this identification—including functorial behaviour across scales and geometric compatibility—are formalized in Appendix C. Consequently, transitions in the growth exponent of $M(r)$ are detected by the discrete curvature statistic $\Delta^2 M_c(k)$ (Corollary 4.21).

The bridge between local curvature (Theorem 3.5) and global dimension (Theorem 4.19) is therefore provided by the scale correspondence (Proposition 4.20), which allows $\Delta^2 M_c(k)$ to detect geometric transitions across all relevant scales.

In the next section we introduce the *Morse Erasure Index* (MEI), a quantitative criterion measuring how effectively a clustering algorithm preserves these geometric transitions.

5 A Morse-Theoretic Metric of Structural Fidelity

We introduce a quantitative metric that evaluates how faithfully a clustering preserves geometric transitions in a dataset. The **Morse Erasure Index (MEI)** measures the fraction of variation in a Morse-type field that remains *visible across cluster boundaries*. Unlike classical external validation indices, the MEI is *intrinsic*: it depends only on the geometry of the data through the graph and the Morse field, not on external labels or ground truth. High MEI indicates that the partition respects geometric structure; low MEI indicates that geometric transitions are absorbed within clusters.

5.1 Morse Field and Graph Structure

Let $G = (V, E, w)$ be the mutual k -nearest neighbour graph on the data, equipped with self-tuning weights of the form

$$w_{ij} = \exp\left(-\frac{d(x_i, x_j)^2}{\sigma_i \sigma_j}\right),$$

where σ_i denotes the distance from x_i to its k -th nearest neighbour (cf. Section 3.2). This produces a locally scaled, density-adaptive geometric structure. Throughout this section, G is treated as an undirected weighted graph, with $w_{ij} = w_{ji} > 0$ for every edge $\{i, j\} \in E$. Unless otherwise stated, we

work on a connected evaluation graph. If the mutual k -nearest neighbour graph has several connected components, the MEI is computed on each connected component or on the largest connected component.

The Morse field is defined as the negative log-density,

$$f(x_i) := -\log \hat{\rho}(x_i),$$

where $\hat{\rho}(x_i)$ is the m_ρ -nearest-neighbour density estimate. The sign convention is immaterial for MEI, since only absolute differences $|f_i - f_j|$ enter the graph total variation. The geometric variability of f along the graph is measured by the total variation

$$\text{TV}_{\text{total}}(f) := \sum_{\{i,j\} \in E} w_{ij} |f_i - f_j|.$$

Given a clustering $\ell : V \rightarrow \{1, \dots, K\}$, the portion of this variation that lies *within* clusters is

$$\text{TV}_{\text{intra}}(f, \ell) := \sum_{\substack{\{i,j\} \in E \\ \ell(i) = \ell(j)}} w_{ij} |f_i - f_j|.$$

5.2 Definition of the MEI

The Morse Erasure Index is defined by

$$\text{MEI}(f, \ell) := 1 - \frac{\text{TV}_{\text{intra}}(f, \ell)}{\text{TV}_{\text{total}}(f)}.$$

Remark. The MEI is well-defined whenever $\text{TV}_{\text{total}}(f) > 0$, which holds whenever f is non-constant on the connected evaluation graph G (cf. the connectivity convention above). Since

$$0 \leq \frac{\text{TV}_{\text{intra}}}{\text{TV}_{\text{total}}} \leq 1,$$

we always have $\text{MEI} \in [0, 1]$.

5.3 Formal Properties

For a split $C = C_1 \sqcup C_2$, define the cut-edge set

$$E_{\text{cut}}(C_1, C_2) := \{\{i, j\} \in E : i \in C_1, j \in C_2\}.$$

Proposition 5.1 (MEI bounds and refinement behavior). *Let $\ell : V \rightarrow \{1, \dots, K\}$ be any clustering with K clusters and let $f : V \rightarrow \mathbb{R}$ be a non-constant Morse field. Then:*

1. $0 \leq \text{MEI}(f, \ell) \leq 1$.
2. $\text{MEI}(f, \ell) = 1$ if and only if $\text{TV}_{\text{intra}}(f, \ell) = 0$.
3. $\text{MEI}(f, \ell) = 0$ if and only if $\text{TV}_{\text{intra}}(f, \ell) = \text{TV}_{\text{total}}(f)$.
4. Let ℓ' be a refinement of ℓ obtained by splitting a single cluster C into C_1, C_2 . Then

$$\text{MEI}(f, \ell') - \text{MEI}(f, \ell) = \frac{1}{\text{TV}_{\text{total}}(f)} \sum_{\{i,j\} \in E_{\text{cut}}(C_1, C_2)} w_{ij} |f_i - f_j|.$$

In particular, the MEI increases if and only if the split separates edges with non-zero weighted variation.

Proof. (1) **Range bounds.**

By definition,

$$\text{MEI}(f, \ell) = 1 - \frac{\text{TV}_{\text{intra}}(f, \ell)}{\text{TV}_{\text{total}}(f)}.$$

Since $\text{TV}_{\text{intra}}(f, \ell)$ sums over a subset of edges in E with non-negative weights and differences, we have

$$0 \leq \text{TV}_{\text{intra}}(f, \ell) \leq \text{TV}_{\text{total}}(f).$$

The upper bound holds because every intra-cluster edge $\{i, j\}$ with $\ell(i) = \ell(j)$ is also counted in the total variation. Since $\text{TV}_{\text{total}}(f) > 0$ (because f is non-constant on the connected evaluation graph), division is well-defined. Therefore,

$$0 \leq \frac{\text{TV}_{\text{intra}}}{\text{TV}_{\text{total}}} \leq 1,$$

which implies

$$0 \leq 1 - \frac{\text{TV}_{\text{intra}}}{\text{TV}_{\text{total}}} \leq 1.$$

(2) Characterization of $\text{MEI} = 1$.

Forward direction ($\Rightarrow$): Suppose $\text{MEI}(f, \ell) = 1$. Then

$$1 - \frac{\text{TV}_{\text{intra}}}{\text{TV}_{\text{total}}} = 1 \quad \Rightarrow \quad \frac{\text{TV}_{\text{intra}}}{\text{TV}_{\text{total}}} = 0 \quad \Rightarrow \quad \text{TV}_{\text{intra}} = 0.$$

Reverse direction ($\Leftarrow$): Suppose $\text{TV}_{\text{intra}}(f, \ell) = 0$. Then

$$\text{MEI}(f, \ell) = 1 - \frac{0}{\text{TV}_{\text{total}}} = 1.$$

(3) Characterization of $\text{MEI} = 0$.

Forward direction ($\Rightarrow$): Suppose $\text{MEI}(f, \ell) = 0$. Then

$$1 - \frac{\text{TV}_{\text{intra}}}{\text{TV}_{\text{total}}} = 0 \quad \Rightarrow \quad \frac{\text{TV}_{\text{intra}}}{\text{TV}_{\text{total}}} = 1 \quad \Rightarrow \quad \text{TV}_{\text{intra}} = \text{TV}_{\text{total}}.$$

Reverse direction ($\Leftarrow$): Suppose $\text{TV}_{\text{intra}}(f, \ell) = \text{TV}_{\text{total}}(f)$. Then

$$\text{MEI}(f, \ell) = 1 - \frac{\text{TV}_{\text{total}}}{\text{TV}_{\text{total}}} = 1 - 1 = 0.$$

(4) Effect of cluster refinement.

Let ℓ be a clustering and let ℓ' be the clustering obtained by splitting cluster C into two non-empty subsets C_1, C_2 such that $C = C_1 \sqcup C_2$ and $C_1 \cap C_2 = \emptyset$. All other clusters remain unchanged. Write $E_{\text{cut}} = E_{\text{cut}}(C_1, C_2)$ for the cut-edge set defined above.

We partition the edge set E into three disjoint subsets:

$$\begin{aligned} E_{\text{old-intra}} &:= \{\{i, j\} \in E : \ell(i) = \ell(j), \{i, j\} \notin E_{\text{cut}}\}, \\ E_{\text{inter}} &:= E \setminus (E_{\text{cut}} \cup E_{\text{old-intra}}). \end{aligned}$$

For the original clustering ℓ :

$$\text{TV}_{\text{intra}}(f, \ell) = \sum_{\{i, j\} \in E_{\text{cut}}} w_{ij} |f_i - f_j| + \sum_{\{i, j\} \in E_{\text{old-intra}}} w_{ij} |f_i - f_j|.$$

For the refined clustering ℓ' , edges in E_{cut} are now inter-cluster (since $\ell'(i) \neq \ell'(j)$ for $i \in C_1, j \in C_2$), so:

$$\text{TV}_{\text{intra}}(f, \ell') = \sum_{\{i, j\} \in E_{\text{old-intra}}} w_{ij} |f_i - f_j|.$$

Thus,

$$\text{TV}_{\text{intra}}(f, \ell) - \text{TV}_{\text{intra}}(f, \ell') = \sum_{\{i,j\} \in E_{\text{cut}}} w_{ij} |f_i - f_j|.$$

Since $\text{TV}_{\text{total}}(f)$ depends only on the graph and the Morse field (not on the clustering), it is the same for both ℓ and ℓ' . Therefore,

$$\begin{aligned} \text{MEI}(f, \ell') - \text{MEI}(f, \ell) &= \left(1 - \frac{\text{TV}_{\text{intra}}(f, \ell')}{\text{TV}_{\text{total}}(f)}\right) - \left(1 - \frac{\text{TV}_{\text{intra}}(f, \ell)}{\text{TV}_{\text{total}}(f)}\right) \\ &= \frac{\text{TV}_{\text{intra}}(f, \ell) - \text{TV}_{\text{intra}}(f, \ell')}{\text{TV}_{\text{total}}(f)} \\ &= \frac{1}{\text{TV}_{\text{total}}(f)} \sum_{\{i,j\} \in E_{\text{cut}}(C_1, C_2)} w_{ij} |f_i - f_j|. \end{aligned}$$

The MEI increases if and only if the right-hand side is strictly positive, which occurs if and only if there exists at least one edge $\{i, j\} \in E_{\text{cut}}(C_1, C_2)$ with $w_{ij} > 0$ and $f_i \neq f_j$. This is precisely the condition that the split separates edges with non-zero weighted variation of the Morse field. $\square$

Corollary 5.2 (Robustness to flat-region splits). *Let ℓ be a clustering and let ℓ' be obtained by splitting a cluster C into C_1, C_2 . Let*

$$E_{\text{cut}}(C_1, C_2) := \{\{i, j\} \in E : i \in C_1, j \in C_2\},$$

and set

$$W_{\text{cut}} := \sum_{\{i,j\} \in E_{\text{cut}}(C_1, C_2)} w_{ij}.$$

If

$$\max_{\{i,j\} \in E_{\text{cut}}(C_1, C_2)} |f_i - f_j| \leq \epsilon,$$

with the convention that the claim is trivial when $E_{\text{cut}}(C_1, C_2) = \emptyset$, then

$$|\text{MEI}(f, \ell') - \text{MEI}(f, \ell)| \leq \frac{\epsilon W_{\text{cut}}}{\text{TV}_{\text{total}}(f)}.$$

Proof. By Proposition 5.1(4),

$$\begin{aligned} |\text{MEI}(f, \ell') - \text{MEI}(f, \ell)| &= \frac{1}{\text{TV}_{\text{total}}(f)} \sum_{\{i,j\} \in E_{\text{cut}}(C_1, C_2)} w_{ij} |f_i - f_j| \\ &\leq \frac{1}{\text{TV}_{\text{total}}(f)} \sum_{\{i,j\} \in E_{\text{cut}}(C_1, C_2)} w_{ij} \cdot \epsilon \\ &= \frac{\epsilon}{\text{TV}_{\text{total}}(f)} \sum_{\{i,j\} \in E_{\text{cut}}(C_1, C_2)} w_{ij} \\ &= \frac{\epsilon \cdot W_{\text{cut}}}{\text{TV}_{\text{total}}(f)}. \end{aligned}$$

$\square$

5.4 Interpretation

Proposition 5.1 establishes the following structural properties of the MEI:

- **MEI close to 1:** geometric transitions of f occur primarily across cluster boundaries (high structural fidelity);
- **MEI close to 0:** geometric variation is absorbed within clusters (structural collapse);
- **Automatic regularization:** refining a clustering by splitting inside regions of low Morse variation produces negligible change in MEI (Corollary 5.2), while splits along high-variation edges increase the MEI proportionally to the separated variation (Proposition 5.1(4)).

The MEI therefore provides a single scalar quantifying how well a partition respects the intrinsic Morse geometry of the data. In all benchmarks (Section 6), algorithms are compared by computing their MEI on a fixed evaluation graph: the clustering with the highest MEI is interpreted as exhibiting the strongest structural fidelity.

5.5 Dependence on the Number of Clusters

The number of clusters K does not appear explicitly in the definition of MEI because it is already encoded in the variational structure of the ratio. Proposition 5.1(4) formalizes the intrinsic balance between granularity and geometric fidelity: refining a clustering (increasing K) reduces TV_{intra} , but only refinements performed at edges where $|f_i - f_j|$ is large produce a meaningful increase in MEI. Splits within flat regions of the Morse field do not alter the index (Corollary 5.2).

The limiting case $K = |V|$ yields $\text{TV}_{\text{intra}} = 0$ and $\text{MEI} = 1$, corresponding to a decomposition that separates all transitions but conveys no macroscopic structure. This is not a degeneracy of the index: it is the maximal-resolution limit of Morse fidelity. At this scale, no transition of the Morse field is erased. The MEI is designed to measure preservation of Morse variation; macroscopic compression belongs to a different layer of the theory.

Conversely, coarse clusterings that merge across transition regions have low MEI, indicating structural loss.

6 Empirical Validation and Benchmarks

This section examines how the theoretical framework developed in Sections 4–5 manifests in computational practice. The goal is not to test continuum theorems directly—which is impossible in principle—but to determine whether empirical behaviour aligns with the structural regime predicted by the Morse-theoretic and fractal layers of RCI.

All evaluations use the *Morse Erasure Index* (MEI) (Section 5.2), an intrinsic, label-free metric defined once the evaluation graph and Morse field have been fixed, which quantifies how much geometric variation of the Morse field remains visible across cluster boundaries. The clustering with the highest MEI is interpreted as exhibiting the strongest structural fidelity to the underlying geometry.

Appendix 12 provides complete reproducibility, including dataset construction, graph building, hyperparameter search for the classical baselines, and the full benchmarking pipeline.

6.1 Experimental Configuration

We evaluate RCI and five classical clustering methods across eight geometric datasets: sphere, saddle surface, torus, dumbbell surface, Hopf link, spiral, Swiss roll, and trefoil knot.

Evaluation graphs and geometric neutrality. The MEI metric requires a graph structure to measure how the Morse field varies across cluster boundaries.

Classical clustering methods (KMeans, DBSCAN, HDBSCAN, Spectral Clustering, and Gaussian Mixture Models) operate directly in the ambient Euclidean metric and return only partition labels; they do not construct graphs during their execution. For MEI evaluation, their partitions are assessed on a *self-tuning* mutual k -NN graph $G_{\text{eval}}(X)$ built post-hoc using the same adaptive kernel scales $\sigma_i = d(x_i, x_{(k)})$ and Gaussian affinity weights $W_{ij} = \exp(-d_{ij}^2/(\sigma_i\sigma_j))$ used internally by RCI. This ensures that all classical methods are evaluated on a common density-adaptive geometric substrate independent of their clustering decisions. RCI is evaluated on its internally produced graph, as described below.

RCI is a purely metric algorithm: its defining quantities—radial profiles $M_c(k)$, normalized geometric density, curvature signals $\Delta^2 M_c(k)$, and boundary indices κ^* —are computed directly from the ambient metric d . In the benchmark implementation used here, RCI is run in its non-spectral configuration. Although the code calls the `SpectralRCI` class, it explicitly sets `use_spectral=False`; no spectral embedding is used in the reported experiments. RCI nevertheless constructs an internal self-tuning mutual k -NN graph. This graph is not a spectral chart; it is the neighbourhood structure produced by the RCI run and is injected into the MEI computation. For MEI evaluation, RCI is therefore assessed on

the graph it constructed during clustering, ensuring consistency between the algorithmic run and the structural metric.

Classical baselines. KMeans, DBSCAN, HDBSCAN, Spectral Clustering, and Gaussian Mixture Models are hyperparameter-tuned via randomized search over standard parameter ranges. The best model under Silhouette and Davies–Bouldin indices is then evaluated structurally via MEI.

RCI configuration. RCI performs no hyperparameter tuning. In the benchmark code, the algorithm is run with `use_spectral=False`; hence no spectral embedding is used for clustering or MEI computation. The value `n_eigenvectors=5` is present only as an inactive constructor argument in this non-spectral configuration. The geometric quantities governing the clustering—profiles, curvature signals, and boundary indices—are computed in the ambient metric as described in Section 3.1.

Density detection vs. geometric transition detection. Density-based algorithms such as DBSCAN and HDBSCAN respond primarily to *sampling density*—the local concentration of points in the dataset. Because this quantity does not distinguish intrinsic geometric structure from variations in sampling, these methods may segment a smooth manifold into many small components when the scale parameter is too small, or merge distinct geometric regions when the scale is too large. This behaviour reflects a structural limitation: sampling density is not a geometric invariant.

RCI approaches the problem from a different perspective. It measures *geometric density* through the Morse profile $M_c(k)$, which accumulates mass radially while normalizing by the intrinsic m -dimensional volume growth $\alpha \log r_k$. This normalization filters out sampling artifacts and produces a scale-aware estimate of how the underlying space expands around c . The second variation $\Delta^2 M_c(k)$ then highlights *transitions* in this normalized field—changes associated with geometric features such as curvature variation, neck regions, or topological constrictions. Because these transitions arise from the geometry rather than the sampling, RCI can identify structurally meaningful boundaries even under uniform or adversarial sampling.

Remark 6.1 (On neutrality, density, and decision regimes). A potential concern is that RCI is evaluated on its internal graph, whereas classical methods are evaluated on a post-hoc mutual k -NN graph. This reflects the benchmark protocol implemented in Appendix 12: RCI produces a graph during its non-spectral run, while classical algorithms return only partition labels and therefore require an evaluation graph to compute MEI.

Both graph constructions use the same self-tuning mutual k -NN principle. The difference lies in their role. For classical methods, the graph is a post-hoc geometric scaffold used only to measure Morse-field variation across the returned partition. For RCI, the graph is part of the neighbourhood structure generated by the algorithmic run and is injected directly into MEI. Thus the comparison is made at the level of structural fidelity: given a partition and the corresponding benchmark evaluation graph, MEI measures how much Morse-field variation remains visible across cluster boundaries. No external labels or ground-truth information are used in either case.

6.2 Structural Scoreboard via MEI

Table 1 reports the empirical MEI across all benchmark datasets. Higher values indicate greater preservation of geometric transitions.

6.3 Discussion

The empirical results reveal a consistent pattern: RCI exhibits substantially higher MEI than classical clustering methods across diverse geometric regimes.

Dominant structural fidelity of RCI. RCI achieves the highest MEI on 7 of 8 datasets, with a mean of 0.57, roughly double that of the second-best performer (HDBSCAN) and an order of magnitude higher than KMeans, Spectral Clustering, and GMM. The advantage is most pronounced for datasets with strong topological or curvature structure (sphere, torus, dumbbell, trefoil), where the Morse transition is clear and compact.

Dataset	RCI	KMeans	DBSCAN	HDBSCAN	Spectral	GMM
Sphere	0.70	0.12	0.56	0.44	0.10	0.13
Saddle	0.03	0.06	0.52	0.41	0.07	0.07
Torus	0.74	0.03	0.00	0.44	0.03	0.03
Dumbbell	0.75	0.00	0.26	0.27	0.00	0.00
Link	0.47	0.04	0.02	0.10	0.03	0.03
Spiral	0.61	0.03	0.03	0.30	0.03	0.03
Swiss Roll	0.61	0.02	0.17	0.04	0.04	0.02
Trefoil	0.68	0.03	0.05	0.37	0.03	0.03
Mean	0.57	0.04	0.20	0.30	0.04	0.04
Std Dev	0.22	0.03	0.21	0.14	0.03	0.04

Table 1: MEI-based structural comparison across all benchmark datasets. RCI attains the highest mean MEI (0.57), winning in 7 of 8 datasets. Bold values indicate the best structural fidelity per dataset. Classical algorithms typically operate in a regime of structural erasure, with mean MEI values near zero.

Structural collapse in classical methods. KMeans, Spectral Clustering, and GMM consistently attain MEI values near zero. This is not a tuning issue: these methods optimise convexity, spectral gaps, or density modes, none of which align with the transitions of the Morse field. The result is *structural erasure*: geometric variation is absorbed within clusters rather than reflected at their boundaries.

DBSCAN and HDBSCAN perform moderately well due to their local density adaptivity, but lack the transition-tracking behaviour characteristic of RCI and do not consistently align boundaries with Morse transitions.

The saddle anomaly. The saddle surface is the single dataset where RCI underperforms. This failure is not an artefact of the MEI metric, but a genuine limitation of the current formulation of RCI.

In negatively curved regions, geodesic balls grow *exponentially* with radius, while the RCI penalty term with $\alpha = m$ cancels only the *polynomial* component of volume growth. As a result, the normalised profile

$$M_c(k) = A_c(k) - m \log |B_k(c)|$$

does not adequately compensate for the underlying hyperbolic expansion. The exponential term dominates, delaying the curvature flip in $\Delta^2 M_c(k)$ and causing RCI to place the boundary substantially farther from the true Morse transition.

Thus, on hyperbolic geometries, the algorithm genuinely fails to detect the transition at the correct scale. This suggests the need for an adaptive or data-driven normalisation scheme, capable of adjusting to exponential volume regimes.

Robustness across geometric complexity. Except for the hyperbolic case, RCI maintains strong and stable structural fidelity across compact manifolds, multi-component structures, and embedded curves. Variation in MEI reflects differences in geometric difficulty rather than algorithmic instability.

Implications for the theoretical framework. The empirical results are consistent with the theoretical motivation behind RCI. In most datasets, the detected boundaries align with regions where the Morse profile changes behaviour, and the method adjusts naturally to the local scale and density of the data without requiring parameter tuning. This supports the idea that the RCI construction captures meaningful geometric transitions in a broad range of settings.

Classical methods behave differently because their objectives—convexity, spectral separation, or density thresholds—do not necessarily reflect the structure encoded in the Morse field. As a result, their MEI scores are lower in cases where geometry plays a central role, even when they perform well under their own internal criteria.

The saddle case highlights a specific limitation: the normalisation $M_c(k) = A_c(k) - m \log |B_k(c)|$ compensates for polynomial volume growth, but not for the exponential growth that appears in strongly negative curvature. In such settings, RCI tends to place boundaries farther from the true transition, which explains its lower MEI on this dataset.

Overall, the method performs as expected across the majority of geometric regimes tested, with the hyperbolic behaviour of the saddle serving mainly as an indicator of where the current normalisation scheme may require refinement.

7 Principled Hyperparameter Specification

Classical clustering algorithms rely on hyperparameters with no intrinsic geometric meaning: KMeans requires the number of clusters K ; DBSCAN requires $(\varepsilon, \text{minPts})$; spectral methods require kernel choices, bandwidths, and an embedding dimension. These parameters are typically selected heuristically and do not arise from analytic or geometric structure.

In contrast, the RCI framework is *parameter-minimal*: nearly all internal quantities are fixed by the analytic structure developed in Sections 3–4 and in Appendix 14. RCI does possess internal parameters, but they belong to two narrow classes: (i) quantities canonically fixed by the theory; and (ii) quantities constrained to lie within a stable theoretical regime rather than freely tunable. Only one degree of freedom remains externally chosen: the observational scale r .

This section specifies the status of each quantity and distinguishes: (i) canonically fixed parameters; (ii) regime-restricted parameters; and (iii) the single operational choice.

7.1 Operational Parameter: Analysis Scale

Resolution scale r . The farthest-point procedure produces coverings $\{B(c_j, r_j^*)\}$ whose nested structure defines the multiscale profile $M(r)$ (Section 4.6). Fixing r selects the geometric resolution at which this profile is evaluated. No theorem identifies a unique canonical value of r ; different scales reveal different geometric regimes. Thus r is the *only* operational parameter chosen by the practitioner.

7.2 Canonically Fixed Structural Quantities

Two quantities are uniquely forced by the analytic structure of the RCI theory.

(S1) Volumetric penalty $\alpha = m$. The Curvature Law (Theorem 3.5) shows that

$$\mathbb{E}[\Delta^2 M_c(k)] \approx -C_m S_g(c) \Delta(r_k^2) \quad \text{if and only if} \quad \alpha = m.$$

Setting $\alpha = m$ cancels the dominant volumetric term $\propto r^m$ and isolates the curvature term $\propto S_g r^{m+2}$. Any other choice mixes curvature with volume. Thus $\alpha = m$ is analytically forced.

(S2) Contrast function $\varphi = \log$. The logarithm is the canonical contrast for density-based analysis because:

- (i) it converts the multiplicative small-ball expansion $\nu(B_r) = \omega_m r^m R(x, r)$ into an additive form suited to discrete differences;
- (ii) it stabilizes variance, enabling the concentration bounds used in Theorem 3.4;
- (iii) it yields the radius–mass C^0 conjugacy (Theorem 4.5) necessary for scale alignment.

Other monotone transforms are possible, but $\varphi = \log$ is the unique choice compatible with the information-theoretic meaning of density and with the Riemannian small-ball expansion.

7.3 Structurally Constrained but Not Uniquely Fixed Quantities

Three quantities must lie in theoretically justified regimes; the theory constrains the admissible range but does not specify a unique value.

(R1) Warm-up neighbourhood size $m_0 \geq 3$. Centred second differences are used (Section 3.1), so $\Delta^2 M_c(k)$ is defined starting at $k = 2$. Requiring additionally $m_0 \geq 3$ allows the simultaneous presence of $\Delta^2 M_c(k)$ and $\Delta^2 M_c(k-1)$ in the concavity-crossing criterion for κ^* .

(R2) Density stencil size m_ρ . Local density is estimated from a fixed neighbourhood of size m_ρ (Section 3.1). The theory requires m_ρ to remain within a fixed small-scale regime (typically $5 \leq m_\rho \leq 20$), ensuring that:

- $r_\rho(i) \sim (m_\rho/N)^{1/m}$ (Lemma 4.3);
- concentration bounds hold uniformly;
- the estimator avoids the asymptotic regime $m_\rho \rightarrow \infty$.

The regime is fixed, but the precise value is not unique. A robust default is $m_\rho = 7$.

(R3) Spectral embedding dimension d . This parameter is active only in the spectral implementation of RCI. In the non-spectral implementation used in the benchmark suite, no spectral embedding is used and d is inactive. The embedding $\Phi_d : X \rightarrow \mathbb{R}^d$ serves two roles: (i) computational acceleration of nearest-neighbour queries; and (ii) theoretical fidelity via local bi-Lipschitz behaviour (Theorem 3.3). Admissible values of d must satisfy a spectral-gap condition

$$\lambda_{d+1}(L) - \lambda_d(L) \geq \delta > 0,$$

which restricts the range of valid choices without fixing a unique d . Standard practice uses either the eigengap heuristic or a held-out structural metric.

7.4 Synthesis: Parameter-Minimality Revisited

RCI contains only one operational degree of freedom: the observational scale r . All remaining quantities fall into two classes:

- **canonically fixed** ($\alpha = m$, $\varphi = \log$);
- **restricted to a stable theoretical regime** ($m_0 \geq 3$, $m_\rho \approx 7$, d chosen via eigengap when the spectral implementation is active).

In contrast, classical clustering algorithms require:

- KMeans: number of clusters K (no geometric basis);
- DBSCAN: $(\varepsilon, \text{minPts})$ controlling a density scale;
- Spectral clustering: affinity kernel, bandwidth, and number of clusters;
- GMM: number of components and covariance model.

Thus RCI is *parameter-minimal* in a rigorous and qualified sense: there are no freely tunable modelling parameters, and the only external selection is the geometric resolution r .

8 Clarifying Objections

This section addresses natural objections that arise when a clustering framework is linked to geometric analysis, discrete Morse theory, and small-ball metric geometry. Each objection is answered strictly on the basis of the formal results proved in Sections 4 and 3, together with the analytic estimates of Appendix 14 and the structural admissibility argument of Appendix 15. We distinguish operational choices from structural quantities that are fixed or regime-constrained by the theory.

8.1 Structural Objections

8.1.1 Objection 1: The theoretical assumptions (smoothness, density regularity, manifold structure) are too strong.

Response. The analytic expansions in Appendix 14 (spectral bi-Lipschitz regularity, curvature expansion, concentration bounds) require smoothness. These hypotheses apply *only* to those expansions.

The foundational metric-measure and Morse results of Section 4 rely primarily on: (i) small-ball regularity, (ii) doubling geometry, and (iii) weak continuity of the mass profile. However, the nerve-theoretic correspondence between drops of $M(r)$ and topological transitions requires *contractible ball intersections* (Assumption 3), a geometric assumption not implied by doubling alone. This assumption is essential for

invoking the Nerve Theorem. Thus the framework avoids full manifold structure but requires contractibility for the topological analysis.

Appendix 15 separates this analytic layer from the structural core. The minimal RCI base $\mathcal{B}_{\text{RCI}}$ is generated from points and distances alone. Smoothness, density regularity, spectral gaps, and probabilistic sampling assumptions are needed for quantitative convergence claims, but not for the existence of the radial base, the restriction structure, the wall of type change, or admissible certification.

8.1.2 Objection 2: The RCI profile and the operator $\Delta^2 M_c(k)$ are ad hoc constructions.

Response. They follow directly from the analytic structure of the small-ball expansion. Appendix 14 shows that

$$\mathbb{E}[M_c(k)] = (\text{volumetric term}) - \alpha \log r_k^{\text{spec}} + (\text{curvature term}) + o(1).$$

Setting $\alpha = m$ cancels all first-order volumetric terms. The discrete second difference $\Delta^2 M_c(k)$ is then the unique second-order operator that isolates the curvature term. Among all finite-difference operators, Δ^2 is selected by the Taylor expansion. Therefore the RCI profile is forced by analysis, not designed ad hoc.

8.1.3 Objection 3: The sheaf-theoretic language is an external decoration.

Response. The sheaf formalism is not assumed by RCI. Appendix 15 shows that RCI first generates the relevant base and restriction structure from the minimal metric data:

$$(X, d) \longrightarrow \mathcal{B}_{\text{RCI}} \longrightarrow \mathcal{I} \longrightarrow \mathcal{W}_n \longrightarrow \mathcal{S}_t.$$

The sheaf-theoretic language enters only after these objects exist, as a way of recognizing that compatible local invariant data glue over admissible sieves. Thus RCI does not invoke sheaves as an external authority; it produces the restriction data that make the sheaf interpretation available.

8.1.4 Objection 4: Focusing on merging transitions is too narrow.

Response. This focus follows from Section 4.5, where drops in the component-count profile $B(r) = \beta_0(U(r))$ correspond to covering-merging events in the induced nerve filtration. The purpose of RCI is to detect these geometric critical transitions and to use them as clustering boundaries. RCI is a clustering algorithm guided by geometric criticality, not a generic geometric probe.

Empirical results (Section 6) confirm that these transitions capture structure across a wide range of datasets.

8.2 Implementation Objections

8.2.1 Objection 5: Implementing RCI on k -NN graphs deviates from the continuum theory.

Response. Appendix 14 provides three forms of discrete control: (i) a bounded-difference inequality for $A_c(k)$, (ii) sub-Gaussian concentration for shell averages, (iii) a bias expansion for the log-density estimator. These ensure that discrete $M_c(k)$ differs from its population analogue by $o(1)$ uniformly in c . Since RCI depends only on metric and small-ball structure, the k -NN graph yields a faithful discrete implementation.

8.2.2 Objection 6: Spectral embeddings may distort geometry and invalidate the Morse interpretation.

Response. RCI is *defined* on the ambient metric (X, d) ; spectral embeddings are optional computational surrogates. The local bi-Lipschitz theorem (theorem 3.3) ensures that small-scale distances are preserved up to $1 \pm \varepsilon_N$, so shell orderings, radii, the profile $M_c(k)$, and the boundary index κ^* are stable under the separation margins assumed in the boundary-stability theorem.

In the benchmark implementation, structural evaluation follows the protocol documented in Appendix 12: RCI is evaluated on the graph produced internally by its non-spectral run, while classical algorithms are evaluated on a freshly constructed mutual k -NN graph built from the original data X . Thus MEI does not use external labels or ground-truth information, and the evaluation graph is fixed by the benchmark protocol.

8.3 Robustness Objections

8.3.1 Objection 7: Noise destroys density information, making $\Delta^2 M_c(k)$ unstable.

Response. Appendix 14 establishes uniform sub-Gaussian concentration for shell averages. The boundary stability theorem (theorem 3.4) states that if the population profile has a non-degenerate curvature gap $\gamma > 0$, then

$$\Pr(\kappa^\star = k^\circ) \geq 1 - N^{-C}.$$

Noise can roughen the Morse profile, but it does not manufacture structural criticality. An RCI boundary is not triggered by an isolated fluctuation; it is produced by a scale-persistent change in the radial profile. Under the non-degenerate gap condition $\gamma > 0$, the concentration bounds in Appendix 14 imply that empirical fluctuations remain below the population transition gap with high probability. Therefore, within the regime of the theorem, noise may blur the detected transition, but it cannot promote arbitrary sampling roughness into a stable RCI boundary.

Appendix 15 gives the structural counterpart of this analytic claim. Noise may move or thicken the empirical wall $\mathcal{W}_n$, and Appendix 14 controls this movement under regularity assumptions. But once an admissible sieve $\mathcal{S}_t$ avoids the wall, the certified object is obtained by restriction:

$$\mathcal{S}_t \cap \mathcal{W}_n = \emptyset \Rightarrow \widehat{\mathcal{I}}_n|_{\mathcal{S}_t} \cong \mathcal{I}|_{\mathcal{S}_t}.$$

8.3.2 Objection 8: Multi-scale variability of $M(r)$ indicates instability.

Response. $M(r)$ is the complexity profile of the covering filtration. Each drop is a genuine merging event (Section 4.5). Multi-scale variability is therefore an intrinsic geometric signature, not instability.

8.4 Synthesis

Operational choices—principally the resolution parameter r —determine the scale at which geometric transitions are examined. The structural quantities $(m_0, m_\rho, d, \alpha, \varphi)$ are either:

- **canonically determined** ($\alpha = m$, $\varphi = \log$), or
- **restricted to stable theoretical regimes** ($m_0 \geq 3$, $m_\rho \in [5, 20]$, spectral dimensions supported by an eigengap).

At any fixed resolution, the geometric features predicted by the analysis (curvature, transition structure, fractal scaling) coincide with the empirical behaviour of the RCI profile and the MEI-based evaluation. Thus the objections above delineate not weaknesses but the precise scope and coherence of the RCI framework: a geometrically principled, analytically controlled, and empirically validated clustering method for metric-measure spaces.

9 Compactification Analogy: Completion Mechanisms in Gauge Theory, RCI, and Fractal Geometry

Having established the metric-measure foundations of RCI (Section 4), its discrete and spectral implementations (Section 3), and the empirical validation (Section 6), we now place the global farthest-point mechanism within a broader geometric framework. The purpose of this section is conceptual: to articulate a shared structural mechanism appearing in three a priori unrelated settings:

- (i) Uhlenbeck’s compactification of the Yang-Mills moduli space,
- (ii) the farthest-point seeding that governs global RCI, and
- (iii) the box-counting completion underlying fractal dimension.

While these constructions live in distinct domains, they resolve the same functional problem: *degeneracies that would otherwise escape the geometric description are systematically reincorporated into a completed object*. The analogy is not an assertion of analytic equivalence; it highlights the common structural role played by these mechanisms.

9.1 Uhlenbeck Compactification: Capturing Singular Limits

The moduli space $\mathcal{M}$ of smooth finite-energy Yang-Mills connections on a compact 4-manifold is not compact: sequences of solutions may develop curvature concentration. Uhlenbeck's compactness theorem [28, 27] establishes the canonical completion mechanism. If A_j is a sequence of Yang-Mills connections with bounded energy, then after gauge transformation there exists a weak limit A_∞ and a finite set of points $\{x_1, \dots, x_k\}$ ("bubble points") such that:

- curvature concentrates at each x_i ,
- each concentration contributes a quantized amount of energy,
- the sequence converges away from $\{x_i\}$.

The compactified moduli space

$$\overline{\mathcal{M}} = \mathcal{M} \cup \{\text{bubble configurations}\}$$

ensures that no curvature loss escapes the geometric description. Every degeneration is represented by adjoining precisely the additional geometrical data required for completeness.

9.2 RCI Seeding as a Discrete Completion Mechanism

In the global RCI procedure, the ambient structure is a metric space (X, d) . The objective is to represent its large-scale geometry via an r -cover. Without constraints, arbitrary center choices may leave regions of X unrepresented. The farthest-point seeding algorithm [13] provides a discrete analogue of a completion operator.

Definition 9.1 (Farthest-Point Seeding). Fix an initial point $c_1 \in X$. Define recursively

$$c_{j+1} = \arg \max_{x \in X} d(x, \{c_1, \dots, c_j\}),$$

and terminate when

$$\max_{x \in X} d(x, \{c_1, \dots, c_M\}) \leq r.$$

Each step inserts a center at the location of maximal coverage deficit, forcing the cover to incorporate the least-represented region of X . The termination condition is equivalent to enforcing that every point of X is within distance r of some center. Thus no geometric region can escape the r -cover: completeness is restored by systematic addition of missing regions. A classical result quantifies the optimality of this mechanism.

Lemma 9.2 (Quasi-Optimal Coverage [13]). *Let (X, d) have doubling dimension D . If $\mathcal{S} = \{c_j\}$ is generated by farthest-point seeding at scale r , then*

$$N(r) \leq |\mathcal{S}| \leq C_D N(r),$$

where $N(r)$ is the minimal r -covering number and C_D depends only on the doubling constant.

Thus farthest-point seeding constructs, up to controlled constants, a canonical representative of the scale- r geometry of X . Moreover, across scales, these insertions record where coverage deficits persist, and changes in their growth rate are reflected in the scale-dependent profile $M(r)$. By Corollary 4.21, transitions in the growth behaviour of $M(r)$ are detected by the second-difference statistics $\Delta^2 M_c(k)$. Therefore the completion mechanism and the Morse-theoretic transitions are structurally aligned.

9.3 Fractal Dimension as Limit Compactification

In fractal geometry, the box-counting dimension is defined via the limit

$$\dim_B(X) = \lim_{\varepsilon \rightarrow 0} \frac{\log N(\varepsilon)}{\log(1/\varepsilon)},$$

where $N(\varepsilon)$ is the optimal covering number at scale ε . This limiting procedure compactifies infinitesimal structure: all arbitrarily small scales are systematically incorporated into a single global invariant. By Lemma 9.2, the RCI global profile satisfies

$$M(r) \lesssim N(r).$$

Thus the sequence of RCI coverings implements the same structural principle: it records the geometry of X at every resolvable scale, and the asymptotics of $M(r)$ encode its dimensional structure.

9.4 Conceptual Synthesis

The three mechanisms above enact the same structural idea:

- **Yang-Mills:** curvature concentration is captured by adjoining bubble configurations.
- **RCI:** uncovered geometric regions are captured by farthest-point insertions enforcing an r -cover.
- **Fractal geometry:** infinitesimal scales are captured via the limiting behavior of $N(\varepsilon)$.

In all cases, *degeneracies do not escape the description*: the structure is completed by explicitly adding the missing geometric components. For RCI, this means that the farthest-point mechanism functions as a discrete completion operator. Every region that lies outside the current representation forces the insertion of a new center, exactly as bubble addition prevents curvature loss in Yang-Mills and as vanishing scales are retained in the box-counting limit. Consequently, the global geometry of X is represented with quantifiable fidelity across scales, specifically via the comparison bounds $N(2r) \leq M(r) \leq C_2 N(r)$, and the merging profile $M(r)$ records the completed structure throughout the resolution range.




Figure 1: **Unified completion mechanism.** Each vertex displays a geometric process that prevents degeneracies from escaping the description: Uhlenbeck adds bubbles at curvature concentration points; RCI inserts farthest-point centers at uncovered regions; fractal geometry retains fine-scale structure through the limit of covering numbers. All three enact the same structural principle of geometric completion.

10 Conclusion and Future Directions

10.1 Conclusion

This work establishes RCI as a unified geometric clustering framework operating across four complementary structural layers:

- Minimal-metric structural:** generation of the radial base, restriction structure, admissible sieves, and certified sections from points and distances alone (Appendix 15);
- Local analytic:** metric finite-difference curvature expansions, together with bi-Lipschitz spectral fidelity when the spectral implementation is used (Section 3);
- Morse-theoretic:** pathwise variation of the log-density field and critical scales of farthest-point growth (Section 4);

- (iv) **Global fractal:** multiscale behavior of quasi-optimal covers and box-counting equivalence (Section 4.6).

Local foundations. The discrete curvature operator $\Delta^2 M_c(k)$ is a statistically stable and geometrically interpretable probe of scalar curvature. Its legitimacy rests on:

- **Bi-Lipschitz fidelity** (Theorem 3.3), controlling local metric distortion when spectral embeddings are used;
- **Boundary stability** (Theorem 3.4), showing that κ^* concentrates on the correct population transition under a curvature gap;
- **Curvature Law** (Theorem 3.5), which connects discrete second differences to scalar curvature via $\mathbb{E}[\Delta^2 M_c(k)] \approx -C_m S_g(c) \Delta(r_k^2)$.

Together with the concentration bounds of Appendix 14, these results establish $\Delta^2 M_c(k)$ as a principled geometric observable.

Structural certification. Appendix 15 establishes a complementary form of robustness. RCI generates its operational base from the minimal metric input (X, d) , then constructs local invariants, restriction maps, and walls of type change. On an admissible sieve avoiding the wall, the empirical invariant is certified by restriction:

$$\mathcal{S}_t \cap \mathcal{W}_n = \emptyset \implies \widehat{\mathcal{I}}_n|_{\mathcal{S}_t} \cong \mathcal{I}|_{\mathcal{S}_t}.$$

This does not replace the analytic estimates of Appendix 14; it explains the structural meaning of robustness once the admissible region has been identified.

Global structure. The multiscale complexity profile $M(r)$ —the number of farthest-point centers at scale r —encodes coarse geometric structure. Two structural results govern its behavior:

- **Fractal completeness** (Theorem 4.19): $M(r)$ is quasi-optimal for the covering number $N(r)$ and recovers the Minkowski dimension;
- **Local–global correspondence** (Proposition 4.20): transitions in the growth exponent of $M(r)$ appear in $\Delta^2 M_c(k)$ at the matching population radius (Corollary 4.21).

The uniform C^0 -conjugacy between the radius potential $u_k(x)$ and the mass profile $v(x)$ (Theorem 4.5) ensures that these transitions reflect actual geometric features of the distribution.

Synthesis. RCI is not a heuristic combination of density estimators, but a coherent geometric mechanism with dual functionality:

- **Locally:** a second-order curvature-sensitive probe of the sampling distribution;
- **Globally:** a Morse-consistent multiscale filtration that recovers Minkowski dimension and structural transitions.

This dual structure induces *parameter minimality*: the only operational choice is the observational scale r , while structural quantities $(m_0, m_\rho, \alpha = m, \varphi = \log)$, together with the spectral dimension d when the spectral implementation is used, are either canonically fixed or restricted to analytically admissible regimes (Section 7).

Empirical evaluation (Section 6) shows that this theoretical design yields high structural fidelity: RCI achieves the highest MEI on 7 of 8 geometric datasets, preserving curvature-related transitions that classical algorithms systematically erase. The framework is therefore metrically generated, structurally certified, statistically controlled, and topologically interpretable, providing a principled solution to geometric clustering on metric-measure spaces.

10.2 Future Directions

(1) Fractal regimes and dimension reconstruction. Because farthest-point covers are quasi-optimal, $M(r)$ provides an explicit approximation to $N(r)$. Refining this correspondence could enable robust estimation of $\dim_{\text{box}}(X)$, detection of multifractal scaling, and characterization of irregular structures in non-smooth or filamentary spaces.

(2) Streaming and temporally evolving datasets. The bounded-difference estimates of Appendix 14 suggest a streaming formulation of RCI capable of tracking geometric transitions under temporal drift, allowing real-time monitoring of curvature, density, and topology in dynamic data.

(3) Connections with optimal-transport curvature. The operator $\Delta^2 M_c(k)$ resembles discrete curvature deficits in Ollivier–Ricci curvature. Establishing a formal correspondence could unify density curvature, transport inequalities, and empirical graph geometry under a single comparison theory.

(4) Anisotropic and adaptive neighbourhoods. The analysis in Appendix 14 assumes isotropic neighbourhoods. Extending the framework to anisotropic or adaptive local neighbourhoods (e.g., PCA-aligned ellipsoids, Mahalanobis metrics, tangent-plane estimates) may broaden the geometric regimes in which the Curvature Law remains valid.

(5) Conjugacy-invariant geometric operators. The uniform C^0 –conjugacy (Theorem 4.5) suggests the existence of a broader family of operators enjoying analogous invariance. Characterizing such operators could lead to a principled geometric learning theory unifying spectral embeddings, density geometry, and discrete regularizers.

Summary. These directions define a coherent research program grounded in the stability, fidelity, and completeness guarantees established in this work. RCI emerges not as a heuristic clustering method but as a principled geometric framework unifying local curvature, global topology, and fractal structure, providing a rigorous foundation for understanding and learning the intrinsic geometry of high-dimensional metric data.

11 Appendix A: Computational Implementation of RCI

This appendix documents the RCI implementation used inside the benchmark pipeline of Section 6. Importantly, the version of RCI used in the benchmarks is *purely ambient*: it operates directly in the original metric space and does not use any spectral embedding. Every quantity appearing in the implementation is an explicit finite-sample object with no asymptotic or geometric assumptions.

11.1 Nearest-Neighbour Structure

Given $X = \{x_i\}_{i=1}^N \subset \mathbb{R}^m$, FAISS is used to compute

$$\text{idx}(i, k), \quad d^2(i, k),$$

the index and squared distance of the k th nearest neighbour of x_i . These arrays are taken exactly as returned by FAISS. All subsequent quantities are defined from these nearest-neighbour lists.

11.2 Self-Tuning Affinity and Mutual Graph

Following standard practice, local scales are defined by

$$\sigma_i = \sqrt{d^2(i, k_\sigma)},$$

where k_σ is the local self-tuning neighbourhood used by the graph construction. The affinity is

$$W(i, j) = \exp\left(-\frac{d^2(i, j)}{\sigma_i \sigma_j}\right)$$

whenever i and j appear in each other's FAISS lists, and is 0 otherwise. The degree is

$$D(i) = \sum_j W(i, j).$$

This graph is exported for MEI evaluation and, in spectral mode, supports the matrix-on-demand Laplacian. In the benchmark path, however, the RCI profile is computed from radial distances, k -NN density estimates, and the ordered neighbourhoods around each centre. The graph is not an additional source of geometric decisions in the profile computation. RCI does *not* use spectral embeddings in the benchmark evaluations.

11.3 Density Estimator

The local density estimator is computed from the original FAISS distances:

$$\rho(i) = \frac{k_\rho}{N V_m R_{k_\rho}(i)^m},$$

where $R_{k_\rho}(i)$ is the k_ρ -th neighbour radius. In the benchmark implementation, unless supplied explicitly, the density stencil is set by

$$k_\rho = \lceil m \log N \rceil,$$

where m is the ambient dimension in the non-spectral benchmark path. No asymptotic interpretation is invoked at the implementation level.

11.4 RCI Profile

For each centre c , the implementation orders all points by increasing distance from c ,

$$j_0 = c, j_1, \dots, j_{N-1}, \quad r_\ell = d(c, j_\ell).$$

For numerical stability the stored radius is $r_\ell + \varepsilon$, with $\varepsilon = 10^{-12}$. The benchmark implementation uses the density-average contrast

$$A_c(k) = \frac{1}{k+1} \sum_{\ell=0}^k \rho(j_\ell), \quad 0 \leq k \leq N-1.$$

The RCI profile is then

$$M_c(k) = A_c(k) - \alpha \log(r_k + \varepsilon),$$

with $\alpha = m$ fixed throughout all experiments, where m is the ambient dimension of the benchmark space $\mathbb{R}^m$.

Remark 11.1 (Density contrast used in the benchmark implementation). The theoretical presentation emphasizes the logarithmic density contrast because it is the canonical contrast associated with the small-ball expansion and the radius–mass conjugacy. The benchmark implementation reported in this paper uses the raw density-average contrast in $A_c(k)$, and then forms

$$M_c(k) = A_c(k) - \alpha \log(r_k + \varepsilon).$$

This implementation choice is reported exactly as used in the experiments. Both contrasts probe the same radial density field, but their full equivalence and stability under monotone density reparametrizations are not claimed here; they are treated as part of the contrast-robustness analysis of RCI in subsequent work.

11.5 Boundary Index

The benchmark implementation uses an operational two-stage extraction. First, it computes

$$\Delta M_c(k) = M_c(k+1) - M_c(k), \quad \Delta^2 M_c(k) = \Delta M_c(k+1) - \Delta M_c(k).$$

A local median smoothing of window size 3 is applied to $\Delta^2 M_c$. The preliminary index is the most negative smoothed second difference. The implementation then refines this index in a local window by selecting the minimum of ΔM_c , followed by endpoint clamping. This is the operational boundary extraction used in the benchmark; the formal well-posedness criterion is given in Section 3.3.

11.6 Spectral Embedding (Used Only for Exploratory Figures)

A spectral embedding appears only in the exploratory script `core.py`, not in the benchmark pipeline. For those figures, we compute eigenpairs of the normalized Laplacian and embed

$$\Phi(i) = (v_1(i), \dots, v_d(i)).$$

The value $d = 5$ used in that script is an empirical choice for visualization. *It plays no role in any result reported in Table 1.*

11.7 Baseline Methods

Baseline clustering methods (KMeans, DBSCAN, HDBSCAN, Spectral Clustering, and GMM) are implemented via `scikit-learn` and `hdbscan`. For Spectral Clustering, the embedding dimension is determined internally by `scikit-learn` from the requested number of clusters and is not chosen by the user.

11.8 Implementation Summary

All computational objects arise from explicit formulas:

- neighbour lists $\text{idx}(i, k)$ and squared distances $d^2(i, k)$;
- weight matrix W and degree function D ;
- normalized Laplacian L defined by

$$(Lu)(i) = u(i) - \frac{1}{D(i)} \sum_j W(i, j) u(j);$$

- sorted radii (r_k) ;
- contrast terms $A_c(k)$;
- RCI profiles $M_c(k)$;
- discrete derivatives $\Delta M_c(k)$, $\Delta^2 M_c(k)$;
- boundary indices κ_c .

Implementation file. The full reference implementation is provided in the file `core.py`, which contains the matrix-on-demand graph/Laplacian construction, optional spectral embedding pipeline, density estimator, and RCI profile machinery. The benchmark path uses the non-spectral configuration, while the spectral routines are retained for exploratory figures and diagnostic experiments.

Note on spectral embedding. Figure 2 shows a spectral embedding computed for visualization purposes. The benchmark evaluations (Section 6) use RCI without spectral embedding, operating directly on the ambient space.

Apart from the explicit local median smoothing used in the operational boundary extraction described above, no post-hoc smoothing, asymptotic rewriting, or geometric interpretation is introduced into the benchmark outputs.

11.9 Cluster Assignment

Given centres $\{c_1, \dots, c_K\}$, the assignment of a point x_i is

$$\ell(i) = \arg \min_{\ell} \left| M_{c_\ell}(\text{rank}_{c_\ell}(i)) - M_{c_\ell}(0) \right|.$$

If the Morse distance is degenerate or the rank is unavailable, a fallback to Euclidean nearest-centre distance in the current working representation is used. The rule is fully deterministic once the centres and profiles have been constructed.

11.10 Empirical Behaviour

Figures in the main text display raw implementation arrays such as:

- $M_c(k)$,
- $\Delta^2 M_c(k)$,
- final labels,
- and, in exploratory spectral figures only, the low-frequency eigenvalue spectrum of L .

No plot is used to infer geometric convergence or curvature properties.




Figure 2: **Spectral RCI analysis on the sphere dataset.** **Top row.** (1) Raw uniformly sampled sphere. (2) RCI labels. (3) Low-frequency spectral embedding (mutual k NN + self-tuning). (4) Farthest-point centres projected back to the sphere. **Bottom row.** (5) Profiles $M_c(k)$ with detected transition scales κ^* . (6) Second differences $\Delta^2 M_c(k)$. (7) Low-frequency eigenvalue spectrum. (8) Residual diagnostics.

11.11 Conclusion

The implementation uses:

- neighbour queries,
- graph/operator construction,
- explicit RCI profile computation,
- operational boundary selection,
- and numerical eigensolvers only in optional spectral or diagnostic routines.

All outputs follow from explicit formulas.

Limitations. The benchmark implementation assumes that FAISS `IndexFlatL2` returns correct nearest-neighbour distances and that floating-point arithmetic remains stable. Eigensolver convergence is relevant only for the optional spectral and diagnostic routines, not for the non-spectral benchmark path. These are standard numerical assumptions but are not covered by the theory.

12 Appendix B: Benchmark Implementation and Reproducibility

This appendix documents the computational framework used to generate the empirical results of Section 6. Its scope is purely methodological: every numerical value in the tables arises directly from the procedures described here, without manual adjustment or interpretive modification.

12.1 Code Structure

All experiments are executed by the script `comparison_suite.py`. The code is modular and mirrors the algorithmic components introduced in the main text.

Synthetic dataset generators. Functions such as `sample_sphere`, `sample_torus`, `sample_dumbbell_surface`, `sample_hopf_link`, and `sample_trefoil_knot` generate finite-sample geometric datasets with known structure. These serve as testbeds for the structural metric evaluated later.

Classical clustering algorithms. KMeans, DBSCAN, HDBSCAN, Spectral Clustering, and Gaussian Mixture Models are applied with randomized hyperparameter search over standard parameter ranges. For each method, the configuration achieving the highest combined internal validity (Silhouette + Davies–Bouldin) is retained. This removes tuning bias as a confounding factor.

RCI implementation. The function `run_rci` applies the full intrinsic RCI pipeline as defined in Sections 3–4. No hyperparameters are tuned. The routine returns

$$(\text{labels}, \text{params}, X_{\text{emb}}, \text{graph}),$$

where `graph` is the adjacency structure produced internally by the algorithm.

12.2 Structural Metric (MEI)

The Morse Erasure Index (Section 5.2) is implemented in `morse_erasure_index`. The function accepts an explicit graph if one is provided. If no graph is passed, a fresh mutual k -NN graph is constructed.

Graph used for RCI. RCI provides its own graph through the `graph_` attribute. MEI evaluates RCI on this graph, ensuring that the score reflects the neighbourhood structure produced by the algorithm itself.

Graphs used for classical algorithms. Classical methods do not construct a graph. For these methods, MEI builds a new mutual k -NN graph from X . Thus each classical algorithm is evaluated on a common baseline adjacency that is independent of its clustering decisions.

MEI computation. Given edges $\{(i, j, w_{ij})\}$ and the Morse field $f = -\log \hat{\rho}$,

$$\text{TV}_{\text{total}}(f) = \sum_{(i,j)} w_{ij} |f_i - f_j|, \quad \text{TV}_{\text{intra}}(f, \ell) = \sum_{\substack{(i,j) \\ \ell(i)=\ell(j)}} w_{ij} |f_i - f_j|,$$

and

$$\text{MEI}(f, \ell) = 1 - \frac{\text{TV}_{\text{intra}}(f, \ell)}{\text{TV}_{\text{total}}(f)}.$$

No smoothing or normalization is applied.

12.3 Execution Pipeline

The function `run_comparison_suite` executes the full benchmark workflow. Its responsibilities include:

- generating all synthetic datasets;
- running RCI and all baseline clustering algorithms;
- performing randomized hyperparameter search for classical methods;

- extracting the internal RCI graph used for MEI;
- evaluating the Morse Erasure Index with the appropriate graph for each method;
- caching labels, embeddings, graphs, and hyperparameters;
- producing the summary CSV used in Table 1.

All artefacts required for replication are written to disk.

12.4 Complete Reference Implementation

The full benchmark implementation—including dataset generators, clustering routines, Laplacian construction, RCI execution, and the MEI metric—is provided in the file `comparison_suite.py`. All dependencies are standard numerical packages: `numpy`, `scikit-learn`, `hdbscan`, `faiss`, and `plotly`.

12.5 Reproducibility Considerations

All sources of randomness (dataset sampling, kNN queries, randomized search, eigenvector computation, and RCI initialization) are controlled by explicit seeds. The pipeline is therefore deterministic.

Each algorithm is evaluated on the neighbourhood model it induces: RCI uses its internal Laplacian graph, while classical methods use a freshly constructed mutual k -NN graph. This ensures that MEI measures the structural behaviour of each method on its own geometry.

12.6 Conclusion

This appendix documents the benchmark pipeline underlying Section 6. All empirical results follow directly from the public implementation, and reflect the finite-sample behaviour of the sampling model and the Morse-field definitions.

13 Appendix C: Structural Homology of RCI

Section 4 established the metric-measure foundations of RCI, showing that the farthest-point filtration induces a multi-scale profile $M(r)$ with controlled topological transitions (Lemmas 4.13–4.15). This appendix formalizes the categorical and topological structure induced by the RCI algorithm under full mathematical rigor.

13.1 Scope and Corrections

This appendix:

- formalizes the sheaf on the *scale poset* of merge times;
- introduces a well-posed cluster adjacency and the induced nerve complexes;
- derives a *linear (type-A) persistence module* on homology;
- provides an explicit morphism of sheaves within the same topos, replacing earlier, stronger categorical claims.

Standing discretization of scales. Let

$$\mathcal{R} = \{r_0 < r_1 < \dots < r_T\}$$

be the finite, totally ordered set of critical RCI scales, or merge times, from finest (r_0) to coarsest (r_T). At each $r \in \mathcal{R}$, RCI returns a finite partition

$$C(r) = \{C_1(r), \dots, C_{M(r)}(r)\}$$

of the sample. For $r_i \leq r_j$, the deterministic RCI merge induces a *surjective* map

$$\pi_{r_j \rightarrow r_i} : C(r_j) \longrightarrow C(r_i), \quad \pi_{r_i \rightarrow r_i} = \text{id}, \quad \pi_{r_k \rightarrow r_i} = \pi_{r_j \rightarrow r_i} \circ \pi_{r_k \rightarrow r_j} \quad (i \leq j \leq k).$$

13.2 The Scale Sheaf (Sheaf Axioms on a Total Order)

Equip $\mathcal{R}$ with the Alexandroff topology: an open set is an *upper set* $U \subset \mathcal{R}$, i.e., $r \in U$ and $r \leq s$ imply $s \in U$. Covers are families of uppers whose union is U . Consider the contravariant functor

$$\mathcal{F}_{\text{RCI}} : \mathcal{R}^{\text{op}} \longrightarrow \mathbf{FinSet}, \quad \mathcal{F}_{\text{RCI}}(r) = C(r), \quad \mathcal{F}_{\text{RCI}}(r \leq s) = \pi_{s \rightarrow r}.$$

We now prove that $\mathcal{F}_{\text{RCI}}$ is a sheaf on $(\mathcal{R}, J_{\uparrow})$.

Lemma 13.1 (Separatedness). *Let $U \subset \mathcal{R}$ be open and let $\{U_{\alpha}\}_{\alpha}$ be an open cover of U . If $s, t \in \prod_{r \in U} C(r)$ satisfy*

$$s|_{U_{\alpha}} = t|_{U_{\alpha}}$$

for all α , then $s = t$.

Proof. Because $\mathcal{R}$ is finite and totally ordered, the family of principal uppers

$$\{\uparrow r := \{s \in \mathcal{R} : r \leq s\}\}_{r \in U}$$

is a cover of U . If $s|_{\uparrow r} = t|_{\uparrow r}$ for all $r \in U$, then evaluating at r gives $s(r) = t(r)$. Hence $s = t$. $\square$

Lemma 13.2 (Gluing). *Let $U \subset \mathcal{R}$ be open, let $\{U_{\alpha}\}_{\alpha}$ be a cover, and suppose we are given*

$$s_{\alpha} \in \prod_{r \in U_{\alpha}} C(r)$$

with

$$s_{\alpha}|_{U_{\alpha} \cap U_{\beta}} = s_{\beta}|_{U_{\alpha} \cap U_{\beta}}$$

for all α, β . Then there exists a unique $s \in \prod_{r \in U} C(r)$ such that $s|_{U_{\alpha}} = s_{\alpha}$ for all α .

Proof. Define $s(r)$ as $s_\alpha(r)$ for any α with $r \in U_\alpha$. This is well-defined by agreement on overlaps. We must check compatibility with the restriction maps. If $r \leq s$ lie in U , choose α such that $r \in U_\alpha$. Since U_α is an upper set and $r \leq s$, we also have $s \in U_\alpha$. The family s_α is compatible with the restriction maps, hence

$$s_\alpha(r) = \pi_{s \rightarrow r}(s_\alpha(s)).$$

Therefore

$$s(r) = \pi_{s \rightarrow r}(s(s)).$$

Uniqueness follows from Lemma 13.1. $\square$

Corollary 13.3. $\mathcal{F}_{\text{RCI}}$ is a sheaf of finite sets on the site $(\mathcal{R}, J_\uparrow)$.

Remark 13.4 (Scale-generated gluing structure). The scale poset $\mathcal{R}$, the transition maps between scales, and the compatible families used in this appendix are generated by the RCI procedure itself. Thus the sheaf structure is not imposed as an external decoration on the data; it formalizes the multiscale gluing structure produced by the algorithm. Throughout this appendix, arrows are written in the restriction direction of the scale site, i.e., contravariantly with respect to the operational merging process.

Remark 13.5 (Why the base must be the scale poset). An alternative approach might attempt to define a sheaf on the manifold $\mathcal{M}$ by assigning clusters to open subsets $U \subset \mathcal{M}$. This fails because:

- (i) restricting the dataset to U changes k -NN relations globally, violating locality;
- (ii) the spectral Laplacian is a *global* operator—its eigenvectors depend on the entire dataset, not just local neighborhoods;
- (iii) separatedness fails: two clusterings that agree on overlapping regions may differ globally due to non-local spectral effects.

In contrast, the scale poset $(\mathcal{R}, J_\uparrow)$ encodes the *deterministic evolution* of the partition under merging, which *is* local in the scale variable. Thus sheaf axioms hold on $\mathcal{R}$ (Lemmas 13.1–13.2) but not, in general, on $\mathcal{M}$.

Remark 13.6 (Canonical partition versus nerve moduli). At the primary RCI output level, the algorithm does not produce a nontrivial Čech nerve. It produces a finite family of locally calibrated cells

$$C^* = \{C_1^*, \dots, C_M^*\},$$

where each cell is generated by an RCI center c_j and its canonical radius r_j^* . Each sampled point is absorbed by exactly one such cell, so the canonical output is a partition of the dataset.

Consequently, the nerve is not the primitive object produced by RCI. A nontrivial nerve appears only after an additional scale operation: one varies, thickens, or radially explores the canonical cells, for instance by considering

$$U_j(\lambda) = B(c_j, \lambda r_j^*), \quad \lambda \geq 1.$$

For each value of λ , the family $\{U_j(\lambda)\}_j$ defines a Čech nerve $N(\lambda)$. Thus the topological object associated with RCI is not a single nerve, but the scale-indexed family

$$\{N(\lambda)\}_{\lambda \in \Lambda},$$

which we call the *RCI nerve moduli*.

Remark 13.7 (Canonical homology versus dynamic homology). The canonical RCI output already carries homological information at the level of the locally calibrated cell system

$$\mathcal{C}^* = \{C_1^*, \dots, C_M^*\}$$

and its canonical radii

$$\{r_1^*, \dots, r_M^*\}.$$

This is the *canonical homology* generated by RCI at its own intrinsic scales.

This canonical homology should not be confused with the Čech nerve layer. At the primary output level, RCI produces a canonical cell system and its homological content, but not a nontrivial Čech nerve. The

nontrivial nerve appears only after an additional scale operation, when the canonical cells are thickened or radially explored.

Thus RCI has two homological levels:

$$\text{canonical RCI cells} \longrightarrow \text{canonical homology},$$

and

$$\text{radial scale exploration of canonical cells} \longrightarrow \text{dynamic homology} \longrightarrow \text{RCI nerve moduli}.$$

13.3 Geometric Spectral Cover and Čech Nerve

Let (X, d) be the ambient metric space and let

$$\Phi_d : X \longrightarrow \mathbb{R}^d$$

denote the diffusion-map spectral embedding of dimension d (Section 3.2). For each scale $r \in \mathcal{R}$ and each cluster $C_j(r) \in C(r)$, choose a *spectral center* $c_j(r) \in X$, for example the farthest-point center selected by RCI. The associated *spectral ball* is

$$U_j(r) := \{y \in \mathbb{R}^d : \|\Phi_d(c_j(r)) - y\| \leq r\}.$$

Each $U_j(r)$ is a closed, convex, and contractible subset of $\mathbb{R}^d$. Write

$$U(r) := \bigcup_{j=1}^{M(r)} U_j(r)$$

for the corresponding geometric union.

Standing geometric compatibility. For every pair of scales $r_i \leq r_j$ and each cluster $C_\ell(r_j) \in C(r_j)$, the spectral balls satisfy

$$U_\ell(r_j) \subseteq U_{\pi_{r_j \rightarrow r_i}(\ell)}(r_i). \quad (\text{C.geom})$$

Justification. The diffusion-map embedding Φ_d is locally bi-Lipschitz, so small-scale distances in $\mathbb{R}^d$ agree with those in (X, d) up to a factor $1 \pm \varepsilon_N$. For each cluster $C_\ell(r_j)$, the RCI radius $r_\ell^*(r_j)$ is chosen so that the ambient metric ball

$$B_d(c_\ell(r_j), r_\ell^*(r_j))$$

covers the entire cluster. Selecting the spectral radius for $U_\ell(r_j)$ compatibly with these bounds ensures

$$\Phi_d(C_\ell(r_j)) \subseteq U_\ell(r_j).$$

Whenever a merge

$$C_\ell(r_j) \subseteq C_{\pi_{r_j \rightarrow r_i}(\ell)}(r_i)$$

occurs at a coarser scale r_i , one may choose the spectral radii so that the ball of the finer cluster is contained within that of its parent. Thus condition (C.geom) is not a geometric consequence of the map Φ_d alone, but an admissible and canonical compatibility choice ensuring functorial coherence across scales in the construction of a spectral cover.

Definition 13.8 (Čech Nerve of the Spectral Cover). For each $r \in \mathcal{R}$, the *RCI Čech nerve* $N(r)$ is the Čech nerve of the spectral cover

$$\{U_j(r)\}_{j=1}^{M(r)}.$$

That is, $N(r)$ is the abstract simplicial complex whose vertex set is $C(r)$ and where a finite set $\{j_1, \dots, j_m\}$ spans a simplex if and only if

$$U_{j_1}(r) \cap \dots \cap U_{j_m}(r) \neq \emptyset.$$

Remark 13.9 (Adjacency graphs as 1-skeletons). The cluster adjacency graph $G(r)$ used in Section 3 can be regarded as the 1-skeleton of the Čech nerve $N(r)$ in the following sense: an edge $\{i, j\}$ in $G(r)$ corresponds to a nonempty intersection

$$U_i(r) \cap U_j(r) \neq \emptyset,$$

i.e., to a 1-simplex of $N(r)$. Empirical adjacency rules, for example those based on $\text{dist}_\wedge$ together with a threshold $\varepsilon(r)$, serve as computational approximations of this geometric condition. While $\text{dist}_\wedge(C_i, C_j) < r$ does not guarantee $U_i(r) \cap U_j(r) \neq \emptyset$ without additional geometric control, it is a numerically stable proxy for intersection of the corresponding spectral balls. The Čech construction formalizes the exact geometric notion.

Lemma 13.10 (Compatibility of the Čech Nerve under Merges). *Let $r_i \leq r_j$ in $\mathcal{R}$. Define a map on vertices*

$$\Phi_{r_j \rightarrow r_i} : N(r_j)_0 \longrightarrow N(r_i)_0, \quad \Phi_{r_j \rightarrow r_i}(j) := \pi_{r_j \rightarrow r_i}(j),$$

and extend it to simplices by

$$\Phi_{r_j \rightarrow r_i}(\{j_1, \dots, j_m\}) := \{\pi_{r_j \rightarrow r_i}(j_1), \dots, \pi_{r_j \rightarrow r_i}(j_m)\}.$$

Then

$$\Phi_{r_j \rightarrow r_i} : N(r_j) \longrightarrow N(r_i)$$

is a well-defined simplicial map, possibly collapsing simplices.

Proof. Let $\sigma = \{j_1, \dots, j_m\}$ be a simplex in $N(r_j)$. By Definition 13.8, this means

$$\bigcap_{p=1}^m U_{j_p}(r_j) \neq \emptyset.$$

By (C.geom), for each p ,

$$U_{j_p}(r_j) \subseteq U_{\pi_{r_j \rightarrow r_i}(j_p)}(r_i).$$

Therefore

$$\bigcap_{p=1}^m U_{\pi_{r_j \rightarrow r_i}(j_p)}(r_i) \supseteq \bigcap_{p=1}^m U_{j_p}(r_j) \neq \emptyset.$$

Hence

$$\{\pi_{r_j \rightarrow r_i}(j_1), \dots, \pi_{r_j \rightarrow r_i}(j_m)\}$$

is again a simplex of $N(r_i)$, and $\Phi_{r_j \rightarrow r_i}$ is simplicial. Functoriality,

$$\Phi_{r_k \rightarrow r_i} = \Phi_{r_j \rightarrow r_i} \circ \Phi_{r_k \rightarrow r_j},$$

follows directly from the functoriality of the merge maps $\pi_{r_\bullet \rightarrow r_\bullet}$. □

13.4 Linear (Type-A) Persistence on Homology

Fix a field $\mathbb{F}$. For each $k \geq 0$ and $r \in \mathcal{R}$, set

$$H_k(r) := H_k(N(r); \mathbb{F}),$$

where $N(r)$ is the Čech nerve of the spectral cover $\{U_j(r)\}_{j=1}^{M(r)}$ defined in Definition 13.8. For adjacent indices $r_i < r_{i+1}$, define the linear map

$$\mathbf{f}_i := H_k(\Phi_{r_{i+1} \rightarrow r_i}) : H_k(r_{i+1}) \longrightarrow H_k(r_i),$$

where $\Phi_{r_{i+1} \rightarrow r_i}$ is the simplicial map of Lemma 13.10. This yields a finite *type-A* persistence module:

$$H_k(r_T) \xrightarrow{\mathbf{f}_{T-1}} H_k(r_{T-1}) \xrightarrow{\mathbf{f}_{T-2}} \dots \xrightarrow{\mathbf{f}_1} H_k(r_1) \xrightarrow{\mathbf{f}_0} H_k(r_0).$$

Theorem 13.11 (Interval decomposition (type- A classification)). *For each $k \geq 0$, the above finite representation over $\mathbb{T}$ decomposes, uniquely up to isomorphism and order, as a finite direct sum of interval modules*

$$I[a, b]$$

supported on contiguous index intervals

$$r_a \leq \cdots \leq r_b.$$

Proof (constructive induction). This is the standard structure theorem for finite-dimensional representations of type- A quivers over a field. Equivalently, it follows from Gabriel’s theorem together with Krull–Schmidt uniqueness: every finite type- A representation decomposes into a direct sum of indecomposable interval representations, and the multiset of intervals is unique up to isomorphism and reordering. $\square$

Remark 13.12 (Sheafified radial barcode). The persistence object constructed here is not the ordinary barcode of the raw point cloud, nor merely the barcode of a single externally prescribed Čech filtration. It is the barcode of the RCI radial nerve system.

More precisely, RCI first produces locally calibrated canonical cells with radii r_j^* . The finite scale poset $\mathcal{R}$ arises only when these canonical cells are radially explored across scales. At each scale $r \in \mathcal{R}$, this exploration induces a partition $C(r)$, a spectral cover, and hence a Čech nerve $N(r)$. The deterministic transition maps between scales then induce simplicial maps between nerves, and applying homology produces a finite type- A persistence module.

Thus the resulting barcode records the homology of a sheafified multiscale radial cover generated by RCI. In degree zero, it records the evolution of connected components of the RCI nerve system under radial merges. It should therefore be read as a barcode of the RCI-generated merge geometry, not as a raw count of clusters and not as a direct barcode of the ambient manifold alone.

Proposition 13.13 ($k = 0$ recovers the merge profile). *At degree $k = 0$, $H_0(N(r_i))$ equals the number of connected components of the Čech nerve $N(r_i)$. Since the nerve is the Čech complex of the spectral cover $\{U_j(r_i)\}$, and since these balls are generated from the RCI clusters at scale r_i , the number of connected components equals the RCI merge profile at scale r_i , namely the component count of the nerve system generated by the spectral cover. Interval deaths in H_0 correspond exactly to component-reducing cluster merges under RCI.*

Proof. $H_0(N(r_i))$ counts connected components of the Čech nerve. By construction (Definition 13.8), two clusters C_j and C_k lie in the same connected component precisely when there exists a path of overlapping spectral balls connecting $U_j(r_i)$ and $U_k(r_i)$.

Under the RCI evolution, a merge identifies the corresponding vertices in the scale system through the map $\pi_{r_{i+1} \rightarrow r_i}$. By Lemma 13.10, this identification induces a simplicial map between the nerves and therefore a linear map on H_0 . Whenever such a merge reduces the number of connected components of the nerve system, the corresponding independent H_0 -generator dies under f_i . Thus deaths in degree zero record exactly the component-reducing merge events of the RCI-generated radial nerve system. $\square$

13.5 Applicability of the Nerve Theorem to RCI Covers

By construction (Definition 13.8), $N(r)$ is precisely the Čech nerve of the spectral cover

$$\{U_j(r)\}_{j=1}^{M(r)}$$

in $\mathbb{R}^d$. Lemma 13.14 below shows that this cover satisfies the standard contractibility assumptions of the Leray–Čech Nerve Theorem.

Lemma 13.14 (Contractibility of finite intersections). *For any nonempty finite collection of indices $\{i_1, \dots, i_m\}$,*

$$U_{i_1}(r) \cap \cdots \cap U_{i_m}(r)$$

is convex and therefore contractible.

Proof. Each $U_j(r)$ is a closed Euclidean ball in $\mathbb{R}^d$. Finite intersections of convex sets are convex, hence contractible. $\square$

Theorem 13.15 (RCI Nerve Theorem). *For every scale $r \in \mathcal{R}$, the geometric realization of the RCI Čech nerve satisfies*

$$|N(r)| \simeq U(r),$$

where $|N(r)|$ denotes geometric realization.

Proof. By Lemma 13.14, all finite intersections of the cover $\{U_j(r)\}$ are either empty or contractible. The Leray–Čech Nerve Theorem therefore applies directly, yielding

$$|N(r)| \simeq \bigcup_j U_j(r) = U(r).$$

$\square$

Interpretation. The persistent homology module of Section 13.5 is computed on the Čech nerves $N(r)$ of the spectral cover $(U_j(r))_j$, whose realizations are homotopy equivalent to the unions $U(r)$. Thus the interval decomposition of Theorem 13.11 encodes topological transitions in the geometry of the spectral balls across scales, not merely combinatorial artifacts of a discrete adjacency relation.

13.6 A Safe Comparative Morphism

We introduce a nontrivial and explicit *morphism of sheaves* within the same topos, providing a rigorous framework for deterministic comparison across scales.

Definition 13.16 (Fixed quotient comparator). Fix an equivalence relation $\sim$ on the point set, for example ground-truth labels. For each r , define $Q(r)$ as the quotient of $C(r)$ obtained by merging clusters whose point sets intersect the same $\sim$ -class. Let

$$\rho_{r_j \rightarrow r_i} : Q(r_j) \longrightarrow Q(r_i)$$

be induced by $\pi_{r_j \rightarrow r_i}$. This defines a sheaf $\mathcal{Q}$ on $(\mathcal{R}, J_\uparrow)$.

Proposition 13.17 (Nontrivial morphism of sheaves). *The natural family of quotient maps*

$$\eta_r : C(r) \longrightarrow Q(r)$$

satisfies

$$\eta_{r_i} \circ \pi_{r_j \rightarrow r_i} = \rho_{r_j \rightarrow r_i} \circ \eta_{r_j} \quad (r_i \leq r_j).$$

Hence

$$\eta : \mathcal{F}_{\text{RCI}} \Rightarrow \mathcal{Q}$$

is a morphism of sheaves.

Proof. Let $x \in C(r_j)$ and let

$$y = \pi_{r_j \rightarrow r_i}(x).$$

By definition of the merge map, the underlying point set of x is contained in the point set of y . Applying the quotient map η before or after the merge therefore identifies the same $\sim$ -equivalence class in $Q(r_i)$:

$$\eta_{r_i}(\pi_{r_j \rightarrow r_i}(x)) = \rho_{r_j \rightarrow r_i}(\eta_{r_j}(x)).$$

Thus the naturality square commutes, and the family η_r defines a natural transformation

$$\eta : \mathcal{F}_{\text{RCI}} \Rightarrow \mathcal{Q}.$$

$\square$

Remark 13.18. This gives a rigorous, nontrivial *comparison* without appealing to geometric morphisms. It quantifies how an externally imposed identification collapses the multiscale information of RCI.




Figure 3: **Structural architecture of the RCI framework.** Information flows from the scale sheaf, which records the deterministic evolution across $\mathcal{R}$, through the Čech nerve of the spectral cover, to the homotopy realization guaranteed by the Nerve Theorem (Theorem 13.15). The resulting spaces feed into the homology persistence module (Theorem 13.11), which is finally compared across scales via the deterministic sheaf morphism $\eta : \mathcal{F}_{\text{RCI}} \Rightarrow \mathcal{Q}$ (Proposition 13.17). Convexity of the spectral balls (Lemma 13.14) ensures that the resulting topological information is geometrically meaningful rather than an artifact of adjacency.

13.7 Summary

- RCI defines a genuine sheaf $\mathcal{F}_{\text{RCI}}$ on the scale poset $(\mathcal{R}, J_{\uparrow})$ (Corollary 13.3).
- The sheaf structure formalizes the multiscale gluing structure generated by the RCI procedure itself (Remark 13.4).
- The canonical RCI output already carries homological information through the locally calibrated cell system and its canonical radii.
- Although the primary RCI output is a canonical partition and not a nontrivial Čech nerve, each scale section of the RCI nerve moduli defines a Čech nerve $N(r)$ of the induced spectral cover, compatible with merges (Lemma 13.10).
- The Leray–Čech Nerve Theorem identifies $|N(r)|$ with the geometric union $U(r)$ (Theorem 13.15).
- Homology across scales then forms a finite type- A persistence module with canonical interval decomposition (Theorem 13.11).
- In degree $k = 0$, the barcode recovers the RCI merge/component profile of the spectral nerve system (Proposition 13.13).

- A canonical quotient map induces a natural transformation $\eta : \mathcal{F}_{\text{RCI}} \Rightarrow \mathcal{Q}$, providing deterministic information collapse (Proposition 13.17).

Together, these results establish a complete algebraic-topological structure for RCI at the level of the scale-generated spectral nerve system.

Computational validation. All theoretical claims in Appendix C were computationally validated on synthetic datasets, including sphere and torus samples. The scale sheaf axioms (separatedness, gluing, and functoriality) passed verification. The Čech nerve construction yielded internally consistent Betti numbers for the spectral covers produced by the validation pipeline:

$$\beta_1 = 1$$

for the sampled sphere cover and

$$\beta_1 = 3$$

for the sampled torus cover in the reported runs. These values refer to the finite spectral nerve at the selected scale, not to a direct computation of the Betti numbers of the ambient smooth manifolds.

The H_0 barcode exhibited internal consistency: the number of initial components minus the number of detected component-reducing merges equals the final component count of the RCI spectral nerve system.

Complete validation code is available in the repository as `homology.py`.




Figure 4: **Geometric visualization of the H_0 -barcode evolution via Čech nerves of the spectral cover.** Each panel displays the 1-skeleton of the Čech nerve $N(r)$ (Definition 13.8) of the RCI spectral cover at scale parameter r . Vertices represent the RCI clusters $C(r)$, and edges indicate nonempty intersections of spectral balls $\{U_j(r)\}_j$ (Section 13.5). The value of H_0 , namely the number of connected components of $N(r)$, is shown above each panel. As r increases, new edges emerge at merge scales detected by the RCI pipeline, producing the local transition $H_0 = 7 \rightarrow 6$ at $r = 0.127$, in agreement with Proposition 13.13. This visualization demonstrates that the H_0 -barcode records the evolution of the RCI-generated Čech nerve system: components are born at fine scales and merge as r increases, with deaths occurring when adjacency edges identify formerly distinct connected components (Lemma 13.10). The local transition $H_0 = 7 \rightarrow 6$ at $r = 0.127$ records one merge event. Across the full run, the arithmetic consistency $H_0(r_{\text{initial}}) - (\text{total detected merges}) = H_0(r_{\text{final}})$, $7 - 4 = 3$, confirms the internal coherence of the type- A persistence module (Theorem 13.11) and validates its computational realization in the RCI pipeline. Dataset: torus with $n = 250$ points.

14 Appendix D: Technical Foundations of Structural Robustness

Objective. The objective of this appendix is to prove the three main theorems stated in Section 3.5: Theorem 3.3 (local bi-Lipschitz regularity of the spectral embedding), Theorem 3.4 (stability of the boundary detector), and Theorem 3.5 (the RCI curvature law). We establish these results through a sequence of intermediate lemmas that control the geometric and statistical behavior of the RCI construction.

Setup and Preliminaries. The argument relies on a set of standard regularity assumptions and several key preliminary lemmas regarding the geometry of the spectral space.

Regularity Assumptions. We assume the following conditions hold.

- (A1) **Bounded-degree oracle graph.** The neighbor list $\mathcal{S}_L(i)$ has size at most a constant L_0 , uniformly in N .
- (A2) **Non-degenerate local scales.** There exist constants $0 < \sigma_{\min} \leq \sigma_{\max} < \infty$ such that $\sigma_i \leq \sigma_{\max}$ for all i .
- (A3) **Spectral gap.** The underlying Laplace–Beltrami operator on the data manifold (M, g) has a spectral gap $\lambda_{d+1}(M) - \lambda_d(M) \geq \delta_0 > 0$. By standard spectral convergence results for graph Laplacians [26], for any $\delta \in (0, \delta_0)$, the discrete gap of L is bounded below by δ with probability at least $1 - e^{-cN}$ for some $c > 0$. Our analysis is conditioned on this high-probability event.
- (A4) **Lipschitz contrast and stable density estimator.** The contrast map ϕ is L_ϕ -Lipschitz. The m -NN radius estimator $r_\rho(y)$ in the spectral space Y is L_ρ -Lipschitz.

Geometric Regularity in the Spectral Space. The following structural properties of the embedded point cloud Y are essential. They are consequences of the local Bi-Lipschitz regularity (Theorem 3.3) established in Appendix D.2.

Lemma 14.1 (Annulus Population and Radial Scaling). *Assume the data X are sampled from a compact C^3 manifold (M, g) of dimension m . With high probability, the embedded cloud $Y = \Phi_d(X)$ satisfies the following properties:*

- (i) *For any center $c_Y \in Y$, radius $r > 0$, and thickness $\tau > 0$, the number of points in the annulus $\mathcal{A}(r, \tau) := \{y \in Y : r \leq \|y - c_Y\| \leq r + \tau\}$ is bounded by*

$$|\mathcal{A}(r, \tau)| \leq C_{\text{ann}} N \tau r^{m-1},$$

for a constant C_{ann} independent of N .

- (ii) *The radius r_k^{spec} of the k -th nearest neighbor ball around any $c_Y \in Y$ satisfies*

$$C_{\min} \left(\frac{k}{N} \right)^{1/m} \leq r_k^{\text{spec}} \leq C_{\max} \left(\frac{k}{N} \right)^{1/m},$$

for constants $0 < C_{\min} \leq C_{\max} < \infty$ independent of N and k .

This follows from the volume preservation properties of the locally bi-Lipschitz spectral embedding established below.

Lemma 14.2 (Uniform Per-Point Displacement Bound via Davis–Kahan). *Under assumptions (A1)–(A3), let X' be a dataset obtained by replacing a single point x_j in X with an arbitrary point x'_j . Let L and L' be the corresponding Laplacians. Then the operator norm of their difference is uniformly bounded, $\|\Delta L\|_{\text{op}} = \|L - L'\|_{\text{op}} \leq C_L$ for a constant C_L independent of N . Consequently, by the Davis–Kahan theorem, the maximal displacement of any embedded point is bounded by a uniform constant:*

$$\max_i \|\Phi_d(x_i) - R\Phi'_d(x_i)\|_2 \leq \frac{2C_L}{\delta} =: \varepsilon,$$

where $R \in O(d)$ is an orthogonal transformation and ε is independent of N .

Proof. Replacing x_j with x'_j affects the affinity matrix K only in the j -th row and column, and in the rows/columns corresponding to the neighbors of x_j and x'_j . By the bounded-degree assumption (A1), the number of affected entries is at most $O(L_0^2)$, a constant. Since each affinity value is in $[0, 1]$, the change in any entry is at most 1. The resulting perturbation matrix $\Delta K = K - K'$ is sparse with a constant number of non-zero entries of constant magnitude. Thus, its operator norm is bounded by a constant independent of N : $\|\Delta K\|_{\text{op}} \leq \|\Delta K\|_{\text{F}} = O(1)$.

The degree matrix perturbation ΔD is also of constant operator norm. The non-degeneracy of local scales (A2) ensures that the entries of D are bounded away from zero, so $D^{-1/2}$ is well-behaved. The perturbation on the normalized Laplacian, $\Delta L = I - D^{-1/2} K D^{-1/2} - (I - (D')^{-1/2} K' (D')^{-1/2})$, can be shown to have an operator norm $\|\Delta L\|_{\text{op}} \leq C_L$ for a constant C_L depending on L_0 , $\sigma_{\min}$, and $\sigma_{\max}$, but not on N .

Given the spectral gap δ from (A3), the Davis–Kahan theorem directly yields a bound on the Frobenius norm of the eigenvector matrix perturbation: $\|V_d - V'_d R\|_{\text{F}} \leq 2\|\Delta L\|_{\text{op}}/\delta \leq 2C_L/\delta$. The maximal per-point displacement is the maximal row-norm of this difference matrix, which is bounded by the Frobenius norm. This establishes the existence of a uniform displacement bound $\varepsilon = O(1)$. $\square$

Concentration via McDiarmid’s Inequality. The bounded-difference property established in the proof allows the application of McDiarmid’s inequality. For a fixed k , suppose the shell-average functional $A_c(k)$ has bounded differences $c_j \leq C_\star(k)$. Then for any $t > 0$,

$$\Pr(|A_c(k) - \mathbb{E}[A_c(k)]| \geq t) \leq 2 \exp\left(-\frac{2t^2}{\sum_{j=1}^N c_j^2}\right) \leq 2 \exp\left(-\frac{2t^2}{NC_\star(k)^2}\right).$$

This inequality is the foundation for the stability result in Theorem 3.4. The term $C_\star(k)$ remains controlled as long as k grows at least polylogarithmically with N . For any polynomial growth rate $k \sim N^\alpha$ with $0 < \alpha \leq 1$, the combinatorial contribution $(N/k)^{1/m} = (N^{1-\alpha})^{1/m}$ is bounded or decays, ensuring robust concentration.

14.1 Local Bi-Lipschitz Regularity of the Spectral Embedding

Objective. The goal of this section is to prove Theorem 3.3, which states that the spectral map Φ_d is locally Bi-Lipschitz with respect to the intrinsic manifold metric d_g . This property provides the formal basis for the geometric fidelity of the RCI framework, ensuring that geometric quantities computed in the spectral space are faithful proxies for their intrinsic counterparts.

Roadmap. The proof proceeds by establishing separate upper and lower Lipschitz bounds. The upper bound (no blow-up) is derived by transferring the smoothness of continuum eigenfunctions to the discrete eigenvectors. The lower bound (no collapse) relies on a quantitative, first-order Taylor expansion of the spectral map, combined with the positive-definiteness of the spectral metric tensor.

The Continuum-to-Discrete Interface. The core of the argument rests on the convergence of the discrete graph Laplacian to its continuum counterpart. We formalize this connection in the following proposition.

Proposition 14.3 (Spectral Approximation). *Let (M, g) be the data manifold and let $\{\lambda_\ell, \varphi_\ell\}$ be the first d eigenpairs of the Laplace–Beltrami operator Δ_g . Let L_N be the discrete Laplacian defined in §3.2. On the high-probability event where a discrete spectral gap $\delta > 0$ exists (Assumption A3), there exists an orthogonal transformation $R_N \in O(d)$ and a sequence $\varepsilon_N \rightarrow 0$ as $N \rightarrow \infty$ such that the matrix of discrete eigenvectors V_d and the matrix of evaluated continuum eigenfunctions $\Phi_d^\star$ satisfy:*

$$\|V_d - \Phi_d^\star R_N\|_{\text{op}} \leq \varepsilon_N.$$

Justification. A full proof is beyond the scope of this appendix, but the result is a direct consequence of established theorems on the consistency of graph Laplacians. The convergence of operators of this class is established in, e.g., [26, 24]. The key hypotheses of these theorems are satisfied by our construction:

- (i) The kernel is positive, localized, and decays rapidly.

- (ii) The bandwidth scaling, managed by the adaptive scales σ_i which are bounded by Assumption (A2), is consistent with the requirements for continuum approximation (e.g., $h_N \rightarrow 0$ and $Nh_N^m/\log N \rightarrow \infty$).
- (iii) The data are sampled from a regular manifold, as per the setup.

Under these conditions, the discrete operator L_N converges to a weighted Laplace–Beltrami operator in a suitable sense, which in turn implies the convergence of the corresponding spectral projectors and eigenspaces. The operator norm bound follows from this convergence combined with the Davis-Kahan theorem. $\square$

From this operator norm bound, we derive the crucial per-point (row-norm) error bound. For any matrix A with at most d non-zero singular values, $\|A\|_F^2 \leq d\|A\|_{\text{op}}^2$. The maximal row norm is bounded by the Frobenius norm. Thus,

$$\max_i \|\Phi_d(x_i) - \Phi_d^*(x_i)R_N\|_2 \leq \|V_d - \Phi_d^*R_N\|_F \leq \sqrt{d}\|V_d - \Phi_d^*R_N\|_{\text{op}} \leq \sqrt{d}\varepsilon_N.$$

Lemma 14.4 (Uniform Upper Lipschitz Bound (No Blow-up)). *There exists a constant $c_2 < \infty$ such that, with high probability, for all $x, y \in X$,*

$$\|\Phi_d(x) - \Phi_d(y)\| \leq c_2 d_g(x, y) + 2\sqrt{d}\varepsilon_N.$$

Proof. On a compact manifold, each eigenfunction φ_ℓ is smooth and thus Lipschitz. Let $c_2^2 = \sum_{\ell=1}^d (\sup_M \|\nabla \varphi_\ell\|_g)^2$, so the continuum map Φ_d^* is c_2 -Lipschitz. By the triangle inequality and the per-point error bound,

$$\begin{aligned} \|\Phi_d(x) - \Phi_d(y)\| &\leq \|\Phi_d(x) - \Phi_d^*(x)R_N\| + \|\Phi_d^*(x)R_N - \Phi_d^*(y)R_N\| + \|\Phi_d^*(y)R_N - \Phi_d(y)\| \\ &\leq \sqrt{d}\varepsilon_N + \|\Phi_d^*(x) - \Phi_d^*(y)\| + \sqrt{d}\varepsilon_N \\ &\leq c_2 d_g(x, y) + 2\sqrt{d}\varepsilon_N. \end{aligned}$$

$\square$

Lemma 14.5 (Positive-Definite Spectral Metric). *Let $G(x) = \sum_{\ell=1}^d \nabla \varphi_\ell(x) \otimes \nabla \varphi_\ell(x)$ be the spectral metric tensor. If Δ_g has a spectral gap at level d , there exists a constant $\beta > 0$ such that $G(x) \succeq \beta I_m$ for all $x \in M$.*

Proof. This is a standard result in spectral geometry. If $G(x_*)$ were singular, a non-zero tangent vector u would exist such that $\langle \nabla \varphi_\ell(x_*), u \rangle_g = 0$ for all $\ell = 1, \dots, d$. This implies the span of the first d eigenfunctions is locally constant, which contradicts non-degeneracy properties guaranteed by the spectral gap and unique continuation principles. Compactness of M ensures the bound is uniform. $\square$

Lemma 14.6 (Quantitative Lower Lipschitz Bound (No Collapse)). *There exist constants $c_1 > 0$ and $r_0 > 0$ such that, with high probability, for all $x, y \in X$ with $d_g(x, y) \leq r_0$,*

$$\|\Phi_d(x) - \Phi_d(y)\| \geq c_1 d_g(x, y) - 2\sqrt{d}\varepsilon_N.$$

Proof. The proof is quantitative. For $x, y \in M$ with $d_g(x, y)$ small, a first-order Taylor expansion of Φ_d^* in normal coordinates at x gives $\Phi_d^*(y) = \Phi_d^*(x) + J_{\Phi_d^*}(x)v + O(d_g(x, y)^2)$, where v is the tangent vector representing the geodesic to y . The squared norm of the linear term is $\|J_{\Phi_d^*}(x)v\|^2 = v^\top G(x)v \geq \beta \|v\|_g^2 = \beta d_g(x, y)^2$, by Lemma 14.5.

By controlling the second-order remainder term, we can find $r_0 > 0$ such that for all $d_g(x, y) \leq r_0$, $\|\Phi_d^*(y) - \Phi_d^*(x)\| \geq (\sqrt{\beta}/2)d_g(x, y)$. Let $c_1 = \sqrt{\beta}/2$.

We transfer this continuum lower bound to the discrete map Φ_d via the reverse triangle inequality:

$$\begin{aligned} \|\Phi_d(y) - \Phi_d(x)\| &\geq \|\Phi_d^*(y)R_N - \Phi_d^*(x)R_N\| - (\|\Phi_d(y) - \Phi_d^*(y)R_N\| + \|\Phi_d(x) - \Phi_d^*(x)R_N\|) \\ &\geq \|\Phi_d^*(y) - \Phi_d^*(x)\| - (\sqrt{d}\varepsilon_N + \sqrt{d}\varepsilon_N) \\ &\geq c_1 d_g(x, y) - 2\sqrt{d}\varepsilon_N. \end{aligned}$$

This holds uniformly for all pairs with $d_g(x, y) \leq r_0$. $\square$

Conclusion. Combining Lemmas 14.4 and 14.6 directly yields Theorem 3.3. The argument is now self-contained, with the critical assumption of spectral convergence (Proposition 14.3) being explicitly stated and justified by reference to the established literature. The analysis confirms that the discrete spectral embedding inherits the local Bi-Lipschitz properties of its continuum counterpart, up to a vanishing, additive error term rigorously controlled by the rate of spectral convergence and the embedding dimension d .

14.2 Stability of the Boundary Detector

Objective. We give a fully rigorous proof of Theorem 3.4. The key point is to formalize the local dependence structure of the shell-average functional $A_c(k)$ and to derive a sub-Gaussian concentration bound with an *effective* sample size of order k , valid with overwhelming probability.

Roadmap. The argument proceeds in four steps: (i) we introduce a high-probability “good geometry” event controlling annulus populations in spectral space; (ii) on this event, we construct a dependency graph for the shell variables and bound its maximal degree; (iii) we apply a Janson–Bousquet-type inequality for weakly dependent variables, obtaining a Bernstein-type concentration for $A_c(k)$; (iv) we combine this with concentration for spectral k -NN radii to control the discrete curvature and prove stability of the boundary detector.

14.2.1 Good Geometry Event and Local Dependence

Fix a center $c_Y = \Phi_d(c) \in Y$ and consider the spectral radius r_k^{spec} and a thickness parameter

$$\tau_N = \tau_0 \left(\frac{k}{N} \right)^{1/m},$$

for some small constant $\tau_0 > 0$ depending only on the manifold model. Define the annulus

$$\mathcal{A}_k(c) = \{y \in Y : r_k^{\text{spec}} - \tau_N \leq \|y - c_Y\| \leq r_k^{\text{spec}} + \tau_N\}.$$

Lemma 14.7 (Annulus population: high-probability bound). *Under the manifold sampling model and the bi-Lipschitz event of Appendix D.2, there exist constants $C_{\text{ann}}, \kappa > 0$ such that for all $k \geq m_0$ and all N sufficiently large,*

$$\Pr \left(|\mathcal{A}_k(c)| \geq C_{\text{ann}} \log N \right) \leq N^{-\kappa}.$$

Proof. By Lemma 14.1, there is a constant $\tilde{C}_{\text{ann}} > 0$ such that

$$\mathbb{E}[|\mathcal{A}_k(c)|] \leq \tilde{C}_{\text{ann}} N \tau_N (r_k^{\text{spec}})^{m-1}.$$

On the manifold model and the scaling of $r_k^{\text{spec}}, r_k^{\text{spec}} \lesssim (k/N)^{1/m}$, so

$$\mathbb{E}[|\mathcal{A}_k(c)|] \leq \tilde{C}'_{\text{ann}} k \tau_0,$$

for another constant $\tilde{C}'_{\text{ann}}$. For fixed k and c_Y , the point cloud in spectral space is i.i.d. up to bi-Lipschitz distortion, hence $|\mathcal{A}_k(c)|$ is stochastically dominated by a binomial variable with mean $\mu_k \leq C_0$ (by choosing τ_0 small). Chernoff’s bound then gives

$$\Pr(|\mathcal{A}_k(c)| \geq L) \leq \exp(-cL)$$

for some $c > 0$ and all $L \geq 2\mu_k$. Taking $L = C_{\text{ann}} \log N$ with C_{ann} large enough yields $\Pr(|\mathcal{A}_k(c)| \geq C_{\text{ann}} \log N) \leq N^{-\kappa}$ for some $\kappa > 0$. $\square$

We now define the high-probability good geometry event:

$$\mathcal{G}_N := \bigcap_{k \geq m_0} \left\{ |\mathcal{A}_k(c)| \leq C_{\text{ann}} \log N \right\}.$$

By a union bound over k in a range of order $O(\log N)$, Lemma 14.7 implies

$$\Pr(\mathcal{G}_N) \geq 1 - N^{-\kappa'}, \tag{15}$$

for some $\kappa' > 0$.

On the event $\mathcal{G}_N$, a perturbation of a single sample point can only affect membership in the k -shell through points lying in the annulus $\mathcal{A}_k(c)$, whose cardinality is now deterministically bounded by $C_{\text{ann}} \log N$.

Definition 14.8 (Local shell variables and dependency graph). For fixed c and k , let $I_k(c)$ be the indices of the k nearest neighbors of c_Y in spectral space, and set

$$Z_j := \log \widehat{\rho}(x_j), \quad A_c(k) := \frac{1}{k} \sum_{j \in I_k(c)} Z_j.$$

We define a dependency graph G_k on vertex set $I_k(c)$ by joining j and j' with an edge whenever a perturbation of some sample point can simultaneously affect both Z_j and $Z_{j'}$.

Lemma 14.9 (Local dependence under good geometry). *On the event $\mathcal{G}_N$, there exists a constant $D_0 > 0$ such that*

$$\Delta(G_k) \leq D_0 \log N$$

for all $k \geq m_0$, where $\Delta(G_k)$ is the maximum degree of G_k .

Proof. A perturbation of a single sample point x_ℓ may: (i) change its own contribution Z_ℓ , and (ii) change the membership of points whose spectral distance to c_Y lies in $\mathcal{A}_k(c)$. On $\mathcal{G}_N$, the number of such indices is at most $C_{\text{ann}} \log N$ for each k . Therefore, each Z_j can be jointly affected with at most $D_0 \log N$ other variables, for some constant D_0 depending only on the manifold model and kernel parameters; this is precisely the statement that $\Delta(G_k) \leq D_0 \log N$. $\square$

14.2.2 Refined Concentration via Dependency Graph

We now apply a McDiarmid–Bernstein inequality for functions of weakly dependent variables indexed by a dependency graph; see, e.g., Janson [18] or Boucheron et al. [2] for concentration methods for weakly dependent variables.

Lemma 14.10 (Refined concentration for shell averages). *Under assumptions (A1)–(A4), Lemma 14.1, and the bi-Lipschitz event of Appendix D.2, there exist constants $c_1, c_2, c > 0$ such that for all $k \geq m_0$, all $t \in (0, 1)$, and all N sufficiently large,*

$$\Pr(|A_c(k) - \mathbb{E}A_c(k)| \geq t) \leq 2 \exp(-c k t^2) + N^{-\kappa'}.$$

In particular, for $k \geq C_0 \log N$ and N large, the $N^{-\kappa'}$ term is dominated, and we have

$$\Pr(|A_c(k) - \mathbb{E}A_c(k)| \geq t) \leq 3 \exp(-c' k t^2).$$

Proof. Condition on the good geometry event $\mathcal{G}_N$. On $\mathcal{G}_N$, changing a single sample point affects at most $O(\log N)$ of the Z_j 's, and each affected Z_j changes by at most a constant (by the Lipschitz assumptions on ϕ and $\widehat{\rho}$). Hence, $A_c(k)$ has one-step differences bounded by

$$|A_c(k)(X) - A_c(k)(X^{(i)})| \leq \frac{C}{k}$$

for some constant $C > 0$, where $X^{(i)}$ denotes a configuration with x_i perturbed.

The effective number of variables on which $A_c(k)$ depends is of order k (points inside the k -ball plus a shell of thickness $O(\tau_N)$), and the dependency graph G_k restricted to this local index set has maximum degree at most $D_0 \log N$ by Lemma 14.9.

The Janson–Bousquet inequality for functions of variables indexed by a dependency graph yields a Bernstein-type bound

$$\Pr(|A_c(k) - \mathbb{E}A_c(k)| \geq t \mid \mathcal{G}_N) \leq 2 \exp\left(-\frac{t^2}{2(v_k + b_k t/3)}\right),$$

with

$$v_k \leq \frac{c_1 \log N}{k}, \quad b_k \leq \frac{c_2}{k}.$$

For fixed $t \in (0, 1)$ and $k \geq C_0 \log N$, the variance proxy satisfies $v_k \leq \tilde{c}_1/k$ for some constant $\tilde{c}_1$, and the denominator in the exponent is of order $1/k$. Thus, there exists $c > 0$ such that

$$\Pr(|A_c(k) - \mathbb{E}A_c(k)| \geq t \mid \mathcal{G}_N) \leq 2 \exp(-c k t^2).$$

Unconditioning and using (15) gives

$$\Pr(|A_c(k) - \mathbb{E}A_c(k)| \geq t) \leq 2 \exp(-c k t^2) + \Pr(\mathcal{G}_N^c) \leq 2 \exp(-c k t^2) + N^{-\kappa'}.$$

For $k \geq C_0 \log N$ and N large, the second term is absorbed into the first, yielding the simplified bound. $\square$

14.2.3 Concentration for Spectral Radii and Curvature

Lemma 14.11 (Concentration for spectral k -NN radii). *Under the manifold sampling model and the local bi-Lipschitz property of Φ_d (Appendix D.2), there exist constants $c_3, c_4 > 0$ such that for all $k \geq m_0$ and all $t > 0$,*

$$\Pr(|\log r_k^{\text{spec}} - \mathbb{E} \log r_k^{\text{spec}}| \geq t) \leq 2 \exp(-c_3 k t^2),$$

and consequently

$$\Pr(|\Delta^2 \log r_k^{\text{spec}} - \mathbb{E} \Delta^2 \log r_k^{\text{spec}}| \geq t) \leq 6 \exp(-c_4 k t^2).$$

Proof. In spectral space, volume growth is m -dimensional up to bi-Lipschitz constants. Counts in metric balls are binomial with mean $\Theta(Nr^m)$; Chernoff bounds give sub-Gaussian fluctuations for the empirical distribution of distances, hence for the k -NN quantile r_k^{spec} . A smooth change of variables $r \mapsto \log r$ transfers this to $\log r_k^{\text{spec}}$. The bound for second differences follows from a union bound over three consecutive indices. $\square$

Combining Lemma 14.10 (for the shell term) and Lemma 14.11 (for the radius term), both components of $M_c(k)$ have deviations with tails of order $\exp(-c k t^2)$, and therefore the discrete curvature inherits the same tail behavior, up to a constant factor.

Corollary 14.12 (Sub-Gaussian curvature fluctuations). *Let*

$$\mathcal{E}(k) := \Delta^2 M_c(k) - \Delta^2 M_c^\circ(k).$$

Then there exists $C_E > 0$ such that for all $k \geq m_0$ and all $t > 0$,

$$\Pr(|\mathcal{E}(k)| \geq t) \leq 6 \exp(-C_E k t^2) + N^{-\kappa'}.$$

In particular, for $k \geq C_0 \log N$ and N large,

$$\Pr(|\mathcal{E}(k)| \geq t) \leq 7 \exp(-C'_E k t^2)$$

for some constant $C'_E > 0$.

14.2.4 Proof of Theorem 3.4

Setup. Assume the curvature gap: there exist $\gamma > 0$ and a unique k° such that

$$\Delta^2 M_c^\circ(k^\circ - 1) \geq \gamma, \quad \Delta^2 M_c^\circ(k^\circ) \leq -\gamma, \quad \Delta^2 M_c^\circ(k) \geq 0 \text{ for } k < k^\circ - 1,$$

and $k^\circ \geq C_0 \log N$.

Local union bound. Define the bad events

$$E_1 = \{\Delta^2 M_c(k^\circ) \geq 0\}, \quad E_2 = \{\Delta^2 M_c(k^\circ - 1) < 0\}.$$

If $\kappa^\star \neq k^\circ$, then either E_1 or E_2 occurs, or a premature sign flip occurs for some $k < k^\circ - 1$.

Since E_1 implies $\mathcal{E}(k^\circ) \geq \gamma$ and E_2 implies $\mathcal{E}(k^\circ - 1) \leq -\gamma$, Corollary 14.12 gives

$$\Pr(E_1) \leq 3 \exp(-C_E k^\circ \gamma^2) + N^{-\kappa'}, \quad \Pr(E_2) \leq 3 \exp(-C_E (k^\circ - 1) \gamma^2) + N^{-\kappa'}.$$

Hence

$$\Pr(E_1 \cup E_2) \leq 6 \exp(-C_E (k^\circ - 1)\gamma^2) + 2N^{-\kappa'} \leq CN^{-C'}$$

for some $C, C' > 0$, using $k^\circ \geq C_0 \log N$.

A similar local union bound over $k < k^\circ - 1$ shows that premature sign flips also occur with probability $O(N^{-C})$. Therefore,

$$\Pr(\kappa^\star = k^\circ) \geq 1 - O(N^{-C}),$$

which completes the proof of Theorem 3.4.

14.3 Geometric Bootstrapping and the Curvature Law

Objective. We prove the RCI Curvature Law (Theorem 3.5). The derivation proceeds in two stages: (i) a continuous expansion in the spectral radius r for the expected shell average and profile, obtained by combining volume and integrand expansions on a Riemannian manifold and transporting them to the spectral chart via the locally bi-Lipschitz spectral embedding; (ii) a discretization step that passes from r to the shell index k , followed by a discrete second-difference computation.

Setup. Let (M, g) be a compact C^3 Riemannian manifold of dimension m , let $f \in C^2(M)$ be a sampling density, and fix a center $c \in M$. All expectations are conditioned on the high-probability bi-Lipschitz event of Appendix D.2 for the spectral embedding Φ_d . Write S_g for the scalar curvature, Δ_g for the Laplace–Beltrami operator, and ω_m for the volume of the Euclidean unit m -ball. Throughout, $r > 0$ is taken below the injectivity radius at c and small enough that the local bi-Lipschitz bounds of Theorem 3.3 hold.

14.3.1 Foundational geometric ingredients

Lemma 14.13 (Gray–Vanhecke volume expansion). *For r sufficiently small,*

$$\text{Vol}_g(B_r^g(c)) = \omega_m r^m \left(1 - \frac{S_g(c)}{6(m+2)} r^2 \right) + O(r^{m+3}).$$

Lemma 14.14 (Log-bias of k -NN density on a manifold). *Let $\hat{f}(x)$ be the k -NN density estimator at $x \in M$ with local scale r . Then*

$$\mathbb{E}[\log \hat{f}(x)] = \log f(x) + \frac{r^2}{2m} \Delta_g(\log f)(x) + \frac{r^2}{6(m+2)} S_g(x) + O(r^3 + r^2 |\nabla_g \log f|^2).$$

Lemma 14.15 (Spectral preimage is a quasi-ball). *Let $B_r^{\text{spec}}(c) \subset Y = \Phi_d(X)$ be the Euclidean ball of radius r centered at $c_Y = \Phi_d(c)$. On the high-probability bi-Lipschitz event of Appendix D.2, there exist constants $c_1, c_2 > 0$, a radius $r_0 > 0$ and a sequence $\varepsilon_N \downarrow 0$ such that for all $0 < r \leq r_0$,*

$$B_{r_-}^g(c) \subset \Phi_d^{-1}(B_r^{\text{spec}}(c)) \subset B_{r_+}^g(c),$$

where

$$r_- := \max \left\{ 0, \frac{r - \varepsilon_N}{c_2} \right\}, \quad r_+ := \frac{r + \varepsilon_N}{c_1},$$

and the Jacobian distortion induced by Φ_d on these sets contributes a multiplicative error $1 + O(\varepsilon_N)$ to all volume integrals.

Proof. By Theorem 3.3 (local bi-Lipschitz regularity), there exist constants $c_1, c_2 > 0$, $r_0 > 0$ and a sequence $\varepsilon_N \rightarrow 0$ such that for all $x, y \in M$ with $d_g(x, y) \leq r_0$,

$$c_1 d_g(x, y) - \varepsilon_N \leq \|\Phi_d(x) - \Phi_d(y)\| \leq c_2 d_g(x, y) + \varepsilon_N. \quad (16)$$

No rescaling of Φ_d is performed; c_1 and c_2 are fixed geometric constants of the embedding.

Left inclusion. Let $x \in M$ satisfy $d_g(x, c) \leq r_- = (r - \varepsilon_N)/c_2$ (we assume $r > \varepsilon_N$, which holds for N large and r fixed). Then, using the upper bound in (16),

$$\|\Phi_d(x) - \Phi_d(c)\| \leq c_2 d_g(x, c) + \varepsilon_N \leq c_2 \frac{r - \varepsilon_N}{c_2} + \varepsilon_N = r.$$

Thus $\Phi_d(x) \in B_r^{\text{spec}}(c)$ and $B_{r_-}^g(c) \subset \Phi_d^{-1}(B_r^{\text{spec}}(c))$.

Right inclusion. Let x satisfy $\|\Phi_d(x) - \Phi_d(c)\| \leq r$. By the lower bound in (16),

$$c_1 d_g(x, c) - \varepsilon_N \leq r, \quad \text{so} \quad d_g(x, c) \leq \frac{r + \varepsilon_N}{c_1} = r_+.$$

Thus $\Phi_d^{-1}(B_r^{\text{spec}}(c)) \subset B_{r_+}^g(c)$.

Jacobian control. On the bi-Lipschitz event, the pullback metric satisfies

$$c_1^2 g \preceq \Phi_d^* \delta \preceq c_2^2 g$$

up to an additive perturbation controlled by ε_N , and therefore the Jacobian determinant obeys

$$c_1^m (1 - O(\varepsilon_N)) \leq J_{\Phi_d}(x) \leq c_2^m (1 + O(\varepsilon_N)).$$

Together with the established set inclusions, this implies

$$\text{Vol}_g(\Phi_d^{-1}(B_r^{\text{spec}}(c))) = (1 + O(\varepsilon_N)) \text{Vol}_g(B_r^g(c)),$$

where $\tilde{r}$ is any radius satisfying $r_- \leq \tilde{r} \leq r_+$. The same multiplicative error transfers to all integrals $\int_{\Phi_d^{-1}(B_r^{\text{spec}}(c))} \psi d\mu_g$ for $\psi \in C^0$, by dominated convergence and the uniform Jacobian bound. $\square$

14.3.2 Expansion of the expected shell average at radius r

For a spectral radius r , define the continuum shell average

$$A_c(r) := \frac{1}{\#\{y \in Y : |y - c_Y| \leq r\}} \sum_{|y - c_Y| \leq r} \log \hat{\rho}(y), \quad M_c(r) := \mathbb{E}[A_c(r)] - \alpha \log r.$$

Equivalently (in the continuum limit), with $x = \Phi_d^{-1}(y)$,

$$\mathbb{E}[A_c(r)] = \frac{\int_{\Phi_d^{-1}(B_r^{\text{spec}}(c))} \mathbb{E}[\log \hat{f}(x)] f(x) d\mu_g(x)}{\int_{\Phi_d^{-1}(B_r^{\text{spec}}(c))} f(x) d\mu_g(x)}, \quad \hat{\rho} = \hat{f}.$$

Proposition 14.16 (Mean shell average; curvature coefficient with metric distortion). *For r sufficiently small, there exist constants $C_{f,m}^{(c_1, c_2)}, C_{g,m}^{(c_1, c_2)} > 0$ depending only on $(m, c_1, c_2, f(c))$ such that*

$$\mathbb{E}[A_c(r)] = \log f(c) + C_{f,m}^{(c_1, c_2)} \Delta_g(\log f)(c) r^2 - C_{g,m}^{(c_1, c_2)} S_g(c) r^2 + O(r^3 + \varepsilon_N).$$

Moreover $C_{f,m}^{(c_1, c_2)} > 0$ and $C_{g,m}^{(c_1, c_2)} > 0$, and in the idealized case $c_1 = c_2 = 1$ they reduce to $C_{f,m}^{(1,1)} = \frac{1}{2m}$ and $C_{g,m}^{(1,1)} = \frac{m}{6(m+2)}$.

Proof. Let

$$\Omega_r := \Phi_d^{-1}(B_r^{\text{spec}}(c)), \quad \mathcal{N}(r) = \int_{\Omega_r} \mathbb{E}[\log \hat{f}(x)] f(x) d\mu_g(x), \quad \mathcal{D}(r) = \int_{\Omega_r} f(x) d\mu_g(x).$$

By Lemma 14.15,

$$B_{r_-}^g(c) \subset \Omega_r \subset B_{r_+}^g(c), \quad r_- = \frac{r - \varepsilon_N}{c_2}, \quad r_+ = \frac{r + \varepsilon_N}{c_1},$$

and the Jacobian distortion induces only a multiplicative $1 + O(\varepsilon_N)$ error. Thus it suffices to compute the expansions of $\mathcal{N}$ and $\mathcal{D}$ for geodesic balls of radii r_- and r_+ and then express the result in terms of r .

Step 1: denominator. For any $a > 0$ fixed, Lemma 14.13 gives

$$\text{Vol}_g(B_{ar}^g(c)) = \omega_m(ar)^m \left(1 - \frac{S_g(c)}{6(m+2)}(ar)^2 \right) + O(r^{m+3}).$$

Expanding f in normal coordinates, we obtain

$$\int_{B_{ar}^g(c)} f(x) d\mu_g(x) = f(c) \omega_m(ar)^m - f(c) \omega_m(ar)^{m+2} \frac{S_g(c)}{6(m+2)} + \frac{\omega_m(ar)^{m+2}}{2m(m+2)} \Delta_g f(c) + O(r^{m+3}).$$

For $a \in \{1/c_2, 1/c_1\}$ this gives two-sided bounds for $\mathcal{D}(r)$ and hence an expansion

$$\mathcal{D}(r) = D_0 r^m + D_2 r^{m+2} + O(r^{m+3} + \varepsilon_N r^m),$$

with $D_0 = \tilde{f}(c) \omega_m$ and $D_2 = \tilde{A}_1 S_g(c) + \tilde{A}_2 \Delta_g f(c)$, where $\tilde{f}(c)$ is comparable to $f(c)$ up to constants depending only on c_1, c_2 and the coefficients $\tilde{A}_1, \tilde{A}_2$ depend only on (m, c_1, c_2) .

Step 2: numerator. By Lemma 14.14,

$$\mathbb{E}[\log \hat{f}(x)] = \log f(x) + \frac{r^2}{2m} \Delta_g(\log f)(x) + \frac{r^2}{6(m+2)} S_g(x) + O(r^3 + r^2 |\nabla \log f|^2).$$

We evaluate the k -NN estimator on a region whose geodesic radius is asymptotically equivalent to r . Therefore the local scale parameter appearing in Lemma 14.14 satisfies $r_{\text{knn}} = \Theta(r)$, and all bias terms depending on the local k -NN radius may be treated, to second order, as functions of r up to multiplicative constants.

Multiplying by $f(x)$ and expanding near c yields

$$\mathbb{E}[\log \hat{f}(x)] f(x) = f(x) \log f(x) + \frac{r^2}{2m} f(x) \Delta_g(\log f)(x) + \frac{r^2}{6(m+2)} S_g(x) f(x) + O(r^3).$$

Integrating over $B_{ar}^g(c)$ and using symmetry and Taylor expansions, we arrive at

$$\int_{B_{ar}^g(c)} f(x) \log f(x) d\mu_g(x) = (f \log f)(c) \omega_m(ar)^m + \left(B_1(a) S_g(c) + B_2(a) \Delta_g(f \log f)(c) \right) r^{m+2} + O(r^{m+3})$$

for explicit coefficients $B_1(a), B_2(a)$ depending only on (m, a) . Similarly,

$$\int_{B_{ar}^g(c)} f(x) \Delta_g(\log f)(x) d\mu_g(x) = f(c) \Delta_g(\log f)(c) \omega_m(ar)^m + O(r^{m+2}),$$

and

$$\int_{B_{ar}^g(c)} S_g(x) f(x) d\mu_g(x) = S_g(c) f(c) \omega_m(ar)^m + O(r^{m+2}).$$

Combining these contributions and sandwiching $\log_r$ between $B_{r_-}^g(c)$ and $B_{r_+}^g(c)$, we obtain

$$\mathcal{N}(r) = N_0 r^m + N_2 r^{m+2} + O(r^{m+3} + \varepsilon_N r^m),$$

with $N_0 = \tilde{f}(c) \log f(c) \omega_m$ and $N_2 = \hat{A}_1 S_g(c) + \hat{A}_2 \Delta_g(f \log f)(c) + \hat{A}_3 f(c) \Delta_g \log f(c)$, for coefficients $\hat{A}_i$ depending only on (m, c_1, c_2) .

Step 3: quotient and identification of coefficients. We write

$$\frac{\mathcal{N}(r)}{\mathcal{D}(r)} = \frac{N_0}{D_0} + \frac{N_2 D_0 - N_0 D_2}{D_0^2} r^2 + O(r^3 + \varepsilon_N).$$

The leading ratio is $\frac{N_0}{D_0} = \frac{\tilde{f}(c) \log f(c)}{\tilde{f}(c)} = \log f(c)$. Using the identity

$$\Delta_g(f \log f) = (\log f) \Delta_g f + 2 \langle \nabla f, \nabla \log f \rangle + f \Delta_g(\log f),$$

the quantity $N_2 D_0 - N_0 D_2$ can be rewritten as a linear combination of $S_g(c)$ and $\Delta_g \log f(c)$, with coefficients depending only on $(m, c_1, c_2, f(c))$. Furthermore, all first-order terms vanish upon integration, since geodesic balls in normal coordinates satisfy

$$\int_{B_r^g(c)} \langle \nabla f, v \rangle d\mu_g = 0 \quad \text{for every fixed vector } v,$$

and the local bi-Lipschitz distortion of Φ_d preserves this symmetry up to a multiplicative $O(\varepsilon_N)$. Consequently,

$$N_2 D_0 - N_0 D_2 = D_0^2 \left(C_{f,m}^{(c_1, c_2)} \Delta_g(\log f)(c) - C_{g,m}^{(c_1, c_2)} S_g(c) \right),$$

and hence

$$\frac{N_2 D_0 - N_0 D_2}{D_0^2} = C_{f,m}^{(c_1, c_2)} \Delta_g(\log f)(c) - C_{g,m}^{(c_1, c_2)} S_g(c),$$

with $C_{f,m}^{(c_1, c_2)}, C_{g,m}^{(c_1, c_2)} > 0$. In the special case $c_1 = c_2 = 1$ these reduce to $C_{f,m}^{(1,1)} = \frac{1}{2m}$ and $C_{g,m}^{(1,1)} = \frac{m}{6(m+2)}$. $\square$

Remark 14.17 (Role of α). The r^2 -coefficient in $\mathbb{E}[A_c(r)]$ is purely geometric/measure-theoretic; the penalty $-\alpha \log r$ affects only lower-order (in k) discrete-curvature terms after discretization. Choosing $\alpha = m$ cancels the leading Euclidean volumetric drift in M_c but does not alter the sign or the qualitative structure of the r^2 -curvature coefficient.

14.3.3 From radius r to shells k and the Curvature Law

Quantile relation and inversion. Let $k(r)$ denote the expected count inside spectral radius r . By Lemmas 14.13 and 14.15,

$$k(r) = N \int_{\Omega_r} f d\mu_g = N \tilde{f}(c) \omega_m \tilde{r}(r)^m (1 - \tilde{\gamma}_g S_g(c) \tilde{r}(r)^2) + \tilde{\gamma}_f \tilde{r}(r)^{m+2} + O(N \tilde{r}(r)^{m+3}),$$

where $\tilde{r}(r)$ is any radius between r_- and r_+ and $\tilde{f}(c)$, $\tilde{\gamma}_g$, $\tilde{\gamma}_f$ depend only on $(m, c_1, c_2, f(c))$. Since $r_{\pm} = a_{\pm} r + O(\varepsilon_N)$ with $a_- = 1/c_2$, $a_+ = 1/c_1$, we have $\tilde{r}(r) = \bar{a} r + O(\varepsilon_N)$ for some $\bar{a} \in [1/c_2, 1/c_1]$. Thus

$$k(r) = \kappa_{\text{eff}} r^m (1 - \gamma_g^{\text{eff}} r^2) + \gamma_f^{\text{eff}} r^{m+2} + O(r^{m+3}),$$

for some effective constants $\kappa_{\text{eff}}, \gamma_g^{\text{eff}}, \gamma_f^{\text{eff}}$ depending only on $(m, c_1, c_2, f(c))$. Solving for r^2 by series inversion gives

$$r^2(k) = \left(\frac{k}{\kappa_{\text{eff}}} \right)^{2/m} \left(1 + \Theta((k/\kappa_{\text{eff}})^{2/m}) \right). \quad (17)$$

Expected profile in k -scale. From Proposition 14.16,

$$M_c(r) = \mathbb{E}[A_c(r)] - \alpha \log r = \text{const} - \alpha \log r + \left(C_{f,m}^{(c_1, c_2)} \Delta_g \log f(c) - C_{g,m}^{(c_1, c_2)} S_g(c) \right) r^2 + O(r^3 + \varepsilon_N).$$

Composing with $r = r(k)$ via (17) yields

$$M_c(k) = \text{const} - \frac{\alpha}{2} \log k + \left(\tilde{C}_{f,m}^{(c_1, c_2)} \Delta_g \log f(c) - \tilde{C}_{g,m}^{(c_1, c_2)} S_g(c) \right) r^2(k) + \tilde{R}(k),$$

where $\tilde{C}_{f,m}^{(c_1, c_2)}, \tilde{C}_{g,m}^{(c_1, c_2)} > 0$ are modified constants (absorbing κ_{eff} and the inversion step) and $\tilde{R}(k) = O(k^{3/m}) + O(\varepsilon_N)$.

Second finite difference. Since $r^2(k) \sim k^{2/m}$, one has

$$\Delta_k^2(\log k) = \Theta(k^{-2}), \quad \Delta_k^2(r^2(k)) = \Theta(k^{2/m-2}),$$

and $\Theta(k^{2/m-2})$ dominates $\Theta(k^{-2})$ for every $m \geq 1$. Therefore the discrete curvature is governed asymptotically by the $r^2(k)$ -term.

Theorem 14.18 (RCI Curvature Law (Proof of Theorem 3.5)). *Under the hypotheses above and with $\alpha = m$, there exist positive constants $K_{f,m}^{(c_1,c_2)}, K_{g,m}^{(c_1,c_2)} > 0$, depending only on $(m, c_1, c_2, f(c))$, such that*

$$\mathbb{E}[\Delta^2 M_c(k)] = \left(K_{f,m}^{(c_1,c_2)} \Delta_g \log f(c) - K_{g,m}^{(c_1,c_2)} S_g(c) \right) \Delta^2(r_k^2) + R_k,$$

where $R_k = o(\Delta^2(r_k^2)) + O(\varepsilon_N)$. In particular, if f is locally constant at c (so that $\Delta_g \log f(c) = 0$),

$$\mathbb{E}[\Delta^2 M_c(k)] \approx -K_{g,m}^{(c_1,c_2)} S_g(c) \Delta^2(r_k^2),$$

so the expected discrete curvature of the RCI profile detects the sign and magnitude of the scalar curvature $S_g(c)$ up to a positive multiplicative factor.

Proof. Apply Δ_k^2 to $M_c(k)$ and use the asymptotics above. The term $-\frac{\alpha}{2} \log k$ contributes $\Theta(k^{-2})$, while the $r^2(k)$ -term contributes $(\tilde{C}_{f,m}^{(c_1,c_2)} \Delta_g \log f(c) - \tilde{C}_{g,m}^{(c_1,c_2)} S_g(c)) \Theta(k^{2/m-2})$, which dominates for all $m \geq 1$. The remainder estimate follows from the composition error in (17) and from $O(r^3) = O(k^{3/m})$. All constants can be absorbed into $K_{f,m}^{(c_1,c_2)}$ and $K_{g,m}^{(c_1,c_2)}$, which are strictly positive because the underlying coefficients in Proposition 14.16 are positive and the inversion step preserves sign. $\square$

Conclusion. The expected discrete curvature of the RCI profile detects an *effective curvature* equal to a linear combination of the intrinsic scalar curvature $S_g(c)$ and the curvature of the sampling density $\Delta_g \log f(c)$, modulated by the universal factor $\Delta^2(r_k^2)$ and by positive geometric constants depending only on $(m, c_1, c_2, f(c))$. This establishes the Curvature Law at the level required for the stability and interpretability of the RCI boundary detector.

15 Appendix E: Minimal-Metric Admissibility and Sheafified Robustness

Objective. Appendix 14 proves analytic stability of RCI under regularity assumptions such as smoothness, local bi-Lipschitz spectral fidelity, concentration of shell averages, and non-degenerate curvature gaps. The purpose of the present appendix is different.

We prove a structural form of robustness that does not begin with smoothness, probability, spectral gaps, or manifold hypotheses. It begins with the minimal data required for any local metric geometry to exist:

$$(X, d), \quad c \in X, \quad x \mapsto d(c, x).$$

From this minimal metric input, RCI generates radial orderings, local windows, cells, invariants, restriction maps, and walls of type change. Once these objects have been generated, admissibility is not an external hypothesis imposed on the algorithm; it is the restriction of the RCI-generated structure to the region where type change has been excluded.

Thus the main point of this appendix is:

$$\text{minimal metric generation} + \text{admissible wall exclusion} \implies \text{certified structural robustness}.$$

15.1 The Minimal Metric Base Generated by RCI

Given (X, d) and a center $c \in X$, RCI defines the radial order

$$(x_{(0)} = c, x_{(1)}, \dots, x_{(N-1)}), \quad d(c, x_{(i)}) \leq d(c, x_{(j)}) \quad (i \leq j).$$

A *radial window* at c is a finite initial segment or admissible subinterval of this radial order. We define the RCI operational base by

$$\mathcal{B}_{\text{RCI}} := \{(c, I) : c \in X, I \subseteq \text{Ord}_c(X) \text{ is a radial window}\}.$$

We partially order $\mathcal{B}_{\text{RCI}}$ by refinement:

$$(c, J) \leq (c, I) \iff J \subseteq I.$$

Thus $\downarrow p$ denotes the family of all refinements of a radial window $p \in \mathcal{B}_{\text{RCI}}$.

Remark 15.1 (Minimality). The construction of $\mathcal{B}_{\text{RCI}}$ requires only points and distances. No manifold structure, density regularity, embedding, spectral gap, smoothness, differentiability, or probability model is required in order to define the base. These additional structures may be used later to obtain quantitative analytic estimates, but they are not needed for the existence of the RCI operational base.

15.2 RCI Invariants and Restriction Structure

At each base point $p = (c, I) \in \mathcal{B}_{\text{RCI}}$, RCI assigns a local invariant

$$\mathcal{I}(p) = (\omega_p, \kappa_p^*, r_p^*, C_p),$$

where ω_p denotes the local radial profile or Morse signal, κ_p^* the boundary index, r_p^* the induced boundary radius, and C_p the corresponding local cell.

If $q \leq p$, the radial window q is a refinement of p . The data computed on p localizes to q through maps

$$\rho_{p \rightarrow q} : \mathcal{I}(p) \longrightarrow \mathcal{I}(q).$$

These maps are induced by the deterministic localization rule of RCI: radial orders and profiles are restricted to smaller windows, while boundary data are localized or recomputed according to the same RCI rule on the refined window. Thus $\rho_{p \rightarrow q}$ should not be read as an arbitrary projection, but as the canonical RCI transition from a window to one of its refinements.

The collection

$$p \mapsto \mathcal{I}(p), \quad (q \leq p) \mapsto \rho_{p \rightarrow q}$$

is therefore a restriction object, or presheaf-like object, on the finite poset $\mathcal{B}_{\text{RCI}}$. Its sheafification, when needed, is not an external structure imposed on RCI. It is the categorical completion of the restriction data already generated by RCI.

Remark 15.2 (Discrete invariant types and sheafification). The RCI invariant types considered in this appendix are treated as objects of a discrete, separated category of local types. Consequently, disagreement of two restricted RCI objects is detected on principal sieves $\downarrow p$.

The passage from the RCI restriction presheaf to its sheafification does not introduce new analytic data. It only enforces the gluing of locally compatible discrete invariant types. In particular, walls of type change are already detected on principal sieves and are unchanged by sheafification.

Remark 15.3 (Sheaf readability). RCI does not assume a sheaf in order to operate. Rather, by constructing a base, local objects, and restriction maps from the metric data itself, RCI produces an object that is naturally readable in sheaf-theoretic language. The sheaf formalism enters as recognition of the generated restriction structure, not as an additional modelling hypothesis.

15.3 Empirical Invariants and Walls of Type Change

Let $\widehat{\mathcal{I}}_n$ denote the empirical RCI invariant computed from a finite sample of size n , and let $\mathcal{I}$ denote the corresponding ideal or limiting invariant on the same RCI-generated base.

A point $p \in \mathcal{B}_{\text{RCI}}$ lies on a wall of type change when the empirical and ideal invariants fail to agree over the local refinement neighbourhood $\downarrow p$. Define

$$\mathcal{W}_n := \left\{ p \in \mathcal{B}_{\text{RCI}} : \widehat{\mathcal{I}}_n|_{\downarrow p} \not\cong \mathcal{I}|_{\downarrow p} \right\}.$$

Thus $\mathcal{W}_n$ is not an arbitrary bad set. It is the RCI-generated locus where the empirical local object changes type relative to the ideal local object.

An admissible sieve is a down-closed subposet,

$$\mathcal{S}_t \subseteq \mathcal{B}_{\text{RCI}},$$

meaning that

$$p \in \mathcal{S}_t, \quad q \leq p \quad \implies \quad q \in \mathcal{S}_t.$$

It is called admissible when it avoids the wall:

$$\mathcal{S}_t \cap \mathcal{W}_n = \emptyset.$$

Equivalently, $\mathcal{S}_t$ is a region of the RCI base on which no type change occurs.

15.4 Structural Robustness by Admissible Restriction

Theorem 15.4 (Structural robustness on admissible sieves). *Let*

$$\mathcal{W}_n = \left\{ p \in \mathcal{B}_{\text{RCI}} : \widehat{\mathcal{I}}_n|_{\downarrow p} \not\cong \mathcal{I}|_{\downarrow p} \right\}.$$

If $\mathcal{S}_t \subseteq \mathcal{B}_{\text{RCI}}$ is an admissible sieve, i.e.

$$\mathcal{S}_t \cap \mathcal{W}_n = \emptyset,$$

then

$$\widehat{\mathcal{I}}_n|_{\mathcal{S}_t} \cong \mathcal{I}|_{\mathcal{S}_t}.$$

Proof. Suppose, for contradiction, that the restrictions do not agree on $\mathcal{S}_t$. Since the invariant category is discrete and separated, disagreement is detected on a principal sieve. Hence there exists $p \in \mathcal{S}_t$ such that

$$\widehat{\mathcal{I}}_n|_{\downarrow p} \not\cong \mathcal{I}|_{\downarrow p}.$$

By definition of the wall $\mathcal{W}_n$, this implies

$$p \in \mathcal{W}_n.$$

But $p \in \mathcal{S}_t$, hence

$$p \in \mathcal{S}_t \cap \mathcal{W}_n,$$

contradicting admissibility:

$$\mathcal{S}_t \cap \mathcal{W}_n = \emptyset.$$

Therefore no such p exists, and the empirical and ideal RCI invariants agree after restriction to the admissible sieve:

$$\widehat{\mathcal{I}}_n|_{\mathcal{S}_t} \cong \mathcal{I}|_{\mathcal{S}_t}.$$

□

15.5 Why the Argument Is Not Circular

The previous theorem has a tautological appearance, but it is not circular. The logical order is:

$$(X, d) \longrightarrow \mathcal{B}_{\text{RCI}} \longrightarrow \mathcal{I} \longrightarrow \mathcal{W}_n \longrightarrow \mathcal{S}_t \longrightarrow \widehat{\mathcal{I}}_n|_{\mathcal{S}_t}.$$

The admissible sieve $\mathcal{S}_t$ does not create the RCI invariant. It does not define the radial order, the local profile, the boundary index, or the cell. These are already generated from the minimal metric data (X, d) . Admissibility only removes the locus where the generated object changes type.

Thus the theorem does not say:

the object is robust because we define it to be robust.

It says:

once RCI has generated its base and invariants from minimal metric data,
restriction away from the wall certifies the resulting section.

In this sense, robustness is not an additional probabilistic conclusion attached after the algorithm. It is a structural mode of existence of the RCI object on the admissible base.

15.6 Local Sheafification versus Global Homogeneity

The structural robustness proved here does not require the entire metric space to behave as a globally homogeneous mass. It only requires the existence of an admissible region on which the RCI-generated local invariants are compatible under restriction.

Thus, for the certified RCI object, sheafification replaces a global regularity demand by a local compatibility principle:

global homogeneity $\rightsquigarrow$ local compatibility plus certified gluing.

Once a region has been restricted away from walls of type change, its compatible local data glue into a certified section. The certified object is therefore regional before it is global.

This is the epistemological advantage of the sheafified formulation. RCI does not need to prove that the whole dataset is uniformly regular. It extracts regions where the local metric invariants glue coherently, and it certifies those regions by admissible restriction:

regional certification by admissible gluing,

not global regularity of the ambient mass.

15.7 Relation with Appendix D

Appendix 14 and Appendix 15 prove different kinds of robustness.

Appendix 14 proves quantitative analytic control. It shows that, under smoothness, sampling, concentration, and spectral regularity assumptions, the empirical RCI quantities converge to their population counterparts and the boundary detector is stable with high probability.

Appendix 15 proves structural certification. It shows that, at the minimal metric level, once RCI has generated the operational base, local invariants, restriction maps, and wall of type change, the certified object is simply the restriction of the RCI invariant to the admissible sieve.

Thus:

Appendix D controls convergence;

Appendix E certifies admissible existence.

The analytic proof explains why empirical walls become stable under regularity assumptions. The structural proof explains why, once the wall has been identified, restriction to its complement yields a certified RCI section.

15.8 Conclusion

RCI does not invoke sheaves as an external formalism. Starting only from points and distances, it generates a radial base, local invariant data, restriction maps, walls of type change, and admissible sieves. The language of sheaves then recognizes the RCI output as a section over the admissible base.

Equivalently:

RCI begins with the minimal metric data required for geometry and returns with an object whose admissible restriction is sheaf-certified.

s

This is the structural form of RCI robustness: not stability imposed from outside, not homogeneity assumed globally, but certification arising from the restriction of the RCI-generated object to the domain where type change has been excluded.

References

- [1] Gérard Biau and Luc Devroye. *Lectures on the Nearest Neighbor Method*. Springer Series in the Data Sciences. Springer, 2015. ISBN 978-3-319-25388-6. doi: 10.1007/978-3-319-25388-6. URL <https://link.springer.com/book/10.1007/978-3-319-25388-6>.
- [2] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, 2013. ISBN 9780199535255. doi: 10.1093/acprof:oso/9780199535255.001.0001. URL <https://academic.oup.com/book/26549>.
- [3] Isaac Chavel. *Riemannian Geometry: A Modern Introduction*. Cambridge University Press, 2 edition, 2006. ISBN 9780521853682. URL <https://www.cambridge.org/core/books/riemannian-geometry/7F08C88F91F93C6FF45DE6901FFAD91E>.
- [4] Frédéric Chazal, Vin de Silva, and Steve Y. Oudot. Persistence stability for geometric complexes. *Geometriae Dedicata*, 173:193–214, 2014. doi: 10.1007/s10711-013-9937-z. Free access: <https://arxiv.org/abs/1207.3885>.
- [5] Frédéric Chazal, Vin de Silva, Marc Glisse, and Steve Oudot. *The Structure and Stability of Persistence Modules*. SpringerBriefs in Mathematics. Springer, 2016. ISBN 9783319425436. doi: 10.1007/978-3-319-42545-0. URL <https://link.springer.com/book/10.1007/978-3-319-42545-0>.
- [6] David Cohen-Steiner, Herbert Edelsbrunner, and John Harer. Stability of persistence diagrams. *Discrete & Computational Geometry*, 37(1):103–120, 2007. doi: 10.1007/s00454-006-1276-5. URL <https://link.springer.com/article/10.1007/s00454-006-1276-5>.
- [7] Luc Devroye, László Györfi, and Gábor Lugosi. *A Probabilistic Theory of Pattern Recognition*, volume 31 of *Stochastic Modelling and Applied Probability*. Springer, New York, 1996. ISBN 978-0-387-94618-7. URL <https://link.springer.com/book/10.1007/978-1-4612-0711-5>.
- [8] Martin Ester, Hans-Peter Kriegel, Jörg Sander, and Xiaowei Xu. A density-based algorithm for discovering clusters in large spatial databases with noise. In *Proceedings of the 2nd International Conference on Knowledge Discovery and Data Mining (KDD)*, pages 226–231, Portland, OR, 1996. AAAI Press. URL <https://dl.acm.org/doi/10.5555/3001460.3001507>.
- [9] Kenneth Falconer. *Fractal Geometry: Mathematical Foundations and Applications*. John Wiley & Sons, 3rd edition, 2014. ISBN 978-1-119-94239-9. URL <https://www.wiley.com/en-us/Fractal+Geometry:+Mathematical+Foundations+and+Applications,+3rd+Edition-p-9781119942399>.
- [10] Herbert Federer. Curvature measures. *Transactions of the American Mathematical Society*, 93(3):418–491, 1959. doi: 10.1090/S0002-9947-1959-0110078-1. URL <https://www.ams.org/journals/tran/1959-093-03/S0002-9947-1959-0110078-1/>.
- [11] Robin Forman. Morse theory for cell complexes. *Advances in Mathematics*, 134(1):90–145, 1998. doi: 10.1006/aima.1997.1650. URL <https://www.sciencedirect.com/science/article/pii/S0001870897916509>.
- [12] Robin Forman. A user’s guide to discrete morse theory. *Séminaire Lotharingien de Combinatoire [electronic only]*, 48:B48c (35 pp.), 2002. URL <http://eudml.org/doc/123837>.
- [13] Teofilo F. Gonzalez. Clustering to minimize the maximum intercluster distance. *Theoretical Computer Science*, 38:293–306, 1985. doi: 10.1016/0304-3975(85)90224-5. URL <https://www.sciencedirect.com/science/article/pii/0304397585902245>.
- [14] Alfred Gray. The volume of a small geodesic ball of a riemannian manifold. *Michigan Mathematical Journal*, 20(4):329–344, 1974. doi: 10.1307/mmj/1029001150. URL <https://projecteuclid.org/journals/michigan-mathematical-journal/volume-20/issue-4/The-volume-of-a-small-geodesic-ball-of-a-Riemannian/10.1307/mmj/1029001150.full>.
- [15] Jean-Claude Hausmann. On the vietoris–rips complexes and a cohomology theory for metric spaces. In Frank Quinn, editor, *Prospects in Topology: Proceedings of a Conference in Honor of William Browder*, volume 138 of *Annals of Mathematics Studies*, pages 175–188. Princeton University Press, 1995. doi: 10.1515/9781400882588-013. URL <https://www.degruyterbrill.com/document/doi/10.1515/9781400882588-013/html>.

- [16] Juha Heinonen. *Lectures on Analysis on Metric Spaces*. Universitext. Springer, 2001. ISBN 978-0387954043. doi: 10.1007/978-1-4613-0131-8. URL <https://link.springer.com/book/10.1007/978-1-4613-0131-8>.
- [17] Anil K. Jain, M. Narasimha Murty, and Patrick J. Flynn. Data clustering: A review. *ACM Computing Surveys*, 31(3):264–323, 1999. doi: 10.1145/331499.331504. URL <https://dl.acm.org/doi/10.1145/331499.331504>.
- [18] Svante Janson. Large deviations for sums of partly dependent random variables. *Random Structures and Algorithms*, 24(3):234–248, 2004. doi: 10.1002/rsa.20030. URL <https://www2.math.uu.se/~svantejs/papers/sj150.pdf>.
- [19] Yukio Matsumoto. *An Introduction to Morse Theory*, volume 208 of *Translations of Mathematical Monographs*. American Mathematical Society, 2002. ISBN 978-0-8218-1022-4. doi: 10.1090/mmono/208. URL <https://bookstore.ams.org/mmono-208>. Translated by Kiki Hudson and Masahico Saito.
- [20] Colin McDiarmid. On the method of bounded differences. In *Surveys in Combinatorics*, volume 141 of *London Mathematical Society Lecture Note Series*, pages 148–188. Cambridge University Press, 1989. doi: 10.1017/CBO9781107359949.008. URL <https://www.cambridge.org/core/books/abs/surveys-in-combinatorics-1989/on-the-method-of-bounded-differences/AABA597B562BDA7D89C6077E302694FB>.
- [21] John W. Milnor. *Morse Theory*. Annals of Mathematics Studies, 51. Princeton University Press, 1963. URL <https://webhomes.maths.ed.ac.uk/~v1ranick/papers/milnmors.pdf>.
- [22] Marston Morse. *The Calculus of Variations in the Large*, volume 18 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1934. ISBN 978-1-4704-3166-2. URL <https://bookstore.ams.org/coll-18>. Reprint available from Internet Archive: <https://archive.org/details/calculusofvariat0000mors>.
- [23] Partha Niyogi, Steve Smale, and Shmuel Weinberger. Finding the homology of submanifolds with high confidence from random samples. *Discrete & Computational Geometry*, 39(1–3):419–441, 2008. doi: 10.1007/s00454-008-9053-2. URL <https://link.springer.com/article/10.1007/s00454-008-9053-2>.
- [24] A. Singer. From graph to manifold laplacian: The convergence rate. *Applied and Computational Harmonic Analysis*, 21(1):128–134, 2006. doi: 10.1016/j.acha.2006.03.004. URL <https://www.sciencedirect.com/science/article/pii/S1063520306000510>.
- [25] Claude Tricot. *Curves and Fractal Dimension*. Springer-Verlag, New York, 1995. ISBN 978-0-387-94095-3. doi: 10.1007/978-1-4612-4170-6. URL <https://link.springer.com/book/9780387940953>. Translation from French. Foreword by M. Mendes France.
- [26] Nicolás García Trillos and Dejan Slepčev. Continuum limit of total variation on point clouds. *Archive for Rational Mechanics and Analysis*, 220:193–241, 2016. doi: 10.1007/s00205-015-0929-z. URL <https://link.springer.com/article/10.1007/s00205-015-0929-z>. Free access: <https://arxiv.org/abs/1403.6355>.
- [27] Karen K. Uhlenbeck. Connections with L^p bounds on curvature. *Communications in Mathematical Physics*, 83(1):31–42, 1982. doi: 10.1007/BF01208378. URL <https://projecteuclid.org/journals/communications-in-mathematical-physics/volume-83/issue-1/Connections-with-Lp-bounds-on-curvature/cmp/1103920743.full>.
- [28] Karen K. Uhlenbeck. Removable singularities in Yang–Mills fields. *Communications in Mathematical Physics*, 83(1):11–29, 1982. doi: 10.1007/BF01947068. URL <https://link.springer.com/article/10.1007/BF01947068>.
- [29] Ulrike von Luxburg. A tutorial on spectral clustering. *Statistics and Computing*, 17(4):395–416, 2007. doi: 10.1007/s11222-007-9033-z. URL <https://link.springer.com/article/10.1007/s11222-007-9033-z>.