



PC1 scores (48.1% variance)

Top loadings: PO4(uM) (+0.46), NO3(uM) (+0.44), NH4(uM) (+0.40)




PC2 scores (20.7% variance)

Top loadings: DO (+0.68), temp (−0.51), Sil(uM) (−0.29)




PC3 scores (12.2% variance)

Top loadings: FIECO-AFL(mg/m^3) (-0.80), depth_m (+0.37), NH4(uM) (+0.27)




PC4 scores (7.2% variance)

Top loadings: depth_m (-0.72), salinity (+0.57), FIECO-AFL(mg/m^3) (-0.33)




PC5 scores (5.5% variance)

Top loadings: Sil(uM) (+0.52), salinity (-0.45), temp (-0.37)




PC6 scores (3.2% variance)

Top loadings: NH4(uM) (+0.68), salinity (-0.45), depth_m (-0.41)




PC7 scores (2% variance)
Top loadings: DO (−0.63), temp (−0.54), NH4(uM) (−0.37)



PC8 scores (0.7% variance)

Top loadings: PO4(uM) (+0.69), Sil(uM) (-0.47), temp (+0.31)



PC9 scores (0.5% variance)

Top loadings: NO3(uM) (+0.78), PO4(uM) (-0.43), SiI(uM) (-0.30)

