

# Ordered fragments of first-order logic

Reijo Jaakkola

University of Helsinki, Tampere University

Funding: Theory of computational logics – Academy of Finland grants 324435  
and 328987

# The satisfiability problem

- ▶ An important invariant of a logic  $\mathcal{L}$  is the complexity of its satisfiability problem, i.e., the problem of determining whether a given sentence of  $\mathcal{L}$  is satisfiable (in other words has a model).

# The satisfiability problem

- ▶ An important invariant of a logic  $\mathcal{L}$  is the complexity of its satisfiability problem, i.e., the problem of determining whether a given sentence of  $\mathcal{L}$  is satisfiable (in other words has a model).
- ▶ A classical result of Church and Turing is that the satisfiability problem for full first-order logic FO is undecidable.

# The satisfiability problem

- ▶ An important invariant of a logic  $\mathcal{L}$  is the complexity of its satisfiability problem, i.e., the problem of determining whether a given sentence of  $\mathcal{L}$  is satisfiable (in other words has a model).
- ▶ A classical result of Church and Turing is that the satisfiability problem for full first-order logic FO is undecidable. This result led researches to focus their study on *fragments* of FO (essentially computable sets  $\mathcal{L} \subseteq \text{FO}$ ) with the hope that expressive logics with decidable satisfiability problem could be identified.

# The satisfiability problem

- ▶ An important invariant of a logic  $\mathcal{L}$  is the complexity of its satisfiability problem, i.e., the problem of determining whether a given sentence of  $\mathcal{L}$  is satisfiable (in other words has a model).
- ▶ A classical result of Church and Turing is that the satisfiability problem for full first-order logic FO is undecidable. This result led researches to focus their study on *fragments* of FO (essentially computable sets  $\mathcal{L} \subseteq \text{FO}$ ) with the hope that expressive logics with decidable satisfiability problem could be identified.
- ▶ An important example of an interesting decidable fragment of FO is the two-variable logic  $\text{FO}^2$  (every sentence can contain at most two variables).

- ▶ An important invariant of a logic  $\mathcal{L}$  is the complexity of its satisfiability problem, i.e., the problem of determining whether a given sentence of  $\mathcal{L}$  is satisfiable (in other words has a model).
- ▶ A classical result of Church and Turing is that the satisfiability problem for full first-order logic FO is undecidable. This result led researches to focus their study on *fragments* of FO (essentially computable sets  $\mathcal{L} \subseteq \text{FO}$ ) with the hope that expressive logics with decidable satisfiability problem could be identified.
- ▶ An important example of an interesting decidable fragment of FO is the two-variable logic  $\text{FO}^2$  (every sentence can contain at most two variables). This logic is decidable because it has the following *bounded model property*: if  $\varphi \in \text{FO}^2$  is satisfiable, then it has a model of size at most  $2^{|\varphi|}$ .

- ▶ Recently there has been an increasing interest in studying fragments that we will refer to as the *ordered fragments* of FO.

- ▶ Recently there has been an increasing interest in studying fragments that we will refer to as the *ordered fragments* of FO. These were originally introduced independently by Quine and Herzig.



- ▶ Recently there has been an increasing interest in studying fragments that we will refer to as the *ordered fragments* of FO. These were originally introduced independently by Quine and Herzog.
- ▶ **Main idea:** Restrict the order in which variables can be quantified, the way variables can be permuted in atomic formulas and the manner in which boolean combinations of formulas can be formed.

- ▶ Recently there has been an increasing interest in studying fragments that we will refer to as the *ordered fragments* of FO. These were originally introduced independently by Quine and Herzog.
- ▶ **Main idea:** Restrict the order in which variables can be quantified, the way variables can be permuted in atomic formulas and the manner in which boolean combinations of formulas can be formed.
- ▶ **This talk:** We will go through the syntax of the two most well-known ordered fragments (ordered logic, fluted logic) and their complexities. In addition, we will take a brief look at some recent results on the complexities of their variants (with respect to the satisfiability problem).

## Definition

Let  $\bar{v}_\omega = (v_1, v_2, \dots)$  be an infinite sequence of variables and let  $\tau$  be a vocabulary. For every  $k \in \mathbb{N}$  we define sets  $OL^k[\tau]$  as follows.

1. Let  $R \in \tau$  be an  $k$ -ary relational symbol and consider the prefix

$$(v_1, \dots, v_k)$$

of  $\bar{v}_\omega$ . Now  $R(v_1, \dots, v_k) \in OL^k[\tau]$ .

2. If  $\varphi, \psi \in OL^k[\tau]$ , then  $\neg\varphi, (\varphi \wedge \psi) \in OL^k[\tau]$ .

3. If  $\varphi \in OL^{k+1}[\tau]$ , then  $\exists v_{k+1}\varphi \in OL^k[\tau]$ .

Finally we define  $OL[\tau] := \bigcup_k OL^k[\tau]$ .

## Definition

Let  $\bar{v}_\omega = (v_1, v_2, \dots)$  be an infinite sequence of variables and let  $\tau$  be a vocabulary. For every  $k \in \mathbb{N}$  we define sets  $OL^k[\tau]$  as follows.

1. Let  $R \in \tau$  be an  $k$ -ary relational symbol and consider the prefix

$$(v_1, \dots, v_k)$$

of  $\bar{v}_\omega$ . Now  $R(v_1, \dots, v_k) \in OL^k[\tau]$ .

2. If  $\varphi, \psi \in OL^k[\tau]$ , then  $\neg\varphi, (\varphi \wedge \psi) \in OL^k[\tau]$ .

3. If  $\varphi \in OL^{k+1}[\tau]$ , then  $\exists v_{k+1}\varphi \in OL^k[\tau]$ .

Finally we define  $OL[\tau] := \bigcup_k OL^k[\tau]$ .

## Example

$\forall v_1 (\neg P(v_1) \wedge \exists v_2 R(v_1, v_2))$  is a sentence of  $OL[\{P, R\}]$ , while  $\exists v_1 \exists v_2 R(v_2, v_1)$ ,  $\exists v_2 \exists v_1 R(v_1, v_2)$  and  $\exists v_1 \exists v_2 (P(v_2) \wedge R(v_1, v_2))$  are not.

# Complexity of ordered logic

- ▶ One can prove that OL has the following bounded model property: if  $\varphi \in \text{OL}$  has a model, then it has one of size at most  $|\varphi|$ .

- ▶ One can prove that OL has the following bounded model property: if  $\varphi \in \text{OL}$  has a model, then it has one of size at most  $|\varphi|$ .
- ▶ **Main idea:** OL can't enforce that there exists more than  $|\varphi|$ -many elements with distinct (quantifier-free) unary types.

- ▶ One can prove that OL has the following bounded model property: if  $\varphi \in \text{OL}$  has a model, then it has one of size at most  $|\varphi|$ .
- ▶ **Main idea:** OL can't enforce that there exists more than  $|\varphi|$ -many elements with distinct (quantifier-free) unary types.

## Theorem (J.)

*Over bounded vocabularies the satisfiability problem of OL is NP-complete.*

- ▶ One can prove that OL has the following bounded model property: if  $\varphi \in \text{OL}$  has a model, then it has one of size at most  $|\varphi|$ .
- ▶ **Main idea:** OL can't enforce that there exists more than  $|\varphi|$ -many elements with distinct (quantifier-free) unary types.

## Theorem (J.)

*Over bounded vocabularies the satisfiability problem of OL is NP-complete.*

## Theorem (Herzig, J.)

*The satisfiability problem of OL is PSPACE-complete.*



## Definition

Let  $\bar{v}_\omega = (v_1, v_2, \dots)$  be an infinite sequence of variables and let  $\tau$  be a vocabulary. For every  $k \in \mathbb{N}$ , we define sets  $\text{FL}^k[\tau]$  as follows.

1. Let  $R \in \tau$  be an  $n$ -ary relation symbol and consider the subsequence

$$(v_{k-n+1}, \dots, v_k)$$

of  $\bar{v}_\omega$ . Now  $R(v_{k-n+1}, \dots, v_k) \in \text{FL}^k[\tau]$ .

2. For every  $\varphi, \psi \in \text{FL}^k[\tau]$ , we have that  $\neg\varphi, (\varphi \wedge \psi) \in \text{FL}^k[\tau]$ .
3. If  $\varphi \in \text{FL}^{k+1}[\tau]$ , then  $\exists v_{k+1}\varphi \in \text{FL}^k[\tau]$ .

Finally, we define  $\text{FL}[\tau] := \bigcup_k \text{FL}^k[\tau]$ .

## Definition

Let  $\bar{v}_\omega = (v_1, v_2, \dots)$  be an infinite sequence of variables and let  $\tau$  be a vocabulary. For every  $k \in \mathbb{N}$ , we define sets  $\text{FL}^k[\tau]$  as follows.

1. Let  $R \in \tau$  be an  $n$ -ary relation symbol and consider the subsequence

$$(v_{k-n+1}, \dots, v_k)$$

of  $\bar{v}_\omega$ . Now  $R(v_{k-n+1}, \dots, v_k) \in \text{FL}^k[\tau]$ .

2. For every  $\varphi, \psi \in \text{FL}^k[\tau]$ , we have that  $\neg\varphi, (\varphi \wedge \psi) \in \text{FL}^k[\tau]$ .
3. If  $\varphi \in \text{FL}^{k+1}[\tau]$ , then  $\exists v_{k+1}\varphi \in \text{FL}^k[\tau]$ .

Finally, we define  $\text{FL}[\tau] := \bigcup_k \text{FL}^k[\tau]$ .

Note that  $\text{OL} \subseteq \text{FL}$ .

## Definition

Let  $\bar{v}_\omega = (v_1, v_2, \dots)$  be an infinite sequence of variables and let  $\tau$  be a vocabulary. For every  $k \in \mathbb{N}$ , we define sets  $\text{FL}^k[\tau]$  as follows.

1. Let  $R \in \tau$  be an  $n$ -ary relation symbol and consider the subsequence

$$(v_{k-n+1}, \dots, v_k)$$

of  $\bar{v}_\omega$ . Now  $R(v_{k-n+1}, \dots, v_k) \in \text{FL}^k[\tau]$ .

2. For every  $\varphi, \psi \in \text{FL}^k[\tau]$ , we have that  $\neg\varphi, (\varphi \wedge \psi) \in \text{FL}^k[\tau]$ .
3. If  $\varphi \in \text{FL}^{k+1}[\tau]$ , then  $\exists v_{k+1}\varphi \in \text{FL}^k[\tau]$ .

Finally, we define  $\text{FL}[\tau] := \bigcup_k \text{FL}^k[\tau]$ .

Note that  $\text{OL} \subseteq \text{FL}$ .

## Example

$\exists v_1 \exists v_2 (P(v_2) \wedge R(v_1, v_2))$  is a sentence of  $\text{FL}[\{P, R\}]$ , while  $\exists v_1 \exists v_2 \exists v_3 (R(v_1, v_2) \wedge R(v_2, v_3))$  is not.

# Complexity of fluted logic

- ▶ FL also has a bounded model property: if  $\varphi \in \text{FL}$  has a model, then it has one of size at most

$$\underbrace{2^{2^{\dots 2}}}_{|\varphi|^k\text{-times}},$$

for some constant  $k$ .

- ▶ FL also has a bounded model property: if  $\varphi \in \text{FL}$  has a model, then it has one of size at most

$$\underbrace{2^{2^{\dots 2}}}_{|\varphi|^k\text{-times}},$$

for some constant  $k$ .

- ▶ **Main idea of proof:** For each  $k \geq 2$  and  $\varphi \in \text{FL}^{(k+1)}$ , there exists  $\varphi' \in \text{FL}^k$  so that  $\varphi$  has a model iff  $\varphi'$  has and if  $\varphi'$  has a model of size  $N$ , then  $\varphi$  has a model of size (roughly)  $2^N$ .

- ▶ FL also has a bounded model property: if  $\varphi \in \text{FL}$  has a model, then it has one of size at most

$$\underbrace{2^{2^{\dots^2}}}_{|\varphi|^k\text{-times}},$$

for some constant  $k$ .

- ▶ **Main idea of proof:** For each  $k \geq 2$  and  $\varphi \in \text{FL}^{(k+1)}$ , there exists  $\varphi' \in \text{FL}^k$  so that  $\varphi$  has a model iff  $\varphi'$  has and if  $\varphi'$  has a model of size  $N$ , then  $\varphi$  has a model of size (roughly)  $2^N$ .

## Theorem (Pratt-Hartmann, Swast, Tendera)

*The satisfiability problem of FL is TOWER-complete.*

# Algebraic view on ordered fragments

- ▶ Originally some of the ordered fragments were discovered by Quine as a by-product of his (eventually successful) attempt at giving a variable-free syntax for FO (predicate functor logic(s)).

# Algebraic view on ordered fragments

- ▶ Originally some of the ordered fragments were discovered by Quine as a by-product of his (eventually successful) attempt at giving a variable-free syntax for FO (predicate functor logic(s)).
- ▶ OL can be seen as consisting of three (relational) algebraic operators: complement  $\neg$ , intersection  $\cap$  and projection  $\exists$ .



# Algebraic view on ordered fragments

- ▶ Originally some of the ordered fragments were discovered by Quine as a by-product of his (eventually successful) attempt at giving a variable-free syntax for FO (predicate functor logic(s)).
- ▶ OL can be seen as consisting of three (relational) algebraic operators: complement  $\neg$ , intersection  $\cap$  and projection  $\exists$ .
- ▶ Similarly, FL consists of  $\neg$ ,  $\exists$  and the so-called suffix intersection  $\dot{\cap}$  (allows one to compute intersections of relations that have different arities).

- ▶ Originally some of the ordered fragments were discovered by Quine as a by-product of his (eventually successful) attempt at giving a variable-free syntax for FO (predicate functor logic(s)).
- ▶ OL can be seen as consisting of three (relational) algebraic operators: complement  $\neg$ , intersection  $\cap$  and projection  $\exists$ .
- ▶ Similarly, FL consists of  $\neg, \exists$  and the so-called suffix intersection  $\dot{\cap}$  (allows one to compute intersections of relations that have different arities).
- ▶ This point of view suggests naturally several syntactical variants of, say, OL and FL (simply add or remove algebraic operators).

# Brief summary of complexity results

- ▶ Adding (restricted) use of equality either to OL or FL does not affect complexity.

# Brief summary of complexity results

- ▶ Adding (restricted) use of equality either to OL or FL does not affect complexity.
- ▶ Adding a swap operator (swap the last two elements in every tuple) increases complexity significantly: OL becomes NEXPTIME-complete while FL becomes undecidable.

# Brief summary of complexity results

- ▶ Adding (restricted) use of equality either to OL or FL does not affect complexity.
- ▶ Adding a swap operator (swap the last two elements in every tuple) increases complexity significantly: OL becomes  $\text{NEXPTIME}$ -complete while FL becomes undecidable.
- ▶ Replacing  $\exists$  with one-dimensional quantification (select at most the first element from every tuple) decreases complexity: OL becomes NP-complete while FL becomes  $\text{NEXPTIME}$ -complete. FL remains  $\text{NEXPTIME}$ -complete even in the presence of equality and swap.

- ▶ Ordered fragments present a fresh viewpoint on the question of what makes satisfiability problems decidable (feasible).

- ▶ Ordered fragments present a fresh viewpoint on the question of what makes satisfiability problems decidable (feasible).
- ▶ Looking at intersections of ordered fragments with *guarded fragments* seems to be a very promising research direction (modal logics often have variable-free syntax).

- ▶ Ordered fragments present a fresh viewpoint on the question of what makes satisfiability problems decidable (feasible).
- ▶ Looking at intersections of ordered fragments with *guarded fragments* seems to be a very promising research direction (modal logics often have variable-free syntax).



- ▶ Ordered fragments present a fresh viewpoint on the question of what makes satisfiability problems decidable (feasible).
- ▶ Looking at intersections of ordered fragments with *guarded fragments* seems to be a very promising research direction (modal logics often have variable-free syntax).
- ▶ How much can we extend the expressive power of the fluted logic while preserving its expressive power?

- ▶ Ordered fragments present a fresh viewpoint on the question of what makes satisfiability problems decidable (feasible).
- ▶ Looking at intersections of ordered fragments with *guarded fragments* seems to be a very promising research direction (modal logics often have variable-free syntax).
- ▶ How much can we extend the expressive power of the fluted logic while preserving its expressive power?

Thanks! :-)