



Transcription factor binding sites detection in

Paramecium

Matthias Grenié *, Jean-François Goût † and Michael Lynch †

*ENS de Lyon, Département de Biologie, and †Indiana University, Biology Department, Lynch Lab

Submitted to Proceedings of the National Academy of Sciences of the United States of America

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Sed vehicula metus sapien. Suspendisse pulvinar, felis ut hendrerit aliquet, dui nisi bibendum erat, fermentum mattis enim nibh id arcu. Vestibulum ultrices eros sed odio tincidunt bibendum. Pellentesque fermentum ante vel nulla commodo fermentum. Vestibulum in augue sit amet libero viverra accumsan eu at magna. Sed at ligula quis nibh pharetra facilisis non eu libero. Suspendisse non quam sit amet massa luctus interdum sit amet in purus. Integer id orci elit, vitae sollicitudin lectus.

Keyword1 | Keyword2 | Keyword3

Abbreviations: SAM, self-assembled monolayer; OTS, octadecyltrichlorosilane

Introduction

S tructure of the introduction

- Whole Genome Duplications background
- WGD influences on Transcription Factor Binding Sites?
- the *Paramecium* project, experimental part, etc. some work is done and random sequences?
- Here, focusing on the computational part, developing pipeline, showed that etc.

Biological questions: How are TFBS affected by WGD? Are they conserved among species, is this linked to expression

- M. Belkin and P. Niyogi, Using manifold structure for partially labelled classification, Advances in NIPS, 15 (2003).
- P. Bérard, G. Besson, and S. Gallot, Embedding Riemannian manifolds by their heat kernel, Geom. and Fun. Anal., 4 (1994), pp. 374–398.
- R.R. Coifman and S. Lafon, Diffusion maps, Appl. Comp. Harm. Anal., 21 (2006), pp. 5-30.
- R.R. Coifman, S. Lafon, A. Lee, M. Maggioni, B. Nadler, F. Warner, and S. Zucker, Geometric diffusions as a tool for harmonic analysis and structure definition of data. Part I: Diffusion maps, Proc. of Nat. Acad. Sci., (2005), pp. 7426–7431.
- P. Das, M. Moll, H. Stamati, L. Kavraki, and C. Clementi, Low-dimensional, freeenergy landscapes of protein-folding reactions by nonlinear dimensionality reduction, P.N.A.S., 103 (2006), pp. 9885–9890.
- D. Donoho and C. Grimes, Hessian eigenmaps: new locally linear embedding techniques for high-dimensional data, Proceedings of the National Academy of Sciences, 100 (2003), pp. 5591–5596.
- D. L. Donoho and C. Grimes, When does isomap recover natural parameterization of families of articulated images?, Tech. Report Tech. Rep. 2002-27, Department of Statistics. Stanford University. August 2002.
- M. Grüter and K.-O. Widman, The Green function for uniformly elliptic equations, Man. Math., 37 (1982), pp. 303–342.

Reserved for Publication Footnotes

level? Conserved among each species? Is there a bias of TFBS usage in certain species?

Results

Discussion

Conservation among species. Major results is that. Divergent resolution of WGD \rightarrow divergence in motifs?

Motifs detection should take phylogeny into account for comparative analysis. Not the same value.

Materials and Methods

developped a whole pipeline (show simple pipeline graph)

- Families upstream sequences extraction
- CDSs extraction and alignment
- CDSs phylogenetic tree
- BigFoot identification (explanation of phylogenetic score and alignment score)
- MEME research
- Comparison MEME and BigFoot
- Identification of given motif in species genome
- Correlation between motifs and expression levels

ACKNOWLEDGMENTS. Thanks to M. Lynch and J.-F. Goût. Funding by NSF.

- R. Hempel, L. Seco, and B. Simon, The essential spectrum of neumann laplacians on some bounded singular domains, 1991.
- Kadison, R. V. and Singer, I. M. (1959) Extensions of pure states, Amer. J. Math. 81, 383-400.
- 11. Anderson, J. (1981) A conjecture concerning the pure states of B(H) and a related theorem. in Topics in Modern Operator Theory, Birkhaüser, pp. 27-43.
- 12. Anderson, J. (1979) Extreme points in sets of positive linear maps on ${\cal B}(H)$. J. Funct. Anal. 31, 195-217.
- 13. Anderson, J. (1979) Pathology in the Calkin algebra. J. Operator Theory 2, 159-167.
- Johnson, B. E. and Parrott, S. K. (1972) Operators commuting with a von Neumann algebra modulo the set of compact operators. J. Funct. Anal. 11, 39-61.
- Akemann, C. and Weaver, N. (2004) Consistency of a counterexample to Naimark's problem. Proc. Nat. Acad. Sci. USA 101, 7522-7525.
- J. Tenenbaum, V. de Silva, and J. Langford, A global geometric framework for nonlinear dimensionality reduction, Science, 290 (2000), pp. 2319–2323.
- Z. Zhang and H. Zha, Principal manifolds and nonlinear dimension reduction via local tangent space alignement, Tech. Report CSE-02-019, Department of computer science and engineering, Pennsylvania State University, 2002.



PNAS | Issue Date | Volume | Issue Number | 1–2





Placeholder

Image

Fig. 1. Figure caption

Table 1. Table caption

Treatments	Response 1	Response 2
Treatment 1	0.0003262	0.562
Treatment 2	0.0015681	0.910
Treatment 3	0.0009271	0.296

2 | www.pnas.org — —



Footline Author