



The AFIT of Today is the Air Force of Tomorrow.



U.S. AIR FORCE

Probability, Statistics, and Calculus Review

LOGM 634

9 October 2014





Statistics & Probability



The AFIT of Today is the Air Force of Tomorrow.

- Statistics
 - Inferring a population's behavior based on the performance of a sample
 - Representative of the population?
 - Sample is unbiased?
- Probability
 - Predicting the future behavior of a single item, based on the population's known performance



Another view



The AFIT of Today is the Air Force of Tomorrow.

- Probability:
 - The study of the laws of chance
- Statistics:
 - Use probability laws on past information to predict future events



Modeling Uncertainty



The AFIT of Today is the Air Force of Tomorrow.

- Basic Method
 - Sample space & events within an experiment
 - Define probabilities of events
 - Compute probabilities of more complex events formed from unions/intersections of the elementary events
- Concept of Random Variables
 - Variable that takes on certain values iaw specific probabilities
 - Specify the probability distribution of a random variable to characterize a random process
- Use both through the course



Key Terms



The AFIT of Today is the Air Force of Tomorrow.


- Probability
- Experiments
- Sample spaces
- Outcomes
- Events
- Random variables



An Experiment

The AFIT of Today is the Air Force of Tomorrow.

- Probability analysis is always made in the context of some *experiment*
 - If I exceed my aircraft's weight and balance limits, can I still take off?
 - I bought a ticket -- will I win the lottery tonight?



I was just handed two dice -- will I roll a seven?




Sample Space


The AFIT of Today is the Air Force of Tomorrow.

- Collection of all possible outcomes of an experiment
 - Mutually exclusive
 - Collectively exhaustive
- Countable (discrete)

We know something will happen in our next experiment, we're just not sure what!



- Uncountable (continuous)





Events



The AFIT of Today is the Air Force of Tomorrow.

- Just a collection (subset) of outcomes contained in the sample space
 - one, a few, or all of 'em
 - Exactly 1 = simple
 - > 1 = compound




A Random Variable

The AFIT of Today is the Air Force of Tomorrow.

- A variable that takes on numerical values iaw some probability distribution
- May be continuous or discrete
- Nothing random about it – it's not really a variable!
- It's a FUNCTION
 - It maps an event (a single outcome or a set of 'em) to the *real number line*,




(runner breaks tape)



(runners total time)




Roll the dice...



The AFIT of Today is the Air Force of Tomorrow.

The sample space





What about the Probability?



The AFIT of Today is the Air Force of Tomorrow.

- Probability is the ***likelihood*** of an outcome
 - Call it “X”
- For each event, X, in the sample space, we assume the probability of X occurring has the following properties:
 - for any specified event X, $0 \leq P(X) \leq 1$
 - If $P(X) = 1$, then X is a *certainty*
 - If $P(X) = 0$, then X is an *impossibility*
 - *Doesn't mean x won't happen – it's just highly unlikely*




What about...

The AFIT of Today is the Air Force of Tomorrow.

- The union:
 - The probability that either x or y or both occur

$$P(X \cup Y) = P(X) + P(Y) - P(X \cap Y)$$



The union

$$P(X \cup Y) = ?$$

The intersection



$$P(X \cap Y) = ?$$



What about.....

The AFIT of Today is the Air Force of Tomorrow.

- Are X and Y *Mutually Exclusive*?



The union



$$P(X \cup Y) = ?$$

The intersection

$$P(X \cap Y) = ?$$



What about.....



The AFIT of Today is the Air Force of Tomorrow.

If X and Y are mutually exclusive then X and Y cannot occur together and $P(X \cap Y) = 0$

So, when X and Y are Mutually Exclusive, then

$$P(X \cup Y) = P(X) + P(Y) - P(X \cap Y)$$

$$P(X \cup Y) = P(X) + P(Y) - 0$$



$$P(X \cup Y) = P(X) + P(Y)$$



What about.....

The AFIT of Today is the Air Force of Tomorrow.

- Are X and Y *Independent*?



The union

$$P(X \cup Y) = ?$$

The intersection


$$P(X \cap Y) = ?$$



Conditional Probabilities

The AFIT of Today is the Air Force of Tomorrow.

- Theorem for Conditional Probability:



Defn: If $P(Y) > 0$, then


$$P(X|Y) = \frac{P(X \cap Y)}{P(Y)} \rightarrow P(X \cap Y) = P(X|Y)P(Y)$$



Conditional & Independent

The AFIT of Today is the Air Force of Tomorrow.

- If X and Y are independent, then neither influences the outcome of the other



then, $P(X|Y) = P(X)$

and $P(X \cap Y) = P(X|Y)P(Y) = P(X)P(Y)$



What about.....

The AFIT of Today is the Air Force of Tomorrow.


- So, back to the Union of X and Y,

If X and Y are Independent, then

$$P(X \cup Y) = P(X) + P(Y) - P(X \cap Y)$$


$$P(X \cup Y) = P(X) + P(Y) - P(X|Y)P(Y)$$

$$P(X \cup Y) = P(X) + P(Y) - P(X)P(Y)$$



When X and Y are Mutually Exclusive,

$$P(X \cup Y) = P(X) + P(Y)$$





Discrete vs Continuous

The AFIT of Today is the Air Force of Tomorrow.

- Discrete: when can *count* stuff
 - We can be **exact**

$$P(X) = P(X = x)$$

- Continuous: used for measuring (other) stuff
 - We're using some tool (clock, meter, etc.)
 - can **never** be exact
 - We thus need to specify a tolerance or confidence

$$P(X) = P(X \leq x)$$



'Formal' Probability


The AFIT of Today is the Air Force of Tomorrow.

- Discrete: use a probability mass function (pmf)
 - We're interested in the *relative frequency* of each event
 - Example: our dice toss
 - can roll a '7' six of 36 ways -- hence $p(7) = 1/6$
 - can roll a '2' only 1 way -- hence $p(2) = 1/36$
 - what about rolling a '13'?
- What's the big assumption?
 - The dice are *fair* -- each face equally likely to appear on each die
- Since the sample space is discrete, we can assign a probability to **every outcome**



A pmf (Our dice toss)

The AFIT of Today is the Air Force of Tomorrow.





Continuous Data


The AFIT of Today is the Air Force of Tomorrow.

- We use a probability density function (pdf)
- Can **not** assign a probability to an exact value -- can only assign probabilities to ranges of values
 - Calculus alert! (the pdf is a *derivative*)
- The shape of the pdf curve shows us what values of a *random variable* are most likely to occur
 - I.e., values of x occur most frequently where $f(x)$ is largest



A pdf

The AFIT of Today is the Air Force of Tomorrow.




The amount of time until a component fails



Graph of a CDF

The AFIT of Today is the Air Force of Tomorrow.





The cdf

The AFIT of Today is the Air Force of Tomorrow.

- The *cumulative distribution function* is based on either the pmf or pdf
 - depends on the data type
 - Gives us the probability of a random variable X taking a value equal to, or less than, some specified value x

$$F(x) = P(X \leq x) \quad \text{for } -\infty < x < \infty$$



cdf Example

The AFIT of Today is the Air Force of Tomorrow.

Let X be the number I'll roll on my next toss of two dice.

The probability I'll roll a 7 or less is

$$P(X \leq 7) = \sum_{k=0}^7 p(x) = 21 / 36$$

It's easy, in the discrete case -- we just sum up all the probabilities less than or equal to the value of interest.



Example

The AFIT of Today is the Air Force of Tomorrow.

Let X be the amount of time to component failure

The probability the component will fail in 225 hours is written as

$$P(T \leq 225 \text{ hrs})$$

Since time is a continuous-valued variable, I would really need to *integrate* some pdf over the interval (0, 225) to compute this probability!



Integrals??!


The AFIT of Today is the Air Force of Tomorrow.

- An integral is a sum of infinitely small things
- Its usually thought of as the area under a curve
- In our case:
 - The *pdf* (a function) defines the shape of the curve
 - A *cdf* value, $F(t)$, represents the *cumulative* area under the pdf curve within the interval $[-\infty, t]$
 - The *total* area under the pdf curve always equals 1
 - To get the area within some interval $[a,b]$, an integral adds all possible pdf values, each evaluated for a small interval dt (for values of t within $[a,b]$)



pdf and cdf


The AFIT of Today is the Air Force of Tomorrow.





Integration

The AFIT of Today is the Air Force of Tomorrow.



An integral as the limit of a sum of rectangles $\rightarrow dt$ wide by $f(t)$ tall.
As dt becomes smaller, the rectangles better approximate the true area.

$$\int_a^b f(t) dt \approx \text{sum} \left(\sum \right) \text{ of the rectangles over the interval } [a, b]$$



The Basic Integral

The AFIT of Today is the Air Force of Tomorrow.

We're integrating with respect to a variable called "x".
We evaluate the integral for values of x within [a,b]

$$\int_a^b dx$$

Indicates the variable involved (integrate wrt)

Integrand: The stuff we insert here—usually a function

- It is the pdf $\rightarrow f(x)$
- Or, the expectation formula $\rightarrow x f(x)$
- If we insert nothing, then we assume the integrand is a *constant*, and equal to 1.0



Integral Examples

The AFIT of Today is the Air Force of Tomorrow.

- The (continuous) cdf is an integral value

$$P(T \leq a) = F(a) = \int_0^a f(t) dt$$

- The probability of failure over a specific time interval [a,b]:

$$P(a \leq T \leq b) = F(b) - F(a)$$

$$= \int_0^b f(t) dt - \int_0^a f(t) dt$$

$$= \int_a^b f(t) dt$$



Some Useful Integrals

The AFIT of Today is the Air Force of Tomorrow.

$$\int k dx \rightarrow kx$$

$$\int kf(x) dx \rightarrow k \int f(x) dx$$

$$\int [f(x) + g(x)] dx \rightarrow \int f(x) dx + \int g(x) dx$$

$$\int x^n dx = \frac{1}{n+1} x^{n+1} \rightarrow \int x^3 dx = \frac{1}{4} x^4$$

$$\int x^{-1} dx = \int \frac{1}{x} dx \rightarrow \ln x$$



Some More Useful Integrals



The AFIT of Today is the Air Force of Tomorrow.

$$\int e^{ax} dx = \frac{1}{a} e^{ax} \quad \rightarrow \quad \int e^{2x} dx = \frac{1}{2} e^{2x}$$

$$\int b e^{ax} dx = \frac{b}{a} e^{ax} \quad \rightarrow \quad \int 3e^{\lambda t} dt = \frac{3}{\lambda} e^{\lambda t}$$

$$\ln e = 1$$

$$\ln e^k = k$$



Applications of the Integral



The AFIT of Today is the Air Force of Tomorrow.

- The mean is an *expectation*
 - Each possible value, t , weighted by its “rectangle area”
 - We use the cdf integrand, but now multiply the integrand by t
 - We integrate over all possible values of t (from $t = 0$ to ∞)

$$E(T) = \int_0^{\infty} t f(t) dt$$

$$MTTF = E(T) = \int_0^{\infty} R(t) dt \quad \text{(A form you'll see soon)}$$



Compare with Discrete Variables



The AFIT of Today is the Air Force of Tomorrow.

- To help see what the integral is doing, recall the average value (expectation) of a discrete RV:

$$E(X) = \sum_{all\ x} x p(x)$$

- In words: the average value of a discrete RV is simply the sum of all possible values of x , each weighted by its probability of occurrence $p(x)$
- The integral is doing the same thing, except we must deal with the “billions and billions” (infinitely many) t values



Variance

The AFIT of Today is the Air Force of Tomorrow.

- The 'spread' or dispersion about the mean

$$Var(X) = E\left[(X - \mu)^2\right]$$

$$Var(X) = \sum_{all\ x} (x - \mu)^2 p(x) \quad (\text{discrete})$$

$$Var(X) = \int_0^{\infty} (x - \mu)^2 f(x) dx \quad (\text{continuous})$$



Variance—An easier way

The AFIT of Today is the Air Force of Tomorrow.

- The 'spread' or dispersion about the mean

$$\text{Var}(X) = E\left[(X - \mu)^2\right] = E\left[X^2\right] - \mu^2 = E\left[X^2\right] - E[X]^2$$

- Note: To find the second moment

$$E\left[X^2\right] = \sum_{\text{all } x} x^2 p(x) \quad (\text{discrete})$$

$$= \int_0^{\infty} x^2 f(x) dx \quad (\text{continuous})$$



Uniform Distribution, $U(a, b)$



The AFIT of Today is the Air Force of Tomorrow.

- Useful when occurrences randomly vary and little else is known about the shape of the distribution

$$pdf : f(t) = \begin{cases} \frac{1}{b-a} & \text{if } a \leq t \leq b \\ 0 & \text{otherwise} \end{cases}$$



Normal Distribution, $N(\mu, \sigma^2)$



The AFIT of Today is the Air Force of Tomorrow.

- Used for errors of various types

pdf:
$$f(t) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(t-\mu)^2}{2\sigma^2}} \quad \text{for } -\infty < t < \infty$$

cdf: no closed form \rightarrow use look-up tables



- Time to perform some task

$$\text{pdf: } f(t) = \begin{cases} \left(\frac{1}{t\sigma\sqrt{2\pi}} \right) e^{-(\ln t - \mu)^2 / (2\sigma^2)} & \text{if } t > 0 \\ 0 & \text{otherwise} \end{cases}$$

cdf: no closed form \rightarrow use look-up tables



- Time to complete some task, time to failure

$$\text{pdf: } f(t) = \begin{cases} \frac{\beta}{\theta} \left(\frac{t}{\theta} \right)^{\beta-1} e^{-(t/\theta)^\beta} & \text{if } t > 0 \\ 0 & \text{otherwise} \end{cases}$$

$$\text{cdf: } F(t) = \begin{cases} 1 - e^{-(t/\theta)^\beta} & \text{if } t > 0 \\ 0 & \text{otherwise} \end{cases}$$



Exponential Distribution

Expon(β)

The AFIT of Today is the Air Force of Tomorrow.

- Constant failure rates, $1/\beta$

$$\text{pdf: } f(t) = \begin{cases} \frac{1}{\beta} e^{-t/\beta} & \text{if } t \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

$$\text{cdf: } F(t) = \begin{cases} 1 - e^{-t/\beta} & \text{if } t \geq 0 \\ 0 & \text{otherwise} \end{cases}$$



Binomial Distribution, $\text{bin}(n, p)$

The AFIT of Today is the Air Force of Tomorrow.

- Discrete distribution representing # of successes in n independent Bernoulli trials where the probability of success for each trial is p
- The binomial (a set of Bernoulli trials)
 - Probability I'll get exactly 3 '7's on my next 5 tosses of two dice
 - Probability of exactly 1 failure among 5 identical components

$$\text{pmf: } p(x) = \binom{n}{x} p^x (1-p)^{n-x} \quad \text{for } x=0,1,\dots,n$$

$$\text{where, } \binom{n}{x} = \frac{n!}{(n-x)!x!} \quad (\text{binomial coefficient})$$



A Poisson Process

The AFIT of Today is the Air Force of Tomorrow.


- A unique relationship between the Poisson pmf and the exponential pdf...
- It's WHEN the time between successive occurrences is exponentially distributed, AND
 - Failure rate = λ
- The number of occurrences in a given time, t , is Poisson distributed
 - Expected # failures over time $t = \lambda t$



Poisson Process – the idea...

The AFIT of Today is the Air Force of Tomorrow.

Time between successive occurrences is exponentially distributed with parameter λ



Number of occurrences in a specified time interval t is Poisson distributed with parameter λt



Poisson Process Assumptions



The AFIT of Today is the Air Force of Tomorrow.

- Things happen *serially* (no stinkin' batches)
- *Stationarity* (parameters are constant w.r.t. time)
- *Independent* increments:
 - What happens in one time period doesn't affect what happens in a different, non-overlapping period



Poisson Distribution

The AFIT of Today is the Air Force of Tomorrow.

- The Poisson is a discrete distribution representing the number of random events that occur in an interval of time, when events occur at a constant rate
 - Probability I'll get exactly 5 emails in the next hour, given they average about λ per hour

$$\text{pmf: } p_n(t) = \frac{e^{-\lambda t} (\lambda t)^n}{n!}$$



Other Useful Things!

The AFIT of Today is the Air Force of Tomorrow.

$$x^m x^n = x^{(m+n)}$$

$$\text{Log}_a(x) = y \iff a^y = x$$

$$\frac{x^m}{x^n} = x^{(m-n)}$$

$$\text{Log}_a(a^x) = x$$

$$(x^m)^n = x^{mn}$$

$$\text{Log}_a(r^x) = x \text{Log}_a(r)$$

$$(xy)^m = x^m y^m$$

$$\text{Log}_a(xy) = \text{Log}_a(x) + \text{Log}_a(y)$$

$$\text{Log}_a(x/y) = \text{Log}_a(x) - \text{Log}_a(y)$$

$$\text{Log}_e(x) = \ln x$$