

SEARCH FOR RESONANT DOUBLE HIGGS PRODUCTION WITH  $b\bar{b}Z\bar{Z}$   
DECAYS IN THE  $b\bar{b}\ell\ell\nu\bar{\nu}$  FINAL STATE IN  $pp$  COLLISIONS AT  $\sqrt{s} = 13$  TeV

by

Rami Kamalieddin

A DISSERTATION

Presented to the Faculty of  
The Graduate College at the University of Nebraska  
In Partial Fulfilment of Requirements  
For the Degree of Doctor of Philosophy

Major: Physics and Astronomy

Under the Supervision of Professor Ilya Kravchenko

Lincoln, Nebraska

July, 2019

SEARCH FOR RESONANT DOUBLE HIGGS PRODUCTION WITH  $b\bar{b}Z\bar{Z}$   
DECAYS IN THE  $b\bar{b}\ell\ell\nu\bar{\nu}$  FINAL STATE IN  $pp$  COLLISIONS AT  $\sqrt{s} = 13$  TeV

Rami Kamalieddin, Ph.D.

University of Nebraska, 2019

Adviser: Ilya Kravchenko

Since the discovery of the Higgs boson in 2012 by the ATLAS and CMS experiments, most of the quantum mechanical properties that describe the long-awaited Higgs boson have been measured. Due to the outstanding work of the LHC, over a hundred of  $\text{fb}^{-1}$  of proton collisions data have been delivered to both experiments. Finally, it became sensible for analyses teams to start working with a very low cross section processes involving the Higgs boson, e.g., a recent success in observing  $t\bar{t}H$  and  $VHbb$  processes. One of the main remaining untouched topics is a double Higgs boson production. However, an additional hundred of  $\text{fb}^{-1}$  per year from the HL-LHC will not necessarily help us much with the SM double Higgs physics, as the process may remain unseen even in the most optimistic scenarios. The solution is to work in parallel on new reconstruction and signal extraction methods as well as new analysis techniques to improve the sensitivity of measurements. This thesis is about both approaches: we have used the largest available dataset at the time the analysis has been performed and developed/used the most novel analysis methods. One such method is the new electron identification algorithm that we have developed in the CMS electron identification group, to which I have had a privilege to contribute during several years of my stay at CERN.

The majority of this thesis is devoted to techniques for the first search at the LHC for double Higgs boson production mediated by a heavy narrow-width resonance in

the  $b\bar{b}ZZ$  channel:  $X \rightarrow HH \rightarrow b\bar{b}ZZ^* \rightarrow b\bar{b}\ell\ell\nu\bar{\nu}$ . The measurement searches for the resonant production of a Higgs boson pair in the range of masses of the resonant parent particle from 250 to 1000 GeV using  $35.9 \text{ fb}^{-1}$  of data taken in 2016 at 13 TeV. Two spin scenarios of the resonance are considered: spin 0 and spin 2. In the absence of the evidence of the resonant double Higgs boson production from the previous searches, we proceed with setting the upper confidence limits.

*“Here will be a quote ”*

*name, year.*

---

## Table of Contents

---

---

## List of Figures

---

---

## List of Tables

---

## CHAPTER 1

---

### *bbZZ* Physics Analysis

---

In this chapter, we report on a search for the resonant production of a double Higgs boson system. We select Higgs boson pairs that subsequently decay through the *bbZZ* channel. The final state consists of two b jets, two charged leptons, and two neutrinos (*2b2l2ν* final state). The analyzed data set was collected in 2016 by the CMS experiment in proton-proton collisions at 13 TeV COM energy, and corresponds to 35.9 fb<sup>-1</sup> of integrated luminosity.

### 1.1 Physics analysis overview

We search for di-Higgs production through the gluon fusion mechanism mediated by two types of possible heavy resonances (separately): a spin-2 Randall-Sundrum (RS1) Kaluza-Klein (KK) graviton and a spin-0 RS1 radion [?, ?]. The width of the graviton and radion is assumed to be negligible with respect to the experimental resolution. We look for decays of WED particles (X) to the *bbZZ* channel with the two b jets, two charged leptons, and two neutrinos in the final state. However, the *bbWW* intermediate channel can also contribute (0(20)%) to the final state of this measurement. Therefore, the full chain of this measurement consists of  $X \rightarrow HH \rightarrow bbZZ/bbWW \rightarrow 2b2l2\nu$ . In the above, one of the Higgs bosons decays to



a pair of b quarks and the other Higgs boson decays to ZZ or WW system. For the purposes of this measurement, we consider only leptonic decays of ZZ and WW. Intermediate taus are allowed to decay to daughter muons and electrons. We explore the invariant masses of WED particles ranging from 250 to 1000 GeV. The result of the measurement is the upper limit on the production cross section of the resonance multiplied by the branching fraction of its decay into the aforementioned  $2b2l2\nu$  final state.

In this data analysis (later referred to as the “analysis”), we will describe first the data sets (often referred to as a “datasets”) and data triggers. The measurement uses the DoubleEG and DoubleMuon PDs defined in the previous chapter. These di-electron and di-muon channels are analyzed separately and the information is combined in the final result. To select the events with two prompt charged leptons, a set of triggers was applied, when the events were recorded. To increase the statistics of the measurement and maximize the number of leptons passing the selection, certain complex trigger strategies were employed.

We then discuss the MC simulation of the signal and background processes (later referred to as “signal” and “background”). The signal MC samples had to be produced for both graviton and radion particles for 16 mass points from 250 to 1000 GeV to cover the whole search range.

The description of the physics object reconstruction and event selection was given in the previous chapter. Physics objects are constructed using the information from the CMS subsystems and the output of the PF algorithm. Then, based on the final state signature, the events containing corresponding physics objects are selected. The construction of the Higgs and Z boson candidates is discussed at length. The characteristics of the signal and background processes are specified and data-MC SFs are discussed.

The MVA technique is used to improve signal-background discrimination. We rely on the BDT classifier (commonly referred to as the BDT for brevity) to reduce the contribution of background processes in the signal-enriched regions. In all physics measurements, there are sources of systematic and statistical uncertainties that affect the final results. We discuss all major uncertainties at length and specify the size of the effect of individual types of uncertainties on the final result.

We present the statistical analysis used to extract the results of the measurements and discuss the CMS statistical package that has been used to extract the final results. Then, we present the results of the measurement and compare them with the theoretical predictions. Finally, we discuss the “grand  $HH$  combination” using measurements of all available  $HH$  channels.

The material in this chapter follows the description of two articles to which the author contributed directly [?, ?].

## 1.2 Data analysis strategy

We form the Lorentz energy-momentum vector associated with the double Higgs system. This vector is constructed as the sum of the Lorentz vectors of the two leptons, two b jets, and the four-vector representing neutrinos ( $\not{p}_T$ ). As the z component of the neutrinos’ momentum is unknown, we form a pseudo-transverse mass:  $\tilde{M}_T(HH) = \sqrt{E^2 - p_z^2}$  (further referred to as “transverse mass” for brevity), where  $E$  and  $p_z$  are the energy and the z-axis component of the Lorentz energy-momentum vector of the di-Higgs candidate.

$\tilde{M}_T(HH)$  distributions are derived for 16 masses and for both spin hypotheses. The  $\tilde{M}_T(HH)$  is constructed using signal and background simulations. The distributions are produced separately for the signal-enriched (SR) and control regions (CR), which

are defined later in this chapter.

To extract the final results, we perform a simultaneous fit of the sum of the signal and background  $\tilde{M}_T(\text{HH})$  distributions in the SR and CRs to the data. The fits are produced for both di-electron and di-muon channels and the results are combined. We obtain 95% CL upper limits on the  $HH$  production cross section multiplied by the BF of the subsequent decay to the final state of this measurement. The employed statistical method is based on the  $\text{CL}_S$  asymptotic procedure [?].

## 1.3 Data and Triggers

### 1.3.1 Data

The search is performed using DoubleMuon and DoubleEG PDs corresponding to an integrated luminosity of  $35.9 \text{ fb}^{-1}$  recorded by the CMS experiment during Run 2 in 2016. The data were produced in proton-proton collisions at the LHC at 13 TeV COM energy. The data were collected in several data taking periods and approximate values of the integrated luminosity of these periods are given in Table ??.

**Table 1.1:** List of data sets collected by the CMS in 2016. Each era contains a unique letter identifier. Also, the date of the data processing is specified. If re-processing was run, it is labeled as “v2”. Corresponding integrated luminosities are shown in the second column.

Dataset	$\int \mathcal{L} \text{ (fb}^{-1}\text{)}$
Run2016B-03Feb2017-v2	$\sim 5.9$
Run2016C-03Feb2017-v1	$\sim 2.7$
Run2016D-03Feb2017-v1	$\sim 4.3$
Run2016E-03Feb2017-v1	$\sim 4.1$
Run2016F-03Feb2017-v1	$\sim 3.2$
Run2016G-03Feb2017-v1	$\sim 3.8$
Run2016H-03Feb2017-v1	$\sim 11.8$
Total Luminosity	$\sim 35.9$


### 1.3.2 Triggers

The data events that are used in this measurement are selected with a set of HLT triggers that required the presence of two muons or two electrons in the event. In the di-electron final state, the trigger requires the leading electron to have  $p_T$  above 23 GeV, and the sub-leading electron to have  $p_T$  above 12 GeV. The latter leg corresponds to a relatively low-efficiency electron. Since the measurement is focused on “golden” electrons coming from on-shell Z boson decays, the offline selection is slightly tighter than the HLT selection and requires the sub-leading electron to have  $p_T$  above 15 GeV.


The final state that contains two prompt muons is selected with an HLT path which is a combination of several HLT paths chained together using the logical “OR” operation. In other words, the muons are required to pass the selection of at least one of the HLT paths. At the HLT level, the leading and sub-leading muons have to pass the 17 and 8 GeV  $p_T$  requirements respectively. The difference among HLT paths has been explained at length in Section ???. The offline analysis increases the  $p_T$  threshold values to 20 and 15 GeV correspondingly. For the offline selection, electrons are selected in the range  $|\eta| < 2.5$  and muons in the range  $|\eta| < 2.4$ . For both channels, the  $|\eta|$  region in the gap (1.4442 to 1.566) between the barrel and endcap is excluded.

The same trigger selections are applied to MC simulated events. The efficiencies are then derived for MC and data, and SFs are determined using the TnP procedure discussed in Section ???. Trigger SFs have been computed for each trigger leg separately since the selection of each leg varies (Fig. ??). Following the recommendations from the CMS Muon POG, scale factors have been calculated separately for two data collecting periods: runs B to G (Fig. ??, and run H separately (Fig. ??), since


the LHC conditions varied significantly for the run H. All eras have slightly different integrated luminosities, see Table ??, so the final SFs are luminosity averaged. Additionally, as was discussed in the chapter on CMS Physics Objects Reconstruction, some triggers did not contain the DZ requirement, while others did. Therefore, scale factors of the DZ selection are also measured, see Fig. ??). Prior to measuring the trigger scale factors, the electrons and muons were required to satisfy ID and ISO selections; more details are given in the subsections ??.




**Figure 1.1:** HLT trigger SFs for electrons approved by the CMS  $e/\gamma$  POG group. SFs are derived for both legs separately: Leg 1 (top) corresponds to the leading electron and Leg 2 (bottom) corresponds to the sub-leading electron. The values of the SFs are shown on the left, and the associated uncertainties with each value are shown on the right. Taken from [?].



**Figure 1.2:** Final HLT SFs for muons as a function of  $p_T$  and  $\eta$ , measured for eras B to G. Left: Scale factors for 8 GeV leg (sub-leading muon). Right: Scale factors for 17 GeV leg (leading muons), provided that the sub-leading muon already passed the 8 GeV  $p_T$  requirement.



**Figure 1.3:** Final HLT SFs for muons as a function of  $p_T$  and  $\eta$ , measured for era H. Left: Scale factors for 8 GeV leg (sub-leading muon). Right: Scale factors for 17 GeV leg (leading muons), provided that the sub-leading muon already passed the 8 GeV  $p_T$  requirement.



**Figure 1.4:** SFs of the dZ requirement for muons as a function of the  $\eta$  values of both muons.

## 1.4 Simulated Samples

The data analysis carried out in this thesis is optimized using the MC simulation. MC samples for signal and background processes have been produced with various HEP software packages (“generators”) that generate the processes of interest: processes that create the final state of this measurement.

### 1.4.1 Signal processes simulation

The signal Monte Carlo (MC) samples have been generated using the MADGRAPH5\_aMC@NLO [?] package. In these samples, the process of the gluon fusion production of WED spin-0 and spin-2 narrow resonances is simulated at leading order (LO). The production of resonances is followed by the decay of the resonances in the double Higgs boson system. All Higgs bosons are assumed to be SM Higgs bosons with an invariant



mass of 125 GeV. Samples are generated for two spin hypotheses and 16 mass values covering the range of heavy resonance masses from 250 to 1000 GeV. Two types of signal samples are present: resonance decaying to  $2b2l2\nu$  final state through the  $X \rightarrow HH \rightarrow bbZZ$  decays and also through the  $X \rightarrow HH \rightarrow bbWW$  decays. In both samples, the first Higgs boson decays to a pair of  $b$  quarks. However, in the first sample, the other Higgs boson decays to  $ZZ$  pair, while in the second sample the other Higgs boson decays to  $WW$  pair. Only  $Z$  boson decays in a dielectron, a dimuon, or a two neutrino state are selected. For  $W$  bosons, the chosen signature is characterized by a  $W$  boson decay to an electron and an anti-electron neutrino or a muon accompanied by an anti-muon neutrino. Intermediate decays to tau leptons with the subsequent decays to electrons and muons are also included.

To compare the expected numbers of events in the simulation to the number of observed events in the data for a given integrated luminosity, the signal production cross section has been normalized to 2 pb. This is a typical theoretical value of the production cross section of the WED particle in the 250-300 GeV range, the range to which the current physics analyses are very sensitive with the available LHC data. Additionally, the computed event rates take into account the branching fractions of the corresponding di-Higgs decay chains to the final state: 0.0012 and 0.0266 for  $HH \rightarrow bbZZ \rightarrow bbl\ell\nu\nu$  and  $HH \rightarrow bbWW \rightarrow bbl\nu\ell\nu$ , respectively [?].

### 1.4.2 Background processes simulation

In this analysis the main background processes are top-antitop production ( $t\bar{t}$ ) and Drell-Yan production in association with jets. Other background processes that contribute to a lesser degree include single top production, diboson production, and the production of a single Higgs boson in association with a  $Z$  boson (“ZH production”),

see Table ???. Other background processes are fully rejected in the event selection and thus are neglected.

**Table 1.2:** Background Monte Carlo samples. First column is the name of the process, the second column is the name of the generator used to simulate the corresponding process.

DY plus 1 Jet	MADGRAPH5_aMC@NLO-PYTHIA
DY plus 2 Jets	MADGRAPH5_aMC@NLO-PYTHIA
DY plus 3 Jets	MADGRAPH5_aMC@NLO-PYTHIA
DY plus 4 Jets	MADGRAPH5_aMC@NLO-PYTHIA
WW	PYTHIA
WZ	PYTHIA
ZZ	PYTHIA
ZH with $H \rightarrow b\bar{b}$ and $Z \rightarrow \ell\ell$	MADGRAPH5_aMC@NLO
$t\bar{t}$	POWHEG-PYTHIA
top quark tW channel	POWHEG-PYTHIA
$\bar{t}$ quark tW channel	POWHEG-PYTHIA
top quark t-channel	POWHEG-PYTHIA
$\bar{t}$ t-channel	POWHEG-PYTHIA
top quark s-channel	MADGRAPH5_aMC@NLO-PYTHIA

The Drell-Yan (DY) process in association with one to four jets is generated at leading order (LO) using MADGRAPH5\_aMC@NLO with the MLM matching scheme [?]. To account for the higher order QCD and electroweak effects in V+jets production (following [?]), DY events are further reweighted according to the dilepton transverse momentum, see ??.

The simulations of the background processes associated with top quark production are generated at next-to-leading order (NLO). The POWHEG [?,?,?,?] generator was used to generate the samples for top quark pair production and single top quark production in the tW and t channels. For the single top s channel production, the MADGRAPH5\_aMC@NLO generator was used. Single top backgrounds have been rescaled to the theoretical values of the next-to-next-to-leading order (NNLO) cross sections [?,?].

PYTHIA 8.212 [?, ?] was used to generate diboson samples at LO. Diboson background yields are normalized to NLO cross sections [?, ?, ?]. The dominant SM Higgs background process, the SM production of a single Higgs boson in the association with a Z boson (ZH), is simulated at NLO using the MADGRAPH5\_aMC@NLO generator with FxFx merging [?]. The SM Higgs background from the ZH process is scaled to NNLO with the MCFM program [?]. All the final cross section values at the NNLO accuracy in perturbative QCD have been computed with the original generators and are found to be in agreement with the values from the LHC Higgs cross section working group [?, ?, ?, ?, ?].

Normalizations for  $t\bar{t}$  and DY background processes are determined from data, as explained later in this chapter.

The NNPDF3.0 [?] Parton Distribution Functions (PDF) set is used for all the LO and NLO samples. POWHEG and MADGRAPH5\_aMC@NLO are interfaced with PYTHIA8.212 for the parton showering and hadronization stages. The description of the underlying event is done using the tune CUETP8M1 derived in [?]. For the simulation of the CMS detector response, GEANT4 [?] was used.

For all MC simulations, further reweighting of events is done using the SFs derived to account for the discrepancies between the data and simulation. This includes SFs related to the lepton efficiencies as well as b tagging efficiencies, see Sections ??, ??.

As discussed in Section ??, multiple overlapping proton-proton interactions occurred in each bunch crossing during data collecting in 2016, with an average of 24 hard scattering vertices per event. To account for this fact, all simulated samples include additional interactions to reproduce the real pileup distribution measured in data.

## 1.5 Physics Objects Selection

During the reconstruction of the collisions at the CMS, data for each event are refined into high-level physics objects that correspond to particles created as a result of a proton-proton interaction. Among the physics objects necessary for this measurement are reconstructed electrons, muons, jets originating from quarks of heavy flavor (b jets), and the missing transverse momentum. The reconstruction details have been given in Section ???. Here we focus on the analysis-specific selection of each of these physics objects.

### 1.5.1 Electrons

Electrons are reconstructed using the GSF algorithm [?]. The electron candidates are then selected by first applying a loose isolation requirement of 0.4, after which they have to pass the Tight ID criterion defined by the CMS  $e/\gamma$  POG, see ??. At the offline level, the analysis applies additionally the loose WP (WP90) [?]. With this WP, one achieves an electron selection efficiency of 90% by utilizing variables such as the agreement between the position of the ECAL cluster and GSF track that forms the electron, the energy of the 3 by 3 crystal core of the electron's cluster, the ratio of electron energy measured in ECAL to the electron's momentum measured in the tracking system, the ratio of the HCAL to ECAL energy,  $p_T$  and  $\eta$  of the electron, etc.

This is a justified level of the desired identification efficiency for the  $Z \rightarrow \ell\ell$  channel since we select on-shell Z boson decays to charged leptons, where produced prompt leptons are very energetic. Finally, the electrons must be isolated from other particles in the event. Numerically, a selection criterion on the isolation is imposed (see Section ??), where the expected contribution of particles from pileup interactions

is  $\rho$ -subtracted using the effective areas method, see Section ?? . In this measurement, the isolation (with the cone size 0.3) for electrons is required to be smaller than 0.06, which means that all other particles together in the isolation cone around the electron cannot have more than 6% of the electron's energy.

### 1.5.2 Muons

Muon candidates are reconstructed using tracker muon and global muon tracks identified by the PF algorithm. The selection procedure is similar to the one of electrons: first, a loose relative isolation selection of 0.4 is applied, then muons have to pass the Tight WP of the muon ID selection. At the offline level, a more stringent set of quality requirements (WP Loose) recommended by the CMS Muon POG is applied to the muon object. The ID definition contains the variables such as muon track segments compatibility, the fraction of valid hits in the inner tracker,  $\chi^2$  variables of the inner tracker and global tracks, position matching variables, energy deposits in the calorimeters, impact parameter variables, etc. Lastly, a relative  $\Delta\beta$ -subtracted PF isolation selection of 0.15 is applied, with the cone size 0.4.

### 1.5.3 Jets

Jets are reconstructed using the anti- $k_T$  jet clustering algorithm. This algorithm clusters PF candidates in a cone with a radius of 0.4. The energy and the resolution of the produced jets (so-called AK4 jets) are further corrected using JEC and JER corrections. These are used to calibrate the energy of the jets and to smear the resolution of jets to match the resolution of jets in the data.

To reject misreconstructed (or misidentified) jets due to detector noise, pileup, etc., a loose jet identification WP is applied following the recommendations from the

CMS JetMET POG. The ID variables include the jet  $p_T$  and  $\eta$ , HCAL and ECAL energy fractions due to charged and neutral hadrons, the jet multiplicity, etc. If jets are found to be overlapping with charged leptons used in the measurement, these jets are not considered by the analysis. We consider jets with a  $p_T$  greater than 30 GeV, which are in the range of  $|\eta| < 2.4$ . At least two jets must be present in the event.

#### 1.5.4 b jets

Each identified jet is further assigned a probability to originate from the b quark using the CMVA b tagging discriminator. This tagger produces as output a continuous discriminator with a value between -1 and 1, that is used to define three WPs depending on the chosen discriminant value. CMS BTag POG provides the CMS with the WP Loose, Medium, and Tight. For our measurement, the WP Medium requirement ( $>0.4432$ ) has led to the best expected (from the simulation) results (the lowest upper limits as will be explained later). Therefore, the WP Medium is chosen in this analysis for the derivation of the final results. Jets passing the WP Medium requirement are classified as b jets. The threshold is chosen such that the misidentification rate for light-quark and gluon jets is about 1% and the b jet tagging efficiency for this working point is about 66%. Two b jets with the largest scores of the CMVA discriminator are used to reconstruct the  $H \rightarrow b\bar{b}$  candidate, which will be explained at length later in this chapter.

#### 1.5.5 Missing transverse momentum

In this measurement two neutrinos are present in the final state. They come from the off-shell Z boson decays to neutrinos and also from leptonic decays of W bosons. Even though the two neutrinos present in the signal events cannot be identified directly

by the CMS detector, their presence can be inferred from the combined transverse momentum vector, using the momentum conservation requirement in the transverse plane for each event. The missing transverse momentum  $\cancel{p}_T$  is computed as the negative vector sum of the transverse momenta of all visible PF objects and is further JEC and JER corrected, see Section ???. The missing transverse energy  $\cancel{E}_T$  is the magnitude of the  $\cancel{p}_T$  vector.

All corrections recommended by the CMS JetMET POG are used in this measurement [?]. Additionally, a set of filters related to the instrumental effects is employed, such as the removal of the misreconstructed signals in the HCAL, noise in the tracker, etc. [?].

## 1.6 Event Selection


The candidate events are selected in the data sample if they contain the following physics objects: 2 b jets, 2 charged leptons, and missing transverse energy. All these objects pass specific kinematic requirements defined in ??, ??, and ??. Then we reconstruct intermediate particles of the decay chain by combining the observed physics objects.  $Z \rightarrow ll$  and  $Z^* \rightarrow \nu\nu$  are reconstructed first. Then  $H \rightarrow ZZ^*$  and  $H \rightarrow b\bar{b}$  are reconstructed. Finally, two Higgs boson candidates are used to form the  $HH$  system that could originate from the narrow resonance we are searching for.

In this chapter, we discuss how  $HH$  candidates are formed, how the selection criteria are applied to reduce contamination from background processes in the signal-enriched region (discussed later), how kinematic quantities used to compute the final results are defined, and how well simulation represents the collected data.

### 1.6.1 Kinematic selection of physics objects

We have described the kinematic selection of leptons and jets (and b jets) in the previous chapter. Because there is another  $bbZZ$  measurement in the CMS Higgs group, our analysis shares some phase space with this other measurement. Therefore, the selected data samples may contain a certain overlap. The other  $bbZZ$  team studies the  $2b2l2q$  final state, where  $q$  stands for any quark initiating the jet. To ensure that both measurements produce statistically independent (“orthogonal”) results, a selection on the MET is introduced. Since the final results - the production cross sections multiplied by the BFs of the final state - depend on the mass point, the MET selection varies with mass. While the derivation of the final results (called “limits”) will be explained later in this chapter, at this point it is worthwhile to point out that both analyses computed limits for all resonance mass hypotheses applying different MET selection criteria. Using a significance-like figure of merit ( $S/(S + B)$ ), see Fig. ??, we derived the set of preliminary MET requirements. The final set of MET requirements that yielded the best simplified combined limits, when the results of both measurements are merged together, is shown in Table ??.





**Figure 1.5:** Optimisation of the MET selection for two  $bbZZ$  analyses. S and B stand for signal and background process yields respectively. A significance-like figure of merit ( $\sqrt{S}/B$ ) as a function of the MET cut is used to identify the best selection for each given mass hypothesis. The green curve shows the significance for this measurement, where events are kept above a given threshold. The red curve shows the significance for the other  $bbZZ$  measurement, where the MET requirement is inverted (events are kept below that threshold). Top: 300 GeV resonance mass. Middle: 600 GeV resonance mass. Bottom: 900 GeV resonance mass. The plots on the left are for the di-electron channel and the plots on the right are for the di-muon channel. The dashed blue lines indicate the values of the MET selection, see Table ??.

**Table 1.3:** The MET requirements as a function of the mass of the  $HH$  candidate. Selection values (the second column) are provided for different mass hypotheses (the first column) of the narrow resonance decaying to the  $HH$  system.

Signal mass, GeV	$\cancel{E}_T$ selection, GeV
260-300	$> 40$
350-600	$> 75$
650-1000	$> 100$

### 1.6.2 Signal candidate construction and selection

Two leptons of the opposite signs and the same flavor with an invariant mass greater than 76 GeV are selected as Z candidates. This mass requirement helps to reject dilepton candidates not corresponding to real Z bosons. Events with a dilepton mass higher than 76 GeV will be used for SR and  $t\bar{t}$  CR. Additional Z mass selection will be discussed later in this chapter. These on-shell Z bosons are assumed to come from the  $H \rightarrow ZZ^*$  decays. The other Z boson, the off-shell Z boson, decays to two neutrinos in our signature and is represented by MET. The Lorentz four-vectors of the dilepton candidate and MET are added and the resulting four-vector represents the first Higgs boson candidate.

The other Higgs boson is formed from the pair of b jets with the highest output value of the CMVA algorithm. No requirement is applied on the existence or absence of other b jets in an event because they may be present due to a b jet misidentification or b jets resulting from pileup interactions or the underlying event. The invariant mass of the  $H \rightarrow b\bar{b}$  candidate is required to be greater than 20 GeV to remove contamination from background events with low mass resonances that decay to two jets, such as events with J/psi, Upsilon or low energy QCD interactions. There is no upper requirement on the  $H \rightarrow b\bar{b}$  invariant mass. Together with the first Higgs boson, the constructed di-Higgs system approximates the double Higgs boson

production that this measurement studies.

The final double Higgs boson candidate ( $HH$  candidate) comprises the  $Z(\ell\ell)$  candidate, the MET representing the  $Z(\nu\nu)$  decay, and the  $H \rightarrow b\bar{b}$  candidate. The four-momentum of this  $HH$  candidate is defined as the sum of the four-momenta of the two leptons and two  $b$  jets in the candidate as well as the four-momentum of the MET, defined in Sec. ??.

$HH$  system decays to a pair of  $b$  quarks and a pair of  $W$  bosons can also result in the same final state. The expected yields for the  $bbWW$  channel with respect to the  $bbZZ$  yields are comparatively small (1 to 4) because of the stringent kinematic selection on the dilepton invariant mass. However, the contribution from the  $HH$  system decaying through the  $bbWW$  intermediate state is still considered to be part of our signal in this measurement. The minimum requirement on the dilepton mass is necessary to ensure that our measurement is orthogonal to the known  $HH$  search from CMS in the  $bbVV$  channel that focused on  $bbWW$  decays [?], where only events with the dilepton mass below 76 GeV were studied.

In collider physics, one of several common definitions of the transverse mass is given by:  $M_T = \sqrt{(E^2 - p_z^2)}$ . This quantity is widely used in CMS searches for new particles. We proceed in the same fashion constructing our final variable -  $\tilde{M}_T(HH)$ , using the di-Higgs four-vector defined in the previous paragraphs.

While up until this moment we had the same requirements for the search of the radion/graviton  $\rightarrow HH$  for any heavy resonance mass, it is worthwhile noting that at this point the analysis becomes mass point specific. This is because the selection on the BDT discriminant (which will be explained later) and the MET requirement differ with the mass hypothesis. When we compute limits for different radion or graviton hypotheses we apply different selection criteria.


Finally, the reader should be informed that if an event does not have enough

physics objects that are suitable (as discussed in the objects selection section) for building the  $Z(\ell\ell)$  or  $H \rightarrow b\bar{b}$  candidates, or if the event does not pass candidate selection requirements on those objects or the full  $HH$  system, the event is discarded.

### 1.6.3 Signal and control kinematic regions

Three regions are defined: signal-enriched or signal region, and two CRs. The signal region is chosen such that the expected signal fraction is the largest there. Two control regions are defined such that they contain predominantly the background events with almost no signal. We define two CRs, one for each of the two main background sources: the CR for  $t\bar{t}$  (CRTT) and the CR for Drell-Yan in association with jets (CRDY). The CRDY contains a high fraction of DY events, and the CRTT contains nothing but  $t\bar{t}$  (more than 95%). This ensures the correctness of the DY and  $t\bar{t}$  normalizations extracted during the fit procedure (explained later).

Two intermediate particles in the decay chain are almost fully reconstructed,  $H \rightarrow b\bar{b}$  and  $Z(\ell\ell)$ . The signal region is defined in the kinematic space of  $H \rightarrow b\bar{b}$  and  $Z(\ell\ell)$  events. This corresponds to an area determined by the mass of the Higgs boson (125 GeV) and the mass of the Z boson (91.2 GeV). To account for detector smearing effects, the mass windows near the pole masses are defined such that we select events within the  $H \rightarrow b\bar{b}$  mass range from 90 to 150 GeV and Z mass range from 76 to 106 GeV, see Fig. ???. In the CMS, these relatively standard mass windows are chosen to take into account the detector resolution effects on a two b jet system and the dilepton mass (and the natural width for a Z boson), containing approximately 95% of true Higgs and Z boson candidates. The proportion of signal events is further increased with respect to background events by applying an additional requirement, the MVA requirement, which will be explained in the next section.



**Figure 1.6:** Signal region, control region  $t\bar{t}$ , and control region Drell-Yan in the phase space of  $Z \rightarrow \ell\ell$  and  $H \rightarrow b\bar{b}$  masses.

CRs are defined by inverting the SR selection. Drell-Yan plus jets (or “DY” later for brevity) background process contains a real Z boson, while the two jets are random and have random invariant mass. We obtain the CRDY by inverting the  $H \rightarrow b\bar{b}$  selection. For CRTT,  $t\bar{t}$  events always have two b quarks whose invariant mass has a broad distribution, while they do not have a real Z boson. Thus, to select CRTT, we invert the  $Z(\ell\ell)$  selection and use only the events in the upper range of the Z mass, see Fig. ??.

CRs are used to fit the event yield of the simulated background processes to the event yield in data and, thus, to determine the relative normalizations of  $t\bar{t}$  and Drell-Yan background processes, which is done using CRTT and CRDY correspondingly.

The efficiency of the candidate selection, up to this point, is summarised in Table ??, where numbers are provided for the SR. The signal contributions are split into two components:  $bbZZ$  and  $bbWW$ , and are shown separately. The corresponding total number of events passing the selection are also given, shown in the fourth column.

**Table 1.4:** Number of events surviving the candidate selection and kinematic requirements. Numbers are given for  $bbZZ$  and  $bbWW$  contributions in the SR. Efficiency values are also provided (the third column) and are normalised to the initial event counts before any selection is applied. A di-muon channel is presented. Numbers for the di-electron channel have the same trend but lower values since in the CMS, efficiencies for electrons are lower than for muons.

Process	Mass, GeV	Efficiency, %	Number of events
bbWW	300	0.2	85
bbZZ	300	10.4	4511
bbWW	900	0.1	23
bbZZ	900	15.1	12963

### 1.6.4 Signal region candidate selection with a multivariate technique

After the SR and CRs have been defined, in the final step of the event selection we require the events in the SR to pass the threshold value of the BDT discriminant. An MVA discriminant that uses a boosted decision trees algorithm is trained considering a number of kinematic quantities of a candidate to be above a certain threshold. As this selection step is complex, it is described in a dedicated chapter (see ??).

### 1.6.5 Event reweighing in simulation

Several factors lead to a difference in the behavior of simulation in comparison to real data. These factors include the difference in the kinematics of simulated events from what it is in nature as well as the differences in the simulation of the detector response that affects reconstruction efficiencies for physics objects that form candidates, as well as tagging and mistagging rates for b jets.

These factors are quantified through per-event SFs that are used as event weights in preparation of all kinematic distributions and computation of the expected event counts. The SFs of all types are produced centrally in CMS by the appropriate

POGs. The full per-event weight for a simulated event is a product of all individual SF discussed below.

- **Lepton efficiency.**

Scale factors that reflect the differences between data and simulation in lepton reconstruction, selection, and triggering, are computed as the ratios of the corresponding efficiencies between data and simulation. Additionally, since the DZ requirement was absent for some triggers, the efficiency of the DZ selection needs to be measured.

The efficiencies and, subsequently, the scale factors are measured separately for the efficiencies of the reconstruction procedure, lepton identification, isolation requirements, and impact parameter requirements. The TnP procedure discussed in Section ?? is applied to measure all of these efficiencies. The scale factors of all listed types for both leptons are used as multiplicative factors to form the per-candidate lepton efficiency SF.

On average, the muon ID scale factors are equal to 0.992 - 1.004, depending on the  $p_t$  and  $\eta$  of leptons. The muon ISO SFs are equal to 0.991-1.004. The electron ID · ISO SFs, computed together, are ranging from 0.832 to 1.073. For the HLT SFs, see Figs. in the subsection ??.

- **b tagging and light flavor mistagging probability scale factors.**

To identify the b jets, the selection on the output of the CMVA tagger is applied. The efficiency of the b jet selection is measured (with the TnP method) in data and in the heavy-flavor enhanced MC sample, the  $t\bar{t}$  process containing top and anti-top quark decays. Since these efficiencies are different, the b tagging SFs are derived. Because this measurement contains two b jets in the final state,

a total per event  $b$  tagging SF is equal to 0.7 - 1.3 for signal and 0.6 - 1.4 for background processes.

- **Kinematic reweighing of the DY plus jets samples.**

DY plus jets samples used in this measurement are produced at LO (see Section ??). The samples are corrected to exhibit NLO kinematic behaviour in the EW interaction by assigning a weight to each event. As the biggest discrepancy between data and simulation for such samples is typically found in the di-jet invariant mass distribution, an LO-to-NLO correction as a function of the  $\Delta\eta_{b\ jets}$  separation is computed following the procedure described in [?]. As stated there, this correction improves the data-MC agreement in many distributions derived from the di-jet objects or based on the  $p_T^{Z(\ell\ell)}$  variable, while has a negligible effect on the remaining kinematic variables. The correction means that a kinematic weight is assigned to each event.

It was found by several Higgs analysis teams that the previous correction does not address the data-MC discrepancies for highly energized Z bosons. The spectrum of the  $p_T$  of the Z boson distribution is harder for DY plus jets MC simulation than for data. This is due to the fact that the simulation does not include higher-order QCD+EW corrections [?]. The DY plus jets simulated samples are reweighed as a function of the  $p_T^{Z(\ell\ell)}$  to update the LO samples to the NLO in QCD+EW. The corrections can be as large as 10% for energetic Z bosons with  $p_T > 400$  GeV.



## 1.7 Multivariate selection in the signal region

To reduce the background contamination in the SR, we employ the standard practice of CMS physics analyses of utilizing the MVA discriminant. In this section, we will discuss which discriminating variables were used, how the MVA discriminant was constructed, and what are the efficiencies of the BDT selection for all major signal and background processes.

### 1.7.1 Kinematic variables of a candidate

A number of kinematic quantities can be constructed out of four-vectors that comprise candidates of our final state. The distributions of these quantities, in general, differ for candidates originating from different background processes as opposed to for candidates from the hypothetical signal.

In this section, we discuss a set of kinematic quantities that are the most discriminating between the expected signal candidates and background candidates. This set of variables is used as an input for computing the multivariate discriminant that is later discussed in this section. In total, nine variables are chosen for this purpose. About 30 - 40 other kinematic variables were considered at the early stages of this data analysis. However, they were discarded as it was found that they do not improve the results of this measurement in a significant way, while at the same time they considerably increase the complexity of the measurement.

Since the MVA selection requirement will be applied on a quantity that was computed based on these variables, the simulation has to reproduce the main variables well. We will define each of the nine variables below. For SR and both CRs, we will show the comparison of distributions from data and simulation for each variable. Since the number of plots would be too large to show for two channels and two spin

hypotheses, not to mention for all masses, we will include in this dissertation only di-muon channel plots for the 300 GeV graviton mass hypothesis. However, all other relevant figures of the nine main variables, including the di-electron channel and other mass points, were examined and discussed with CMS collaborators during the review of the measurement in CMS. They show similar behavior and agreement as those included here.

As we outlined before, even for simple physics objects the distributions from MC simulations and collected data have discrepancies. For more complex objects and candidates, these discrepancies become even larger. In general, MC simulations are not supposed to reproduce very narrow regions of the kinematic phase space well, or align with the data perfectly at the high order of the QCD. Therefore, one normally adjusts the event yields (rates) of background processes using the normalization values determined in the fit to data in the control regions. The distributions as they are, without the extra normalizations from the fit being applied to the event yields of MC simulated background processes, are called pre-fit distributions. After the normalizations from the fit are applied to the rates of background processes, post-fit distributions are obtained.

The list below summarises the set of variables used as input to the MVA discriminant:

- **$\Delta R$  separation between two b jets.** The  $\Delta R$  (defined in Section ??) separation between the b jets ( $\Delta R_{b \text{ jets}}$ ) gradually decreases as we start considering larger graviton or radion masses. This fact is explained by the Lorentz boost of the  $H \rightarrow b\bar{b}$  system. When the heavy resonance produces two SM Higgs bosons, they are highly boosted. For the Higgs boson, which will further decay to a pair of b quarks ( $H \rightarrow b\bar{b}$ ), the b quarks will come out of the decay almost collinear


to each other with a minimal value of  $\Delta R_{b \text{ jets}}$ . For background processes, this separation does not depend on the graviton or radion mass hypothesis.

- **$\Delta R$  separation between two leptons.** The distance between two leptons in the  $\eta - \varphi$  space ( $\Delta R_{leptons}$ ) tends to be smaller for the dilepton system coming from the real Z bosons originated from the  $H \rightarrow ZZ$  decays, in comparison to the  $\Delta R_{leptons}$  distance between two random OSSF leptons produced by background processes.
- **Missing transverse momentum.**  $\cancel{p}_T$  distribution coming from the off-shell Z bosons is constrained by the invariant mass of the parent Higgs boson and has, thus, a more narrow distribution. Conversely, the MET coming from  $t\bar{t}$  background process can produce smaller and larger values of the MET outside of constraintment.
- **Invariant mass of the  $H \rightarrow b\bar{b}$  candidate.**  $H \rightarrow b\bar{b}$  mass for two b jets coming from a real Higgs boson tends to be close to the SM Higgs boson invariant mass smeared by the b jet energy-momentum resolution, while the background candidates from unrelated b jets can have smaller and larger masses, especially if they come from the top quarks of the  $t\bar{t}$  background.
- **Transverse momentum of the  $H \rightarrow b\bar{b}$  candidate.** The transverse momentum of the  $H \rightarrow b\bar{b}$  candidate ( $p_T^{Hbb}$ ) tends to have a relatively narrow and peaky distribution for candidates coming from the hypothetical signal in comparison to the broad distributions produced by background processes.
- **Invariant mass of the  $ZZ^*$  system.**  $ZZ^*$  mass of two Z bosons coming from the Higgs boson decay ( $H \rightarrow ZZ$ ) tends to be close to the SM Higgs boson mass smeared by the  $\cancel{p}_T$  resolution, while the background candidates from DY


production in association with jets can have random masses. These can be particularly large if they come from the DY production in association with several jets in addition to some instrumental MET present in the event.

- **Transverse momentum of the  $ZZ^*$  system.** The transverse momentum of the  $H \rightarrow ZZ$  candidate ( $p_T^{H_{zz}}$ ) tends to have a relatively narrow and peaky distribution for candidates coming from the hypothetical signal in comparison to broad distributions produced by the background processes.
- **Invariant mass of the Z boson candidate.** Z bosons coming from the Higgs boson have the invariant mass near the pole mass, while Z bosons from DY plus jets production can have a broader peak.
- **Transverse momentum of the Z boson candidate.** The transverse momentum of the Z candidate ( $p_T^Z$ ) tends to have a relatively narrow distribution for candidates coming from the real Z boson produced in the decays of the hypothetical signal in comparison to broad distributions of the  $p_T^{\ell\ell}$ , when dilepton systems are produced by background processes.


At this stage of the analysis description, we would like to refer the reader to the pre-fit plots (figures in the left columns). The figures are shown in the following order: all nine distributions in CRDY (see Figs. ??, ??), in SR (see Figs. ??, ??), and in CRTT (see Figs. ??, ??). The simulated distributions are found in an acceptable agreement with those from the data and are judged to be adequate for use in a multivariate discriminant construction discussed next.




**Figure 1.7:** Comparison of data and MC samples. 300 GeV, CRDY region, mm channel. Prefit plot on the left; Full Postfit plot on the right.



**Figure 1.8:** Comparison of data and MC samples. 300 GeV, CRDY region, mm channel. Profits plot on the left; Full Postfit plot on the right.




**Figure 1.9:** Comparison of data and MC samples. 300 GeV, SR region, mm channel. Profit plot on the left; Full Postfit plot on the right.




**Figure 1.10:** Comparison of data and MC samples. 300 GeV, SR region, mm channel. Profit plot on the left; Full Postfit plot on the right.





**Figure 1.11:** Comparison of data and MC samples. 300 GeV, CRTT region, mm channel. Profit plot on the left; Full Postfit plot on the right.



**Figure 1.12:** Comparison of data and MC samples. 300 GeV, CRTT region, mm channel. Profit plot on the left; Full Postfit plot on the right.

### 1.7.2 Multivariate discriminant: a BDT classifier

In many areas of HEP, such as object identification, a reduction of the background contribution in the signal region, etc., there are needs to discriminate a “signal” from one or several “backgrounds.” The discriminating algorithms can use a single physical variable for signal-background separation, or several variables. The way an algorithm uses the input variables can be of a sequential nature, or through some non-linear dependence. The Higgs boson discovery, in part, was possible due to an employed BDT approach, where several variables are combined into an artificial BDT score or output. The BDT output, as a higher-level construct, has no direct physical interpretation. However, due to the shown gain in signal-background separation performance, the BDT has been successfully used for years for the final classification decision.

Multivariate discriminants (or models or classifiers) based on the BDT or other algorithms are constructed using machine learning (ML) techniques. Now, at almost every stage of the HEP data analysis, starting from the track reconstruction, through the object identification and the signal region purification, and up to the construction of the final discriminating MVA variable, physicists rely on the ML methods.

In simple terms, the ML procedure consists of the training and testing stages, and sometimes a validation stage is added. First, the data is split into two non-overlapping sets: train and test data. Then the classifier is trained when the model is learning the patterns in the train data. After that, the performance of the model is tested using independent test data.

After the training procedure, the ML algorithm produces a discriminator, which is a function that is related to the likelihood of an event to belong to a signal or to a background category. Here and in other parts of the thesis, we are describing the BDT procedure, which is a non-parametric supervised learning method used for

classification; other MVA methods that are not related to this measurement, are well-explained in [?].

Every analyst prefers a classifier with low bias and low variance. A bias is the difference between the average prediction of our model and the correct value, the latter of which we are trying to predict. A classifier with a high bias oversimplifies the model and does not take into account all the nuances of a given train dataset. Therefore, analysts prefer low bias models, which are flexible and perform well both on the train and test data, and always provide a stable separation between the signal and background hypotheses. A variance is the variability of model prediction for a given data point and is related to the spread of our data. A model with high variance learns almost every single pattern in the train data and, therefore, will not generalize well when given new unseen data. As a result, models with a high variance achieve the best scores using the train data but perform poorly on the test data. Thus, low variance models are preferable, which are robust when applying the trained classifiers on a dataset that is statistically independent from the one used for training.

One of the most popular ML algorithms in HEP is based on the boosted decision trees (BDT). A single decision tree is a sequence of simple binary splits on each of the input variables. The tree, thus, can be depicted as a multidimensional set of selections on input space. The tree assigns a score of +1 or -1 if a given event is signal or background-like. The key elements in each step of the BDT splits are the variables used for the selection and the positions of the selections. Both are determined using an impurity criterion  $I(p_n)$ , which is a function of the signal purity  $p_n$  in a tree node  $n$ . The signal purity is defined as a ratio of the signal contribution to the total number of signal and background events,  $p_n = S/(S + B)$ . The impurity criterion is chosen such that it is at the minimum when  $p_n=0$  or  $p_n=1$ . These values would give perfect signal-background discrimination. The criterion is at the maximum, when

$p_n=0.5$ , which means no discrimination is achieved. Each split maximises the gain function  $G = I(p_n) - f_1 I(p_{c1}) - f_2 I(p_{c2})$ . In this formulation of  $G$ , the labels  $c1$  and  $c2$  refer to child nodes of the parent node, and  $f1$  and  $f2$  are the fractions of events of the parent node that are found in child nodes  $c1$  and  $c2$ , respectively. As a definition of the impurity criterion for this measurement, the Gini coefficient is used:  $I_G(p_n) = 2p_n \cdot (1 - p_n)$ .

The tree keeps applying the selection on the input variables until either a maximum depth is reached (a number of consecutive splits), or a minimum number of events in the final child nodes are attained. The problem with such a tree growing procedure is that it is prone to overtraining, which means the model will not generalize well and have a high variance.


One way to deal with BDT overtraining is boosting. The idea behind boosting is that signal events from the training sample that end up in a background node or vice versa are given intentionally large weights (a penalty). These weights are normally significantly larger than the weights for events with the correct leave node prediction. Such a penalty procedure re-weights the train data and then new decision trees are created. The boosting is applied hundreds of times, which produces a set of new reweighed trees (a forest). Intuitively, boosting is a method in which a large number of shallow trees, which have only a few splits, are combined by taking a weighted average of their output scores. The discriminating power of a single shallow tree is usually poor, but such a tree is less prone to overtraining, therefore when combined, the ensemble of these trees produces a model with high and stable performance.

For the boosting algorithm, we chose the gradient boosting method, as is implemented in the popular MVA package for HEP - TMVA [?]. Gradient boosting can be thought of as a function expansion approach. In this case, each tree corresponds to a summand. In this approach, the parameters for each summand (tree) are determined


by minimizing an error function. In the TMVA implementation, the binomial log-likelihood loss function is used. The adopted procedure follows the greedy algorithm approach - only one tree is modified at a time, while the other trees remain invariable.

Several decision trees of the BDT procedure, which were employed in the analysis, are shown below. This training was done in the d-electron channel. In all figures produced by the TMVA, index 0 refers to  $H \rightarrow ZZ$  candidate and index 1 refers to  $H \rightarrow b\bar{b}$  candidate. For example, “hmass1” denotes the mass of the  $H \rightarrow ZZ$  candidate, “hpt0” denotes the  $p_T$  of the  $H \rightarrow b\bar{b}$  candidate, etc. For the Z boson candidate, the on-shell Z boson decaying to charged leptons,  $Z(\ell\ell)$ , has either an index 0 or the index is dropped. The closer the node color is to blue, the more the event is classified as signal-like. The same idea is illustrated with the red color for background events.

Di-electron and di-muon channels have been trained separately and the BDT metrics (discussed later in this chapter) show similar performance. To save space, in this chapter we show mostly figures related to the di-electron channel.




**Figure 1.13:** BDT trees 1 and 2 (indexing starts from 0). These trees relied on: the mass and the  $p_T$  of the  $Z(\ell\ell)$  candidate,  $\Delta R$  separation between two b jets, the mass of the  $H \rightarrow b\bar{b}$  candidate, and the  $p_T$  of the  $ZZ^*$  system. The TMVA notation is explained in ??.



**Figure 1.14:** BDT trees 400 and 401 (indexing starts from 0). These trees relied on: the  $p_T$  of the  $Z(\ell\ell)$  candidate,  $\Delta R$  separation between two b jets,  $\Delta R$  separation between two charged leptons, the  $p_T$  of the  $H \rightarrow b\bar{b}$  candidate, and the mass and the  $p_T$  of the  $ZZ^*$  system. The TMVA notation is explained in ??.





**Figure 1.15:** BDT trees 799 and 800 (indexing starts from 0). These trees relied on: the  $p_T$  of the  $Z(\ell\ell)$  candidate,  $\Delta R$  separation between two b jets,  $\Delta R$  separation between two charged leptons, the mass of the  $H \rightarrow b\bar{b}$  candidate, and the mass of the  $ZZ^*$  system. The TMVA notation is explained in ??.

The BDT discriminant is trained on a pure sample of signal and background events from simulation, using the train data. The properties of the discriminant are studied

on an independent testing sample of pure signal and background events, using the test data. The Deep Neural Network (DL) approach has also been studied - however, due to the lack of event statistics, the DL approach could not offer a performance higher than the one achieved with the BDT, and thus was abandoned. Training 16 BDTs per the spin hypothesis seemed impractical since our data analysis is dominated by statistical errors. Instead, we follow the approach of other  $HH$  analyses [?]: we split the range from 250 to 1000 into two regions: low mass (250 to 450 GeV) and high mass (above 450 GeV up to 1 TeV) regions. This separation boundary at 450 GeV was optimized first using ROC curves (described later in this section), then by running the whole analysis chain all the way to the final limits. This procedure not only saves CPU cycles by training two BDTs instead of sixteen, but mostly increases the statistics and improves the robustness of the trained BDT.

During the training procedure some variables provide more discriminating power and on average are used more often than others. The more often the variable is used, the higher its importance is. The tables below show the ranking and average importance of each of the BDT training. For the low and high mass regions, and di-electron and di-muon channels, see Tables ??, ??, ??, ??.

Rank	Variable	Importance, %
1	$\Delta R_{b \text{ jets}}$	13.9
2	MET	12.1
3	Mass of $H \rightarrow b\bar{b}$ candidate	11.9
4	$p_T^{ZZ^*}$	11
5	$\Delta R_{leptons}$	10.9
6	$p_T$ of $H \rightarrow b\bar{b}$ candidate	10.7
7	$p_T$ of Z boson candidate	10.2
8	Mass of $ZZ^*$ system	10.1
9	Mass of Z boson candidate	9.26

**Table 1.5:** Di-electron channel. Relative importance of the input variables in the low mass BDT training.

Rank	Variable	Importance, %
1	$\Delta R_{leptons}$	14.1
2	Mass of $H \rightarrow b\bar{b}$ candidate	13.7
3	$\Delta R_{b\ jets}$	13.2
4	$p_T$ of $H \rightarrow b\bar{b}$ candidate	12.1
5	$p_T$ of Z boson candidate	11.5
6	$p_T^{ZZ^*}$	11.3
7	MET	10.3
8	Mass of $ZZ^*$ system	7.7
9	Mass of Z boson candidate	6.1

**Table 1.6:** Di-electron channel. Relative importance of the input variables in the high mass BDT training.

Rank	Variable	Importance, %
1	Mass of $H \rightarrow b\bar{b}$ candidate	13.8
2	$\Delta R_{b\ jets}$	13.1
3	$\Delta R_{leptons}$	12.9
4	$p_T$ of $H \rightarrow b\bar{b}$ candidate	11.7
5	$p_T^{ZZ^*}$	11.3
6	$p_T$ of Z boson candidate	11.1
7	MET	11
8	Mass of $ZZ^*$ system	8.8
9	Mass of Z boson candidate	6.2


**Table 1.8:** Di-muon channel. Relative importance of the input variables in the high mass BDT training.

Rank	Variable	Importance, %
1	$\Delta R_{b\ jets}$	13
2	MET	12.2
3	Mass of $H \rightarrow b\bar{b}$ candidate	11.9
4	$p_T$ of $H \rightarrow b\bar{b}$ candidate	11.3
5	$p_T$ of Z boson candidate	11.1
6	$p_T^{ZZ^*}$	10.9
7	$\Delta R_{leptons}$	10.5
8	Mass of $ZZ^*$ system	9.7
9	Mass of Z boson candidate	9.5


**Table 1.7:** Di-muon channel. Relative importance of the input variables in the low mass BDT training.

The correlations among the input variables are shown in the Figs. ??, ?. For low mass training, signal samples for mass hypotheses from 250 to 450 GeV are combined together and represent the “signal”. This is done similarly for high mass training - it is a mix of signal samples for mass hypotheses from 500 to 1000 GeV. For background samples, the “background” is a mix of samples of two main background processes: DY in association with jets and  $t\bar{t}$  production, weighted by the cross section value.


Below we include figures of the BDT input variables. They are the same variables previously shown, but are produced by the TMVA package. In this style they allow for an easier visual judgment of which variables behave with greater difference between the signal and the background, see Figs. ??, ?.




**Figure 1.16:** Input variable correlations of the di-electron channel, low mass training. Top: signal sample mix. Bottom: background sample mix. The TMVA notation is explained in ??.



**Figure 1.17:** Input variable correlations of the di-electron channel, high mass training. Top: signal sample mix. Bottom: background sample mix. The TMVA notation is explained in ??.



**Figure 1.18:** Variables used in the low mass training for di-electron channel. Index '1' refers to  $H \rightarrow b\bar{b}$  decay and index '0' refers to  $H \rightarrow ZZ$  decay. The TMVA notation is explained in ??.



**Figure 1.19:** Variables used in the high mass training for di-electron channel. Index '1' refers to  $H \rightarrow b\bar{b}$  decay and index '0' refers to  $H \rightarrow ZZ$  decay. The TMVA notation is explained in ??.


After the training, the BDT model is created. This model produces the BDT distribution (BDT response), see Fig. ??. This figure overlays train and test parameters of the BDT training of the di-electron channel. The parameters of the model, such as the number of trees, the allowed number of splits per variable, the maximum tree



depth and others, have been thoroughly studied. The default TMVA parameters are found to provide a good performance with this measurement.

It is difficult to get high performance during low mass training, since this region corresponds to the range of  $HH$  masses where all the background processes are concentrated (Figs. ??, ??). The event yield of background in this region is very high and most variables have similar distributions for signal and background processes. The BDT performance is noticeably better than what can be achieved using a simple linear discriminant method (LD), see Figs. ??, ??.

Performance of the high mass training can be considered perfect, see Fig. ?. Most background processes concentrate in the low mass region; therefore, for the high mass region even the linear discriminant is performing well, see Figs. ??, ??.




**Figure 1.20:** BDT discriminants for di-electron channel. Top: low mass training. Bottom: high mass training.

In machine learning, performance measures are needed to compare several models. Usually for a classification problem, one relies on the Receiver Operating Characteristic curve (ROC) and its Area Under The Curve (AUC). These two are the most im-

portant evaluation metrics for checking the performance for any classification model. The ROC is a curve plotted in the space of signal efficiency and background rejection efficiency, where background rejection is equal to one minus the background efficiency. On the 0 to 1 ranges for axes (values of efficiencies), a diagonal line represents the efficiency of the random guess, which is 0.5. A perfect ROC would bend very close to the top right corner. Such ROC has the AUC almost equal to 1. The ROC curves for di-electron training in low and high mass regions are shown in Fig. ??.

ROC can be regarded as a probability curve and, in this case, the AUC represents the degree or measure of signal-background separability. It tells how much the model is capable of distinguishing between two given classes. The higher the AUC is, the better the model is at predicting signal as a signal-like event and background as a background-like event.




**Figure 1.21:** ROC curves for di-electron channel. Top: low mass training. Bottom: high mass training.

For two given signal efficiencies, one would prefer the model, which has a lower background efficiency; or in other words, a higher background rejection efficiency. In practice, when AUCs are close to 1 and ROCs are similar, one compares AUCs

directly and decides which model to use based on the AUC value.

Finally, we provide the figures of the BDT discriminant in the SR. The plots are shown for the 300 GeV radion decaying in the di-electron and di-muon channels. Both figures are of the post-fit type.



**Figure 1.22:** BDT distributions for the radion case, electron(muon) channel is shown at the top (bottom). Signal region is presented for the 300 GeV mass hypothesis. For electrons the selection is at 0.4, for muons at 0.7, as is described in Section ??.

**Table 1.9:** The BDT selection values used in this measurement.

Channel	260 and 270 GeV	300 and 350 GeV	400 and 450 GeV	600 to 1000 GeV
Di-muon	0.1	0.7	0.7	0.99
Di-electron	0.4	0.4	0.925	0.99

### 1.7.3 BDT selection requirement in the signal region

To remove the background contribution in the SR, we apply a BDT requirement to di-Higgs candidates. The requirement is specific to the mass hypothesis and specific to the lepton channel; however, the requirement does not depend on the resonance nature. A simple grid search method was adopted to find the best BDT selection for each mass point and channel. During the optimization procedure, we produced the final limits for each value of the considered BDT requirement. Then, we determine the value of the BDT requirement that corresponds to the best expected limit. The optimized values of the BDT selection that are used in this physics analysis are summarised in the Table ???. The BDT selection is not applied in the CRs. The BDT selection achieves about a 90 % efficiency for the signal while reducing the background contribution by a factor of few at the low mass region to more than 100 for the high mass region.

The corresponding efficiencies of the BDT selection are presented in Table ??. These values are derived for the graviton signal hypothesis and the main background processes for two channels separately.

## 1.8 Uncertainties of the measurement

The outcome of the statistical analysis (or StatAn, which is discussed in the next section) is the limit on the  $HH$  production cross section multiplied by the BF of the

**Table 1.10:** Efficiency of the BDT selection requirement in two mass regions and for two channels:  $ee$  channel (top) and  $\mu\mu$  channel (bottom).

sample	Efficiency at 300 GeV, [%]	Efficiency at 900 GeV, [%]
signal (bbZZ)	89.2	94.9
signal (bbWW)	75.0	88.4
$t\bar{t}$	28.8	0.2
Drell-Yan	74.2	1.2
Single top	33.1	1.1
ZH	88.8	10.7
Dibosons	90.0	5.0

sample	Efficiency at 300 GeV, [%]	Efficiency at 900 GeV, [%]
signal (bbZZ)	58.1	91.1
signal (bbWW)	25.9	96.3
$t\bar{t}$	13.6	0.2
Drell-Yan	39.0	0.8
Single top	13.0	0.2
ZH	56.0	8.4
Dibosons	51.4	6.2

final state (or just the limit). The StatAn uses the predicted event counts for different processes as well as the shape information of kinematic distributions for those events. The predicted event counts are computed by multiplying the number of events in simulated samples of different processes by the best-known cross sections for those processes, while also assigning event weights based on several other factors. However, the results of the analysis are highly affected by systematic and statistical uncertainties. We first discuss the types of uncertainties, how they affect the final limits, and also provide the size of this impact for dominant sources of the uncertainties. Then, we will discuss the fit procedure and the variable that was used to extract the limits.

Uncertainties that affect the sensitivity of the search come from a variety of sources: statistical uncertainties due to the limited size of the sample statistics, experimental uncertainties related to the accuracy of the detector response, the amount of collected data (uncertainty on the Luminosity), differences of simulated samples



from real data, and the theoretical uncertainties on cross sections or proton structure. Following the CMS practices, below we will divide all uncertainties into two broad classes: the “normalization” and the “shape” uncertainties. The former modifies only the yields (or “normalizations” for brevity) of selected events from different processes. The latter may also distort the shape of the distribution of the final kinematic variable that is used in the extraction of the limits.

To compute the effect of a particular source of the systematic uncertainty on the final limit, we use the up/down method. This is a procedure of varying a factor affecting the expected yields up or down by the amount of the estimated uncertainty on the effect and examining how the yields change.

### 1.8.1 Normalisation uncertainties

The sources of systematic uncertainties that affect normalization are discussed below. The sizes of certain types of uncertainties vary depending on the resonance mass hypothesis and the type of the channel (di-electron or di-muon). In this case, ranges of the uncertainty values are listed. Normalization uncertainties listed in this section modify all background processes but  $t\bar{t}$  and DY. The levels of those backgrounds are determined from data during the statistical analysis that uses two control regions dominated by these two background processes.

- **Luminosity.** The CMS estimated the uncertainty on the integrated luminosity of the 2016 data set to be 2.5%, see [?]. This uncertainty directly affects the expected event yields for both signal and background processes (excluding two dominant ones: DY and  $t\bar{t}$ ).
- **Pileup.** How well the PU interactions are replicated in the MC simulation affects the signal and background event yields. The effect of this uncertainty

on normalizations is approximately 6%. The effect is estimated by varying the number of pileup interactions in simulated samples up and down within the range, reflecting the limited knowledge of the total inelastic proton-proton interaction cross section at 13 TeV.

- Proton PDF.** The impossibility to know precisely the structure of the interacting protons is evaluated using an ensemble of PDF replicas from the NNPDF set [?] following the PDF4LHC prescription [?,?]. The root mean square value of the expected event yields in simulated samples, computed with the PDF replicas from this ensemble, is taken as a measure of the bias. The size of the effect is found to be of order 5%.
- Theoretical uncertainties of the QCD scales.** The bias associated with the theoretical uncertainties in the QCD factorization and renormalization scales is evaluated. This uncertainty affects the expected yield of the signal and background events (except the  $t\bar{t}$  and DY processes). This uncertainty is computed by independently varying the factorization and renormalization scales, used as parameters in the Pythia event generator. The variation is done in the MC simulation following a standard CMS prescription given by the PDF4LHC [?]. The scales in signal samples are varied by a factor of 2 up and down - multiplying the original nominal scale values by two corresponding factors: 0.5 and 2. The unphysical cases, when one of the scales fluctuates up, while the other fluctuates down, are not considered. In each bin of the final kinematic variable that is used in the StatAn, the maximum and minimum variations of the yields are used to build a bias region around the nominal shape. The size of the envelope indicates that these scales affect the yields of the processes at a level of 4-6%.

- **Theoretical values of the cross sections.** The uncertainty on the theoretical cross section value of the single top production is propagated to the background yields of the single top background processes and used to normalize their event yield. The uncertainty is found to be 5–7%.
- **Missing transverse momentum.** During the clustering of the jets and a subsequent application of JEC and JER corrections, neutral hadrons and photons that do not belong to any jet (“unclustered energy”) and jets with transverse momenta below 10 GeV - lack such corrections. This affects the MET reconstruction and results in a small systematic bias. The effect of JEC on the unclustered energy and subsequently on the magnitude of the  $\cancel{p}_T$  is studied. The effect is propagated to the final kinematic variable, which contains the  $\cancel{p}_T$ . The effect of this bias is studied by shifting the energy of each particle not contained in jets or contained in low- $p_{T,jets}$  by its uncertainty, which is estimated during particle’s reconstruction. These variations are found to affect the event yields of signal and background processes at the level of 3%; however, they do not show a visible effect on the shape of the final kinematic variable. This bias is categorized as a “normalization” systematic source.

### 1.8.2 Shape uncertainties

Certain types of systematic uncertainties (later referred to as “shape uncertainties”) distort the shapes of some kinematic distributions and BDT discriminant distributions. As a consequence, the shape of the final discriminating variable is modified. The parameters defining each source of the bias are varied within one standard deviation up and down. The effect is propagated through all variables included in the construction of the final kinematic variable used in the StatAn procedure. This

method prepares three shapes: in addition to the nominal shape of the final variable, it also produces two modified shapes corresponding to the up and down variations of the parameters. All these shapes are then used in the statistical analysis of the data, see ???. Below we discuss each source of the shape uncertainty individually.

- Lepton efficiency.** Data-MC simulation differences in the lepton reconstruction in the tracker, identification and isolation selection criteria, and in the efficiencies of the HLT trigger requirements are computed and used to correct the MC samples. The final statistical analysis relies on the MC correctly simulating the efficiency for an event to be reconstructed and to pass online and offline selection requirements. The lepton scale factors are derived from high-statistics samples of Z boson decays. This simulation is not perfect, thus, corrections are applied to candidates as per-candidate (or per-event) weights, as explained later in this chapter. These corrections, though, are also known with limited accuracy. The uncertainty of lepton efficiency corrections as a function of lepton  $p_T$  and  $\eta$  is propagated to the final kinematic distribution. The effect of these uncertainties is sub-percent for the muon channel and up to 6% for the electron channel.
- Jet energy scale.** The JEC directly affects the invariant mass and  $p_T$  of the  $H \rightarrow b\bar{b}$  candidates and the magnitude of the  $\cancel{p}_T$ . Both objects are used in the construction of the final kinematic variable; therefore the JEC effect is thoroughly studied. The JEC is varied within one standard deviation of its uncertainty as a function of jet  $p_T$  and  $\eta$ , and the effect on the b jet kinematics and on the  $\cancel{p}_T$  is calculated. This effect is propagated through the steps of the measurement yielding the variation of the final variable shape. The JEC uncertainty has an effect on the yields of the signal and background processes

at a level of 5 to 10%.

- Jet energy resolution.** The shapes of the final kinematic distributions are affected by the difference between the jet energy resolution in data and simulation, because this bias modifies the mass and the  $p_T$  of  $H \rightarrow b\bar{b}$  candidates and the magnitude of the  $\cancel{p}_T$ . The JER is varied in simulation by one standard deviation as a function of jet  $p_T$  and  $\eta$ , and the effect is propagated through the steps of the measurement. This bias affects the final distribution at the order of 0.5%.
- b tagging.** The data-mc difference in the efficiency to tag a  $b$ -jet (and the probability to misidentify a light flavor or a gluon jet as a  $b$  jet) is calculated and MC samples are corrected by the corresponding SFs, see Section ?? . They are derived using heavy-flavor enhanced jet MC samples. The uncertainties on these SFs are propagated through the measurement steps to the final variable. The effect of the  $b$ -tagging efficiency (flavor misidentification) is the highest for DY process, at about 5% (7–10%), and is at the sub-percent level for other processes (7–10%).
- Bin-by-bin uncertainties.** This is a statistical uncertainty. Out of hundreds of uncertainty sources in this measurement, the most dominant ones are related to the limited size of simulated samples. In certain cases, these statistical uncertainties can produce sizeable fluctuations of the bin content of the shape of the final variable. Because of this, we add an individual nuisance parameter for each bin of the final variable distribution. For each bin in the bins of the final variable, the corresponding bin-by-bin uncertainty, defined as a single Gaussian-constrained Barlow-Beeston-lite parameter [?, ?], is created and used to scale the total yield in a given bin.

## 1.9 Statistical Analysis

As was mentioned in Section ??, the final variable ( $\tilde{M}_T(\text{HH})$ ) is constructed as the sum of the Lorentz vectors of the  $Z(\ell\ell)$  candidate, the  $\text{H} \rightarrow b\bar{b}$  candidate, and  $\cancel{p}_T$ . This variable, a pseudo-transverse mass of the double Higgs system, is used to extract the results in this measurement. In this section, we discuss how the distributions of the  $\tilde{M}_T(\text{HH})$  variable are used in the statistical analysis to obtain the limits on the  $HH$  production multiplied by the BFs of the final state.

### 1.9.1 The likelihood function

The results in this measurement are obtained with the maximum likelihood fit. We perform a simultaneous fit of the SR and both CRs for both di-electron and di-muon channels using the likelihood function constructed as a product of Poisson terms over all bins of the input  $\tilde{M}_T(\text{HH})$  distributions in the three regions (SR, CRDY, CRTT) with Gaussian terms to constrain the nuisance parameters:

$$L(r_{\text{signal}}, r_k | \text{data}) = \prod_{i=1}^{N_{\text{bins}}} \frac{\mu_i^{n_i} \cdot e^{-\mu_i}}{n_i!} \cdot \prod_{j=1}^{N_{\text{nuisances}}} e^{-\frac{1}{2}\theta_j^2}$$

where the product index  $i$  refers to the bin of the input distributions, the product index  $j$  refers to uncertainties accounted for by the fit model, and  $n_i$  is the number of observed data events in the bin  $i$ . The mean value for each of the Poisson distributions is computed as:

$$\mu_i = r_{\text{signal}} \cdot S_i + \sum_k r_k \cdot B_{k,i},$$

where  $k$  refers to the background process  $k$ , and  $B_{k,i}$  is the content of the bin  $i$  of the background shape for a process  $k$ , while  $S_i$  is the content of the bin  $i$  of the signal shape. The  $S$  and  $B_k$  distributions are prepared using MC simulations with the best-known cross section values. This is done to have the number of expected MC events equal to the number of events in the analyzed data set (with the corresponding integrated luminosity). When filling the distributions for  $S$  and  $B_k$ , we also use a set of weights composed of scale factors of various types that depend on kinematic parameters of the candidates.

The parameter  $r_k$  sets the normalization of the background process  $k$  while  $r_{signal}$  is the signal strength parameter. All  $r$  parameters are floating freely in the fit. Two values of the signal strength parameter are of special interest:  $r_{signal} = 0$  describes the background-only hypothesis, while  $r_{signal} = 1$  corresponds to the case where the  $HH$  cross-section matches the cross-section used for the initial signal normalization inspired by BSM models, 2 pb in our case, which is a typical value for predictions of WED models (e.g., at 300 GeV). The terms  $\theta_j$  represent the set of nuisance parameters that are introduced into the likelihood function as Gaussian constraints. In the final likelihood function, we have a nuisance parameter for each source of systematic uncertainty associated with each physics object and for each normalization and shape uncertainty for a given process. Each normalization nuisance parameter describes a relative difference in the event yield of the process before and after the fit to data; and each shape nuisance parameter quantifies the difference in shape of the  $\tilde{M}_T(HH)$  distribution obtained with interpolation or extrapolation of the nominal, up, and down shapes (see Section ??) corresponding to the variation of a given systematic effect to achieve the best fit to the data.


The fit is performed separately for each of the 16 resonance mass points in the range 250 to 1000 GeV for a radion and a graviton resonance hypothesis individually.

During the fit, the values of the  $r_{signal}$  and  $r_k$ , along with the uncertainties, are determined.


The implementation of the maximum likelihood fit is done in the Higgs Combine package, a software framework supported by the CMS experiment [?]. The framework is based on the RooStats package [?] that is widely used in the HEP community.

Figure ?? shows the result of the fit for one mass point - 300 GeV. These  $HH$  transverse mass distributions for the graviton hypothesis are of the post-fit type - the normalizations and shapes of all components were adjusted according to the best-fit values. A similar figure in the case of the radion hypothesis is shown in Fig. ?. The signal component is very close to zero, it would not be visible on the figures. Therefore, the signal event yield is further scaled by a certain high factor for visual purposes, to make the signal contribution clearly visible.





**Figure 1.23:** Transverse mass of the reconstructed  $HH$  candidates for data, the simulated signal for the 300 GeV mass hypothesis of the graviton, and simulated background processes. Event yields of the MC simulations are scaled according to the fit results. The top row shows the figures for the di-muon channel, while the bottom row is for the di-electron channel. For each row, the left plot is for the CRDY, the middle is for the CRTT, and the right is for SR. Signal scaling choice is discussed in the text. The crosshatched area represents the sum of statistical and systematic uncertainties.



**Figure 1.24:** Transverse mass of the reconstructed  $HH$  candidates for data, the simulated signal for the 300 GeV mass hypothesis of the radion, and simulated background processes. Event yields of the MC simulations are scaled according to the fit results. The top row shows the figures for the di-muon channel, while the bottom row is for the di-electron channel. For each row, the left plot is for the CRDY, the middle is for the CRTT, and the right is for SR. Signal scaling choice is discussed in the text. The crosshatched area represents the sum of statistical and systematic uncertainties.

## 1.9.2 The fit results

Previous  $HH$  measurements provided no evidence for  $HH$  production through a narrow width resonance in the mass range from 250 GeV to 1 TeV. Thus, we proceed setting upper 95 % confidence level limits on the  $HH$  production cross-section, using the modified frequentist  $CL_s$  approach (asymptotic  $CL_s$ ) [?, ?, ?, ?].

It is worthwhile letting the reader know that before the real data is used to derive observed limits, the toy data set is first used to optimize the expected limits. This toy data set, called the Asimov data set, is commonly used by CMS and ATLAS experi-

ments for StatAn procedures. The Asimov data set (see [?]) represents the alternative hypothesis, which in most cases is the expected background with no fluctuations. The signal strength is set to zero, which, in a simple counting experiment [?] corresponds to the total number of events equal to the number of background events (no signal events).

As this measurement is a search, we use a standard CMS practice that helps experimenters avoid bias in the measurement - a blinding procedure. First, the measurement is performed only with a simulation. The expected sensitivity and the expected limits (with the Asimov data set) on the quantity being measured are determined, and the procedure is frozen. Second, the likelihood fit is performed and a set of diagnostic figures is prepared with real data, including all kinematic distributions shown in Section ??, the BDT outputs ??, and  $\tilde{M}_T(\text{HH})$  distributions in the signal region ??, ??. The figures are examined by the relevant CMS groups for data and simulation agreement. In our case, they were the CMS Higgs Physics Analysis group (PAG), CMS  $\text{H} \rightarrow \text{ZZ}$  PAG subgroup, and the CMS double Higgs PAG subgroup. At this stage, the results are kept blind and no real data is used. Finally, all outputs of the fit are examined, and the mentioned CMS PAGs give the “green light” for the unblinding procedure. Then, if the result obtained with the real data show no signs of experimental errors, the analysis reaches its final state - it is given into the hands of the Analysis Review Committee (ARC). The ARC serves several purposes: once again it verifies that the whole analysis chain is correct and helps lead the analysis towards the publication. During interactions with the ARC, additional studies and checks are performed to prove the correctness of the results.


The observed and expected 95% upper CL limits for the full mass range and both resonances are listed in Table ??. We produce the standard CMS-style figures of limits, shown in Fig. ??. The green and yellow bands correspond to one and two

standard deviations around the expected limit respectively. Since 450 GeV is the separation boundary between two mass regions, low mass and high mass - in which different BDT discriminants are applied - the limit calculation is performed with both of the BDTs at 450 GeV, where the discontinuity is seen in the figure. The figure also shows the expected production cross section for an RS1 KK graviton and RS1 radion in WED models. These cross sections are computed in [?]. As seen from the figure, this measurement is not sensitive by itself to probe the chosen model of  $HH$  production. The combination of this measurement with other similar measurements is necessary, as discussed in Section ??.

**Table 1.11:** The expected and observed  $HH$  production cross section upper limits at 95% CL for different narrow resonance graviton (top) and radion (bottom) mass hypotheses for both dielectron and dimuon channels combined.

Mass, GeV	Observed Limit, pb	Expected Limit, pb
250	253.5	589.1
260	272.2	585.9
270	274.4	537.5
300	380.0	434.4
350	330.6	309.4
400	90.4	119.9
450	59.8	63.3
500	31.0	36.6
550	14.5	20.2
600	9.8	12.7
650	18.5	11.1
700	16.1	10.1
750	13.7	8.8
800	10.1	6.5
900	8.1	4.8
1000	5.8	4.2

Mass, GeV	Observed Limit, pb	Expected Limit, pb
250	107.3	297.7
260	170.8	410.9
270	207.0	470.3
300	451.7	496.9
350	532.6	496.9
400	155.7	171.1
450	89.3	82.0
500	36.0	54.4
550	18.7	28.5
600	13.2	19.6
650	24.6	17.2
700	16.4	12.0
750	13.9	10.4
800	12.6	9.8
900	6.9	5.6
1000	5.7	4.5



**Figure 1.25:** Expected (dashed line) and observed (solid line) limits on the cross section of a resonant  $HH$  production as a function of the mass of the narrow resonance for both leptonic channels combined. Graviton case is shown at the top and radion case at the bottom. The red line shows a theoretical prediction for the production of a WED particle with certain model assumptions [?].

## 1.10 Discussion

In searches for a process, when intermediate particles can decay through a variety of channels, a common experimental approach is to search for multiple final states in parallel. Then the results are combined into a single result. This is a common method in the measurement of limits on a cross-section of a physics process such as the production of a heavy resonance.


At CMS, the Higgs boson program, endorsed by the CMS Higgs PAG, searches for  $HH$  production, both of the SM and BSM types. These searches cover multiple channels and are published as individual papers. For di-Higgs measurements, the individual channels include many of the channels seen in Fig. ???. This is usually followed by the final paper that presents the combined result of all individual channels, the “grand combination”.

This measurement, the search for a resonant production of the  $HH$  system decaying through the intermediate  $bbZZ$  state into  $2b2\ell2\nu$  final state is completed, approved by the CMS collaboration [?], and is shown at conferences [?]. This was the first measurement of the  $HH$  production in the  $bbZZ$  intermediate state that was performed at CERN using the LHC data.

This data analysis, at the time of this writing, is being combined with a similar channel where the  $HH$  system decays through the same intermediate state  $bbZZ$  into the final state of  $2b2\ell2q$ , and a paper will be submitted to the Physical Review D shortly after the defence of this dissertation.

Once the full 13 TeV dataset is fully analyzed, the CMS plans to release the grand combination paper for the  $HH$  production search that is based on the  $bbZZ$  channel discussed here as well as many other channels. As of this writing, the best available grand combination from the CMS experiment is based on a partial 13 TeV data set

analysis, see Fig. ??, published in [?] that includes a subset of all channels. That paper does not include the present measurement, which would be found just above the  $bbWW$  graph, which is the curve shown in orange. As can be seen from the figure, the most sensitive channels are the  $bb\gamma\gamma$  for the mass region below 500 GeV and the  $bbbb$  channel for higher masses.



**Figure 1.26:** Combination of  $HH$  channels using 2016 data. Expected (dashed) and observed (solid line) 95% CL exclusion limits are shown. The results describe the production cross section of a narrow width spin 0 (top) and spin 2 (bottom) resonance decaying into a pair of SM Higgs bosons.



While no individual channel measurement is sensitive enough to see a resonant  $HH$  production at the level predicted by the WED theory, the grand combination for the full data set may be able to see the process, if it exists, and approach the sensitivity of the order of picobarns. With more data taken in Run 3 and the HL-LHC period, the BSM theories discussed here will be probed more stringently. However, the ability of a particle collider experiment to detect the SM production of the  $HH$  system is further off into the future.

A projected sensitivity for a combined  $HH$  measurement based on the full data set expected to be collected with the HL-LHC (almost  $3\text{ ab}^{-1}$  of data), shows that this amount of data is not enough to reach the discovery sensitivity [?] : “the statistical combination of the five decay channels results in an expected significance for the standard model  $HH$  signal of  $2.6\sigma$ ”. This is a clear sign that more data are needed for both BSM and SM searches of the  $HH$  production. However, many Higgs PAG analysts would agree that new statistical and MVA tools should be developed and employed. Thus, the next versions of the  $HH$  analyses will most likely use a sophisticated neural network not only for signal-background separation but also for lepton reconstruction and more. At the moment, there is a strong effort in the CMS community to develop such tools and methods.

This concludes the discussion of analysis details. In the next chapter, we will summarise the main ideas that have been covered throughout this thesis.

---

## References

---

- [1] Alexandra Oliveira. Gravity particles from Warped Extra Dimensions, predictions for LHC. 2014.
- [2] Lisa Randall and Raman Sundrum. Large mass hierarchy from a small extra dimension. *Phys. Rev. Lett.*, 83:3370–3373, Oct 1999.
- [3] Alexandra Oliveira. Gravity particles from Warped Extra Dimensions, predictions for LHC. 2014.
- [4] Michele de Gruttola, Rami Kamalieddin, Ilya Kravchenko, Lesya Shchutska. Search for resonant diHiggs production with  $bbZZ$  decays with the  $2b2l2\nu$  signature using 35.9/fb data of 2016 pp collisions at the LHC. [http://cms.cern.ch/iCMS/jsp/openfile.jsp?tp=draft&files=AN2017\\_198\\_v17.pdf](http://cms.cern.ch/iCMS/jsp/openfile.jsp?tp=draft&files=AN2017_198_v17.pdf).
- [5] Search for resonant double Higgs production with  $bbZZ$  decays in the  $b\bar{b}l\ell\nu\nu$  final state. Technical Report CMS-PAS-HIG-17-032, CERN, Geneva, 2018.
- [6] Gunter Zech. Upper Limits in Experiments with Background Or Measurement Errors. *Nucl. Instrum. Meth.*, A277:608, 1989.
- [7] Michele de Gruttola, Caterina Vernieri, Pierluigi Bortignon, David Curry, Ivan Furic, Jacobo Konigsberg, Sean-Jiun Wang, Paolo Azzurri, Tommaso Boccali,

- Andrea Rizzi, Silvio Donato, Stephane Brunet Cooperstein , James Olsen, Christopher Palmer, Lorenzo Bianchini, Christoph Grab, Gael Ludovic Perrin, and Luca Perrozzi. Search for the Standard Model Higgs Boson Produced in Association with W and Z and Decaying to Bottom Quarks. [http://cms.cern.ch/iCMS/jsp/db\\_notes/noteInfo.jsp?cmsnoteid=CMS%20AN-2015/168](http://cms.cern.ch/iCMS/jsp/db_notes/noteInfo.jsp?cmsnoteid=CMS%20AN-2015/168).
- [8] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro. The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations. *JHEP*, 07:079, 2014.
- [9] SM Higgs Branching Ratios and Total Decay Widths (update in CERN Report4 2016). [https://twiki.cern.ch/twiki/bin/view/LHCPhysics/CERNYellowReportPageBR#Higgs\\_2\\_g](https://twiki.cern.ch/twiki/bin/view/LHCPhysics/CERNYellowReportPageBR#Higgs_2_g)
- [10] Johan Alwall et al. Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions. *Eur. Phys. J. C*, 53:473–500, 2008.
- [11] Stefan Kallweit, Jonas M. Lindert, Philipp Maierhofer, Stefano Pozzorini, and Marek Schonherr. NLO QCD+EW predictions for V + jets including off-shell vector-boson decays and multijet merging. *JHEP*, 04:021, 2016.
- [12] Simone Alioli, Paolo Nason, Carlo Oleari, and Emanuele Re. NLO single-top production matched with shower in POWHEG: s- and t-channel contributions. *JHEP*, 09:111, 2009. [Erratum: JHEP02,011(2010)].
- [13] Paolo Nason. A New method for combining NLO QCD with shower Monte Carlo algorithms. *JHEP*, 11:040, 2004.

- [14] Stefano Frixione, Paolo Nason, and Carlo Oleari. Matching NLO QCD computations with Parton Shower simulations: the POWHEG method. *JHEP*, 11:070, 2007.
- [15] Simone Alioli, Paolo Nason, Carlo Oleari, and Emanuele Re. A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX. *JHEP*, 06:043, 2010.
- [16] Nikolaos Kidonakis. Differential and total cross sections for top pair and single top production. Technical Report arXiv:1205.3453, Kennesaw State University, May 2012. Comments: 4 pages, 10 figures; presented at XX International Workshop on Deep-Inelastic Scattering and Related Subjects (DIS2012), Bonn, Germany, March 26-30, 2012.
- [17] Michal Czakon, Paul Fiedler, and Alexander Mitov. Total Top-Quark Pair-Production Cross Section at Hadron Colliders Through  $O(\alpha_s^4)$ . *Phys. Rev. Lett.*, 110:252004, 2013.
- [18] Torbjorn Sjostrand, Stephen Mrenna, and Peter Z. Skands. A Brief Introduction to PYTHIA 8.1. *Comput. Phys. Commun.*, 178:852–867, 2008.
- [19] Torbjorn Sjostrand, Stefan Ask, Jesper R. Christiansen, Richard Corke, Nishita Desai, Philip Ilten, Stephen Mrenna, Stefan Prestel, Christine O. Rasmussen, and Peter Z. Skands. An Introduction to PYTHIA 8.2. *Comput. Phys. Commun.*, 191:159–177, 2015.
- [20] Measurements of the  $pp \rightarrow WZ$  inclusive and differential production cross section and constraints on charged anomalous triple gauge couplings at  $\sqrt{s} = 13$  TeV. Technical Report CMS-PAS-SMP-18-002, CERN, Geneva, 2018.

- [21] Measurement of the WW cross section pp collisions at  $\sqrt{s}=13$  TeV. Technical Report CMS-PAS-SMP-16-006, CERN, Geneva, 2016.
- [22] Vardan Khachatryan et al. Measurement of the ZZ production cross section and  $Z \rightarrow \ell^+ \ell^- \ell'^+ \ell'^-$  branching fraction in pp collisions at  $\sqrt{s}=13$  TeV. *Phys. Lett. B*, 763:280, 2016. [Erratum: *Phys. Lett. B* 772, 884 (2017)].
- [23] Rikkert Frederix and Stefano Frixione. Merging meets matching in MC@NLO. *JHEP*, 12:061, 2012.
- [24] John M. Campbell and R. K. Ellis. MCFM for the Tevatron and the LHC. *Nucl. Phys. Proc. Suppl.*, 205-206:10, 2010.
- [25] The LHC Higgs cross-section working group. <https://twiki.cern.ch/twiki/bin/view/LHCPhysics/LHCHXSWG>.
- [26] SM Higgs production cross sections at  $\sqrt{s} = 13$  TeV. <https://twiki.cern.ch/twiki/bin/view/LHCPhysics/CERNYellowReportPageAt13TeV#ZH1H>
- [27] NNLO+NNLL top-quark-pair cross sections. [https://twiki.cern.ch/twiki/bin/view/LHCPhysics/TtbarNNLO#Top\\_quark\\_pair\\_cross\\_se](https://twiki.cern.ch/twiki/bin/view/LHCPhysics/TtbarNNLO#Top_quark_pair_cross_se)
- [28] Single Top Cross sections. <https://twiki.cern.ch/twiki/bin/viewauth/CMS/SingleTopSigma>.
- [29] CMS GEN XSEC Task Force. <https://twiki.cern.ch/twiki/bin/viewauth/CMS/GenXsecTaskF>
- [30] Richard D. Ball et al. Parton distributions for the LHC Run II. *JHEP*, 04:040, 2015.
- [31] Vardan Khachatryan et al. Event generator tunes obtained from underlying event and multiparton scattering measurements. *Eur. Phys. J. C*, 76(3):155, 2016.

- [32] S. Agostinelli et al. GEANT4—a simulation toolkit. *Nucl. Instrum. Meth. A*, 506:250, 2003.
- [33] Wolfgang Adam, R Frühwirth, Are Strandlie, and T Todor. Reconstruction of Electrons with the Gaussian-Sum Filter in the CMS Tracker at the LHC. Technical Report CMS-NOTE-2005-001, CERN, Geneva, Jan 2005.
- [34] CMS MET group. MET Corrections and Uncertainties for Run-II. <https://twiki.cern.ch/twiki/bin/viewauth/CMS/MissingETRun2Corrections>.
- [35] CMS MET group. MET Filter Recommendations for Run II. <https://twiki.cern.ch/twiki/bin/view/CMS/MissingETOptionalFiltersRun2>.
- [36] Albert M Sirunyan et al. Search for resonant and nonresonant Higgs boson pair production in the  $b\bar{b}\mu\mu$  final state in proton-proton collisions at  $\sqrt{s} = 13$  TeV. 2017.
- [37] A.M. Sirunyan, R. Kamalieddin, and et al I. Kravchenko. Evidence for the higgs boson decay to a bottom quark–antiquark pair. *Physics Letters B*, 780:501 – 532, 2018.
- [38] S. Kallweit, J. M. Lindert, P. Maierhöfer, S. Pozzorini, and M. Schönherr. Nlo qcd+ew predictions for  $v + \text{jets}$  including off-shell vector-boson decays and multijet merging. *Journal of High Energy Physics*, 2016(4):21, Apr 2016.
- [39] Geoffrey I. Webb Claude Sammut. *Encyclopedia of Machine Learning*. Springer, 1st edition. edition, 2011.
- [40] Helge Voss, Andreas Höcker, Jörg Stelzer, and Frerik Tegenfeldt. TMVA, the toolkit for multivariate data analysis with ROOT. In *XIth International Work-*

*shop on Advanced Computing and Analysis Techniques in Physics Research (ACAT)*, page 40, 2007.

- [41] Combination of searches for Higgs boson pair production in proton-proton collisions at  $\sqrt{s} = 13$  TeV. Technical Report CMS-PAS-HIG-17-030, CERN, Geneva, 2018.
- [42] CMS Collaboration. CMS Luminosity Measurements for the 2016 Data Taking Period. CMS Physics Analysis Summary CMS-PAS-LUM-17-001, CERN, Geneva, 2017.
- [43] Michiel Botje et al. The PDF4LHC Working Group Interim Recommendations. 2011.
- [44] Sergey Alekhin et al. The PDF4LHC Working Group Interim Report. 2011.
- [45] Jon Butterworth et al. PDF4LHC recommendations for LHC Run II. *J. Phys.*, G43:023001, 2016.
- [46] Roger Barlow and Christine Beeston. Fitting using finite monte carlo samples. *Computer Physics Communications*, 77(2):219 – 228, 1993.
- [47] J. S. Conway. Incorporating Nuisance Parameters in Likelihoods for Multisource Spectra. In *Proceedings, PHYSTAT 2011 Workshop on Statistical Issues Related to Discovery Claims in Search Experiments and Unfolding*, CERN, Geneva, Switzerland 17-20 January 2011, pages 115–120, 2011.
- [48] CMS Higgs WG. Documentation of the RooStats -based statistics tools for Higgs PAG. <https://twiki.cern.ch/twiki/bin/view/CMS/SWGuideHiggsAnalysisCombinedLimit>.

- [49] L. Moneta, K. Belasco, K. S. Cranmer, A. Lazzaro, D. Piparo, G. Schott, W. Verkerke, and M. Wolf. The RooStats Project. In *13<sup>th</sup> International Workshop on Advanced Computing and Analysis Techniques in Physics Research (ACAT2010)*. SISSA, 2010. PoS(ACAT2010)057.
- [50] Thomas Junk. Confidence level computation for combining searches with small statistics. *Nucl.Instrum.Meth.*, A434:435, 1999.
- [51] A. L. Read. Presentation of search results: the CLs technique. *J. Phys. G: Nucl. Part. Phys.*, 28, 2002.
- [52] SM Higgs Combination. Technical Report CMS-PAS-HIG-11-011, CERN, Geneva, 2011.
- [53] Glen Cowan, Kyle Cranmer, Eilam Gross, and Ofer Vitells. Asymptotic formulae for likelihood-based tests of new physics. *Eur. Phys. J.*, C71:1554, 2011. [Erratum: *Eur. Phys. J.*C73,2501(2013)].
- [54] Eilam Gross. Practical statistics for high energy physics. *CERN Yellow Reports: School Proceedings*, 4(0):165, 2017.
- [55] Andres Tiko. HH resonance searches at ATLAS and CMS. Tokyo, Japan, November 2018. "Higgs Couplings 2018" conference.
- [56] A. M. Sirunyan, A. Tumasyan, and W. et al. Adam. Combination of searches for higgs boson pair production in proton-proton collisions at  $\sqrt{s} = 13$  TeV. *Phys. Rev. Lett.*, 122:121803, Mar 2019.
- [57] Prospects for HH measurements at the HL-LHC. Technical Report CMS-PAS-FTR-18-019, CERN, Geneva, 2018.



- [58]
- [59] S. Frixione, P. Nason, and C. Oleari. Matching nlo qcd computations with parton shower simulations: the powheg method. *JHEP*, 11:070, 2007.
- [60] Gionata Luisoni, Paolo Nason, Carlo Oleari, and Francesco Tramontano.  $HW^\pm/HZ + 0$  and 1 jet at NLO with the POWHEG BOX interfaced to GoSam and their merging within MiNLO. *JHEP*, 10:083, 2013.
- [61] Comparison of nuisances for background only case, 350 GeV mass hypothesis.  
[http://rkamalie.web.cern.ch/rkamalie/feb12/Comparison\\_of\\_nuisances\\_expectedSignal0\\_3](http://rkamalie.web.cern.ch/rkamalie/feb12/Comparison_of_nuisances_expectedSignal0_3)
- [62] Comparison of nuisances for s+b case, 350 GeV mass hypothesis.  
[http://rkamalie.web.cern.ch/rkamalie/feb12/Comparison\\_of\\_nuisances\\_expectedSignal1\\_3](http://rkamalie.web.cern.ch/rkamalie/feb12/Comparison_of_nuisances_expectedSignal1_3)
- [63] k-factor for DY/Z. <https://twiki.cern.ch/twiki/bin/viewauth/CMS/SummaryTable1G25ns#D>
- [64] Standard Model Cross Sections for CMS at 13 TeV.  
<https://twiki.cern.ch/twiki/bin/viewauth/CMS/StandardModelCrossSectionsat13TeVInclusi>
- [65] Ryan Gavin, Ye Li, Frank Petriello, and Seth Quackenbush. W Physics at the LHC with FEWZ 2.1. *Comput. Phys. Commun.*, 184:208, 2013.
- [66] Stefano Frixione, Paolo Nason, and Giovanni Ridolfi. A Positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction. *JHEP*, 09:126, 2007.
- [67] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro. The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations. *JHEP*, 07:079, 2014.

- [68] Rikkert Frederix, Emanuele Re, and Paolo Torrielli. Single-top t-channel hadroproduction in the four-flavour scheme with POWHEG and aMC@NLO. *JHEP*, 09:130, 2012.
- [69] Serguei Chatrchyan et al. Determination of jet energy calibration and transverse momentum resolution in CMS. *JINST*, 6:P11002, 2011.
- [70] Emanuele Re. Single-top Wt-channel production matched with parton showers using the POWHEG method. *Eur. Phys. J.*, C71:1547, 2011.
- [71] Gael L. Perrin, Pedro Fernandez Manteca. Muon Identification and Isolation Scale-Factors on 2016 Dataset. [https://indico.cern.ch/event/611558/contributions/2465881/attachments/1407735/2151747/TnP\\_06\\_02\\_2017.pdf](https://indico.cern.ch/event/611558/contributions/2465881/attachments/1407735/2151747/TnP_06_02_2017.pdf).
- [72] CMS Muon POG. Tracking SFs on the full 2016 data. [https://twiki.cern.ch/twiki/bin/view/CMS/MuonWorkInProgressAndPagResults#Results\\_on\\_the\\_full\\_2016\\_data](https://twiki.cern.ch/twiki/bin/view/CMS/MuonWorkInProgressAndPagResults#Results_on_the_full_2016_data).
- [73] Gael L. Perrin. Double Muon trigger efficiency per-leg approach. [https://indico.cern.ch/event/636555/contributions/2577291/attachments/1453162/2241537/TnP\\_DoubleMuSF\\_03\\_05\\_17.pdf](https://indico.cern.ch/event/636555/contributions/2577291/attachments/1453162/2241537/TnP_DoubleMuSF_03_05_17.pdf).
- [74] Grace Dupuis. Collider Constraints and Prospects of a Scalar Singlet Extension to Higgs Portal Dark Matter. *JHEP*, 07:008, 2016.
- [75] Shinya Kanemura, Kunio Kaneta, Naoki Machida, Shinya Odori, and Tetsuo Shindou. Single and double production of the Higgs boson at hadron and lepton colliders in minimal composite Higgs models. *Phys. Rev.*, D94(1):015028, 2016.

- [76] Oleg Antipin, David Atwood, and Amarjit Soni. Search for RS gravitons via  $W(L)W(L)$  decays. *Phys. Lett.*, B666:155–161, 2008.
- [77] A. Liam Fitzpatrick, Jared Kaplan, Lisa Randall, and Lian-Tao Wang. Searching for the Kaluza-Klein Graviton in Bulk RS Models. *JHEP*, 09:013, 2007.
- [78] Kaustubh Agashe, Hooman Davoudiasl, Gilad Perez, and Amarjit Soni. Warped Gravitons at the LHC and Beyond. *Phys. Rev.*, D76:036006, 2007.
- [79] CMS Higgs PAG. List of question for the preapproval checks. <https://twiki.cern.ch/twiki/bin/viewauth/CMS/HiggsWG/HiggsPAGPreapprovalChecks>.
- [80] CMS BTV POG. Supported Algorithms and Operating Points. [https://twiki.cern.ch/twiki/bin/viewauth/CMS/BtagRecommendation80XReReco#Supported\\_Algorithms\\_and\\_Operati](https://twiki.cern.ch/twiki/bin/viewauth/CMS/BtagRecommendation80XReReco#Supported_Algorithms_and_Operati).
- [81] Abdus Salam and John Clive Ward. On a gauge theory of elementary interactions. *Nuovo Cim.*, 19:165–170, 1961.
- [82] Steven Weinberg. A model of leptons. *Phys.Rev.Lett.*, 19:1264–1266, 1967.
- [83] Albert M Sirunyan et al. Evidence for the Higgs boson decay to a bottom quark-antiquark pair. 2017.
- [84] Albert M Sirunyan et al. Particle-flow reconstruction and global event description with the CMS detector. *JINST*, 12(10):P10003, 2017.
- [85] Aruna Kumar Nayak. Reconstruction of physics objects in the CMS detector. *PoS*, CHARGED2012:010, 2012.
- [86] Serguei Chatrchyan et al. Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC. *Phys. Lett.*, B716:30–61, 2012.

- [87] Georges Aad et al. Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Phys. Lett.*, B716:1–29, 2012.
- [88] Giuliano Panico. Prospects for double Higgs production. *Frascati Phys. Ser.*, 61:102, 2016.
- [89] I. Belotelov, I. Golutvin, D. Bourilkov, A. Lanyov, E. Rogalev, M. Savina, and S. Shmatov. Search for ADD extra dimensional gravity in di-muon channel with the CMS detector. CMS Note 2006/076, 2006.
- [90] M. Aldaya, P. Arce, J. Caballero, B. de la Cruz, P. Garcia-Abia, J. M. Hernandez, M. I. Josa, and E. Ruiz. Discovery potential and search strategy for the standard model Higgs boson in the  $H \rightarrow ZZ^* \rightarrow 4\mu$  decay channel using a mass-independent analysis. CMS Note 2006/106, 2006.
- [91] A. Brandt et al. Measurements of single diffraction at  $\sqrt{s} = 630$  GeV: Evidence for a non-linear  $\alpha(t)$  of the pomeron. *Nucl. Phys. B*, 514:3, 1998.
- [92] W. Buchmüller and D. Wyler. Constraints on SU(5)-type leptoquarks. *Phys. Lett. B*, 177:377, 1986.
- [93] CMS Collaboration. CMS technical design report, volume II: Physics performance. *J. Phys. G*, 34:995, 2007.
- [94] CMS Collaboration. Jet performance in pp collisions at  $\sqrt{s}=7$  TeV. CMS Physics Analysis Summary CMS-PAS-JME-10-003, 2010.
- [95] S. Chatrchyan et al. The CMS experiment at the CERN LHC. *JINST*, 3:S08004, 2008.

- [96] Particle Data Group, J. Beringer, et al. Review of Particle Physics. *Phys. Rev. D*, 86:010001, 2012.
- [97] ALEPH, CDF, D0, DELPHI, L3, OPAL, SLD Collaborations, the LEP Electroweak Working Group, the Tevatron Electroweak Working Group, and the SLD Electroweak and Heavy Flavour Groups. Precision electroweak measurements and constraints on the Standard Model. 2010.
- [98] I. Bertram, G. Landsberg, J. Linnemann, R. Partridge, M. Paterno, and H. B. Prosper. A recipe for the construction of confidence limits. Technical Report TM-2104, Fermilab, 2000.
- [99] Vardan Khachatryan et al. Search for the standard model Higgs boson produced through vector boson fusion and decaying to  $b\bar{b}$ . *Phys. Rev.*, D92(3):032008, 2015.
- [100] Performance of muon identification in pp collisions at  $\sqrt{s}=7$  TeV. Technical Report CMS-PAS-MUO-10-002, CERN, Geneva, 2010.
- [101] Serguei Chatrchyan et al. Performance of CMS muon reconstruction in  $pp$  collision events at  $\sqrt{s}=7$  TeV. *JINST*, 7:P10002, 2012.
- [102] CMS COLLABORATION. Particle-flow event reconstruction in CMS and performance for jets, taus, and  $E_T^{miss}$ . CMS Physics Analysis Summary CMS-PAS-PFT-09-001, CERN, 2009.
- [103] CMS COLLABORATION. Commissioning of the particle-flow event reconstruction with the first lhc collisions recorded in the cms detector. CMS Physics Analysis Summary CMS-PAS-PFT-10-001, CERN, 2010.

- [104] Vardan Khachatryan et al. Performance of Electron Reconstruction and Selection with the CMS Detector in Proton-Proton Collisions at  $\sqrt{s} = 8$  TeV. *JINST*, 10(06):P06005, 2015.
- [105] CMS Collaboration. Search for pair production of first-generation scalar leptquarks in pp collisions at  $\sqrt{s} = 7$  TeV. Submitted to *Phys. Rev. Lett.*, 2010.
- [106] CMS Collaboration. Performance of cms muon reconstruction in pp collision events at  $\sqrt{s} = 7$  TeV. Submitted to *J. Inst.*, 2012.
- [107] ATLAS Collaboration. Search for the Higgs boson in the  $H \rightarrow WW(*) \rightarrow \ell^+ \nu \ell^- \bar{\nu}$  decay channel in pp collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector. Submitted to *Phys. Rev. Lett.*, 2011.
- [108] Matteo Cacciari and Gavin P. Salam. Dispelling the  $N^3$  myth for the  $k_t$  jet-finder. *Phys. Lett. B*, 641:57, 2006.
- [109] CMS Luminosity Measurements for the 2016 Data Taking Period. Technical Report CMS-PAS-LUM-17-001, CERN, Geneva, 2017.
- [110] CMS Muon POG. Reference muon id, isolation and trigger efficiencies for Run-II. <https://twiki.cern.ch/twiki/bin/viewauth/CMS/MuonReferenceEffsRun2>.
- [111] John M. Campbell, R. Keith Ellis, Paolo Nason, and Emanuele Re. Top-pair production and decay at NLO matched with parton showers. *JHEP*, 04:114, 2015.
- [112] Ryan Gavin, Ye Li, Frank Petriello, and Seth Quackenbush. FEWZ 2.0: A code for hadronic Z production at next-to-next-to-leading order. *Comput. Phys. Commun.*, 182:2388, 2011.

- [113] Ye Li and Frank Petriello. Combining QCD and electroweak corrections to dilepton production in FEWZ. *Phys. Rev. D*, 86:094034, 2012.
- [114] Nikolaos Kidonakis. Differential and total cross sections for top pair and single top production. In *Proceedings, 20th International Workshop on Deep-Inelastic Scattering and Related Subjects (DIS 2012): Bonn, Germany, March 26-30, 2012*, page 831, 2012. [831(2012)].
- [115] Chris Palmer. VHbb Electron Trigger and ID+ISO SFs for 2016 data. [https://indico.cern.ch/event/604949/contributions/2543520/attachments/1439974/2216426/VHbb\\_TnP\\_SF\\_s\\_egamma\\_april.pdf#search=vhbb%20AND%20cerntaxonomy%3A%22Indico%2FExperiments%2FCMS%20meetings%2FPH%20%2D%20Physics%2FEgamma%22](https://indico.cern.ch/event/604949/contributions/2543520/attachments/1439974/2216426/VHbb_TnP_SF_s_egamma_april.pdf#search=vhbb%20AND%20cerntaxonomy%3A%22Indico%2FExperiments%2FCMS%20meetings%2FPH%20%2D%20Physics%2FEgamma%22).
- [116] CMS JetMET group. Jet Energy Resolution. <https://twiki.cern.ch/twiki/bin/viewauth/CMS/JetResolution>.
- [117] CMS EGM POG. Multivariate Electron Identification for Run2. <https://twiki.cern.ch/twiki/bin/viewauth/CMS/MultivariateElectronIdentificationRun2>.
- [118] CMS Higgs WG. Binned shape analysis with the Higgs Combination Tool. [https://twiki.cern.ch/twiki/bin/view/CMS/SWGuideHiggsAnalysisCombinedLimit#Binned\\_shape\\_analysis](https://twiki.cern.ch/twiki/bin/view/CMS/SWGuideHiggsAnalysisCombinedLimit#Binned_shape_analysis).
- [119] bbbb team. Search for resonant pair production of Higgs bosons decaying to bottom quark-antiquark pairs in proton-proton collisions at 13 TeV. <http://cms.cern.ch/iCMS/analysisadmin/get?analysis=HIG-17-009-pas-v5.pdf>.

- [120] Matteo Cacciari, Gavin P. Salam, and Gregory Soyez. The anti- $k_t$  jet clustering algorithm. *JHEP*, 04:063, 2008.
- [121] Torbjörn Sjöstrand, Stephen Mrenna, and Peter Skands. PYTHIA 6.4 physics and manual. *JHEP*, 05:026, 2006.
- [122] C. Giunti and M. Laveder. Neutrino mixing. In F. Columbus and V. Krasnoholovets, editors, *Developments in Quantum Physics*. Nova Science Publishers, Inc., 2004.
- [123] Richard Phillips Feynman, Robert Benjamin Leighton, and Matthew Sands. *The Feynman lectures on physics; New millennium ed.* Basic Books, New York, NY, 2010. Originally published 1963-1965.
- [124] Savas Dimopoulos, Stuart Raby, and Frank Wilczek. Proton decay in supersymmetric models. *Physics Letters B*, 112(2):133 – 136, 1982.
- [125] David J Griffiths. *Introduction to elementary particles; 2nd rev. version.* Physics textbook. Wiley, New York, NY, 2008.
- [126] M. Della Negra, P. Jenni, and T. S. Virdee. Journey in the search for the higgs boson: The atlas and cms experiments at the large hadron collider. *Science*, 338(6114):1560–1568, 2012.
- [127] Jennifer Ouellette. Einstein’s quest for a unified theory. *APS*, 2015.
- [128] E A Davis and Isabel Falconer. *J.J. Thompson and the discovery of the electron.* Taylor and Francis, Hoboken, NJ, 2002.
- [129] Oreste Piccioni. *The Discovery of the Muon*, pages 143–162. Springer US, Boston, MA, 1996.



- [130] G. Danby, J-M. Gaillard, K. Goulianos, L. M. Lederman, N. Mistry, M. Schwartz, and J. Steinberger. Observation of high-energy neutrino reactions and the existence of two kinds of neutrinos. *Phys. Rev. Lett.*, 9:36–44, Jul 1962.
- [131] M. L. Perl, G. S. Abrams, A. M. Boyarski, et al. Evidence for anomalous lepton production in  $e^+ - e^-$  annihilation. *Phys. Rev. Lett.*, 35:1489–1492, Dec 1975.
- [132] K. Kodama et al. Observation of tau neutrino interactions. *Phys. Lett.*, B504:218–224, 2001.
- [133] S. M. Bilenky. Neutrino in Standard Model and beyond. *Phys. Part. Nucl.*, 46(4):475–496, 2015.
- [134] S Chandrasekhar. *Newton's principia for the common reader*. Oxford Univ., Oxford, 2003. The book can be consulted by contacting: PH-AID: Wallet, Lionel.
- [135] Hanoeh Gutfreund and Jurgen Renn. *The road to relativity: the history and meaning of Einstein's "The foundation of general relativity" : featuring the original manuscript of Einstein's masterpiece*. Princeton University Press, Princeton, NJ, Apr 2015.
- [136] J. Butterworth. *Smashing Physics*. Headline Publishing Group, 2014.
- [137] W N Cottingham and D A Greenwood. *An Introduction to the Standard Model of Particle Physics; 2nd ed*. Cambridge Univ. Press, Cambridge, 2007.
- [138] Eric W. Weisstein. Fundamental forces.
- [139] Carl Bender. Mathematical physics.

- [140] Andrew Wayne. QED and the Men Who Made It: Dyson, Feynman, Schwinger, and Tomonaga by Silvan S. Schweber. *The British Journal for the Philosophy of Science*, 46(4):624–627, 1995.
- [141] C. Patrignani et al. Review of Particle Physics. *Chin. Phys.*, C40(10):100001, 2016.
- [142] Michelangelo L Mangano. Introduction to QCD. (CERN-OPEN-2000-255), 1999.
- [143] Matt Strassler. Of particular significance: Conversations about science with theoretical physicist matt strassler.
- [144] S. L. Glashow. Partial Symmetries of Weak Interactions. *Nucl. Phys.*, 22:579–588, 1961.
- [145] F. Englert and R. Brout. Broken symmetry and the mass of gauge vector mesons. *Phys. Rev. Lett.*, 13:321–323, Aug 1964.
- [146] Peter W. Higgs. Broken symmetries and the masses of gauge bosons. *Phys. Rev. Lett.*, 13:508–509, Oct 1964.
- [147] G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble. Global conservation laws and massless particles. *Phys. Rev. Lett.*, 13:585–587, Nov 1964.
- [148] Pauline Gagnon. *Who cares about particle physics? : making sense of the Higgs boson, the Large Hadron Collider and CERN*. Oxford University Press, 2016.
- [149] Matthias U. Mozer. Electroweak Physics at the LHC. *Springer Tracts Mod. Phys.*, 267:1–115, 2016.

- [150] Roger Wolf. *The Higgs Boson Discovery at the Large Hadron Collider*, volume 264. Springer, 2015.
- [151] Steven Weinberg. The Making of the Standard Model. *Eur. Phys. J. C*, 34(hep-ph/0401010):5–13. 21 p. ; streaming video, 2003.
- [152] Jose Andres Monroy Montanez, Kenneth Bloom, and Aaron Dominguez. Search for production of a Higgs boson and a single Top quark in multilepton final states in pp collisions at  $\sqrt{s} = 13$  TeV, Jul 2018. Presented 23 Jul 2018.
- [153] Albert M Sirunyan et al. Search for Higgs boson pair production in the  $\gamma\gamma b\bar{b}$  final state in pp collisions at  $\sqrt{s} = 13$  TeV. 2018.
- [154] Matthew J. Dolan, Christoph Englert, and Michael Spannowsky. New Physics in LHC Higgs boson pair production. *Phys. Rev.*, D87(5):055002, 2013.
- [155] Peisi Huang, Aniket Joglekar, Min Li, and Carlos E. M. Wagner. Corrections to di-Higgs boson production with light stops and modified Higgs couplings. *Phys. Rev.*, D97(7):075001, 2018.
- [156] Lisa Randall and Raman Sundrum. A Large mass hierarchy from a small extra dimension. *Phys. Rev. Lett.*, 83:3370–3373, 1999.
- [157] Kunihiro Uzawa, Yoshiyuki Morisawa, and Shinji Mukohyama. Excitation of Kaluza-Klein gravitational mode. *Phys. Rev.*, D62:064011, 2000.
- [158] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo. Phenomenology of the Randall-Sundrum Gauge Hierarchy Model. *Phys. Rev. Lett.*, 84:2080, 2000.
- [159] Michael Forger and Hartmann Romer. Currents and the energy momentum tensor in classical field theory: A Fresh look at an old problem. *Annals Phys.*, 309:306–389, 2004.

- [160] Chuan-Ren Chen and Ian Low. Double take on new physics in double Higgs boson production. *Phys. Rev.*, D90(1):013018, 2014.
- [161] Roberto Contino, Margherita Ghezzi, Mauro Moretti, Giuliano Panico, Fulvio Piccinini, and Andrea Wulzer. Anomalous Couplings in Double Higgs Production. *JHEP*, 08:154, 2012.
- [162] Gennadi Sardanashvily. *Noether's theorems: applications in mechanics and field theory*. Atlantis studies in variational geometry. Springer, Paris, 2016.
- [163] Thomas Schörner-Sadenius. *The Large Hadron Collider: harvest of run 1*. Springer, Cham, 2015.
- [164] CERN. *Large Hadron Collider in the LEP Tunnel*, Geneva, 1984. CERN.
- [165] Lyndon R Evans and Philip Bryant. LHC Machine. *JINST*, 3:S08001. 164 p, 2008. This report is an abridged version of the LHC Design Report (CERN-2004-003).
- [166] Karsten Eggert, K Honkavaara, and Andreas Morsch. Luminosity considerations for the LHC. Technical Report CERN-AT-94-04-DI. CERN-LHC-Note-263. LHC-NOTE-263, CERN, Geneva, Feb 1994.
- [167] Oswald Gröbner. The LHC Vacuum System. (LHC-Project-Report-181. CERN-LHC-Project-Report-181):5 p, May 1998.
- [168] Thomas Lenzi. Development and Study of Different Muon Track Reconstruction Algorithms for the Level-1 Trigger for the CMS Muon Upgrade with GEM Detectors. Master's thesis, U. Brussels (main), 2013.

- [169] Combination of searches for Higgs boson pair production in proton-proton collisions at  $\sqrt{s} = 13$  TeV. Technical Report CMS-PAS-HIG-17-030, CERN, Geneva, 2018.
- [170] The CMS collaboration. Missing transverse energy performance of the cms detector. *Journal of Instrumentation*, 6(09):P09001, 2011.
- [171] Erwin Schrödinger. *Statistical thermodynamics; 2nd ed.* Cambridge Univ. Press, Cambridge, 1952.
- [172] Precise determination of the mass of the Higgs boson and studies of the compatibility of its couplings with the standard model. Technical Report CMS-PAS-HIG-14-009, CERN, Geneva, 2014.
- [173] R. P. Feynman. The theory of positrons. *Phys. Rev.*, 76:749–759, Sep 1949.
- [174] Werner Herr and B Muratori. Concept of luminosity. 2006.
- [175] Charles W. Misner, Kip S. Thorne, and John Archibald Wheeler. *Gravitation*. W. H. Freeman San Francisco, 1973.
- [176] Charles W. Misner, K. S. Thorne, and J. A. Wheeler. *Gravitation*. W. H. Freeman, San Francisco, 1973.
- [177] Cecile Noels. Literature in focus - The Large Hadron Collider: A Marvel of Technology. Literature in focus - The Large Hadron Collider: A Marvel of Technology. (BUL-NA-2009-414. 51/2009):11, Dec 2009.
- [178] John Hauptman. *Particle physics experiments at high energy colliders*. 2011.
- [179] CMS Collaboration. The Phase-2 Upgrade of the CMS Tracker. Technical Report CERN-LHCC-2017-009. CMS-TDR-014, CERN, Geneva, Jun 2017.

- [180] Philippe Bloch, Robert Brown, Paul Lecoq, and Hans Rykaczewski. *Changes to CMS ECAL electronics: addendum to the Technical Design Report*. Technical Design Report CMS. CERN, Geneva, 2002.
- [181] *The CMS hadron calorimeter project: Technical Design Report*. Technical Design Report CMS. CERN, Geneva, 1997.
- [182] G Baatian, Albert M Sirunyan, Virgil E Emeliantchik, Igor Barnes, Alvin T Laasanen, and Arnold Pompos. Design, Performance, and Calibration of CMS Hadron-Barrel Calorimeter Wedges. Technical Report CMS-NOTE-2006-138. 1, CERN, Geneva, May 2007.
- [183] G L Bayatian, S Chatrchyan, H Hmayakyan, A Poblaguev, M E Zeller, and B S Yuldashev. *CMS Physics: Technical Design Report Volume 1: Detector Performance and Software*. Technical Design Report CMS. CERN, Geneva, 2006. There is an error on cover due to a technical problem for some items.
- [184] CMS Collaboration. The Phase-2 Upgrade of the CMS Muon Detectors. Technical Report CERN-LHCC-2017-012. CMS-TDR-016, CERN, Geneva, Sep 2017. This is the final version, approved by the LHCC.
- [185] G. Bauer et al. The CMS data acquisition system software. *J. Phys. Conf. Ser.*, 219:022011, 2010.
- [186] Vardan Khachatryan et al. The CMS trigger system. *JINST*, 12(01):P01020, 2017.
- [187] A. M. Sirunyan et al. Particle-flow reconstruction and global event description with the CMS detector. *JINST*, 12(10):P10003, 2017.

- [188] Serguei Chatrchyan et al. Description and performance of track and primary-vertex reconstruction with the CMS tracker. *JINST*, 9(10):P10009, 2014.
- [189] Giuseppe Cerati et al. Parallelized and Vectorized Tracking Using Kalman Filters with CMS Detector Geometry and Events. 2018.
- [190] Serguei Chatrchyan et al. Description and performance of track and primary-vertex reconstruction with the CMS tracker. *JINST*, 9(10):P10009, 2014.
- [191] E Chabanat and N Estre. Deterministic Annealing for Vertex Finding at CMS. 2005.
- [192] R Frühwirth, Wolfgang Waltenberger, and Pascal Vanlaer. Adaptive Vertex Fitting. Technical Report CMS-NOTE-2007-008, CERN, Geneva, Mar 2007.
- [193] A. M. Sirunyan et al. Particle-flow reconstruction and global event description with the CMS detector. *JINST*, 12(10):P10003, 2017.
- [194] Andreas Hoecker, Peter Speckmayer, Joerg Stelzer, Jan Therhaag, Eckhard von Toerne, and Helge Voss. TMVA: Toolkit for Multivariate Data Analysis. *PoS, ACAT:040*, 2007.
- [195] Matteo Cacciari, Gavin P. Salam, and Gregory Soyez. The anti- $k_t$  jet clustering algorithm. *JHEP*, 04:063, 2008.
- [196] Lior Rokach. Ensemble-based classifiers. *Artificial Intelligence Review*, 33(1):1–39, Feb 2010.
- [197] Identification of b quark jets at the CMS Experiment in the LHC Run 2. Technical Report CMS-PAS-BTV-15-001, CERN, Geneva, 2016.

- [198] A. M. Sirunyan et al. Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV. *JINST*, 13(05):P05011, 2018.
- [199] A Zee. *Quantum Field Theory in a Nutshell*. Nutshell handbook. Princeton Univ. Press, Princeton, NJ, 2003.
- [200] Francis Halzen and Alan Douglas Martin. *Quarks and leptons: an introductory course in modern particle physics*. Wiley, New York, NY, 1984.
- [201] Gregory Soyez. Pileup mitigation at the LHC: a theorist’s view. *Phys. Rept.*, 803:1–158, 2019.
- [202] N. Adam, J. Berryhill, V. Halyo, A. Hunt, K. Mishra. Generic Tag and Probe Tool for Measuring Efficiency at CMS with Early Data . [http://cms.cern.ch/iCMS/jsp/openfile.jsp?tp=draft&files=AN2009\\_111\\_v1.pdf](http://cms.cern.ch/iCMS/jsp/openfile.jsp?tp=draft&files=AN2009_111_v1.pdf).
- [203] Walter D. Goldberger and Mark B. Wise. Modulus stabilization with bulk fields. *Phys. Rev. Lett.*, 83:4922–4925, 1999.
- [204] Piotr Traczyk and Grzegorz Wrochna. Search for Randall-Sundrum graviton excitations in the CMS experiment. 2002.
- [205] Walter D. Goldberger and Mark B. Wise. Bulk fields in the Randall-Sundrum compactification scenario. *Phys. Rev.*, D60:107505, 1999.
- [206] Sreerup Raychaudhuri and K Sridhar. *Particle physics of brane worlds and extra dimensions*. Cambridge monographs on mathematical physics. Cambridge University Press, Cambridge, 2016.
- [207] Sebastien Wertz and Vincent Lemaître. Search for Higgs boson pair production in the  $b\bar{b}\ell\nu\ell\nu$  final state with the CMS detector, 2018.



- [208] Luca Cadamuro, Yves Sirois, and Roberto Salerno. Search for Higgs boson pair production in the  $b\bar{b}\tau^+\tau^-$  decay channel with the CMS detector at the LHC. Recherche de la production de paires de bosons de Higgs dans le canal de désintégration  $b\bar{b}\tau^+\tau^-$  avec le détecteur CMS auprès du LHC, Sep 2017. Presented 05 Oct 2017.
- [209] CMS e-Lab. Energy, Mass and Momentum in High-Energy Physics. <https://www.i2u2.org/elab/cms/library/ref-mass-energy-momentum.jsp>.
- [210] The CMS collaboration. Performance of CMS muon reconstruction in pp collision events at  $\sqrt{s} = 7$  TeV. *Journal of Instrumentation*, 7(10):P10002–P10002, oct 2012.