

SEARCH FOR RESONANT DOUBLE HIGGS PRODUCTION WITH BBZZ  
DECAYS IN THE  $b\bar{b}\ell\ell\nu\bar{\nu}$  FINAL STATE IN pp COLLISIONS AT  $\sqrt{s} = 13$  TeV

by

Rami Kamalieddin

A DISSERTATION

Presented to the Faculty of  
The Graduate College at the University of Nebraska  
In Partial Fulfilment of Requirements  
For the Degree of Doctor of Philosophy

Major: Physics and Astronomy

Under the Supervision of Professor Ilya Kravchenko

Lincoln, Nebraska

May, 2019

SEARCH FOR RESONANT DOUBLE HIGGS PRODUCTION WITH BBZZ  
DECAYS IN THE  $b\bar{b}\ell\ell\nu\bar{\nu}$  FINAL STATE IN pp COLLISIONS AT  $\sqrt{s} = 13$  TeV

Rami Kamalieddin, Ph.D.

University of Nebraska, 2019

Advisers: Ilya Kravchenko

Since the discovery of the Higgs boson in 2012 by the ATLAS and CMS experiments, most of the quantum mechanical properties that describe the long-awaited Higgs boson have been measured. Due to an impeccable work of the LHC, dozens of  $fb^{-1}$  of data have been delivered to both experiments. Finally, it became possible for analyses that have a very low cross section to observe rare decay modes of the Higgs boson, as was done successfully recently in  $t\bar{t}H$  and  $VHbb$  channels. The only untouched territory is a double Higgs boson production. Data will not help us much either at the HL-LHC, the process will remain unseen even in the most optimistic scenarios, so one has to rely solely on new reconstruction methods as well as new analysis techniques. This thesis is addressing both goals. I have been blessed by an opportunity to work in the CMS electron identification group, where we have developed new electron identification algorithms. The majority of this thesis, however, will be devoted to the second goal of HL-LHC. We establish the techniques for the first ever analysis at the LHC that searches for the double Higgs production mediated by a heavy narrow-width resonance in the  $b\bar{b}ZZ$  channel:  $X \rightarrow HH \rightarrow b\bar{b}ZZ^* \rightarrow b\bar{b}\ell\ell\nu\bar{\nu}$ . The analysis searches for a resonant production of a Higgs boson pair in the range of masses of the resonant parent particle from 250 to 1000  $GeV$ . Both spin scenarios of the resonance are considered: spin 0 (later called "graviton") and spin 2 (later called "radion"). In the absence of the

evidence of the resonant double Higgs boson production from the previous searches, we set upper confidence limits. When combined with other search channels, this analysis will contribute to the discovery of the double Higgs production and we would be able to finally probe the Higgs boson potential using its self-coupling.

*“... a place for a smart quote!”*

*Lenin, 1922.*

## ACKNOWLEDGMENTS

This will be a longgggg list!

---

## Table of Contents

---

<b>List of Figures</b>	<b>vii</b>
<b>List of Tables</b>	<b>viii</b>
<b>1 Theoretical approach</b>	<b>1</b>
1.1 The Standard Model . . . . .	1
1.2 Beyond the Standard Model . . . . .	4
<b>Bibliography</b>	<b>9</b>
<b>References</b>	<b>10</b>

---

## List of Figures

---

1.1	SSB Potential form . . . . .	4
1.2	SM double Higgs boson production. . . . .	6
1.3	Left: comparison of the double Higgs boson mass distribution at the LO at 14 and 100 TeV center-of-mass energy, Right: the total SM HH cross section and the box and the triangular contributions ("box" and "tri" at the plots). "nl" denotes the contribution from the new non-linear $t\bar{t}HH$ Higgs interaction if this new coupling exists [15]. . . . .	7
1.4	BSM (WED) double Higgs boson production. . . . .	8
1.5	Double Higgs decay in the 2 b, 2 lepton, and 2 neutrino final state. . . .	9
1.6	Double Higgs decay channels according to the SM branching fractions. . .	9

---

## List of Tables

---



## CHAPTER 1

---

### Theoretical approach

---

## 1.1 The Standard Model

The SM uses the Lagrangian formalism to describe the interactions of elementary particles and fields. The SM Lagrangian can be split into four main terms [1]:

- Gauge bosons and their interactions
- Fermions and their interactions with the gauge bosons
- Higgs boson, its self-interaction, and interaction with the gauge bosons to give them mass, which is not possible solely by the  $\mathcal{L}_{YM}$
- Fermions and their interactions with the Higgs boson, which through the Yukawa mechanism gives mass to fermions

or equivalently:

$$\mathcal{L}_{SM} = \mathcal{L}_{YM} + \mathcal{L}_{ferm} + \mathcal{L}_H + \mathcal{L}_{Yuk} \quad (1.1)$$

The first term in the SM Lagrangian:

$$\mathcal{L}_{YM} = -\frac{1}{4}W_{\mu\nu}^i(x)W_i^{\mu\nu}(x) - \frac{1}{4}B_{\mu\nu}(x)B^{\mu\nu}(x) - \frac{1}{4}G_{\mu\nu}^a(x)G_a^{\mu\nu}(x) \quad (1.2)$$

where

$$B_{\mu\nu}(x) \equiv \partial_\mu B_\nu - \partial_\nu B_\mu \quad (1.3)$$

$$W_{\mu\nu}^i(x) \equiv \partial_\mu W_\nu^i(x) - \partial_\nu W_\mu^i(x) - g\varepsilon^{ijk}W_\mu^j W_\nu^k \quad (1.4)$$

$$G_{\mu\nu}^a(x) \equiv \partial_\mu G_\nu^a(x) - \partial_\nu G_\mu^a(x) - g_s f^{abc}G_\mu^b G_\nu^c \quad (1.5)$$

$$(1.6)$$

with  $i, j, k = 1, 2, 3$  and  $a, b, c = 1, \dots, 8$ . The fields have the following connections to their underlying symmetries:  $B_{\mu\nu}$  corresponds to  $U(1)$  symmetry of the weak hypercharge  $Y_k$  and "B" term is simply a kinematic term while "W" and "G" terms describe interactions among the corresponding bosons, where  $W_{\mu\nu}^i$  corresponds to  $SU(2)_I$  symmetry of the weak isospin  $I_w^i$ , and  $G_{\mu\nu}^a$  corresponds to  $SU(3)_c$  symmetry of the QCD color charge.  $g$  and  $\varepsilon$  are  $SU(2)$  coupling and structure constants, while  $g_s$  and  $f$  are coupling and structure constants for  $SU(3)$ .

The next term in the SM Lagrangian shows how fermions interact with the gauge bosons. Notice, that the mass terms are still absent:

$$\mathcal{L}_{ferm} = i\bar{\Psi}_L \not{D} \Psi_L + i\bar{\psi}_{lR} \not{D} \psi_{lR} + i\bar{\Psi}_Q \not{D} \Psi_Q + i\bar{\psi}_{uR} \not{D} \psi_{uR} + i\bar{\psi}_{dR} \not{D} \psi_{dR} \quad (1.7)$$

Above  $\Psi$  represents a doublet of a charged lepton and a corresponding neutral lepton within the same lepton family of  $SU(2)_L$ , a letter Q is reserved for a family of quarks, and  $\psi_R$  describes a right-handed leptonic singlet.

Gauge bosons interactions are present due to the derivative term:

$$D_\mu = \partial_\mu + igI_w^i W_\mu^i + ig'Y_w B_\mu + ig_s T_c^a G_\mu^a \quad (1.8)$$

$$(1.9)$$

Physical fields in this notation are represented by a linear combination of W and B fields:

$$A_\mu = B_\mu \cos \theta_W + W_\mu^3 \sin \theta_W \quad (1.10)$$

$$Z_\mu = -B_\mu \sin \theta_W + W_\mu^3 \cos \theta_W$$

where  $\theta_W$  is known as the *Weinberg angle* [2].


These two first terms of the SM Lagrangian is enough to have a theory of fermions and bosons, but they have no mass [3]. As discussed before, to ensure that weak bosons are massive, we need a Higgs term. Higgs mechanism enters the SM Lagrangian through the corresponding Higgs Lagrangian term given by

$$\mathcal{L}_H = (D_\mu \Phi)^\dagger (D^\mu \Phi) - V(\Phi), \quad V(\Phi) = -\mu^2 (\Phi^\dagger \Phi) + \frac{\lambda}{4} (\Phi^\dagger \Phi)^2 \quad (1.11)$$

where

$$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 = (v + H + i\chi)/\sqrt{2} \end{pmatrix} \quad \text{and} \quad v = 2\sqrt{\frac{\mu^2}{\lambda}} \quad (1.12)$$

Here  $\mu$  and  $\lambda$  are parameters of the Higgs potential. The discovery of the Higgs boson and the measurement of its mass studies the  $\mu$  parameter, double Higgs boson non-resonant searches target  $\lambda$  parameter to know more precisely what is the shape of Higgs potential. After the SSB, the value of the Higgs field vacuum expectation value can be expressed in term of  $\mu$  and  $\lambda$  and is usually denoted by  $v$  [4].



**Figure 1.1:** Shape of the Higgs potential before and after SSB that is determined at the leading orders by  $\mu$  and  $\lambda$  parameters.

I After adding  $\mathcal{L}_H$  and rearranging term, bosons have masses given by:

$$M_W = \frac{gv}{2}, \quad M_Z = \frac{M_W}{\cos \theta_W}, \quad M_H = \sqrt{2\mu^2} \quad (1.13)$$

The final piece of the SM Lagrangian is the Yukawa term, which Lagrangian is given by:

$$\mathcal{L}_{Yuk} = -i\bar{\Psi}_L G_l \psi_{l_R} \Phi - i\bar{\Psi}_Q G_u \psi_{u_R} \tilde{\Phi} - i\bar{\Psi}_Q G_d \psi_{d_R} \Phi + h.c. \quad (1.14)$$

where  $\tilde{\Phi} = i\sigma^2 \Phi^*$

The masses of fermions enter the equation through the  $3 \times 3$  matrices  $G$ , which are not known from the theory and are the parameters of the SM. The mass of the fermion is proportional to the Yukawa coupling of the corresponding fermion to the Higgs boson, which has already been mentioned when we discussed the Fig. ??.

## 1.2 Beyond the Standard Model

Several BSM theories [5–7] predict a resonant production of the double Higgs boson events through a heavy narrow width resonance ( $\sim O(1 - 10)$  GeV), which could be

spin 0 or spin 2 particle [8]. In this particular analysis data is compared with respect to predictions from the Warped Extra Dimensions theory (WED) [9]. WED theory addressing the hierarchy problem adds additional fifth dimension to the 4-dimensional (4D) space-time. In the framework that Randall and Sundrum (RS) [10] followed 4D space then is nothing but an EFT approximation, where the radion or graviton may exist as Kaluza-Klein (KK) [11] excitation modes at the TeV scale. Since LHC had provided us with no evidence of the SM particles interacting with the additional RS dimensions, it is postulated that they are confined to 3-brane, or a wall. At the same time, gravity, which is not in the SM, can propagate freely in the full higher-dimensional space, so-called bulk. If/when the bulk is compactified, it may produce KK modes of the gravitons. In this analysis RS model with parameter  $k$  of the order of Planck scale and  $\bar{M}_{Pl}$ , a reduced 4D  $M_{Pl}$  which is a function of the 5D Planck scale  $M$  and a parameter  $k$  with  $k < M$ , are assumed to satisfy the constraint  $0.01 \leq k/\bar{M}_{Pl} \leq 1$ , because values outside of this range are not applicable/or complicate the theory [12]. Considered in this search graviton and radion are thus RS KK graviton and RS radion particles that emerge in RS scenario with the excitation or a KK state mass of the order of TeV.

If we denote a part of the KK 5D wave function, often called a profile, as  $f_X^{(n)}(\phi)$ , where  $n$  is referred to the KK mode, then the graviton 4D profile wave-function can be formulated as  $h_{\mu\nu}^{(n)}(x_\mu)(f_X^{(n)}(\phi))$  and the zero-th mode of this function would correspond to the graviton that is a gravity mediator. Its effective mass is of the order of TeV. The Lagrangian describing the interaction of the graviton with the SM fields is given then by

$$\mathcal{L}_{graviton} = -\frac{x_1 \tilde{k}}{m_G} h^{\mu\nu(1)} \times d_i T_{\mu\nu}^{(i)}, \quad (1.15)$$


where  $T_{\mu\nu}^{(i)}$  is a 4D canonical energy-momentum tensor [13] for the SM field  $i$  and  $d_i$  is an integral of the profiles of the SM fields and KK gravitons.  $\tilde{k}$  is a free parameter inversely proportional to the Planck mass and varies from 0.01 to 1 when  $M_{graviton}$  is from 100 to 1500 GeV.

For radion the Lagrangian is similar and is given by:

$$\mathcal{L}_{radion} = -\frac{r}{\Lambda_R} \times a_i T_{\mu}^{\mu(i)}, \quad (1.16)$$


where  $\Lambda_R$  is a scale parameter proportional to the Planck mass and  $r$  is a 5D Radion field. If we make an assumption that the profiles of the graviton and radion are localised at the TeV scale, then the coupling of them to the massive SM fields is of the order of 1. Throughout this thesis, theory curves contain model results for the case  $\tilde{k} = 0.1$  and  $\Lambda_R = 3$  TeV.

In this analysis the gravitons/radions in the search are produced by a gluon fusion mechanism. Five Feynman diagrams describe this process, two of which are present in the SM (a "box" and a "triangular" diagrams on Fig. 1.2), and the other three are a BSM extension of the SM (BSM contact interaction diagrams on Fig. 1.4).




**Figure 1.2:** SM double Higgs boson production.

In the SM two main diagrams interfere destructively and the total cross section is thus lowered (Fig. 1.3 on the right). The box diagrams dominates the double Higgs boson production and peaks near 400 GeV. An extensive study has been performed by theorist for the future 100 TeV collider [?, 14]. However, since the kinematic distribution of the double Higgs mass remains unchanged to a high degree (see Fig. 1.3 on the left), we can extrapolate 100 TeV results to the current LHC machine.



**Figure 1.3:** Left: comparison of the double Higgs boson mass distribution at the LO at 14 and 100 TeV center-of-mass energy, Right: the total SM HH cross section and the box and the triangular contributions ("box" and "tri" at the plots). "nl" denotes the contribution from the new non-linear  $t\bar{t}HH$  Higgs interaction if this new coupling exists [15].


It is also interesting to measure BSM contact interaction couplings and a future non-resonant version of this analysis will target that. In this case,  $c_2$ , the coupling of two heavy quarks with two Higgs bosons,  $c_{2g}$ , the coupling of two gluons with two Higgs bosons, and  $c_g$ , the direct coupling of the gluons to the Higgs boson will be studied (see Fig. 1.4).



**Figure 1.4:** BSM (WED) double Higgs boson production.


Now it is time to discuss the decay of the double Higgs system. This analysis considers separately graviton and radion decays into two SM Higgs bosons with the subsequent decays of the Higgs boson to W or Z boson pairs. W bosons are allowed to decay only leptonically. For Z boson decays, the signature is characterised by the on-shell Z boson decaying into a lepton pair and an off-shell Z boson decaying to invisible (see Fig. 1.5).





**Figure 1.5:** Double Higgs decay in the 2 b, 2 lepton, and 2 neutrino final state.

All the decay channels of the double Higgs system to the SM particles are summarised in the Fig. 1.6. Combined branching fraction of the double Higgs boson decay in the two b quarks, two leptons, and two neutrinos final state is approximately 2.78%.



**Figure 1.6:** Double Higgs decay channels according to the SM branching fractions.

---

## References

---

- [1] Matthias U. Mozer. Electroweak Physics at the LHC. *Springer Tracts Mod. Phys.*, 267:1–115, 2016.
- [2] Steven Weinberg. The Making of the Standard Model. *Eur. Phys. J. C*, 34(hep-ph/0401010):5–13. 21 p. ; streaming video, 2003.
- [3] Roger Wolf. *The Higgs Boson Discovery at the Large Hadron Collider*, volume 264. Springer, 2015.
- [4] Jose Andres Monroy Montanez, Kenneth Bloom, and Aaron Dominguez. Search for production of a Higgs boson and a single Top quark in multilepton final states in pp collisions at  $\sqrt{s} = 13$  TeV, Jul 2018. Presented 23 Jul 2018.
- [5] Peisi Huang, Aniket Joglekar, Min Li, and Carlos E. M. Wagner. Corrections to di-Higgs boson production with light stops and modified Higgs couplings. *Phys. Rev.*, D97(7):075001, 2018.
- [6] Matthew J. Dolan, Christoph Englert, and Michael Spannowsky. New Physics in LHC Higgs boson pair production. *Phys. Rev.*, D87(5):055002, 2013.

- [7] Shinya Kanemura, Kunio Kaneta, Naoki Machida, Shinya Odori, and Tetsuo Shindou. Single and double production of the Higgs boson at hadron and lepton colliders in minimal composite Higgs models. *Phys. Rev.*, D94(1):015028, 2016.
- [8] Albert M Sirunyan et al. Search for Higgs boson pair production in the  $\gamma\gamma b\bar{b}$  final state in pp collisions at  $\sqrt{s} = 13$  TeV. 2018.
- [9] Alexandra Oliveira. Gravity particles from Warped Extra Dimensions, predictions for LHC. 2014.
- [10] Lisa Randall and Raman Sundrum. A Large mass hierarchy from a small extra dimension. *Phys. Rev. Lett.*, 83:3370–3373, 1999.
- [11] Kunihiro Uzawa, Yoshiyuki Morisawa, and Shinji Mukohyama. Excitation of Kaluza-Klein gravitational mode. *Phys. Rev.*, D62:064011, 2000.
- [12] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo. Phenomenology of the Randall-Sundrum Gauge Hierarchy Model. *Phys. Rev. Lett.*, 84:2080, 2000.
- [13] Michael Forger and Hartmann Romer. Currents and the energy momentum tensor in classical field theory: A Fresh look at an old problem. *Annals Phys.*, 309:306–389, 2004.
- [14] Chuan-Ren Chen and Ian Low. Double take on new physics in double Higgs boson production. *Phys. Rev.*, D90(1):013018, 2014.
- [15] Roberto Contino, Margherita Ghezzi, Mauro Moretti, Giuliano Panico, Fulvio Piccinini, and Andrea Wulzer. Anomalous Couplings in Double Higgs Production. *JHEP*, 08:154, 2012.

- [16] Richard Phillips Feynman, Robert Benjamin Leighton, and Matthew Sands. *The Feynman lectures on physics; New millennium ed.* Basic Books, New York, NY, 2010. Originally published 1963-1965.
- [17] Savas Dimopoulos, Stuart Raby, and Frank Wilczek. Proton decay in supersymmetric models. *Physics Letters B*, 112(2):133 – 136, 1982.
- [18] David J Griffiths. *Introduction to elementary particles; 2nd rev. version.* Physics textbook. Wiley, New York, NY, 2008.
- [19] M. Della Negra, P. Jenni, and T. S. Virdee. Journey in the search for the higgs boson: The atlas and cms experiments at the large hadron collider. *Science*, 338(6114):1560–1568, 2012.
- [20] Jennifer Ouellette. Einstein’s quest for a unified theory. *APS*, 2015.
- [21] E A Davis and Isabel Falconer. *J.J. Thompson and the discovery of the electron.* Taylor and Francis, Hoboken, NJ, 2002.
- [22] Oreste Piccioni. *The Discovery of the Muon*, pages 143–162. Springer US, Boston, MA, 1996.
- [23] G. Danby, J-M. Gaillard, K. Goulianos, L. M. Lederman, N. Mistry, M. Schwartz, and J. Steinberger. Observation of high-energy neutrino reactions and the existence of two kinds of neutrinos. *Phys. Rev. Lett.*, 9:36–44, Jul 1962.
- [24] M. L. Perl, G. S. Abrams, and et al Boyarski. Evidence for anomalous lepton production in  $e^+ - e^-$  annihilation. *Phys. Rev. Lett.*, 35:1489–1492, Dec 1975.
- [25] K. Kodama et al. Observation of tau neutrino interactions. *Phys. Lett.*, B504:218–224, 2001.

- [26] S. M. Bilenky. Neutrino in Standard Model and beyond. *Phys. Part. Nucl.*, 46(4):475–496, 2015.
- [27] S Chandrasekhar. *Newton's principia for the common reader*. Oxford Univ., Oxford, 2003. The book can be consulted by contacting: PH-AID: Wallet, Lionel.
- [28] Hanoch Gutfreund and Jürgen Renn. *The road to relativity: the history and meaning of Einstein's "The foundation of general relativity" : featuring the original manuscript of Einstein's masterpiece*. Princeton University Press, Princeton, NJ, Apr 2015.
- [29] J. Butterworth. *Smashing Physics*. Headline Publishing Group, 2014.
- [30] W N Cottingham and D A Greenwood. *An Introduction to the Standard Model of Particle Physics; 2nd ed.* Cambridge Univ. Press, Cambridge, 2007.
- [31] Eric W. Weisstein. Fundamental forces.
- [32] Carl Bender. Mathematical physics.
- [33] Andrew Wayne. QED and the Men Who Made It: Dyson, Feynman, Schwinger, and Tomonaga by Silvan S. Schweber. *The British Journal for the Philosophy of Science*, 46(4):624–627, 1995.
- [34] C. Patrignani et al. Review of Particle Physics. *Chin. Phys.*, C40(10):100001, 2016.
- [35] Michelangelo L Mangano. Introduction to QCD. (CERN-OPEN-2000-255), 1999.
- [36] Matt Strassler. Of particular significance: Conversations about science with theoretical physicist matt strassler.

- [37] S. L. Glashow. Partial Symmetries of Weak Interactions. *Nucl. Phys.*, 22:579–588, 1961.
- [38] F. Englert and R. Brout. Broken symmetry and the mass of gauge vector mesons. *Phys. Rev. Lett.*, 13:321–323, Aug 1964.
- [39] Peter W. Higgs. Broken symmetries and the masses of gauge bosons. *Phys. Rev. Lett.*, 13:508–509, Oct 1964.
- [40] G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble. Global conservation laws and massless particles. *Phys. Rev. Lett.*, 13:585–587, Nov 1964.
- [41] Pauline Gagnon. *Who cares about particle physics? : making sense of the Higgs boson, the Large Hadron Collider and CERN*. Oxford University Press, 2016.