Effat College of Engineering · Computer Science

CS3081 Artificial Intelligence
Assignment: Uncertainty and Bayesian Networks

Instructor: Dr. Naila Marir **Total:** 50 marks (5 exercises × 10 marks)

Name: *Remas Rawas* Student ID: *S24109161*

Instructions

1. Show all your steps. If you write only the final number, you get at most half of the marks.
2. Write probabilities with 4 decimal places (for example 0.5367), unless the question says something else.
3. Exercises 1 to 4, and Exercise 5 parts (a) and (b), are solved by hand.
4. Exercise 5 part (c) is solved in Python with pgmpy in VS Code, like in the lab.
5. Submit two files on Blackboard: these pages with your answers written in the boxes, scanned as one PDF, and your notebook `assignment_bn.ipynb` with all outputs.
6. This is individual work. You can discuss ideas with classmates, but you must write your own answers.

Exercise 1 — Joint distribution (10 marks)

The Effat shuttle bus is either **on time** or **late**. The weather in Jeddah is either **clear** or **dusty**. The table gives the probability of each pair. For example, the probability that the weather is clear **and** the bus is late is 0.14.

Weather	Shuttle = on time	Shuttle = late
clear	0.56	0.14
dusty	0.12	0.18

1. **(2 marks)** Show that the four numbers in the table make a valid probability distribution.

All four probabilities are between 0 and 1.
0.56 + 0.14 + 0.12 + 0.18 = 1.0000
So this is a valid probability distribution.

2. **(2 marks)** Find P(dusty), the probability that the weather is dusty. Then find P(late), the probability that the bus is late.

P(dusty) = 0.12 + 0.18 = 0.3000
P(late) = 0.14 + 0.18 = 0.3200

3. **(2 marks)** Find P(late | dusty) and P(late | clear).

P(late | dusty) = 0.18 / 0.30 = 0.6000
P(late | clear) = 0.14 / (0.56 + 0.14) = 0.2000

4. **(2 marks)** Find P(dusty | late). Is it the same as P(late | dusty)? Explain your answer in one sentence.

P(dusty | late) = 0.18 / 0.32 = 0.5625
They are different because one divides by P(late) and the other divides by P(dusty).

5. **(2 marks)** Are the weather and the bus independent? Use numbers to explain your answer.

The weather and the bus are dependent.
P(late | dusty) = 0.6000, but P(late) = 0.3200.
The chance of being late changes when we know it is dusty.

Exercise 2 — Bayes' rule: a spam filter (10 marks)

An email program wants to find spam emails. It knows these facts:

- 30% of all emails are spam. So P(spam) = 0.3 and P(normal) = 0.7.
- In spam emails, the word "free" appears 60% of the time: P(free | spam) = 0.6.
- In normal emails, the word "free" appears 5% of the time: P(free | normal) = 0.05.

1. **(3 marks)** Find P(free), the probability that any email contains the word "free". Write the name of the rule you use.

I used the law of total probability.
P(free) = P(free | spam) P(spam) + P(free | normal) P(normal)
= (0.60 × 0.30) + (0.05 × 0.70)
= 0.18 + 0.035 = 0.2150

2. **(3 marks)** A new email contains the word "free". Use Bayes' rule to find P(spam | free).

By Bayes' rule,
P(spam | free) = P(free | spam) P(spam) / P(free)
= (0.60 x 0.30) / 0.215 = 0.8372

3. **(3 marks)** The program also checks the word "winner": P(winner | spam) = 0.4 and P(winner | normal) = 0.02. Assume that, once we know if an email is spam or normal, the two words appear independently of each other. A new email contains **both** "free" and "winner". Find P(spam | free, winner).

Spam with both words = 0.30 x 0.60 x 0.40 = 0.0720
Normal with both words = 0.70 x 0.05 x 0.02 = 0.0007
P(both words) = 0.0720 + 0.0007 = 0.0727
P(spam | free, winner) = 0.0720 / 0.0727 = 0.9904

4. **(1 mark)** Compare your answers to parts 2 and 3. Did the probability of spam go up or down when we added the word "winner"? Explain why in one sentence.

It went up from 0.8372 to 0.9904 because "winner" is more common in spam emails than in normal emails.

Exercise 3 — Bayesian network: structure and joint probability (10 marks)

In Jeddah summers, a **heatwave** (H) can cause an **AC overload** (A) in a campus building. A heatwave and an AC overload can both cause a **power cut** (C). A power cut can cause the **lab session to be cancelled** (L). Each variable is yes or no.

The network has 4 arrows: H → A, H → C, A → C, and C → L. The tables give only the probability of "yes". The probability of "no" is 1 minus the probability of "yes".


P(H = yes)
0.3

H	P(A = yes | H)
yes	0.6
no	0.1

H	A	P(C = yes | H, A)
yes	yes	0.7
yes	no	0.2
no	yes	0.5
no	no	0.05

C	P(L = yes | C)
yes	0.8
no	0.1

1. **(2 marks)** Draw the network. For each node, write its parents.



H has no parents. A's parent is H.
C's parents are H and A. L's parent is C.

2. **(2 marks)** Write P(H, A, C, L) as a product of the probabilities from the network's tables.

$P(H, A, C, L) = P(H) \times P(A \mid H) \times P(C \mid H, A) \times P(L \mid C)$

3. **(2 marks)** A full joint table for 4 yes/no variables needs 15 numbers ($2^4 - 1$). How many numbers do the 4 tables of this network need? Count only the "yes" probabilities, and show how you counted.

H has 1 entry and A has 2 (one per value of H).
C has 4 entries (all H, A pairs) and L has 2 (one per value of C).
Total = 1 + 2 + 4 + 2 = 9 numbers.

4. **(2 marks)** Find P(H = yes, A = no, C = yes, L = yes).

$P(A = no \mid H = yes) = 1 - 0.60 = 0.4000$
$P(H=yes, A=no, C=yes, L=yes) = 0.30 \times 0.40 \times 0.20 \times 0.80$
$= 0.0192$

5. **(2 marks)** Find P(H = yes | C = yes). Then give P(H = yes | C = yes, L = yes) without doing a new calculation, and explain why the two answers are the same.

$P(H=yes, C=yes) = 0.30 \times (0.60 \times 0.70 + 0.40 \times 0.20) = 0.1500$
$P(C=yes) = 0.15 + 0.70 \times (0.10 \times 0.50 + 0.90 \times 0.05) = 0.2165$
$P(H=yes \mid C=yes) = 0.15 / 0.2165 = 0.6928$
$P(H=yes \mid C=yes, L=yes) = 0.6928$ too.
Once we know C, knowing L gives no extra information about H.

Exercise 4 — Inference by enumeration, challenge question (10 marks)

This is the **challenge question** of the assignment. Show every step.

Use the network and tables from Exercise 3. Today the lab session **was cancelled** (L = yes).

1. **(4 marks)** Use inference by enumeration to find P(H = yes | L = yes), the probability that there was a heatwave. Write the query variable, the evidence and the hidden variables, then show all sums and the normalization.

   Query: H. Evidence: L = yes. Hidden variables: A and C.
   For each H and A pair, I add the C=yes and C=no cases.
   H=yes, A=yes: 0.30 x 0.60 x (0.70 x 0.80 + 0.30 x 0.10) = 0.1062
   H=yes, A=no: 0.30 x 0.40 x (0.20 x 0.80 + 0.80 x 0.10) = 0.0288
   H=no, A=yes: 0.70 x 0.10 x (0.50 x 0.80 + 0.50 x 0.10) = 0.0315
   H=no, A=no: 0.70 x 0.90 x (0.05 x 0.80 + 0.95 x 0.10) = 0.08505
   P(H=yes, L=yes) = 0.1062 + 0.0288 = 0.1350
   P(H=no, L=yes) = 0.0315 + 0.08505 = 0.11655
   P(L=yes) = 0.1350 + 0.11655 = 0.25155
   Normalize: P(H=yes | L=yes) = 0.1350 / 0.25155 = 0.5367
   I kept extra digits in the working to avoid rounding early.

2. **(3 marks)** Now find P(A = yes | L = yes), the probability that there was an AC overload. Which variables are hidden this time?

   Query: A. Evidence: L = yes. Hidden variables: H and C.
   I use the same sums over C from part 1 and group them by A.
   P(A=yes, L=yes) = 0.1062 + 0.0315 = 0.1377
   P(A=no, L=yes) = 0.0288 + 0.08505 = 0.11385
   P(L=yes) = 0.1377 + 0.11385 = 0.25155
   P(A=yes | L=yes) = 0.1377 / 0.25155 = 0.5474

3. **(3 marks)** A technician checks the building and reports that there was **no** AC overload (A = no). The lab is still cancelled (L = yes). Find P(H = yes | L = yes, A = no). Compare it with your answer to part 1: did the probability of a heatwave go up or down? Use the network to explain why in two or three sentences.

Now A = no is evidence, so I only use those two cases from part 1.
H=yes: 0.30 x 0.40 x (0.20 x 0.80 + 0.80 x 0.10) = 0.0288
H=no: 0.70 x 0.90 x (0.05 x 0.80 + 0.95 x 0.10) = 0.08505
P(H=yes | L=yes, A=no) = 0.0288 / (0.0288 + 0.08505)
= 0.2530
It went down from 0.5367 to 0.2530. A heatwave makes an overload more likely, so seeing no overload is evidence against a heatwave.
A heatwave can still cause a power cut directly.

Exercise 5 — Approximate inference by sampling (10 marks)

Now we find the same probability, P(H = yes | L = yes), by **sampling**. Sampling gives an estimate, not the exact answer.

(a) Rejection sampling (3 marks). Reminder: in rejection sampling, we keep only the samples that agree with the evidence (here L = yes), and we throw away the others. Below are 12 samples from the network of Exercise 3.

#	H	A	C	L
1	no	no	no	no
2	yes	yes	yes	yes
3	no	no	no	no
4	no	yes	yes	yes
5	yes	yes	no	no
6	no	no	no	yes
7	yes	no	yes	yes
8	no	no	no	no
9	no	no	no	no
10	yes	yes	yes	no
11	no	no	no	no
12	no	no	yes	yes

1. Write the numbers of the samples we keep and the samples we throw away. Then estimate P(H = yes | L = yes).

Keep: 2, 4, 6, 7, 12 (all have L = yes).
Throw away: 1, 3, 5, 8, 9, 10, 11.
Only 2 and 7 have H = yes, so the estimate is 2/5 = 0.4000.

2. What fraction of the 12 samples did we throw away? Why is this a problem for rejection sampling?

We threw away 7/12 = 0.5833 of the samples.
This leaves fewer samples to estimate the probability.
If the evidence is rare, most samples get thrown away.

(b) Likelihood weighting (3 marks). Reminder: in likelihood weighting, we do not sample the evidence. We fix it (here L = yes), we sample the other variables, and we give each sample a **weight**. The weight is the probability of the evidence given the sample's values: here, P(L = yes | C). Below are 6 samples.

#	H	A	C	L
1	no	no	no	yes (fixed)
2	yes	yes	yes	yes (fixed)
3	no	no	no	yes (fixed)
4	yes	no	no	yes (fixed)
5	no	yes	yes	yes (fixed)
6	yes	yes	yes	yes (fixed)

1. Write the weight of each sample.

If C = yes, the weight is 0.8000. If C = no, it is 0.1000.
1: 0.1000 2: 0.8000 3: 0.1000
4: 0.1000 5: 0.8000 6: 0.8000

2. Estimate P(H = yes | L = yes) using the weights: add the weights of the samples where H = yes, then divide by the sum of all weights.

Weight for H = yes (samples 2, 4, 6) = 0.8 + 0.1 + 0.8 = 1.7
Total weight = 0.1 + 0.8 + 0.1 + 0.1 + 0.8 + 0.8 = 2.7
Estimate = 1.7 / 2.7 = 0.6296

3. Compare your estimates from (a) and (b) with the exact answer from Exercise 4. Why are they different? How can we make them closer to the exact answer?

Rejection sampling gave 0.4000, below the exact answer 0.5367.
Likelihood weighting gave 0.6296, above the exact answer.
The estimates vary because we used small random samples.
Using more samples usually brings them closer to the exact answer.

(c) Python with pgmpy (4 marks). Answer this part in your notebook, not on paper. In VS Code, create `assignment_bn.ipynb`.

1. Build the network of Exercise 3 with `DiscreteBayesianNetwork` and `TabularCPD`, like in the lab. Name the nodes `heatwave`, `overload`, `powercut` and `cancel`, and use the values `yes` and `no`. Print `model.check_model()`. It must print `True`.
2. Use `VariableElimination` to find P(H | L = yes). Your answer must be the same as in Exercise 4.
3. Use rejection sampling with 10,000 samples to estimate P(H = yes | L = yes). Hint: import `BayesianModelSampling` from `pgmpy.sampling` and `State` from `pgmpy.factors.discrete`. Then use `BayesianModelSampling(model).rejection_sample(evidence=[State("cancel", "yes")], size=10000)`. This gives a table of samples. Count the fraction of rows where `heatwave` is `yes`.
4. In a Markdown cell, write two sentences that compare the exact answer and the sampling estimate.