

Advanced Dynamics Notes

Jed Rembold

September 24, 2009

Contents

1	Survey of Elementary Particles	5
1.1	Mechanics of a Particle	5
1.2	Mechanics of a System of Particles	7
1.3	Constraints	11
1.4	D'Alembert's Principle and Lagrange's Equations	13
1.5	Velocity Dependent Potentials and Dissipation Functions	15
1.6	Simple Applications of Lagrangian Formalism	15
2	Variational Principles and Lagrange's Equations	19
2.1	Hamilton's Principle and Calculus of Variations	19
2.2	The 3 Classic Problems of the Calculus of Variations	21
2.3	Calculus of Variations and Lagrange's Equations	21
2.4	Non-Holonomic Systems and Lagrange Multipliers	24
2.5	Conservation Theorems and Symmetry Properties	29
2.5.1	Energy Conservation and Time Homogeneity	30
2.5.2	Momentum Conservation and Space Homogeneity	31
2.5.3	Angular Momentum Conservation and Rotational Invariance	32
3	The Central Force Problem	35
3.1	2 body to 1 body problem and Reduced Mass	35

Chapter 1

Survey of Elementary Particles

1.1 Mechanics of a Particle

We begin with a review of common expressions. Recall that

\mathbf{r} = position vector of a particle from a given origin
 \mathbf{v} = particle's vector velocity

We thus know that

$$\mathbf{v} = \frac{d\mathbf{r}}{dt} \quad \text{and} \quad \mathbf{p} = m\mathbf{v}$$

where

\mathbf{p} = linear momentum of a particle
 m = the mass of a particle
 \mathbf{F} = total force exerted on the particle

The mechanics or motion of the particle are given by Newton's 2nd Law of Motion:

$$\mathbf{F} = \frac{d\mathbf{p}}{dt} = \dot{\mathbf{p}}$$

For an inertial or Galilean reference frame, this can be expressed as

$$\mathbf{F} = \frac{d}{dt}(m\mathbf{v})$$

For a particle with constant mass, this simplifies to:

$$\mathbf{F} = m\mathbf{a} = m\frac{d\mathbf{v}}{dt} = m\frac{d^2\mathbf{r}}{dt^2}$$


where \mathbf{a} is the particle's acceleration. Now, if $\mathbf{F} = 0 = \dot{\mathbf{p}}$, then \mathbf{p} equals a constant, and thus we have Conservation of Linear Momentum of the particle.

We also know that

\mathbf{L} = angular momentum about a point O

and that

$$\mathbf{L} = \mathbf{r} \times \mathbf{p}$$



The change of the angular momentum with respect to time is given by

$$\begin{aligned} \frac{d}{dt}\mathbf{L} &= \frac{d}{dt}(\mathbf{r} \times \mathbf{p}) = \frac{d}{dt}(\mathbf{r} \times m\mathbf{v}) \\ &= \underbrace{\frac{d\mathbf{r}}{dt} \times m\mathbf{v}}_0 + \mathbf{r} \times \frac{d}{dt}(m\mathbf{v}) \\ &= \mathbf{r} \times \mathbf{F} \\ &= \mathbf{N} \end{aligned}$$

which is called the moment of force or, more commonly, the torque. Note here that \mathbf{L} and \mathbf{N} depend on O, the point about which the moments were taken. Now, if $\mathbf{N} = \dot{\mathbf{L}} = 0$, then \mathbf{L} is a constant, and thus we have Conservation of Angular Momentum.


The Work done on a particle by force \mathbf{F} is

$$W_{AB} = \int_A^B \mathbf{F} \cdot d\mathbf{s}$$

where \mathbf{s} is the trajectory. Assuming that the mass is constant:

$$\begin{aligned} W_{AB} &= m \int_A^B \frac{d\mathbf{v}}{dt} \cdot \mathbf{v} dt \\ &= \frac{m}{2} \int_A^B \frac{d}{dt}(v^2) \\ &= \frac{m}{2} (v_B^2 - v_A^2) \\ &\equiv T_B - T_A \end{aligned}$$

where $T = \frac{1}{2}mv^2$ is the kinetic energy. Thus we see evidence of the Work-Energy Theorem, which states that the work done is equal to the change in kinetic energy. Now, if W_{AB} depends only on the endpoints A and B (i.e. is path independent), then the force responsible is conservative.



Thus we see that, in a conservative system, the work around a closed path is

$$\oint \mathbf{F} \cdot d\mathbf{s} = 0$$

Remark 1: Dissipative forces such as friction are *not* conservative since

$$\mathbf{F} \cdot d\mathbf{s} > 0$$

for work done *by* the particle.

For a conservative force (W_{AB} path independent), we can write

$$\mathbf{F} = -\nabla V(\mathbf{r})$$

where V is the potential or potential energy. Thus we have

$$\mathbf{F} \cdot d\mathbf{s} = -dV \quad \Rightarrow \quad \mathbf{F}_s = -\frac{dV}{ds}$$

Remark 2: Note that

$$\mathbf{F} = -\nabla (V(\mathbf{r}) + \text{constant}) = -\nabla V(\mathbf{r})$$

and thus the zero of $V(\mathbf{r})$ is arbitrary.

Remark 3: For a conservative system, $W_{AB} = V_A - V_B$. Thus we have that

$$V_A - V_B = T_B - T_A \quad \Rightarrow \quad V_A + T_A = V_B + T_B$$

which is our expression for the Conservation of Total Energy!

1.2 Mechanics of a System of Particles

The goal here is to generalize Newton's 2nd law to a system of particles. Starting with the equation of motion for the i^{th} particle:

$$\sum_j \mathbf{F}_{ji} + \mathbf{F}_i^{(e)} = \dot{\mathbf{p}}_i$$

where

\mathbf{F}_{ji} = the internal force on particle i by particle j and

$\mathbf{F}_i^{(e)}$ = an external force

Now, assuming that $\mathbf{F}_{ij} = -\mathbf{F}_{ji}$ (Newton's 3rd Law, or the Weak Law of Action and Reaction), then

$$\sum_i \mathbf{F}_i = \frac{d^2}{dt^2} \sum_i m_i \mathbf{r}_i = \sum_i \mathbf{F}_i^{(e)} + \underbrace{\sum_{i \neq j} \mathbf{F}_{ji}}_0$$

We define \mathbf{R} as:

$$\mathbf{R} = \frac{\sum_i m_i \mathbf{r}_i}{\sum_i m_i} = \frac{\sum_i m_i \mathbf{r}_i}{M}$$

where \mathbf{R} is the center of mass and M is the total mass. Thus we have that

$$M \frac{d^2}{dt^2} \mathbf{R} = \sum_i \mathbf{F}_i^{(e)} \equiv \mathbf{F}^{(e)}$$

Consequently, purely internal forces (forces on particles from other particles) vanish by Newton's 3rd law. The total linear momentum can thus be expressed as

$$\mathbf{P} = \sum_i m_i \frac{d\mathbf{r}_i}{dt} = M \frac{d\mathbf{R}}{dt}$$

Thus total linear momentum is conserved when the total external forces are equal to zero.

The total angular momentum is written as

$$\mathbf{L} = \sum_i \mathbf{L}_i = \sum_i \mathbf{r}_i \times \mathbf{p}_i$$

Hence,

$$\begin{aligned} \frac{d\mathbf{L}}{dt} &= \dot{\mathbf{L}} = \sum_i \frac{d}{dt} (\mathbf{r}_i \times \mathbf{p}_i) \\ &= \sum_i \mathbf{r}_i \times \dot{\mathbf{p}}_i \\ &= \sum_i \mathbf{r}_i \times \mathbf{F}_i^{(e)} + \sum_{i \neq j} \mathbf{r}_i \times \mathbf{F}_{ji} \end{aligned}$$

but

$$\begin{aligned} \sum_{i \neq j} \mathbf{r}_i \times \mathbf{F}_{ji} &= \frac{1}{2} \sum_{i \neq j} [\mathbf{r}_i \times \mathbf{F}_{ji} + \mathbf{r}_j \times \mathbf{F}_{ij}] \\ &= \frac{1}{2} \sum_{i \neq j} (\mathbf{r}_i - \mathbf{r}_j) \times \mathbf{F}_{ji} \end{aligned}$$

If we define

$$\mathbf{r}_{ij} = \mathbf{r}_i - \mathbf{r}_j$$

then we can write

$$\sum_{i \neq j} \mathbf{r}_i \times \mathbf{F}_{ji} = \frac{1}{2} \sum_{i \neq j} \mathbf{r}_{ij} \times \mathbf{F}_{ji}$$


Note that if \mathbf{F}_{ji} is parallel to \mathbf{r}_{ij} , then $\mathbf{r}_{ij} \times \mathbf{F}_{ji} = 0$, which is called the Strong Law of Action and Reaction. If this is true, then

$$\frac{d\mathbf{L}}{dt} = \sum_i \mathbf{r}_i \times \mathbf{F}_i^{(e)} + \underbrace{\frac{1}{2} \sum_{i \neq j} \mathbf{r}_{ij} \times \mathbf{F}_{ji}}_0 = \mathbf{N}^{(e)}$$

Thus, if the applied external torque, $\mathbf{N}^{(e)}$, equals 0, then \mathbf{L} is constant in time and we have Conservation of Angular Momentum.

Remark 1: The Strong Law of Action and Reaction requires the internal forces to be central.

We can also describe the particle position with respect to the center of mass:



So we have

$$\begin{aligned} \mathbf{r}_i &= \mathbf{R} + \mathbf{r}'_i \\ \mathbf{v}_i &= \mathbf{V} + \mathbf{v}'_i \end{aligned}$$

where

$$\begin{aligned} \mathbf{V} &= \frac{d\mathbf{R}}{dt} = \text{velocity of center of mass} \\ \mathbf{v}'_i &= \frac{d\mathbf{r}'_i}{dt} = \text{velocity about the center of mass} \end{aligned}$$

Then

$$\begin{aligned}
 \mathbf{L} &= \sum_i \mathbf{r}_i \times \mathbf{p}_i = \sum_i (\mathbf{R} + \mathbf{r}'_i) \times (m_i \mathbf{V} + m_i \mathbf{v}'_i) \\
 &= \sum_i \mathbf{R} \times m_i \mathbf{V} + \sum_i \mathbf{R} \times \underbrace{m_i \mathbf{v}'_i}_0 + \sum_i \underbrace{\mathbf{r}'_i \times m_i \mathbf{V}}_0 + \sum_i \mathbf{r}'_i \times m_i \mathbf{v}'_i \\
 &= \mathbf{R} \times M \mathbf{V} + \sum_i \mathbf{r}'_i \times m_i \mathbf{v}'_i
 \end{aligned}$$

So the total angular momentum is:

$$\begin{aligned}
 \mathbf{L} &= \mathbf{R} \times M \mathbf{V} + \sum_i \mathbf{r}'_i \times m_i \mathbf{v}'_i \\
 &= \text{angular momentum of the CM} + \text{angular momentum about the CM}
 \end{aligned}$$

Now,

$$\begin{aligned}
 W_{AB} &= \sum_i \int_A^B \mathbf{F}_i \cdot d\mathbf{s}_i = \sum_i \int_A^B \mathbf{F}_i^{(e)} \cdot d\mathbf{s}_i + \sum_{i \neq j} \int_A^B \mathbf{F}_{ji} \cdot \mathbf{s}_i \\
 &= \sum_i \int_A^B m_i \dot{\mathbf{v}}_i \cdot \mathbf{v}_i dt \\
 &= \sum_i \int_A^B d\left(\frac{1}{2} m_i v_i^2\right) \\
 &= T_B - T_A
 \end{aligned}$$

with total kinetic energy $T = \frac{1}{2} \sum_i m_i v_i^2$. We can also write

$$\begin{aligned}
 T &= \frac{1}{2} \sum_i m_i v_i^2 = \frac{1}{2} \sum_i m_i (\mathbf{V} + \mathbf{v}'_i) \cdot (\mathbf{V} + \mathbf{v}'_i) \\
 &= \frac{1}{2} M V^2 + \sum_i \underbrace{m_i \mathbf{v}'_i \cdot \mathbf{V}}_0 + \frac{1}{2} \sum_i m_i \mathbf{v}'_i{}^2 \\
 &= \frac{1}{2} M V^2 + \frac{1}{2} \sum_i m_i \mathbf{v}'_i{}^2 \\
 &= \text{KE of CM} + \text{KE about CM}
 \end{aligned}$$

We now go back to

$$W_{AB} = \sum_i \int_A^B \mathbf{F}_i^{(e)} \cdot d\mathbf{s}_i + \sum_{i \neq j} \int_A^B \mathbf{F}_{ji} \cdot \mathbf{s}_i$$

For conservative *external* forces, we have that:

$$\sum_i \int_A^B \mathbf{F}_i^{(e)} \cdot d\mathbf{s}_i = - \sum_i \int_A^B \nabla_i V_i \cdot d\mathbf{s}_i = - \sum_i V_i \Big|_A^B$$

For conservative *internal* forces:

$$\mathbf{F}_{ji} = -\nabla_j V_{ji}$$

where $V_{ji} = V_{ji}(|\mathbf{r}_i - \mathbf{r}_j|)$ to satisfy the Strong Law of Action and Reaction

$$\begin{aligned} &= \nabla_i V_{ij} \\ &= -\mathbf{F}_{ij} \end{aligned}$$

where there is no implied sum over repeated indices above. Therefore

$$\begin{aligned} W_{AB} &= - \sum_i V_i \Big|_A^B + \frac{1}{2} \sum_{i \neq j} \int_A^B (\mathbf{F}_{ji} \cdot d\mathbf{s}_i + \mathbf{F}_{ij} \cdot d\mathbf{s}_j) \\ &= - \sum_i V_i \Big|_A^B + \frac{1}{2} \sum_{i \neq j} \int_A^B \mathbf{F}_{ji} \cdot (d\mathbf{s}_i - d\mathbf{s}_j) \\ &= - \sum_i V_i \Big|_A^B + \frac{1}{2} \sum_{i \neq j} \int_A^B \mathbf{F}_{ji} \cdot d\mathbf{s}_{ij} \\ &= - \sum_i V_i \Big|_A^B - \frac{1}{2} \sum_{i \neq j} \int_A^B \nabla_{ji} V_{ji} \cdot d\mathbf{s}_{ij} \\ &= - \sum_i V_i \Big|_A^B - \frac{1}{2} \sum_{i \neq j} V_{ji} \Big|_A^B \end{aligned}$$

Thus the total potential energy is

$$V = \sum_i V_i - \frac{1}{2} \sum_{i \neq j} V_{ij}$$

Remark 2: $T + V$ is conserved for T, V the total kinetic and potential energy.

Remark 3: For a rigid body, internal potential energy is constant, and thus internal forces do no work.

1.3 Constraints


Constraints limit the motion of a system: ie, beads on a string, ball on a circular track, or gas molecules in a container.

Holonomic Constraints: A Holonomic constraint is one where the constraint can be expressed as an equation connecting the space and time coordinates of a particle with the form:

$$f(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \dots, \mathbf{r}_N, t) = 0$$

Examples include rigid bodies (which have equations of the form $(\mathbf{r}_i - \mathbf{r}_j)^2 - C_{ij}^2 = 0$) and particles constrained to move on a curve or surface.

Non-Holonomic Constraints: A Non-Holonomic Constraint is one where the constraint can not be expressed as above. Examples include gas molecules in a box or a particle placed on a sphere surface (but *not* stuck to it)



Rheonomous Constraints: A Rheonomous constraint is one which depends explicitly on time

Scleronomous Constraints: A Scleronomous constraint is one which has no explicit time dependence.

Remark 1: Constraints are equivalent to saying that there are forces that can not be specified explicitly; only by their effect on the systems motion.

Now we will consider Holonomic constraints and introduce generalized coordinates. A system of N particles has $d \cdot N$ degrees of freedom (independent coordinates) in d spacial dimensions. If the holonomic constraints are expressed in K equations, then we have $d \cdot N - K$ degrees of freedom expressed in terms of the generalized coordinates $q_1, q_2, q_3, \dots, q_{dN-K}$.

Transformation Equations relate the original variables $\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N$ in terms to q_i via

$$\mathbf{r}_j = \mathbf{r}_j(q_1, q_2, q_3, \dots, q_{dN-K}, t)$$

Examples would include a particle constrained to move on a sphere of fixed radius, a pendulum with a sliding attach-point, or a double pendulum.

1.4 D'Alembert's Principle and Lagrange's Equations

Let

$$\underbrace{\mathbf{F}_i}_{\substack{\text{total force} \\ \text{on } i^{\text{th}} \\ \text{particle}}} = \underbrace{\mathbf{F}_i^{(a)}}_{\substack{\text{applied} \\ \text{forces}}} + \underbrace{\mathbf{f}_i}_{\substack{\text{force of} \\ \text{constraint}}}$$


By Newton's 2nd law:

$$\mathbf{F}_i = \dot{\mathbf{p}}_i \Rightarrow \mathbf{F}_i - \dot{\mathbf{p}}_i = 0$$

which gives

$$\sum_i (\mathbf{F}_i - \dot{\mathbf{p}}_i) \cdot \underbrace{\delta \mathbf{r}_i}_{\substack{\text{infinitesimal} \\ \text{displacement}}} = 0 = \sum_i \left(\mathbf{F}_i^{(a)} - \dot{\mathbf{p}}_i \right) \cdot \delta \mathbf{r}_i + \sum_i (\mathbf{f}_i - \dot{\mathbf{p}}_i) \cdot \delta \mathbf{r}_i$$

Consider the case where $\sum_i \mathbf{f}_i \cdot \delta \mathbf{r}_i = 0$. This is akin to



i.e. the forces of constraint do no work. This leaves us with:

$$\sum_i \left(\mathbf{F}_i^{(a)} - \dot{\mathbf{p}}_i \right) \cdot \delta \mathbf{r}_i = 0 \quad \Rightarrow \text{D'Alembert's Principle} \quad (\dagger)$$

Note that the above equation contains no constraints, so we'll drop the "(a)" designation unambiguously in the future.

For $\mathbf{r}_i = \mathbf{r}_i(q_1, q_2, q_3, \dots, q_n, t)$, we will write (\dagger) in terms of the generalized coordinates q_i :

$$\begin{aligned} \delta \mathbf{r}_i &= \sum_j \frac{\partial \mathbf{r}_i}{\partial q_j} \delta q_j \\ \Rightarrow \sum_i \mathbf{F}_i \cdot \delta \mathbf{r}_i &= \sum_{i,j} \mathbf{F}_i \cdot \frac{\partial \mathbf{r}_i}{\partial q_j} \delta q_j \equiv \sum_j Q_j \delta q_j \end{aligned}$$

where

$$Q_j = \sum_i \mathbf{F}_i \cdot \frac{\partial \mathbf{r}_i}{\partial q_j} = \text{the generalized force}$$

Working on the other half of the equation, we have:

$$\begin{aligned}\Rightarrow \sum_i \dot{\mathbf{p}}_i \cdot \delta \mathbf{r}_i &= \sum_{i,j} m_i \dot{\mathbf{v}}_i \cdot \frac{\partial \mathbf{r}_i}{\partial q_j} \delta q_j \\ &= \sum_{i,j} \left[\frac{d}{dt} \left(m_i \mathbf{v}_i \cdot \frac{\partial \mathbf{r}_i}{\partial q_j} \right) - m_i \mathbf{v}_i \cdot \frac{d}{dt} \left(\frac{\partial \mathbf{r}_i}{\partial q_j} \right) \right] \delta q_j\end{aligned}$$

Using the fact that

$$\begin{aligned}\dot{\mathbf{v}}_i &= \frac{d\mathbf{r}_i}{dt} = \frac{\partial}{\partial t} \mathbf{r}_i + \sum_k \dot{q}_k \frac{\partial \mathbf{r}_i}{\partial q_k} \quad \text{by chain rule} \\ \Rightarrow \quad \frac{\partial \mathbf{v}_i}{\partial \dot{q}_j} &= \frac{\partial \mathbf{r}_i}{\partial q_j}\end{aligned}$$

Thus:

$$\begin{aligned}\sum_i \dot{\mathbf{p}}_i \cdot \delta \mathbf{r}_i &= \sum_{i,j} \left[\frac{d}{dt} \left(m_i \mathbf{v}_i \cdot \frac{\partial \mathbf{v}_i}{\partial \dot{q}_j} \right) - m_i \mathbf{v}_i \cdot \frac{\partial \mathbf{v}_i}{\partial q_j} \right] \delta q_j \\ &= \sum_j \left\{ \frac{d}{dt} \left[\frac{\partial}{\partial \dot{q}_j} \left(\sum_i \frac{1}{2} m_i v_i^2 \right) \right] - \frac{\partial}{\partial q_j} \left(\sum_i \frac{1}{2} m_i v_i^2 \right) \right\} \delta q_j\end{aligned}$$

And D'Alembert's principle becomes:

$$\sum_i (\mathbf{F}_i - \dot{\mathbf{p}}_i) \cdot \delta \mathbf{r}_i = \sum_j \left\{ Q_j - \frac{d}{dt} \frac{\partial T}{\partial \dot{q}_j} + \frac{\partial T}{\partial q_j} \right\} \delta q_j = 0$$

with $T = \sum_i \frac{1}{2} m_i v_i^2$. Now if the system is Holonomic, then it is possible to find q_j such that δq_j is independent of δq_k for all $j \neq k$. Thus, the individual coefficients must vanish:

$$\frac{d}{dt} \frac{\partial T}{\partial \dot{q}_j} - \frac{\partial T}{\partial q_j} = Q_j$$

If $\mathbf{F}_i = -\nabla U$ (conservative force), then

$$Q_j = \sum_i \mathbf{F}_i \cdot \frac{\partial \mathbf{r}_i}{\partial q_j} = - \sum_i \nabla_i U \cdot \frac{\partial \mathbf{r}_i}{\partial q_j} = - \frac{dU}{dq_j} = - \frac{dU}{dq_j} + \underbrace{\frac{d}{dt} \left(\frac{dU}{d\dot{q}_j} \right)}_0$$

where the last term is 0 since we are taking U to be velocity independent. Thus we have

$$\frac{d}{dt} \frac{\partial}{\partial \dot{q}_j} (T - U) - \frac{\partial}{\partial q_j} (T - U) = 0$$

Define the Lagrangian to be $\mathcal{L} = T - U$. Thus Lagrange's equations become:

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}_j} - \frac{\partial \mathcal{L}}{\partial q_j} = 0$$

Note that the Lagrangian is *not* unique! (See HW)

1.5 Velocity Dependent Potentials and Dissipation Functions

If $U = U(q_j, \dot{q}_j)$, then

$$Q_j = -\frac{\partial U}{\partial q_j} + \frac{d}{dt} \left(\frac{\partial U}{\partial \dot{q}_j} \right) \quad \text{and} \quad \mathcal{L} = T - U$$

An example would be a charged particle in an EM field (See HW). If frictional forces are present, then we'll have

$$\frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}_j} \right) - \frac{\partial \mathcal{L}}{\partial q_j} = \tilde{Q}_j = \text{forces not arising from a potential}$$

Example: Consider Rayleigh's dissipation function:

$$\mathcal{F} = \frac{1}{2} \sum_i (k_x v_{ix}^2 + k_y v_{iy}^2 + k_z v_{iz}^2)$$

And

$$F_{fx} = -\frac{\partial \mathcal{F}}{\partial v_x}; \quad \mathbf{F}_f = -\nabla_v \mathcal{F}$$

where F_{fx} is the x component of the frictional force and the sum over i is over all particles. Thus we have that \tilde{Q}_j (the generalized force due to friction) is

$$\begin{aligned} \tilde{Q}_j &= \sum_i \mathbf{F}_{fi} \cdot \frac{\partial \mathbf{r}_i}{\partial q_j} = - \sum_i \nabla_v \mathcal{F} \cdot \frac{\partial \mathbf{r}_i}{\partial q_j} \\ &= - \sum_i \nabla_v \mathcal{F} \cdot \frac{\partial \mathbf{v}_i}{\partial \dot{q}_j} = - \frac{\partial \mathcal{F}}{\partial \dot{q}_j} \\ \Rightarrow \quad \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}_j} \right) - \frac{\partial \mathcal{L}}{\partial q_j} + \frac{\partial \mathcal{F}}{\partial \dot{q}_j} &= 0 \end{aligned}$$

1.6 Simple Applications of Lagrangian Formalism

Example 1: Consider a single free particle in space:

$$T = \frac{1}{2} m (\dot{x}_1^2 + \dot{x}_2^2 + \dots + \dot{x}_d^2) \quad \text{in d-dimensional space}$$

$$U = 0$$

$$Q_i = \sum_j \mathbf{F}_j \cdot \frac{\partial \mathbf{r}_j}{\partial q_i}$$

where, in cartesian coordinates, $\frac{\partial x_j}{\partial q_i} = \delta_{ij}$

$$\mathcal{L} = T$$

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{x}_i} - \frac{\partial \mathcal{L}}{\partial x_i} = 0$$

$$\Rightarrow m\ddot{x}_i = F_{x_i} = 0$$

Example 2: Consider a particle constrained on a sphere. Using polar coordinates, we have


$$x_1 = R \sin \theta \cos \phi$$

$$x_2 = R \sin \theta \sin \phi$$

$$x_3 = R \cos \theta$$

Calculate $\dot{x}_1, \dot{x}_2, \dot{x}_3$ and substitute into T to get the kinetic energy for the system whose natural coordinates are spherical.

Example 3: Here we will consider a pendulum with a sliding pivot:



So we have that

$$x = x' + l \sin \phi$$

$$y = l - l \cos \phi$$

Thus we have:

$$\begin{aligned}
 T &= \frac{1}{2}m (\dot{x}^2 + \dot{y}^2) \\
 &= \frac{1}{2}m \left(\left(\dot{x}' + l \cos \phi \dot{\phi} \right)^2 + \left(l \sin \phi \dot{\phi} \right)^2 \right) \\
 &= \frac{1}{2}m \left(\dot{x}'^2 + 2l \cos \phi \dot{x}' \dot{\phi} + l^2 \cos^2 \phi \dot{\phi}^2 + l^2 \sin^2 \phi \dot{\phi}^2 \right) \\
 &= \frac{1}{2}m \left(\dot{x}'^2 + 2l \cos \phi \dot{x}' \dot{\phi} + l^2 \dot{\phi}^2 \right)
 \end{aligned}$$

$$U = mgy = mg(l - l \cos \phi)$$

$$\mathcal{L} = \frac{1}{2}m \left(\dot{x}'^2 + 2l \cos \phi \dot{x}' \dot{\phi} + l^2 \dot{\phi}^2 \right) - mgl(1 - \cos \phi)$$

Remark 1: Generalized coordinates are not necessarily orthogonal!

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}_i} - \frac{\partial \mathcal{L}}{\partial q_i}$$

$$\begin{aligned}
 \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{x}'} - \frac{\partial \mathcal{L}}{\partial x'} &= \frac{d}{dt} (m\dot{x}' + ml \cos \phi \dot{\phi}) - 0 = 0 \\
 \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{\phi}} - \frac{\partial \mathcal{L}}{\partial \phi} &= \frac{d}{dt} (ml \cos \phi \dot{x}' + ml^2 \dot{\phi}) + ml \sin \phi \dot{x}' \dot{\phi} + mgl \sin \phi = 0
 \end{aligned}$$

To find the equilibrium points:

$$U = mgl - mgl \cos \phi$$

$$\begin{aligned}
 \frac{\partial U}{\partial \phi} &= mgl \sin \phi = 0 \quad \Rightarrow \quad \phi = 0, \pi \\
 \frac{\partial^2 U}{\partial \phi^2} &= mgl \cos \phi|_{\phi=0, \pi} = \begin{cases} mgl > 0, & \phi = 0 \\ -mgl < 0, & \phi = \pi \end{cases}
 \end{aligned}$$


So we have a stable equilibrium at $\phi = 0$ and an unstable equilibrium at $\phi = \pi$ (as we'd expect).

Chapter 2

Variational Principles and Lagrange's Equations

2.1 Hamilton's Principle and Calculus of Variations

Consider a system with n generalized coordinates: q_1, q_2, \dots, q_n . These coordinates make up an n -dimensional configuration space with each q corresponding to an axis in the space.




1. t is a parameter of curve \mathcal{C} with $t \in [t_-, t_+] = \text{interval}$
2. $q_- = q(t_-)$, $q_+ = q(t_+)$
3. $\frac{d}{dt}q(t) \equiv \dot{q}(t)$ is the tangent vector
4. If $L(q(t), \dot{q}(t), t)$ is a function of the q 's and their tangents, then we define a number that characterizes the path:

$$S(\mathcal{C}) = \sum_{t_-}^{t_+} L(q(t), \dot{q}(t), t) dt$$

Note that this is more general than $t = \text{time}$ and $L = \text{Lagrangian}$. The above is true of *any* function of q, \dot{q} and t .

5. S is called a functional (a functional is an animal that eats a function and spits out a number).
6. If $\mathcal{C} \rightarrow \mathcal{C} + \delta\mathcal{C}$, then $S \rightarrow S + \delta S$ continuously

Example 1: Let $n = 2$ and we'll work in 2-d Euclidean space. We have that $q = q(x_1, x_2)$ and \mathcal{C} is a curve with parameter t .



We can find the length of \mathcal{C} by:

$$\begin{aligned} l(\mathcal{C}) &= \lim_{N \rightarrow \infty} \sum_{i=1}^N \sqrt{(q(t_{i+1}) - q(t_i))^2} \\ &= \lim_{N \rightarrow \infty} \sum_{i=1}^N \sqrt{\left(\frac{q(t_{i+1}) - q(t_i)}{t_{i+1} - t_i} \right)^2} (t_{i+1} - t_i) \\ &= \int_{t_1}^{t_2} \sqrt{(\dot{q}(t))^2} dt \end{aligned}$$

$$\begin{aligned} L(q, \dot{q}, t) &= L(\dot{q}) = \sqrt{\dot{q}^2} \\ &\Rightarrow S = l \end{aligned}$$

Example 2: Now we'll look at a point moving on curve \mathcal{C} with velocity $v(q)$ that takes time T to go from q_1 to q_2 .

$$\begin{aligned} T &= \lim_{N \rightarrow \infty} \sum_{i=1}^n \frac{1}{v(q(t_i))} \sqrt{(q(t_{i+1}) - q(t_i))^2} \\ &= \int_{t_1}^{t_2} \frac{1}{v(q(t))} \sqrt{(\dot{q})^2} \\ &\Rightarrow L(q, \dot{q}, t) = L(q, \dot{q}) = \frac{\sqrt{\dot{q}^2}}{v(q)} \quad \text{and} \quad S = T \end{aligned}$$

Example 3: Again, we have a moving particle with time = t and position = q . We can write the kinetic and potential energy as:

$$T = \frac{\dot{q}^2}{2m}, \quad U(q)$$

Thus:

$$L(q, \dot{q}, t) = \mathcal{L}(q, \dot{q}) = T(\dot{q}) - U(q)$$


$$S = \int_{t_1}^{t_2} \mathcal{L}(q(t), \dot{q}(t), t) dt$$

So S is the Action. Hamilton's principle says that S has a stationary value for the actual path of motion:

$$\Rightarrow \delta S = \delta \int_{t_1}^{t_2} \mathcal{L}(q, \dot{q}, t) dt = 0$$

2.2 The 3 Classic Problems of the Calculus of Variations

The Brachistochrone Problem: Brachistochrone means “short time”, so these type of problems are attempting to minimize the time. A massive particle moves from A to B under the force



of gravity along path \mathcal{C} . Which \mathcal{C} gives the shortest travel time?

The Geodesics Problem: A ship is traveling from Portland, OR to Hawaii along the surface of a sphere. Which route is the shortest?

The Isoperimetric Problem (Dido's Problem): For a curve \mathcal{C} with given length l , which form gives the maximum area?


2.3 Calculus of Variations, Hamilton's Principle, and Lagrange's Equations

Consider a curve \mathcal{C}_ε in configuration space.

$$\mathcal{C}_\varepsilon : \quad q_\varepsilon(\tau) = q_\varepsilon(q_1(\tau, \varepsilon), \dots, q_n(\tau, \varepsilon))$$

Some things to note:

1. $q_\varepsilon(\tau)$ is a path for every fixed ε .
2. $q_{\varepsilon=0}(\tau) = q(\tau)$



3.

$$\left. \begin{aligned} q_{\varepsilon}(\tau_-) &= q(\tau_-) \\ q_{\varepsilon}(\tau_+) &= q(\tau_+) \end{aligned} \right\} \text{ for all } \varepsilon$$

4. $q_{\varepsilon}(\tau)$ are continuously differentiable with respect to ε (fixed τ).Consider a function L and functional S :

$$S_L(\mathcal{C}) = \int_{\tau_-}^{\tau_+} L(q(\tau), \dot{q}(\tau), \tau) d\tau$$

Then, under variations of the path \mathcal{C} :

1. $S(\mathcal{C})$ is minimal if $S(\mathcal{C}_{\varepsilon}) > S(\mathcal{C})$ for all ε in the neighborhood of \mathcal{C}
2. $S(\mathcal{C})$ is maximal if $S(\mathcal{C}_{\varepsilon}) < S(\mathcal{C})$ for all ε in the neighborhood of \mathcal{C}
3. $S(\mathcal{C})$ is stationary, and \mathcal{C} is an extremal of S if

$$\begin{aligned} \delta S &= \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} [S(\mathcal{C}_{\varepsilon}) - S(\mathcal{C})] = 0 \quad \forall \mathcal{C}_{\varepsilon} \text{ in nbhd} \\ &= \text{variation of functional } S \end{aligned}$$

Remark 1: We find the extremals to solve classical problems.

$$\begin{aligned} \delta S &= \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} [S(\mathcal{C}_{\varepsilon}) - S(\mathcal{C})] \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\tau_-}^{\tau_+} \frac{1}{\varepsilon} [L(q_{\varepsilon}, \dot{q}_{\varepsilon}, \tau) - L(q, \dot{q}, \tau)] d\tau \end{aligned}$$

We then Taylor expand $S(\mathcal{C}_{\varepsilon})$ for small ε . Let $q_{\varepsilon}(\tau) = q(\tau) + \varepsilon \eta(\tau)$. Then

$$\delta S = \lim_{\varepsilon \rightarrow 0} \int_{\tau_-}^{\tau_+} \frac{1}{\varepsilon} \left[L(q, \dot{q}, \tau) + \sum_{i=1}^n \left(\frac{\partial L}{\partial q_{\varepsilon,i}} \frac{\partial q_{\varepsilon,i}}{\partial \varepsilon} + \frac{\partial L}{\partial \dot{q}_{\varepsilon}} \frac{\partial \dot{q}_{\varepsilon}}{\partial \varepsilon} \right) \varepsilon + \mathcal{O}(\varepsilon^2) - L(q, \dot{q}, \tau) \right] d\tau$$

Note that, since $q_\varepsilon(\tau) = q(\tau) + \varepsilon\eta(\tau)$, we have

$$\begin{aligned} \frac{\partial q_\varepsilon}{\partial \varepsilon} &= \eta(\tau) \\ \Rightarrow \quad \frac{\partial L}{\partial q_\varepsilon} &= \frac{\partial L}{\partial q} \underbrace{\frac{\partial q}{\partial q_\varepsilon}}_1 = \frac{\partial L}{\partial q} \end{aligned}$$

Thus

$$\begin{aligned} \delta S &= \lim_{\varepsilon \rightarrow 0} \int_{\tau_-}^{\tau_+} \frac{1}{\varepsilon} \left[\sum_{i=1}^n \left(\frac{\partial L}{\partial q_i} \eta_i + \frac{\partial L}{\partial \dot{q}_i} \dot{\eta}_i \right) \varepsilon + \mathcal{O}(\varepsilon^2) \right] d\tau \\ &= \int_{\tau_-}^{\tau_+} \sum_{i=1}^n \frac{\partial L}{\partial q_i} \eta_i d\tau + \sum_{i=1}^n \left[\underbrace{\frac{\partial L}{\partial \dot{q}_i} \eta_i}_{0} \Big|_{\tau_-}^{\tau_+} - \int_{\tau_-}^{\tau_+} \eta_i \frac{d}{dt} \frac{\partial L}{\partial \dot{q}_i} d\tau \right] \\ &= \int_{\tau_-}^{\tau_+} \sum_{i=1}^n \left[\frac{\partial L}{\partial q_i} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}_i} \right] \eta_i(\tau) d\tau \\ &= 0 \text{ since } S \text{ is stationary.} \end{aligned}$$

Now, applying the fundamental lemma of the calculus of variations:

Lemma: Let $f(\tau)$ be continuous for $\tau \in [\tau_-, \tau_+]$ and

$$\int_{\tau_-}^{\tau_+} f(\tau) \eta(\tau) d\tau = 0 \quad \forall \eta$$

which are 2 times differentiable and obey

$$\eta(\tau_+) = \eta(\tau_-) = 0$$

Then

$$f(\tau) = 0$$

since $\eta_i(\tau)$ is arbitrary and vanishes at the endpoints.

$$\Rightarrow \quad \frac{\partial L}{\partial q_i} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}_i} = 0 = \text{Euler Equations} \quad (\dagger)$$

Remark 2: When $L = T - U$ in (\dagger) , these are the Euler-Lagrange equations.

Remark 3: The functional $S = \int L(q(\tau), \dot{q}(\tau), \tau) d\tau$ is stationary only if the functional L obeys the Euler-Lagrange equations.

Remark 4: (\dagger) implies that

$$\begin{aligned} \frac{\partial L}{\partial q_i} - \sum_k \frac{\partial}{\partial \dot{q}_k} \left(\frac{\partial L}{\partial \dot{q}_i} \right) \ddot{q}_k - \sum_k \frac{\partial}{\partial q_k} \left(\frac{\partial L}{\partial \dot{q}_i} \right) \dot{q}_k - \frac{\partial}{\partial t} \left(\frac{\partial L}{\partial \dot{q}_i} \right) &= 0 \\ \Rightarrow \sum_k L_{ik} \ddot{q}_k &= \frac{\partial L}{\partial q_i} - \frac{\partial^2 L}{\partial \dot{q}_i \partial \tau} - \sum_k \left(\frac{\partial^2 L}{\partial q_k \partial \dot{q}_i} \right) \dot{q}_k \end{aligned} \quad (\ddagger)$$

and

$$L_{ik} = \frac{\partial^2 L}{\partial \dot{q}_i \partial \dot{q}_k} \quad \leftarrow \text{symmetric } n \times n \text{ matrix}$$

So the Euler-Lagrange equations (\ddagger) are equivalent to n -coupled ODE's of second order for $q(\tau)$. Therefore, the initial point of $q(\tau_-)$ and initial tangent $\dot{q}(\tau_-)$ completely determine the path.

Remark 5: (\ddagger) has the structure of Newton's Second Law.

Remark 6: Lagrange's equations of motion follow naturally from Hamilton's Principle.

Example: Geodesics in d-dimensional Euclidean space: Recall that the length of curve \mathcal{C} is given by

$$l(\mathcal{C}) = \int_{\tau_i}^{\tau_+} \sqrt{(\dot{q}(\tau))^2} d\tau = L(q, \dot{q}, \tau) = \sqrt{\dot{q}^2} = \left[\sum_{j=1}^d (q_j(\tau))^2 \right]^{1/2}$$

(\dagger) then implies that

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_i} - \frac{\partial L}{\partial q_i} = \frac{d}{dt} \left(\frac{1}{2} \frac{1}{\sqrt{\dot{q}^2}} \cdot 2\dot{q}_i \right) = 0 \quad \forall i$$

$$\Rightarrow \frac{\dot{q}_i}{\sqrt{\dot{q}^2}} = \text{constant}$$

$$\Rightarrow \dot{q} = a = \text{constant} \quad \Rightarrow \quad q(\tau) = a\tau + b = \text{straight lines}$$

So the geodesics must run thru 2 points to specify a and b , implying that we require a 2-d condition.

2.4 Extensions of Hamilton's Principle to Non-Holonomic Systems (Lagrange Multipliers)

$$\delta S = \delta \int_{t_-}^{t_+} L(q, \dot{q}, t) dt = 0 = \int_{\tau_-}^{\tau_+} \sum_i \left(\frac{\partial}{\partial q_i} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}_i} \right) d\tau \delta q_i = 0$$

Non-Holonomic constraints imply that the generalized coordinates are *not* independent. This implies that displacements of the path may or may not satisfy the constraints. If displacements satisfy constraints, then the constraints are holonomic. If the displacements do *not* satisfy the constraints, then we want to eliminate the constraints by means of Lagrange Multipliers. Lagrange multipliers work for “semi-holonomic” constraints, which can be put in the form:

$$f_\alpha(q_1, \dots, q_n, \dot{q}_1, \dots, \dot{q}_n) = 0 \quad (\dagger)$$

where $\alpha = 1, 2, \dots, n$.

Remark 1: Semi-holonomic differs from holonomic in that the latter can be expressed in terms of *one* constraint equation (function of generalized coordinates only), whereas the former can be more than one (function of tangents as well).

Remark 2: In terms of path displacements, the semi-holonomic constraints can be expressed:

$$\sum_k a_{ik} dq_k + a_{it} dt = 0 \quad \text{where } i = 1, \dots, m$$

This is more restrictive than (\dagger) .

Remark 3: (\dagger) implies that:

$$\sum_{\alpha=1}^m \lambda_\alpha f_\alpha = 0$$

where

$$\lambda_\alpha = \lambda_\alpha(q_1, \dots, q_n, \dot{q}_1, \dots, \dot{q}_n, t)$$

are undetermined functions.

Recall Hamilton's Principle:

$$\delta \int_{t_1}^{t_2} \mathcal{L} dt = 0$$

which implies

$$\int_{t_1}^{t_2} \sum_k \left(\frac{\partial \mathcal{L}}{\partial q_k} - \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}_k} \right) dt \delta q_k = 0$$

And δq_k are no longer independent (if we have non-holonomic constraints). But if the constraints are semi-holonomic, then:

$$\delta \int_{t_1}^{t_2} \left(\mathcal{L} + \sum_{\alpha=1}^m \lambda_\alpha f_\alpha \right) dt = 0$$

Changing notation slightly, let

$$L = \mathcal{L}_1, \quad \underbrace{\sum_{\alpha=1}^m \lambda_\alpha f_\alpha}_{\star} = \lambda \mathcal{L}_2$$

where (\star) in the piece that will make the δq_k 's independent. Then

$$\begin{aligned}\mathcal{L}_3 &= \mathcal{L}_1 + \lambda \mathcal{L}_2 \\ \Rightarrow \delta \int_{t_1}^{t_2} (\mathcal{L}_1 + \lambda \mathcal{L}_2) dt &= \delta \int_{t_1}^{t_2} \mathcal{L}_3 dt = 0 \\ \Rightarrow \int_{t_1}^{t_2} \sum_k \left(\frac{\partial \mathcal{L}_3}{\partial q_i} - \frac{d}{dt} \frac{\partial \mathcal{L}_3}{\partial \dot{q}_i} \right) dt \delta q_i &= 0\end{aligned}$$

And \mathcal{L}_3 obeys Lagrange's equations:

$$\frac{d}{dt} \frac{\partial \mathcal{L}_3}{\partial \dot{q}_i} - \frac{\partial \mathcal{L}_3}{\partial q_i} = 0$$

Also

$$\begin{aligned}\frac{d}{dt} \frac{\partial \mathcal{L}_3}{\partial \dot{q}_i} - \frac{\partial \mathcal{L}_3}{\partial q_i} &= \frac{d}{dt} \frac{\partial}{\partial \dot{q}_i} (\mathcal{L}_1 + \lambda \mathcal{L}_2) - \frac{\partial}{\partial q_i} (\mathcal{L}_1 + \lambda \mathcal{L}_2) \\ &= \frac{d}{dt} \frac{\partial}{\partial \dot{q}_i} \mathcal{L}_1 - \frac{\partial \mathcal{L}_1}{\partial q_i} + \frac{d}{dt} \frac{\partial}{\partial \dot{q}_i} (\lambda \mathcal{L}_2) - \frac{\partial}{\partial q_i} (\lambda \mathcal{L}_2) \\ &= 0\end{aligned}$$

Now,

$$\begin{aligned}\frac{d}{dt} \frac{\partial \mathcal{L}_1}{\partial \dot{q}_i} - \frac{\partial \mathcal{L}_1}{\partial q_i} &= \frac{\partial}{\partial q_i} (\lambda \mathcal{L}_2) - \frac{d}{dt} \frac{\partial (\lambda \mathcal{L}_2)}{\partial \dot{q}_i} \equiv Q_i \\ \Rightarrow \frac{d}{dt} \frac{\partial \mathcal{L}_1}{\partial \dot{q}_i} - \frac{\partial \mathcal{L}_1}{\partial q_i} &= Q_i = \text{forces of constraint}\end{aligned}$$

with

$$Q_i = \frac{\partial}{\partial q_i} (\lambda \mathcal{L}_2) - \frac{d}{dt} \frac{\partial}{\partial \dot{q}_i} (\lambda \mathcal{L}_2)$$

Remark 4: Lagrange multipliers allow us to map a semi-holonomic system of n -generalized coordinates and m -constraints to a holonomic one with $n + m$ generalized coordinates.

Example:

$$\mathcal{L} = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2) - U(x, y) \equiv \mathcal{L}_1$$

$$\text{Constraint: } f(\dot{x}, \dot{y}, y) = \dot{x}\dot{y} + ky = 0 \equiv \mathcal{L}_2 \quad \text{where } k = \text{const} \quad (1)$$

$$\Rightarrow \mathcal{L}_3 = \mathcal{L}_1 + \lambda \mathcal{L}_2 \quad \text{and} \quad \frac{d}{dt} \frac{\partial \mathcal{L}_3}{\partial \dot{q}_i} - \frac{\partial \mathcal{L}_3}{\partial q_i} = 0$$

$$\Rightarrow \frac{d}{dt} \frac{\partial \mathcal{L}_3}{\partial \dot{x}} - \frac{\partial \mathcal{L}_3}{\partial x} = \frac{d}{dt} (m\dot{x} + \lambda\dot{y}) + \frac{\partial U}{\partial x} = 0 \quad (2)$$

$$\frac{d}{dt} \frac{\partial \mathcal{L}_3}{\partial \dot{y}} - \frac{\partial \mathcal{L}_3}{\partial y} = \frac{d}{dt} (m\dot{y} + \lambda\dot{x}) + \frac{\partial U}{\partial y} - k\lambda = 0 \quad (3)$$


Thus we have 3 unknowns: x, y, λ , and 3 equations: (1), (2), (3).

Remark 5: Physically, the λ 's represent forces of constraints.

Remark 6: Forces of constraint do no work in (virtual) displacements (See §2.4 of Goldstein).

Remark 7: Lagrange multipliers can be used for holonomic constraints ($\alpha = 1$) as well.

Example: Hoop Rolling without Slipping Down an Inclined Plane: We are using the



generalized coordinates x, θ . Our constraint in the rolling constraint:

$$\begin{aligned} r\dot{\theta} &= \dot{x} \\ \Rightarrow r\theta &= x \\ \Rightarrow r\theta - x &= 0 \end{aligned} \tag{3}$$

$$\begin{aligned} T &= \frac{1}{2}m\dot{x}^2 + \frac{1}{2}mr^2\dot{\theta}^2 \\ U &= mg(l - x)\sin\phi \end{aligned}$$

$$\begin{aligned} \Rightarrow \mathcal{L}_1 &= \frac{1}{2}m\dot{x}^2 + \frac{1}{2}mr^2\dot{\theta}^2 - mg(l - x)\sin\phi \\ \mathcal{L}_2 &= r\theta - x \\ \mathcal{L}_3 &= \mathcal{L}_1 + \lambda\mathcal{L}_2 \\ &= \frac{1}{2}m\dot{x}^2 + \frac{1}{2}mr^2\dot{\theta}^2 - mg(l - x)\sin\phi + \lambda(r\theta - x) \end{aligned}$$

$$\Rightarrow \frac{d}{dt} \frac{\partial \mathcal{L}_3}{\partial \dot{x}} - \frac{\partial \mathcal{L}_3}{\partial x} = m\ddot{x} - mg\sin\phi + \lambda = 0 \tag{1}$$

$$\frac{d}{dt} \frac{\partial \mathcal{L}_3}{\partial \dot{\theta}} - \frac{\partial \mathcal{L}_3}{\partial \theta} = mr^2\ddot{\theta} - \lambda r = 0 \tag{2}$$

$$\begin{aligned} \Rightarrow m\ddot{x} &= mg\sin\phi - \lambda \\ &= mg\sin\phi - mr\ddot{\theta} \quad \text{from (2)} \\ &= mg\sin\phi - m\ddot{x} \quad \text{by (3)} \end{aligned}$$


$$\Rightarrow 2m\ddot{x} = mg \sin \phi \quad \Rightarrow \quad \ddot{x} = \frac{1}{2}g \sin \phi$$

So rolling has half the acceleration as the hoop sliding on a frictionless plane!

$$\ddot{\theta} = \frac{g}{2r} \sin \phi$$

$$\lambda = \frac{mg}{2} \sin \phi \rightarrow \text{“frictional force” of constraint}$$

Example: Bead on a Wire Loop:



The normal way we'd solve this problem would be:

$$T = \frac{1}{2}mR^2\dot{\phi}^2, \quad U = mgy = mgR \sin \phi$$

$$\mathcal{L} = \frac{1}{2}mR^2\dot{\phi}^2 - mgR \sin \phi$$

$$\Rightarrow \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{\phi}} - \frac{\partial \mathcal{L}}{\partial \phi} = mR^2\ddot{\phi} + mgR \cos \phi = 0$$

$$\Rightarrow \ddot{\phi} = -\frac{g}{R} \cos \phi$$

Now, using Lagrange multipliers:

$$T = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2), \quad U = mgy \quad \Rightarrow \quad \mathcal{L}_1 = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2) - mgy$$

Our constraint is:

$$x^2 + y^2 = R^2 \quad \Rightarrow \quad x^2 + y^2 - R^2 = 0 = \mathcal{L}_2$$

which implies

$$\Rightarrow \mathcal{L}_3 = \mathcal{L}_1 - \lambda \mathcal{L}_2 = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2) - mgy + \lambda(x^2 + y^2 - R^2)$$

$$\begin{aligned} \frac{d}{dt} \frac{\partial \mathcal{L}_3}{\partial \dot{x}} - \frac{\partial \mathcal{L}_3}{\partial x} &= m\ddot{x} - 2\lambda x = 0 \\ \frac{d}{dt} \frac{\partial \mathcal{L}_3}{\partial \dot{y}} - \frac{\partial \mathcal{L}_3}{\partial y} &= m\ddot{y} + mg - 2\lambda y = 0 \end{aligned}$$

So our 3 equations are:

$$\begin{aligned} m\ddot{x} &= 2\lambda x \\ m\ddot{y} &= -mg + 2\lambda y \\ R^2 &= x^2 + y^2 \end{aligned}$$

Letting

$$\begin{aligned} x &= R \cos \phi & y &= R \sin \phi \\ \dot{x} &= -R \sin \phi \dot{\phi} & \dot{y} &= R \cos \phi \dot{\phi} \\ \ddot{x} &= -R \cos \phi \dot{\phi}^2 - R \sin \phi \ddot{\phi} & \ddot{y} &= -R \sin \phi \dot{\phi}^2 + R \cos \phi \ddot{\phi} \end{aligned}$$

Taking $m\ddot{x} = 2\lambda x$ and plugging in \ddot{x} implies:

$$\begin{aligned} m \left(-R \cos \phi \dot{\phi}^2 - R \sin \phi \ddot{\phi} \right) &= 2\lambda R \cos \phi \\ \Rightarrow \quad \lambda &= -\frac{m}{2} \dot{\phi}^2 - \frac{m \sin \phi}{2 \cos \phi} \ddot{\phi} \end{aligned}$$

Taking $m\ddot{y} = -mg + 2\lambda y$:

$$\begin{aligned} \Rightarrow \quad m \left(-R \sin \phi \dot{\phi}^2 + R \cos \phi \ddot{\phi} \right) &= -mg + 2 \left(-\frac{m}{2} \dot{\phi}^2 - \frac{m}{2} \tan \phi \ddot{\phi} \right) R \sin \phi \\ \Rightarrow \quad (mR \cos \phi + mR \tan \phi \sin \phi) \ddot{\phi} &= -mg \\ \Rightarrow \quad \ddot{\phi} &= -\frac{g}{R} \cos \phi \end{aligned}$$

2.5 Conservation Theorems and Symmetry Properties

Consider a function $f(q, \dot{q}, t)$. If

$$\frac{d}{dt} (f(q, \dot{q}, t)) = 0$$

then

$$f(q, \dot{q}, t) = \text{const} \tag{\dagger}$$

where (\dagger) is called the

- first integral of the equations of motion
- integrals of motion
- constant of motion

Remark 1: Conservation laws are an example of (\dagger) .

Nöther's Theorem: Symmetries in a system are related to conservation laws.

2.5.1 Energy Conservation and Time Homogeneity

Consider the Lagrangian:

$$\mathcal{L} = \mathcal{L}(q(\tau), \dot{q}(\tau), \tau)$$

Time homogeneity (time-translational invariance) means that \mathcal{L} does not depend explicitly on time.

$$\Rightarrow \mathcal{L} = \mathcal{L}(q(\tau), \dot{q}(\tau))$$

Then

$$\frac{d}{dt}\mathcal{L} = \sum_i \frac{\partial \mathcal{L}}{\partial q_i} \dot{q}_i + \sum_i \frac{\partial \mathcal{L}}{\partial \dot{q}_i} \ddot{q}_i + \underbrace{\frac{\partial \mathcal{L}}{\partial t}}_0$$

Recall Lagrange's equations of motion:

$$\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}_j} = \frac{\partial \mathcal{L}}{\partial q_j}$$

These imply

$$\begin{aligned} \frac{d}{dt}\mathcal{L} &= \sum_i \left(\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}_j} \right) \dot{q}_j + \sum_i \frac{\partial \mathcal{L}}{\partial \dot{q}_j} \ddot{q}_j \\ &= \sum_i \frac{d}{dt} \left(\dot{q}_j \frac{\partial \mathcal{L}}{\partial \dot{q}_j} \right) \\ &\Rightarrow \frac{d}{dt} \left[\sum_i \dot{q}_j \frac{\partial \mathcal{L}}{\partial \dot{q}_j} - \mathcal{L} \right] = 0 \\ &\Rightarrow \sum_i \dot{q}_j \frac{\partial \mathcal{L}}{\partial \dot{q}_j} - \mathcal{L} \equiv H = E \end{aligned}$$

Then $H = E = \text{const}$ is the energy and energy is conserved in a system whose Lagrangian has no explicit time dependence.

Example:

$$\mathcal{L} = T(\dot{q}) - U(q), \quad T = \frac{1}{2} \sum_j m_j \dot{q}_j^2$$

$$\begin{aligned} \Rightarrow H &= \sum_i \dot{q}_i m_i \dot{q}_i - \frac{1}{2} \sum_i m_i \dot{q}_i^2 + U(q) \\ &= \frac{1}{2} \sum_i m_i \dot{q}_i^2 + U(q) = T(\dot{q}) + U(q) = \text{total energy} \end{aligned}$$

Remark 1: Here we have assumed that $U = U(q)$, so there does not exist dissipative forces (which dissipate energy).

Remark 2: If \mathcal{L} is an explicit function of time:

$$\frac{dH}{dt} = -\frac{\partial \mathcal{L}}{\partial t}$$

Remark 3: H is sometimes called the energy function, and its form coincides with the Hamiltonian.

Remark 4: For frictional forces that are derivable from a dissipative function \mathcal{F} :

$$\text{From §1.5: } \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}_j} \right) - \frac{\partial \mathcal{L}}{\partial q_j} + \frac{\partial \mathcal{F}}{\partial \dot{q}_j} = 0$$

Then

$$\begin{aligned} \frac{d}{dt} \mathcal{L} &= \sum_i \frac{\partial \mathcal{L}}{\partial q_i} \dot{q}_i + \sum_i \frac{\partial \mathcal{L}}{\partial \dot{q}_i} \ddot{q}_i + \frac{\partial \mathcal{L}}{\partial t} \\ &= \sum_i \dot{q}_i \left(\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}_i} + \frac{\partial \mathcal{F}}{\partial \dot{q}_i} \right) + \sum_i \frac{\partial \mathcal{L}}{\partial \dot{q}_i} \ddot{q}_i + \frac{\partial \mathcal{L}}{\partial t} \\ &= \sum_i \frac{d}{dt} \left(\dot{q}_i \frac{\partial \mathcal{L}}{\partial \dot{q}_i} \right) + \underbrace{\sum_i \dot{q}_i \frac{\partial \mathcal{F}}{\partial \dot{q}_i}}_{=2\mathcal{F} \text{ by §1.5}} + \frac{\partial \mathcal{L}}{\partial t} \\ &\Rightarrow \frac{dH}{dt} = -\frac{\partial \mathcal{L}}{\partial t} - 2\mathcal{F} \end{aligned}$$

for $\mathcal{L} = \mathcal{L}(q, \dot{q})$ and $H = E$ then this gives the dissipative rate.

2.5.2 Momentum Conservation and Space Homogeneity

This will correspond to space translational invariance. This implies that the Lagrangian is invariant under $\mathbf{r}_i \rightarrow \mathbf{r}_i + \boldsymbol{\varepsilon}$ (every particle moves by some same displacement). For now, we'll only consider shifts in coordinates, not velocities.

$$\delta \mathcal{L} = \sum_i \frac{\partial \mathcal{L}}{\partial \mathbf{r}_i} \delta \mathbf{r}_i = 0 = \sum_i \frac{\partial \mathcal{L}}{\partial q_i} \underbrace{\delta q_i}_{\star}$$

where (\star) is arbitrary, and only for holonomic constraints. This implies that

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial q_i} &= 0 = \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}_i} \\ \Rightarrow \frac{\partial \mathcal{L}}{\partial \dot{q}_i} &= \text{const} \end{aligned}$$

If $U = U(q)$ only, then

$$\frac{\partial \mathcal{L}}{\partial \dot{q}_i} = m\dot{q}_i = p_i$$

$$\Rightarrow p_i \equiv \frac{\partial \mathcal{L}}{\partial \dot{q}_i} = \text{canonical (or conjugate) momentum}$$

p also generalizes to include $U = U(q, \dot{q})$. Now consider $\mathcal{L} = T(\dot{q}) - U(q)$:


$$\frac{d}{dt} \frac{\partial T}{\partial \dot{q}_i} + \frac{\partial U}{\partial q_i} = 0 \quad \Rightarrow \quad \frac{d}{dt} \frac{\partial T}{\partial \dot{q}_i} = \dot{p}_i = -\frac{\partial U}{\partial q_i} \equiv Q_i$$

$$\Rightarrow p_i = Q_i = \text{generalized force}$$

2.5.3 Angular Momentum Conservation and Rotational Invariance

Let $\delta\phi$ be an infinitesimal rotation and

$$\delta\phi = \delta\phi \hat{n}$$



$$\delta\mathcal{L} = \sum_i \frac{\partial \mathcal{L}}{\partial \mathbf{r}_i} \cdot \delta\mathbf{r}_i + \sum_i \frac{\partial \mathcal{L}}{\partial \mathbf{v}_i} \cdot \delta\mathbf{v}_i = 0$$

Note that

$$\delta\mathbf{r}_i = \delta\phi \times \mathbf{r}_i$$

$$\delta\mathbf{v}_i = \delta\phi \times \mathbf{v}_i \quad \text{assuming } \delta\dot{\phi} = 0$$

So

$$\delta\mathcal{L} = \sum_i \left[\frac{\partial \mathcal{L}}{\partial \mathbf{r}_i} \cdot (\delta\phi \times \mathbf{r}_i) + \frac{\partial \mathcal{L}}{\partial \mathbf{v}_i} \cdot (\delta\phi \times \mathbf{v}_i) \right]$$

but

$$\mathbf{p}_i = \frac{\partial \mathcal{L}}{\partial \mathbf{v}_i} \quad \text{and} \quad \frac{\partial \mathcal{L}}{\partial \mathbf{r}_i} = \frac{d}{dt} \mathbf{p}_i$$

$$\Rightarrow \delta\mathcal{L} = \sum_i \left[\frac{d}{dt} \mathbf{p}_i \cdot (\delta\boldsymbol{\phi} \times \mathbf{r}_i) + \mathbf{p}_i \cdot (\delta\boldsymbol{\phi} \times \mathbf{v}_i) \right]$$

Now, recall that $a \cdot (b \times c) = b \cdot (c \times a) = c \cdot (a \times b)$.

$$\begin{aligned} \Rightarrow \delta\mathcal{L} &= \sum_i [\delta\boldsymbol{\phi} \cdot (\mathbf{r}_i \times \dot{\mathbf{p}}_i) + \delta\boldsymbol{\phi} \cdot (\mathbf{v}_i \times \mathbf{p}_i)] \\ &= \delta\boldsymbol{\phi} \cdot \sum_i \frac{d}{dt} (\mathbf{r}_i \times \mathbf{p}_i) = 0 \end{aligned}$$

But $\delta\boldsymbol{\phi}$ is arbitrary, so

$$\frac{d}{dt} \sum_i (\mathbf{r}_i \times \mathbf{p}_i) = 0$$

Now $\mathbf{M}_i = \mathbf{r}_i \times \mathbf{p}_i$ = angular momentum, so we have that

$$\sum_i \mathbf{M}_i = \text{const}$$

Remark 1: The components of angular momentum along any axis (e.g. the z-axis) are given by

$$M_z = \sum_i \frac{\partial \mathcal{L}}{\partial \dot{\phi}_i} \quad \text{where } \boldsymbol{\phi} = \phi \hat{\mathbf{z}}$$

Example: We'll take a look at how this works in cylindrical coordinates:

$$M_z = \sum_i (\mathbf{r}_i \times \mathbf{p}_i)_z = \sum_i m_i (x_i \dot{y}_i - y_i \dot{x}_i)$$

$$\begin{aligned} x &= r \cos \phi & \dot{x} &= \dot{r} \cos \phi - r \sin \phi \dot{\phi} \\ y &= r \sin \phi & \dot{y} &= \dot{r} \sin \phi + r \cos \phi \dot{\phi} \end{aligned}$$

$$\begin{aligned} \Rightarrow M_z &= \sum_i m_i \left[r_i \cos \phi_i \left(\dot{r}_i \sin \phi_i + r_i \cos \phi_i \dot{\phi}_i \right) + r_i \sin \phi_i \left(\dot{r}_i \cos \phi_i - r_i \sin \phi_i \dot{\phi}_i \right) \right] \\ &= \sum_i m_i r_i^2 \dot{\phi}_i \end{aligned}$$

Comparing with

$$\begin{aligned} \sum_i \frac{\partial \mathcal{L}}{\partial \dot{\phi}_i} &= \sum_i \frac{\partial}{\partial \dot{\phi}_i} \left[\frac{1}{2} m_i \left(\dot{r}_i^2 + r_i^2 \dot{\phi}_i^2 + \dot{z}^2 \right) - U(r_i, \phi, z) \right] \\ &= \sum_i m_i r_i^2 \dot{\phi}_i \end{aligned}$$

Chapter 3

The Central Force Problem

3.1 2 Body to 1 Body Problem and Reduced Mass

The two body problem: