

# 1 Quotient spaces

Let  $X$  be a Banach space and  $M \subset X$  a closed linear subspace. Let  $\sim$  be the equivalence relation defined through  $x \sim y$  if and only if  $x - y \in M$  and let  $X/M$  be the quotient space with respect to  $\sim$ .

a) For  $[x] \in X/M$  let

$$\|[x]\|_{X/M} := \inf_{y \in [x]} \|y\|_X = \inf_{y \in M} \|x - y\|_X$$

Show that  $\|\cdot\|_{X/M}$  is a norm on  $X/M$ .

*Proof.* Indeed we have that  $\|\cdot\|_{X/M}$  is well-defined, to see this let  $[x], [y] \in X/M$  be such that  $[x] = [y]$ , i.e.  $x \sim y$  which means that we find  $m_0 \in M$  such that  $x - y = m_0$ , thus

$$\begin{aligned} \|[x]\|_{X/M} &= \inf_{m \in M} \|x - m\|_X = \inf_{m \in M} \|y - (m - m_0)\| \\ &= \inf_{\tilde{m} \in M} \|y - \tilde{m}\| = \|[y]\|_{X/M} \end{aligned}$$

since we know that  $M$  is a subspace. Also since  $\|\cdot\|$  is a norm on  $X$  we know that for all  $[x] \in X/M$  we have that  $\|[x]\|_{X/M} < \infty$ . Moreover

$$\|[x]\|_{X/M} = \inf_{y \in M} \|x - y\| = 0$$

Thus we can construct a sequence  $y_n$  in  $M$  such that  $\|x - y_n\| \rightarrow 0$  as  $n \rightarrow \infty$ . But since  $M \subset X$  is closed we have that  $y_n \rightarrow x \in M$ , henceforth  $[x] = 0$ .

Next we have

$$\begin{aligned} \|[x] + [y]\|_{X/M} &= \|[x + y]\|_{X/M} = \inf_{v \in M} \|x + y - v\| \\ &= \inf_{v \in M} \|x - v + y - v + v\| \\ &\leq \inf_{v \in M} \|x - v\| + \inf_{v \in M} \|y - v\| + \inf_{v \in M} \|v\| \\ &= \inf_{v \in M} \|x - v\| + \inf_{v \in M} \|y - v\| \\ &= \|[x]\|_{X/M} + \|[y]\|_{X/M} \end{aligned}$$

Finally we have for all  $\lambda \in \mathbb{K}$

$$\begin{aligned}
\|\lambda[x]\|_{X/M} &= \|[\lambda x]\|_{X/M} = \inf_{y \in M} \|\lambda x - y\| \\
&= \inf_{\lambda y \in M} \|\lambda x - \lambda y\| \\
&= \inf_{u \in M} \|\lambda x - \lambda u\| \\
&= \inf_{u \in M} |\lambda| \|x - u\| = |\lambda| \|x\|_{X/M}
\end{aligned} \tag{*}$$

Where we used in (\*) that  $\varphi_\lambda : M \rightarrow M$  given by  $\varphi_\lambda(x) = \lambda x$  is a bijection for all  $\lambda \in \mathbb{K}$ . Thus we have shown that  $\|\cdot\|_{X/M}$  is a norm on  $X/M$ .  $\square$

**b)** Show that the projection  $\pi : X \rightarrow X/M, x \mapsto [x]$  is continuous.

*Proof.* Evidently, the function  $\pi$  is linear because we have by definition that

$$\begin{aligned}
[x + y] &= [x] + [y] \\
\lambda[x] &= [\lambda x]
\end{aligned}$$

For all  $x, y \in X, \lambda \in \mathbb{K}$ . Moreover we have

$$\|\pi(x)\|_{X/M} = \|[x]\|_{X/M} = \inf_{y \in M} \|x - y\| \leq \inf_{y \in M} \|x\| + \underbrace{\inf_{y \in M} \|y\|}_{=0} = \|x\|_X$$

That is  $\pi$  is a bounded linear operator and thus continuous.  $\square$

c) Show that  $X/M$  is complete.

*Proof.* Recall from Exercise Sheet 1 Exercise 4: Normed spaces and Banach spaces where we have shown

**Theorem:** Let  $(X, \|\cdot\|)$  be a normed space.  $(X, \|\cdot\|)$  is a Banach space if and only if, all absolutely convergent series are also convergent, i.e. if

$$\sum_{n=0}^{\infty} \|x_n\| < \infty \implies \sum_{n=0}^{\infty} x_n < \infty$$

Thanks to the above theorem in order to prove that  $X/M$  is a Banach space it is enough to show that every series in  $X/M$  that converges absolutely also converges in  $X/M$ .

To this extent let us take an arbitrary sequence  $([x_n])_{n \in \mathbb{N}}$  in  $X/M$  such that its series is absolutely convergent, i.e. we have

$$\sum_{n=0}^{\infty} \|[x_n]\|_{X/M} < \infty$$

By definition of the quotient norm (via the inf) we have for all  $n \in \mathbb{N}$  the existence of some  $y_n \in M$  such that  $\|x_n - y_n\| \leq \|[x_n]\|_{X/M} + 1/2^n$ . Since we've already established the convergence of the right hand side, the convergence of the left hand side is immediate by domination. But  $X$  is a Banach space, thus it follows from the Theorem above that

$$\sum_{n=0}^{\infty} (x_n - y_n) = x < \infty$$

We now claim that the series over the sequence  $([x_n])_{n \in \mathbb{N}}$  in  $X/M$  converges to  $[x] \in X/M$ .

Indeed, let us consider for  $N \in \mathbb{N}$

$$\left\| [x] - \sum_{n=0}^N [x_n] \right\|_{X/M} = \left\| [x - \sum_{n=0}^N x_n] \right\|_{X/M} \leq \left\| x - \sum_{n=0}^N x_n \right\|_X \rightarrow 0$$

as  $N \rightarrow \infty$ . Where the last inequality just follows by the very definition of the quotient norm.  $\square$

**d)** Let  $X$  and  $Y$  be Banach spaces and  $T \in \mathcal{L}(X, Y)$ .

We define the kernel of  $T$ ,  $\ker T := \{x \in X : Tx = 0\}$  and the range of  $T$ ,  $\text{ran}(T) := \{Tx : x \in X\} \subset Y$ . Let  $\iota : \text{ran} T \rightarrow Y, x \mapsto x$  be the inclusion map. Show that then there exists a bijective operator  $\hat{T} \in \mathcal{L}(X/\ker T, \text{ran} T)$  with  $T = \iota \hat{T} \pi$  and  $\|\hat{T}\| = \|T\|$ .

*Proof.* We have indeed that  $\ker T \subset X$  is a closed linear subspace. Now we define

$$\hat{T} : X/\ker T \rightarrow \text{ran} T, [x] \mapsto T(x)$$

We notice that that  $\hat{T}$  is well-defined. Indeed, let  $[x], [y] \in X/\ker T$  be such that  $[x] = [y]$ , this is the case if and only if  $x \sim y$  i.e.  $x - y \in \ker T$ , or equivalently  $x \in y + \ker T$ . This entails that we can write  $x = y + t$  for some  $t \in \ker T$ . Thus we get

$$\hat{T}([x]) = T(x) = T(y + t) = T(y) + \underbrace{T(t)}_{=0} = T(y) = T([y])$$

Which shows that  $\hat{T}$  is well-defined. Moreover,  $\hat{T}$  is by definition (defined through  $T$ ) surjective onto the range of  $T$ . Furthermore  $\hat{T}$  is injective, let  $[x], [y] \in X/\ker T$  be such that

$$\begin{aligned} \hat{T}([x]) = \hat{T}([y]) &\iff T(x) = T(y) \implies T(x - y) = 0 \\ &\iff x - y \in \ker T \\ &\iff x \sim y \\ &\iff [x] = [y] \end{aligned}$$

The linearity of  $\hat{T}$  follows immediatly by the linearity of  $\pi$  and the linearity of  $T$ . Finally, we have that  $\pi$  is continuous, and by assumption  $T$  is continuous too.

We have seen that  $\|[x]\|_{X/\ker T} \leq \|x\|_X$  and by the continuity of  $T$  we obtain for some  $C > 0$  that

$$\|\hat{T}([x])\|_Y = \|T(x)\|_Y \leq C\|x\|_X$$

let us now set  $\tilde{C} := \max(1, C)$ . This yields that

$$\|\hat{T}([x])\|_Y \leq C\|x\|_X \leq \tilde{C}\|x\|_X = \tilde{C}\|x - 0\|_X$$

Since  $0 \in \ker T$  we conclude that

$$\|\hat{T}([x])\|_Y \leq \inf_{t \in \ker T} \|x - t\| = \|x\|_{X/\ker T}$$

showing that indeed  $\hat{T} \in \mathcal{L}(X/\ker T, \text{ran} T)$ .

Further, by the very definition of  $\hat{T}$  we have that  $T = \iota \hat{T} \pi$ .

Lastly, we observe that since  $0 \in \ker T$  we have that if  $\|x\|_X \leq 1$  for an arbitrary  $x \in X$ , then  $\|x - 0\|_X = \|x\|_X \leq 1$  and thus by definition of the infimum as the greatest lower bound we must also have  $\inf_{y \in \ker T} \|x - y\| \leq 1$ .

Conversely, if  $\inf_{y \in \ker T} \|x - y\| \leq 1$ . Then again by definition of the greatest upper bound we must also have that  $\|x - 0\| = \|x\| \leq 1$ . This shows that

$$\begin{aligned} \|\hat{T}\| &= \sup_{\substack{[x] \in X/\ker T \\ \|[x]\|_{X/\ker T} \leq 1}} \|\hat{T}([x])\|_Y = \sup_{\substack{[x] \in X/\ker T \\ \|[x]\|_{X/\ker T} \leq 1}} \|T(x)\|_Y \\ &= \sup_{\substack{x \in X \\ \|x\|_X \leq 1}} \|T(x)\|_Y = \|T\| \end{aligned}$$

□

## 2 Reflexivity and weak convergence

**a)** Let  $X$  be a normed space and  $X^*$  its dual. Let  $(L_n)_{n \in \mathbb{N}}$  be a sequence in  $X^*$  and  $(x_n)_{n \in \mathbb{N}}$  a sequence in  $X$ . Let  $L_n \xrightarrow{*} L$  in  $X^*$  with respect to the weak-\* topology and  $x_n \rightarrow x$  with respect to the norm in  $X$ . Show that if  $X$  is reflexive, then  $L_n(x_n) \rightarrow L(x)$  as  $n \rightarrow \infty$ .

*Proof.* We recall that the space  $X^{**}$  is always a Banach space, since by assumption  $X$  is reflexive we have  $X \cong X^{**}$ , i.e.  $X$  is a Banach space as well. Thus we know that the reflexivity of  $X$  is equivalent to saying that  $X^*$  is reflexive.

**Claim:** Let  $X$  be a reflexive Banach space, then weak-\* convergence of a sequence  $(f_n)_{n \in \mathbb{N}}$  in  $X^*$  implies weak convergence.

**Proof of Claim:** Let  $(f_n)_{n \in \mathbb{N}}$  be a sequence in  $X^*$  that converges towards  $f \in X^*$  w.r.t.  $\tau_W^*$ , this is the case if and only if  $f_n(x) \rightarrow f(x)$  for all  $x \in X$  as  $n \rightarrow \infty$ .

We want to show that  $f_n \rightharpoonup f$  in  $X^*$ , that is by **Lemma 6.2.1.** equivalent to showing that  $\lambda(f_n) \rightarrow \lambda(f)$  as  $n \rightarrow \infty$  for all  $\lambda \in X^{**}$ .

Since  $X$  is reflexive, the canonical inclusion  $J_X : X \rightarrow X^{**}$  is surjective, that is to say that for every  $\lambda \in X^{**}$  we find an  $x \in X$  such that  $J_X(x) = \lambda$ . But then by the definition of the canonical inclusion we have

$$\lambda(f_n) = J_X(x)(f_n) = f_n(x) \rightarrow f(x) = J_X(x)(f) = \lambda(f)$$

which gives the weak convergence.  $\square$

Next we recall that weakly convergent sequences are always bounded. By our efforts above we now have for the sequence  $(L_n)_{n \in \mathbb{N}}$  in  $X^*$  that  $L_n \xrightarrow{*} L$  in  $X^*$  implies that  $L_n \rightharpoonup L$  in  $X^*$  and the sequence  $(L_n)_{n \in \mathbb{N}}$  is bounded. Let  $M > 0$  be a constant such that  $\|L_n\|_{X^*} \leq M$  for all  $n \in \mathbb{N}$ , then

$$\begin{aligned} |L_n(x_n) - L(x)| &= |L_n(x_n) - L_n(x) + L_n(x) - L(x)| \\ &\leq |L_n(x_n - x)| + |L_n(x) - L(x)| \\ &\leq \|L_n\|_{X^*} \|x_n - x\|_X + |L_n(x) - L(x)| \\ &\leq M \|x_n - x\|_X + |L_n(x) - L(x)| \end{aligned}$$

Letting  $n \rightarrow \infty$  the right hand side converges to zero since  $x_n \rightarrow x$  with respect to the norm in  $X$  and  $L_n$  converges in the sense of weak-\* converges towards  $L$  (see characterization at begin of proof of claim).  $\square$

b) Let  $X$  and  $Y$  be normed vector spaces and let  $T : X \rightarrow Y$  be linear. Furthermore let  $T$  be such that

$$x_n \rightharpoonup 0 \text{ in } X \implies Tx_n \rightharpoonup 0 \text{ in } Y$$

Let  $X$  be reflexive, show that then  $T$  is bounded.

*Proof.* For the sake of contradiction, assume that  $T$  is not bounded, this means that for every  $n \in \mathbb{N}$  there exists  $x_n \in X$  such that  $\|Tx_n\| \geq n^3\|x_n\|$ . We can easily choose  $x_n$  to be normed to 1. Next we define  $\tilde{x}_n := x_n/n$  and notice that since  $x_n$  is normed that  $\|\tilde{x}_n\|_X \rightarrow 0$  as  $n \rightarrow \infty$ .

Moreover we have

$$\|T\tilde{x}_n\| = \frac{1}{n}\|T(x_n)\| \geq \frac{1}{n} \cdot n^3\|x_n\| = n^2$$

Thus we have shown that we can easily construct a sequence  $(x_n)_{n \in \mathbb{N}}$  in  $X$  that converges in norm to 0 but for which we have

$$\|Tx_n\| \geq n^2 \text{ for all } n \in \mathbb{N}$$

Let us therefore take such a sequence  $(x_n)_{n \in \mathbb{N}}$  as above. We then have that  $x_n \rightarrow 0$  in the strong (i.e. norm) sense and thus in particular  $x_n \rightharpoonup 0$ . By our initial assumption this gives that  $T(x_n) \rightharpoonup T(0) = 0$  in  $Y$ . This means that for all  $f \in Y^*$  we have that

$$f(T(x_n)) \rightarrow f(0) = 0$$

Furthermore we can view  $T(x_n)$  as an element  $\Psi_n$  of  $Y^{**}$  by using the canonical embedding  $J_Y : Y \rightarrow Y^{**}$ , setting  $\Psi_n := J_Y(T(x_n)) \in Y^{**}$ . By definition of the canonical embedding we then have

$$\Psi_n(f) := J_Y(T(x_n))(f) = f(T(x_n)) \rightarrow 0 \text{ for all } f \in Y^*$$

This yields that we have

$$\sup_{n \in \mathbb{N}} |\Psi_n(f)| \leq c_f$$

Applying the Banach-Steinhaus Theorem to  $Y^{**}$  we further get that there exists a constant  $c \geq 0$  such that

$$\sup_{n \in \mathbb{N}} \|\Psi_n\|_{Y^{**}} \leq c$$

But we recall that the canonical embedding  $J_Y$  is an isometry between  $Y$  and  $Y^{**}$ . Thus our derived bound yields

$$\sup_{n \in \mathbb{N}} \|\Psi_n\|_{Y^{**}} = \sup_{n \in \mathbb{N}} \|J_Y(T(x_n))\|_{Y^{**}} = \sup_{n \in \mathbb{N}} \|T(x_n)\|_Y \leq c$$

Which is absurd because we have constructed our sequence to satisfy

$$\|T(x_n)\|_Y \geq n^2 \rightarrow \infty \text{ as } n \rightarrow \infty$$

arriving at a contradiction, we conclude that  $T$  must be indeed bounded.  $\square$

**Remark 2.1.** We did not make use of the fact that  $X$  is reflexive, maybe it is possible to show the above in a more elegant way when tightening the conditions for  $X$  to be reflexive.



### 3 Eigenvalues of the Laplace operator

Let  $\Omega \subset \mathbb{R}^n$  be open, bounded and non-empty. We call  $\lambda \in \mathbb{C}$  an eigenvalue of  $-\Delta$  with zero boundary values, if there exists  $u \in H_0^{1,2}(\Omega)$  such that  $\|u\|_{L^2(\Omega)} = 1$  and  $-\Delta u = \lambda u$  in a weak sense, i.e. such that

$$\int_{\Omega} \nabla u \nabla \varphi dx = \lambda \int_{\Omega} u \varphi dx \text{ for all } \varphi \in H_0^{1,2}(\Omega)$$

Then we call  $u$  an eigenfunction to the eigenvalue  $\lambda$ .

**a)** Let  $\lambda_1$  and  $\lambda_2$  be eigenvalues of  $-\Delta$  with eigenfunctions  $u_1$  and  $u_2$ . Show that if  $\lambda_1 \neq \lambda_2$ , then  $u_1$  and  $u_2$  are orthogonal in  $L^2$ , i.e.  $\langle u_1, u_2 \rangle_{L^2(\Omega)} = 0$

*Proof.* We have

$$\langle u_1, u_2 \rangle_{L^2} = \int_{\Omega} u_1(x) u_2(x) dx$$

Since  $u_1, u_2$  are both eigenvalues of  $-\Delta$  with eigenvalues  $\lambda_1$  respectively  $\lambda_2$  we can say that

$$\lambda_1 \int_{\Omega} u_1(x) u_2(x) dx = \lambda_1 \langle u_1, u_2 \rangle_{L^2}$$

using  $\varphi = u_2 \in H_0^{1,2}(\Omega)$  as a test function in the above, similarly we have

$$\lambda_2 \int_{\Omega} u_1(x) u_2(x) dx = \lambda_2 \langle u_1, u_2 \rangle_{L^2}$$

using  $\varphi = u_1 \in H_0^{1,2}(\Omega)$  as a test function in the above definition. We thus obtain that

$$\underbrace{(\lambda_1 - \lambda_2)}_{\neq 0} \langle u_1, u_2 \rangle_{L^2} = 0 \implies \langle u_1, u_2 \rangle_{L^2} = 0$$

since by assumption  $\lambda_1 \neq \lambda_2$ . □