

No Uniform Interpolation in Linear Logic

Rémi Di Guardia   

Université Paris Cité, CNRS, Inria, IRIF, F-75013, Paris, France

Abstract

The literature on uniform interpolation in linear logic is very scarce: the sole known result is that the multiplicative-additive fragment of linear logic has the uniform interpolation property, and so do its intuitionistic and affine variants [1]. We exhibit a simple formula of the unit-free multiplicative-exponential fragment of linear logic that cannot have any uniform interpolant in propositional linear logic. This example shows not only that propositional linear logic does not have the uniform interpolation property, but also that its intuitionistic and affine variants neither do. Furthermore, building on this example, we show the undecidability of the following problem in both classical and intuitionistic linear logic: given a formula, does it have a uniform interpolant?

2012 ACM Subject Classification Theory of computation → Linear logic

Keywords and phrases Linear Logic, Uniform Interpolation, Sequent Calculus, Geometry of Interaction

Funding This work is supported by the European Research Council (ERC) under the European Union's Ninth Framework Programme Horizon Europe (ERC Synergy Project Malinca, Grant Agreement n. 101167526).

Acknowledgements We are thankful to Olivier Laurent for pointing out similar formulas to our counter-example in [15, 4] and a simpler treatment of the intuitionistic case. We also thank Hugo Herbelin and Alexis Saurin for insightful discussions around interpolation.

1 Introduction

Looking at known results for interpolation in linear logic and its fragments, a reader can find a modicum of work, chiefly on Craig's and Lyndon's interpolation and with very little on uniform interpolation. *Craig's interpolation* states that if $A \vdash B$ is provable, then there exists a formula C built only from variables occurring in both A and B and such that both $A \vdash C$ and $C \vdash B$ are provable. *Lyndon's interpolation* is a strengthening of this property taking account the polarity of variables. These two properties have been proved in first-order linear logic and studied in several fragments of this logic. Notably, Roorda [22] proved and disproved Lyndon's interpolation in various fragments of propositional linear logic; Brotherston and Rajeev [2] used display logic to obtain Craig's interpolation in the multiplicative and multiplicative-additive fragments of linear logic, as well as in classical logic (they then formalised these results working with Dawson [5]); Strassburger [25] provides in the calculus of structures for multiplicative-exponential linear logic a result corresponding to Craig's interpolation; a proof of Craig's and Lyndon's interpolation in first-order linear logic, along with links to cut-elimination, has been developed by Saurin [23], who then considered with Fiorillo and Osorio-Valencia [7] this question in the setting of the proof-net syntax of multiplicative linear logic. Let us also mention that Rasga, Carnielli and Sernadas have results for variations of exponential-free linear logic using translations between different logics [21], and that Fussner and Santschi [8] researched Craig's interpolation in extensions of classical and intuitionistic propositional linear logic using algebraic techniques.

The literature is even more scarce on uniform interpolation in linear logic. *Uniform interpolation*, introduced by Pitts [20] in the setting of intuitionistic logic, is a stronger version of Craig's interpolation where the interpolant depends only on one of the two formulas under

consideration. More precisely, a uniform interpolant for a formula A , with respect to a variable X , is a formula C such that:

- the variables of C are included in those of A , except X must not appear in C ;
- $A \vdash C$ is provable;
- for all formulas B such that X does not appear in B and $A \vdash B$ is provable, $C \vdash B$ is provable too.

In particular, uniform interpolants are an interpretation of first-order quantifiers in the logic under consideration, since the formula C above behaves like $\exists X A$ (at least for provability purposes). Not every logic has uniform interpolation, for instance the modal logic S4 does not [10] even though it has Craig’s and Lyndon’s interpolations [18, 9]. It is easy to check that classical logic has this property: a uniform interpolant for a formula A with respect to X is simply $A[\top/X] \vee A[\perp/X]$, that is the disjunction of substituting X by true or false. The foundational work of Pitts [20] proves that intuitionistic logic has uniform interpolation. The algorithm to compute an interpolant is more complex than in the classical case, and the proof technique is based on a syntactic approach. This method by Pitts was then adapted to many other logics. In particular, it was used by Alizadeh, Derakhshan and Ono [1] to prove uniform interpolation in the Lambek calculus, as well as in classical and intuitionistic multiplicative-additive linear logic – and also in the affine variants of these two. This is, at the best of the author’s knowledge, the *sole* result on uniform interpolation in linear logic to this day. Pitts’ method does not extend easily to full propositional linear logic because of the contraction rule, that has a premise sequent “bigger” than its conclusion sequent, and that necessitated a particular treatment in Pitts’ paper [20] so as to remove it from the calculus.

Nonetheless, since classical logic and intuitionistic logic both have uniform interpolation, one could expect linear logic to have it too. We show it is not the case by exhibiting a counter-example. Actually, we present formulas A and $(B_n)_{n \in \mathbb{N}}$ such that:

1. an atom X appears in A but not in any of the B_n ;
2. $A \vdash B_n$ is provable for all number n ;
3. for all formulas C such that X does not appear in C , for any number $n \in \mathbb{N}$, if $A \vdash C$ and $C \vdash B_n$ are both provable, then C is of size at least n .

The result immediately follows: A has no uniform interpolant with respect to X , because such an interpolant would have an unbounded size. Furthermore, the counter-example we exhibit is not only one for uniform interpolation in linear logic, but also in its intuitionistic and affine variants. A subtle use of the exponential modality is mandatory inside the formula A , since uniform interpolation holds with no exponential (case of multiplicative-additive linear logic) and with exponentials everywhere (case of classical logic). This complexity of A entails difficulties to prove some results for sequents of the shape $A \vdash C$. In particular, the most challenging part in this paper is the proof of property 3 above, that involves some technical tools from the literature of linear logic, among which geometry of interaction.

Once proved that uniform interpolation does not hold in a given logic, a natural follow-up question is: given a formula, does it have a uniform interpolant? We show that this decision problem is undecidable in both classical linear logic and intuitionistic linear logic. This is done by reducing provability in these logics to the existence, or not, of a uniform interpolant. Strikingly, this reduction follows by a modification of our counter-example for a formula A with no uniform interpolant, keeping in particular the “shape” of A .

Outline. We first recall the sequent calculus syntax of classical linear logic in Section 2, along with a definition of uniform interpolation in this setting. Then, we introduce some useful but technical tools in Section 3. The main part, Section 4, consists of a presentation

$$\begin{array}{c}
\frac{}{\vdash X^+, X^-} (ax) \quad \frac{\vdash A, \Gamma \quad \vdash A^\perp, \Delta}{\vdash \Gamma, \Delta} (cut) \\
\\
\frac{\vdash A, B, \Gamma}{\vdash A \wp B, \Gamma} (\wp) \quad \frac{\vdash A, \Gamma \quad \vdash B, \Delta}{\vdash A \otimes B, \Gamma, \Delta} (\otimes) \quad \frac{\vdash \Gamma}{\vdash \perp, \Gamma} (\perp) \quad \frac{}{\vdash 1} (1) \\
\frac{\vdash A, \Gamma \quad \vdash B, \Gamma}{\vdash A \& B, \Gamma} (\&) \quad \frac{\vdash A, \Gamma}{\vdash A \oplus B, \Gamma} (\oplus_1) \quad \frac{\vdash B, \Gamma}{\vdash A \oplus B, \Gamma} (\oplus_2) \quad \frac{}{\vdash \top, \Gamma} (\top) \\
\frac{\vdash A, \Gamma}{\vdash ?A, \Gamma} (?d) \quad \frac{\vdash ?A, ?A, \Gamma}{\vdash ?A, \Gamma} (?c) \quad \frac{\vdash \Gamma}{\vdash ?A, \Gamma} (?w) \quad \frac{\vdash A, ?\Gamma}{\vdash !A, ?\Gamma} (!)
\end{array}$$

■ **Figure 1** Inference Rules of propositional Linear Logic

of our counter-example and a proof it cannot have a uniform interpolant using the tools from the preceding section. It is followed in Section 5 by a proof that our counter-example is also one for uniform interpolation in *intuitionistic* linear logic and in *affine* logic (both classical and intuitionistic). Finally, our counter-example is adapted in Section 6 so as to obtain undecidability of whether a uniform interpolant exists, in classical linear logic as well as in its intuitionistic variant.

2 Linear Logic & Uniform Interpolation

We present here linear logic in a standard way as a *unilateral* sequent calculus [11]. In particular, a proof is a *derivation* (tree) whose conclusion sequent is of the shape $\vdash \Gamma$.

Formulas are defined by the following grammar, with X an atom in some countable set:

$$A, B ::= \underbrace{X^- \mid X^+}_{\text{atom}} \mid \underbrace{A \wp B \mid A \otimes B \mid \perp \mid 1}_{\text{multiplicative}} \mid \underbrace{A \& B \mid A \oplus B \mid \top \mid 0}_{\text{additive}} \mid \underbrace{?A \mid !A}_{\text{exponential}}$$

Orthogonality $(\cdot)^\perp$ (*a.k.a.* negation, duality) is an involution defined inductively by:

$$\begin{array}{llll}
(X^-)^\perp \stackrel{\text{def}}{=} X^+ & (A \wp B)^\perp \stackrel{\text{def}}{=} A^\perp \otimes B^\perp & (A \& B)^\perp \stackrel{\text{def}}{=} A^\perp \oplus B^\perp & (?A)^\perp \stackrel{\text{def}}{=} !A^\perp \\
(X^+)^\perp \stackrel{\text{def}}{=} X^- & (A \otimes B)^\perp \stackrel{\text{def}}{=} A^\perp \wp B^\perp & (A \oplus B)^\perp \stackrel{\text{def}}{=} A^\perp \& B^\perp & (!A)^\perp \stackrel{\text{def}}{=} ?A^\perp \\
\perp^\perp \stackrel{\text{def}}{=} 1 & \top^\perp \stackrel{\text{def}}{=} 0 & & \\
1^\perp \stackrel{\text{def}}{=} \perp & 0^\perp \stackrel{\text{def}}{=} \top & &
\end{array}$$

In addition to the connectives in the above grammar, we define another binary connective known as **linear implication** by $A \multimap B \stackrel{\text{def}}{=} A^\perp \wp B$. **Rules** of the logic are given on Figure 1, where A and B stand for arbitrary formulas, Γ and Δ for sets of (occurrences of) formulas, and $?\Gamma$ for a set of (occurrences of) $?$ -formulas. Remark we restrict the *ax*-rule to *atomic formulas*, a well-known simplification. This sequent calculus is equipped with a **cut-elimination** procedure, denoted by $\rightarrow$, that is weakly normalising and that we recall in Appendix A. Let us also mention that linear logic has the **sub-formula property** [11]: given a *cut-free* derivation π of conclusion sequent $\vdash \Gamma$, all the formulas appearing in all sequents of π are sub-formulas of the formulas in Γ .

Uniform interpolation in linear logic can be stated as follows. We denote by $\mathcal{V}(A)$ the set of atoms occurring in the formula A – e.g. $\mathcal{V}((X^+ \wp X^-) \& Y^-) = \{X, Y\}$.

► **Definition 1** (Uniform Interpolant). *Given a formula A and an atom X , a **uniform interpolant** for A with respect to X is a formula C such that:*

- $\mathcal{V}(C) \subseteq \mathcal{V}(A) \setminus \{X\}$;
- $\vdash A^\perp, C$ is provable;

120 ■ for all formulas B , if $X \notin \mathcal{V}(B)$ and $\vdash A^\perp, B$ is provable then $\vdash C^\perp, B$ is provable.
 121 The **uniform interpolation property** holds when every formula admits a uniform inter-
 122 polant with respect to every atom.

123 3 Toolbox

124 We present here two tools used to prove that our counter-example has no uniform interpolant.
 125 The first is a variant of *slices*, the second of *Geometry Of Interaction*. They are needed
 126 solely to prove our most technical statement in Section 4, corresponding to property 3 in the
 127 introduction, as well as the look-alike statement in Section 6.

128 3.1 A variant of Slices

129 The $\&$ -rule stands apart in linear logic since it is the sole rule with some sharing of the
 130 context Γ . A useful tool when considering $\&$ -rules is the notion of a *slice* [11, 13] which is a
 131 derivation missing some additive component: a slice is obtained from a derivation by deleting
 132 one sub-tree of each $\&$ -rule. In a slice, no rule duplicates a context. As our framework has
 133 not only additive but also exponential rules, the well-behaved concept is the one of a *0-slice*,
 134 meaning we only slice the $\&$ -rules that are not above a $!$ -rule, *i.e.* those at depth 0. This
 135 notion of a slice already appeared in the literature, see *e.g.* [19].

136 ► **Definition 2** (0-slice). For π a derivation, a *0-slice* of π is a (usually non-derivation) tree
 137 π' obtained by deleting one of the two sub-trees of each $\&$ -rule of π that is not above a $!$ -rule.
 138 Thus, in π' some (but maybe not all) $\&$ -rules are unary, yielding $\&_1$ - and $\&_2$ -rules:

$$139 \quad \frac{\vdash A, \Gamma}{\vdash A \& B, \Gamma} (\&_1) \quad \frac{\vdash B, \Gamma}{\vdash A \& B, \Gamma} (\&_2)$$

140 ► **Example 3.** Here is a derivation followed by its two 0-slices (note it has three slices):

$$141 \quad \frac{\frac{\frac{\overline{\vdash 1}^{(1)} \quad \overline{\vdash 1}^{(1)}}{\vdash 1 \& 1} (\&) \quad \frac{\vdash 1 \& 1}{\vdash !(1 \& 1)} (!)}{\vdash \top, !(1 \& 1)} (\top) \quad \frac{\vdash \top, !(1 \& 1)}{\vdash \top \& \perp, !(1 \& 1)} (\&_1) \quad \frac{\frac{\frac{\overline{\vdash 1}^{(1)} \quad \overline{\vdash 1}^{(1)}}{\vdash 1 \& 1} (\&) \quad \frac{\vdash 1 \& 1}{\vdash !(1 \& 1)} (!)}{\vdash \perp, !(1 \& 1)} (\perp)}{\vdash \top \& \perp, !(1 \& 1)} (\&_2)$$

142 **Cut-elimination** can be defined in the natural way for 0-slices, taking the definition for
 143 derivations and extending the $\& - \oplus$ key case with

$$144 \quad \frac{\frac{\vdash A, \Gamma}{\vdash A \& B, \Gamma} (\&_1) \quad \frac{\vdash A^\perp, \Delta}{\vdash A^\perp \oplus B^\perp, \Delta} (\oplus_1)}{\vdash \Gamma, \Delta} (cut) \longrightarrow \frac{\vdash A, \Gamma \quad \vdash A^\perp, \Delta}{\vdash \Gamma, \Delta} (cut)$$

$$145 \quad \frac{\frac{\vdash B, \Gamma}{\vdash A \& B, \Gamma} (\&_2) \quad \frac{\vdash B^\perp, \Delta}{\vdash A^\perp \oplus B^\perp, \Delta} (\oplus_2)}{\vdash \Gamma, \Delta} (cut) \longrightarrow \frac{\vdash B, \Gamma \quad \vdash B^\perp, \Delta}{\vdash \Gamma, \Delta} (cut)$$

146 as well as replacing $\&$ -rules with $\&_1$ - or $\&_2$ -rules during a $?d - !$ key case (where a $!$ -rule is
 147 removed). Observe it is not possible to eliminate *cut*-rules of the two following shapes:

$$148 \quad \frac{\frac{\vdash A, \Gamma}{\vdash A \& B, \Gamma} (\&_1) \quad \frac{\vdash B^\perp, \Delta}{\vdash A^\perp \oplus B^\perp, \Delta} (\oplus_2)}{\vdash \Gamma, \Delta} (cut) \quad \frac{\frac{\vdash B, \Gamma}{\vdash A \& B, \Gamma} (\&_2) \quad \frac{\vdash A^\perp, \Delta}{\vdash A^\perp \oplus B^\perp, \Delta} (\oplus_1)}{\vdash \Gamma, \Delta} (cut)$$

149 The only property of 0-slices we need is their behaviour with respect to cut-elimination.

150 ► **Lemma 4.** *Let π and ϕ be derivations such that $\pi \longrightarrow \phi$. For each 0-slice ϕ' of ϕ , there*
 151 *exists a 0-slice π' of π such that $\pi' \longrightarrow \phi'$ or $\pi' = \phi'$.*

152 **Proof.** We can check that each cut-elimination step respects this property, with the equality
 153 case coming from a reduction on rules not in the considered 0-slice. Some cases are detailed
 154 in Appendix C. ◀

155 ► **Remark 5.** Observe a corresponding result for slices does not hold, because a formula of the
 156 shape $?(A \oplus B)$ can be associated to both a $\oplus_1$ - and a $\oplus_2$ -rule. This is a phenomenon at play,
 157 e.g., in the Seely isomorphism $?(A \oplus B) \simeq ?A \wp ?B$. The cut-elimination case responsible for
 158 this failure is the $?c - !$ key case, because the two copies of the duplicated derivation (the “!
 159 box”) may not delete the same sub-trees for all of their corresponding $\&$ -rules.

160 3.2 A variant of Geometry Of Interaction

161 We define here the key tool for proving our counter-example has no uniform interpolant,
 162 previously used in the restricted setting of unit-free multiplicative-additive linear logic [6]. It
 163 falls within the spirit of *Geometry Of Interaction*, shortened as *GOI* [12]. The quite unusual
 164 part is that in the next section we will look at *connectives* of the *cut*-formulas along a path
 165 from GOI, whereas usually one ignores those.

166 ► **Definition 6** (GOI projection graph). *The **GOI projection graph** $\mathcal{G}_\pi$ of a derivation, or*
 167 *of a 0-slice, π of conclusion sequent $\vdash \Gamma$ is the undirected edge-bicoloured graph defined as*
 168 *follows, colors being **ax** and **cut**.*


169 **Vertices:** *All occurrences of atomic formulas X^+ and X^- (for all atoms) from the formulas*
 170 *of Γ and from the cut-formulas A and $A^\perp$ in π , with each cut-rule yielding a pair of*
 171 *cut-formulas giving separate and dual occurrences.*

172 **ax-edges:** *For each pair of occurrences X^+ and X^- such that π contains an ax-rule*
 173 $\frac{}{\vdash X^+, X^-} (ax)$, *there is an ax-edge between X^+ and X^- .*

174 **cut-edges:** *For each cut-rule $\frac{\vdash A, \Delta \quad \vdash A^\perp, \Sigma}{\vdash \Delta, \Sigma} (cut)$ in π , there is a cut-edge between dual*
 175 *occurrences from the two formulas A and $A^\perp$.*

176 In our figures, **ax**-edges are coloured in **azure** and depicted above the vertices, whereas
 177 **cut**-edges are coloured in **crimson** and depicted below the vertices.

178 ► **Example 7.** Here is a derivation π followed by its GOI projection graph $\mathcal{G}_\pi$:




► Remark 8. Observe they are two ways to obtain several ax -edges on a same occurrence: this occurrence can be duplicated using a $?c$ -rule on one of its ancestor (in the formula tree), or by being in the context of a $\&$ -rule. Both cases can be observed in Example 7.

We will consider **alternating** paths in a GOI projection graph, which are paths¹ where consecutive edges are coloured differently – *e.g.* we start with an ax -edge, then take a cut -edge, then an ax -edge, and so on. We have a typical result of GOI about *persistent* paths (see *e.g.* [3]): some paths are preserved through cut-elimination. What is relevant for our purposes is that this holds not only for cut-elimination between derivations but also between 0-slices.

► **Lemma 9.** *Consider a cut-elimination step $\pi \rightarrow \phi$ between two derivations, or between two 0-slices, of conclusion $\vdash \Gamma$. Let u and v be occurrences of atoms from Γ . If there is an alternating path between u and v in $\mathcal{G}_\phi$, then there is an alternating path between them in $\mathcal{G}_\pi$.*

Proof. By inspection on the kind of the cut-elimination step. Since we only care about ax -rules and atoms, the non-immediate cases are either trivial with $\mathcal{G}_\phi \subseteq \mathcal{G}_\pi$ ($\& - \oplus$ key cases, $?d - !$ key case, $?w - !$ key case, $\top - cut$ commutative case) or easy to check (ax key case, $?c - !$ key case, $\& - cut$ commutative case). A few of these cases are given in Appendix C. ◀

► Remark 10. It is easy to find a counter-example to the converse of Lemma 9. One could be tempted to consider not only ax -rules but also $\top$ -rules in a GOI projection graph, also adding vertices for all occurrences of units. Such an altered setting does not suit our purposes for Lemma 9 would not hold – consider the following, with the occurrences of $\top$ and X^+ :



4 A Counter-Example to Uniform Interpolation

Let us fix two atoms X and Y . Here is our counter-example formula A that has no uniform interpolant in linear logic, along with an infinite family of formulas B_n used to prove it:

$$A \stackrel{\text{def}}{=} X^+ \otimes ! (X^+ \multimap (X^+ \otimes Y^+)) \otimes (X^+ \multimap Y^+) \quad \begin{cases} B_0 & \stackrel{\text{def}}{=} Y^+ \otimes Y^+ \\ B_{n+1} & \stackrel{\text{def}}{=} B_n \otimes Y^+ \end{cases} \quad (1)$$

The main intuition is the following. The formula A is made of three parts: X^+ , a “reusable machine” taking as input a X^+ and returning a X^+ and a Y^+ , and a “one-use machine” taking as input a X^+ and returning a Y^+ . Thence, A can “produce” any non-null number of Y^+ , so that $A \vdash B_n$ is provable for all $n \in \mathbb{N}$. Therefore, one may think that $!Y^+ \otimes Y^+$ is a uniform interpolant for A , and indeed we do have $!Y^+ \otimes Y^+ \vdash B_n$ provable; however, $A \vdash !Y^+ \otimes Y^+$ cannot be proved. This generalizes to all “uniform interpolant candidates”.

¹ A path is a finite (and non-empty) alternating sequence of vertices and edges $(v_0, e_1, v_1, e_2, v_2, \dots, e_n, v_n)$ such that for all $i \in \{1, \dots, n\}$, the two endpoints of e_i are v_{i-1} and v_i .

Let us point out that A can be viewed as some “pseudo ! exponential” of Y^+ , since the following dereliction and selection rules for Y^- are admissible:

$$\frac{\vdash Y^-, \Gamma}{\vdash A^\perp, \Gamma} \quad \frac{\vdash Y^-, A^\perp, \Gamma}{\vdash A^\perp, \Gamma}$$

The formula A is reminiscent of two other “pseudo ! exponentials” of Y^+ from the literature:

■ $Y^+ \otimes !(Y^+ \multimap (Y^+ \otimes Y^+)) \otimes !(Y^+ \multimap 1)$ used by Yves Lafont to discuss some non-admissible !-introduction rule [15, page 14];

■ $X^+ \otimes !(X^+ \multimap (X^+ \otimes X^+)) \otimes !(X^+ \multimap 1) \otimes !(X^+ \multimap Y^+)$ used by Anupam Das to prove different the exponentials of linear logic and their encoding in multiplicative-additive linear logic with least and greatest fixed points [4] – actually the formula in the reference is not exactly the previous one but an isomorphic formula.

The later formula by Anupam Das could also be used here in place of our A . Nonetheless, we preferred the definition in Equation (1) because it is smaller and because we do not need the full power of an exponential – in particular not the $?w$ -rule.

To obtain that there is no uniform interpolant for A , it stays to check the three properties claimed in the introduction: $X \notin \mathcal{V}(B_n)$; and $A \vdash B_n$ is provable; and for all formulas C such that $X \notin \mathcal{V}(C)$, if $A \vdash C$ and $C \vdash B_n$ are both provable, then C is of size at least n . Note that the first property holds by definition. The second is easy to prove.

► **Lemma 11.** *For all $n \in \mathbb{N}$, there is a derivation of $\vdash A^\perp, B_n$.*

Proof. We build a derivation π_n of $\vdash X^-, ?(X^+ \otimes (X^- \wp Y^-)), X^+ \otimes Y^-, B_n$ by induction on $n \in \mathbb{N}$. The result then follows by exhibiting this derivation:

$$\frac{\frac{\frac{\vdash X^-, ?(X^+ \otimes (X^- \wp Y^-)), X^+ \otimes Y^-, B_n}{\vdash X^-, ?(X^+ \otimes (X^- \wp Y^-)) \wp (X^+ \otimes Y^-), B_n} (\wp)}{\vdash X^- \wp ?(X^+ \otimes (X^- \wp Y^-)) \wp (X^+ \otimes Y^-), B_n} (\wp)$$

We define the wanted derivations π_n as:

$$\left\{ \begin{array}{l} \pi_0 \stackrel{\text{def}}{=} \frac{\frac{\frac{\frac{\frac{\vdash X^-, X^+}{\vdash X^-, X^+} (ax)}{\vdash X^-, X^+ \otimes (X^- \wp Y^-), X^+ \otimes Y^-, Y^+ \otimes Y^+} (\otimes)}{\vdash X^-, X^+ \otimes (X^- \wp Y^-), X^+ \otimes Y^-, Y^+ \otimes Y^+} (\otimes)}{\vdash X^-, ?(X^+ \otimes (X^- \wp Y^-)), X^+ \otimes Y^-, Y^+ \otimes Y^+} (?d)} \\ \pi_{n+1} \stackrel{\text{def}}{=} \frac{\frac{\frac{\frac{\frac{\frac{\frac{\vdash X^-, X^+}{\vdash X^-, X^+} (ax)}{\vdash X^-, X^+ \otimes (X^- \wp Y^-), ?(X^+ \otimes (X^- \wp Y^-)), X^+ \otimes Y^-, B_n \otimes Y^+} (\otimes)}{\vdash X^-, X^+ \otimes (X^- \wp Y^-), ?(X^+ \otimes (X^- \wp Y^-)), X^+ \otimes Y^-, B_n \otimes Y^+} (\otimes)}{\vdash X^-, ?(X^+ \otimes (X^- \wp Y^-)), ?(X^+ \otimes (X^- \wp Y^-)), X^+ \otimes Y^-, B_n \otimes Y^+} (?d)}{\vdash X^-, ?(X^+ \otimes (X^- \wp Y^-)), X^+ \otimes Y^-, B_n \otimes Y^+} (?c)} \end{array} \right.$$

234

235 The third wished property is more complex to prove: we use the tools from Section 3. It
 236 also needs an intermediate result, that is not surprising, albeit a little knotty to prove.

237 ► **Lemma 12.** *Let C be a formula such that $X \notin \mathcal{V}(C)$ and $\vdash A^\perp, C$ is provable. There*
 238 *exists a cut-free derivation of $\vdash A^\perp, C$ in which no $!$ -rule has $?(X^+ \otimes (X^- \wp Y^-))$ in its*
 239 *conclusion sequent.*

240 **Proof.** Consider a cut-free derivation π of $\vdash A^\perp, C$ with a minimal number of $!$ -rules having
 241 $?(X^+ \otimes (X^- \wp Y^-))$ in their conclusion sequents. We prove this number is null.

242 By the sub-formula property and the constraint on the context in a $!$ -rule, any given
 243 $!$ -rule in π is of the shape

$$\frac{\phi \quad \vdash (? (X^+ \otimes (X^- \wp Y^-)))^i, D, ?\Delta}{\vdash (? (X^+ \otimes (X^- \wp Y^-)))^i, !D, ?\Delta} (!)$$

245 with a conclusion sequent containing $i \geq 0$ copies of $? (X^+ \otimes (X^- \wp Y^-))$ for some $i \in \mathbb{N}$,
 246 and where $!D, ?\Delta$ are sub-formulas of C . Towards a contradiction, assume $i \geq 1$. Substituting
 247 X by 0 in ϕ yields a derivation $\phi[0/X]$ of $\vdash (? (0 \otimes (\top \wp Y^-)))^i, D, ?\Delta$ since, by hypothesis,
 248 there is no X in $D, ?\Delta$. Consider any result τ of fully eliminating cut-rules in

$$\frac{\frac{\frac{\vdash \top, 0 \otimes Y^+}{\vdash \top \wp (0 \otimes Y^+)} (\top) \quad \frac{\vdash \top \wp (0 \otimes Y^+)}{\vdash !(\top \wp (0 \otimes Y^+))} (\wp) \quad \frac{\vdash !(\top \wp (0 \otimes Y^+))}{\vdash !(\top \wp (0 \otimes Y^+))} (!)}{\vdash !(\top \wp (0 \otimes Y^+))} (!) \quad \frac{\phi[0/X] \quad \vdash (? (0 \otimes (\top \wp Y^-)))^i, D, ?\Delta}{\vdash (? (0 \otimes (\top \wp Y^-)))^i, D, ?\Delta} (?c)}{\vdash (? (0 \otimes (\top \wp Y^-)))^i, D, ?\Delta} (cut)}{\vdash D, ?\Delta} \quad \frac{\vdash D, ?\Delta}{\vdash !D, ?\Delta} (!)}{\vdash (? (X^+ \otimes (X^- \wp Y^-)))^i, !D, ?\Delta} (?w)$$

250 where doubled inference rules mean we apply the rule several times depending on i . Then,
 251 τ has no $!$ -rule having $? (X^+ \otimes (X^- \wp Y^-))$ in its conclusion sequent by the sub-formula
 252 property. Therefore, the derivation obtained from π by substituting with τ the sub-tree

$$\frac{\phi \quad \vdash (? (X^+ \otimes (X^- \wp Y^-)))^i, D, ?\Delta}{\vdash (? (X^+ \otimes (X^- \wp Y^-)))^i, !D, ?\Delta} (!)$$

253 contradicts the minimality assumption on π . ◀

254 ► **Remark 13.** The above lemma is the part of the demonstration that fails in presence
 255 of quantifiers – remember that uniform interpolation is trivial with first-order quantifiers.
 256 Indeed, setting $C \stackrel{\text{def}}{=} \exists X A$, the sequent $\vdash A^\perp, C$ is provable but all its cut-free derivations
 257 contain some $!$ -rule having $? (X^+ \otimes (X^- \wp Y^-))$ in its conclusion sequent – simply because
 258 this is the only $?$ -formula and $\vdash ! (Z^+ \multimap (Z^+ \otimes Y^+))$ is not provable for any atom Z . The
 259 reason our proof of Lemma 12 fails with quantifiers is that the sub-formula property no
 260 longer holds as stated in this setting: even if $X \notin \mathcal{V}(C)$, an existential quantifier in C may
 261 be instantiated with X .

262 Using this intermediate result, we now proceed to the core but technical part of the
 263 demonstration: any interpolant between A and B_n must have size at least n .

264 ► **Proposition 14.** *Let C be a formula such that $X \notin \mathcal{V}(C)$. If $\vdash A^\perp, C$ and $\vdash C^\perp, B_n$ are*
 265 *both provable for some number $n \in \mathbb{N}$, then C contains at least $n + 2$ occurrences of Y^+ .*

Proof. Let us call π and ϕ derivations of respectively $\vdash A^\perp, C$ and $\vdash C^\perp, B_n$. We assume them *cut*-free without any loss of generality. We will use the GOI projection graph of (some 0-slice of) their composition by *cut* to identify Y^- 's in $C^\perp$ linked to the Y^+ 's of B_n . Thanks to alternating paths, we will prove that these occurrences of Y^- are pairwise distinct. This yields one Y^- in $C^\perp$ for each Y^+ in B_n , concluding the proof.

Finding an alternating path in some 0-slice. Notice that $\frac{\frac{\pi}{\vdash A^\perp, C} \quad \frac{\phi}{\vdash C^\perp, B_n}}{\vdash A^\perp, B_n} (cut)$ reduces to a *cut*-free derivation of $\vdash A^\perp, B_n$, that we call τ . This τ is its own unique 0-slice, since there is no $\&$ -connective in $A^\perp$ nor in B_n . By repeated applications of Lemma 4, there is a 0-slice μ of $\frac{\frac{\pi}{\vdash A^\perp, C} \quad \frac{\phi}{\vdash C^\perp, B_n}}{\vdash A^\perp, B_n} (cut)$ such that $\mu \rightarrow^* \tau$.

Consider some occurrence Y_α^+ in B_n , with the index α used to distinguish it from other occurrences. Since τ is a *cut*-free derivation of $\vdash A^\perp, B_n$, there must be an *ax*-rule between this occurrence Y_α^+ and one of the two occurrences of Y^- in $A^\perp$: call it Y_1^- . Hence, there is an alternating path (made of one *ax*-edge) between these two occurrences in the GOI projection graph $\mathcal{G}_\tau$ of τ . By repeated applications of Lemma 9, there also is an alternating path p between Y_1^- and Y_α^+ in $\mathcal{G}_\mu$. Thus, there is an occurrence Y_β^- and an *ax*-rule $\frac{}{Y_\beta^-, Y_\alpha^+} (ax)$ in μ and $Y_\beta^- - Y_\alpha^+$ is the last edge of p . This occurrence Y_β^- can only be in $C^\perp$.

We use the alternating path p to show there is no sub-formula $?D$ of $C^\perp$ containing Y_β^- – say differently, there is no $?$ between the root of the formula $C^\perp$ and Y_β^- . Once this claim is proved, we are done: each occurrence of Y^+ in B_n is linked by an *ax*-edge to an occurrence of Y^- in $C^\perp$. All these occurrences of Y^- are pairwise distinct (recall Remark 8): there is no sub-formula $?D$ of $C^\perp$ containing such an occurrence, so they cannot be contracted; and there is no $\&$ -rule that can share the identified *ax*-rules since we are in a 0-slice, and these *ax*-rules cannot be above a $!$ -rule for there is no $?$ -connective nor $!$ -connective below the Y^+ of B_n . Therefore, we have found $n + 2$ occurrences of Y^- in $C^\perp$, as wished.

Exploiting the alternating path. We prove our claim by showing a stronger result:


- for all occurrences of Y^- in C (respectively in $C^\perp$) belonging to p , there is no $?$ between the root of C (respectively of $C^\perp$) and these occurrences;
- for all occurrences of Y^+ in C (respectively in $C^\perp$) belonging to p , there is no $!$ between the root of C (respectively of $C^\perp$) and these occurrences.

We proceed by following the edges of p starting from its endpoint Y_1^- in $A^\perp$.

Exploiting the alternating path: base case. Consider the first edge $Y_1^- - Y_2^+$ of p , with Y_2^+ necessarily in C . Using Lemma 12, without any loss of generality, there is no $!$ -connective between the root of C and Y_2^+ . Otherwise, since there is an *ax*-rule on $\vdash Y_1^-, Y_2^+$, there would be a $!$ -rule (associated to this $!$ -connective) with in its conclusion sequent a $?$ -sub-formula of $A^\perp$ containing Y_1^- , meaning with $?(X^+ \otimes (X^- \wp Y^-))$: contradiction.

Exploiting the alternating path: inductive case. See Figure 2 for an illustration of our present reasoning. Assume the results holds for a prefix of p , and let us consider its last edge.

1. Suppose this prefix ends with an *ax*-edge $Y_k^- - Y_{k+1}^+$ with Y_{k+1}^+ in C . We know there is no $!$ between the root of C and Y_{k+1}^+ . As p is alternating, the subsequent edge is a *cut*-edge



■ **Figure 2** Alternating path p in the GOI projection graph $\mathcal{G}_\mu$ in the proof of Proposition 14

- 305 $Y_{k+1}^+ - Y_{k+2}^-$ with Y_{k+2}^- in $C^\perp$. Since Y_{k+1}^+ and Y_{k+2}^- are dual occurrences by definition of
 306 a **cut**-edge, there is no ? between the root of $C^\perp$ and Y_{k+2}^- .
- 307 2. Suppose this prefix ends with a **cut**-edge $Y_k^+ - Y_{k+1}^-$ with Y_{k+1}^- in $C^\perp$. We know there is
 308 no ? between the root of $C^\perp$ and Y_{k+1}^- . As p is alternating, the subsequent edge is an
 309 **ax**-edge $Y_{k+1}^- - Y_{k+2}^+$ with Y_{k+2}^+ either in B_n or in $C^\perp$. In the first case, we are done: we
 310 have $Y_{k+2}^+ = Y_\alpha^+$ since there is no **cut**-edge with endpoint Y_{k+2}^+ to continue our path, and
 311 p is supposed to end in Y_α^+ . In the second case, since there is an **ax**-rule $\frac{}{Y_{k+1}^-, Y_{k+2}^+}^{(ax)}$
 312 by definition of an **ax**-edge, it follows there is no ! between the root of $C^\perp$ and Y_{k+2}^+ –
 313 because the context condition of a corresponding !-rule would prevent the presence of
 314 Y_{k+1}^- in all sequents above it.
- 315 3. Suppose this prefix ends with an **ax**-edge $Y_k^- - Y_{k+1}^+$ with Y_{k+1}^+ in $C^\perp$. This case is identical
 316 to case 1 up to switching C and $C^\perp$.
- 317 4. Suppose this prefix ends with a **cut**-edge $Y_k^+ - Y_{k+1}^-$ with Y_{k+1}^- in C . This case is identical
 318 to case 2, with no need to consider Y_{k+2}^+ in $A^\perp$ since $A^\perp$ has no Y^+ .
- 319 This proves our claim, and concludes the proof. ◀

- 320 ► **Remark 15.** By going in the other direction of the path p , we can similarly show that:
 321 ■ for all occurrences of Y^- in C (respectively in $C^\perp$) belonging to p , there is no ! between
 322 the root of C (respectively of $C^\perp$) and these occurrences;
 323 ■ for all occurrences of Y^+ in C (respectively in $C^\perp$) belonging to p , there is no ? between
 324 the root of C (respectively of $C^\perp$) and these occurrences.
- 325 This means the “useful” part of C as identified by p is exponential-free.

326 Now that we have all needed properties, we easily obtain our main result.

327 ► **Theorem 16.** *Linear logic does not have the uniform interpolation property.*

328 **Proof.** The formula A cannot have a uniform interpolant C with respect to X , as such a C
 329 would have an unbounded number of occurrences of Y^+ by Lemma 11 and Proposition 14. ◀

330 5 Adaptation to the Intuitionistic and Affine variants

331 Our counter-example to the uniform interpolation property in linear logic is also a counter-
 332 example to it in *intuitionistic* linear logic [14]. This can be directly obtained through a result
 333 of Harold Schellinx [24]. Formulas of propositional intuitionistic linear logic are given by:

$$334 \quad A, B ::= X^+ \mid A \multimap B \mid A \otimes B \mid 1 \mid A \& B \mid A \oplus B \mid \top \mid 0 \mid !A$$

335 ► **Proposition 17** ([24, Proposition 3.8]). *Consider a formula D of intuitionist linear logic
 336 which does not contain 0 or does not contain $\multimap$. Then $\vdash D$ is provable in linear logic if and
 337 only if it is provable in intuitionistic linear logic.*

338 ► **Theorem 18.** *Intuitionistic linear logic does not have the uniform interpolation property.*

339 **Proof.** The formula A from Equation (1) has no uniform interpolant with respect to X . In-
 340 deed, $\vdash A \multimap B_n$ is provable intuitionistically for all $n \in \mathbb{N}$ using Lemma 11 and Proposition 17
 341 – for 0 is not in $A \multimap B_n$. This yields an intuitionistic derivation of $A \vdash B_n$:

$$\frac{\frac{\overline{A \vdash A}^{(ax)} \quad \overline{B_n \vdash B_n}^{(ax)}}{A, A \multimap B_n \vdash B_n}^{(\multimap)} \quad \vdash A \multimap B_n}{A \vdash B_n}^{(cut)}$$

343 Hence, we conclude as in the classical case: an interpolant for A would have an unbounded
 344 number of occurrences of Y^+ by Proposition 14 and conservativity – *i.e.* a sequent provable
 345 in intuitionistic linear logic is *a fortiori* provable in (classical) linear logic. ◀

346 Our counter-example is also one to the uniform interpolation property in *affine* linear
 347 logic, both classical and intuitionistic. Affine logic is defined as linear logic where the
 348 $?w$ -rule is replaced by the more general $\frac{\vdash \Gamma}{\vdash A, \Gamma}^{(w)}$. It is equipped with a weakly normalising
 349 cut-elimination procedure – see Appendix B. We need here one more intermediate result.

350 ► **Lemma 19.** *In affine logic, there is no derivation of conclusion $\Gamma \vdash \perp$ where the formulas of*
 351 *Γ are included in $\{X^+, !(X^+ \multimap (X^+ \otimes Y^+)), X^+ \multimap (X^+ \otimes Y^+), X^+ \otimes Y^+, Y^+, X^+ \multimap Y^+\}$,*
 352 *with a same formula appearing possibly several times in Γ .*

353 **Proof.** By contradiction, consider a smallest *cut*-free derivation of $\Gamma \vdash \perp$ with Γ a sequent
 354 of the aforementioned shape. Its last rule is either a $\otimes$ - $\wp$ - $?d$ - $?c$ - or w -rule. In each case,
 355 one of the premises of this rule is a sequent of the aforementioned shape, which contradicts
 356 the minimality hypothesis. ◀

357 ► **Theorem 20.** *Affine logic and intuitionistic affine logic do not have the uniform interpola-*
 358 *tion property.*

359 **Proof.** We proceed almost exactly as for linear logic, the sole difference being in the beginning
 360 of the proof of Proposition 14. There, that in the normalised derivation τ of $\vdash A^\perp, B_n$ (or
 361 $A \vdash B_n$) there is an ax -rule on each Y^+ of B_n is no longer immediate: we invoke Lemma 19,
 362 which implies no sub-formula of B_n can be weakened in τ – this lemma holding in (classical)
 363 affine logic, so *a fortiori* in intuitionistic affine logic by conservativity. ◀

364 6 Undecidability of the Existence of a Uniform Interpolant

365 A natural question following the failure of the uniform interpolation property is to study the
 366 complexity of the subsequent decision problem:

367 **Uniform interpolant:** Given a formula A and an atom X , does A have a uniform interpolant
 368 with respect to X ?

369 In this section, we show this problem is undecidable in both classical and intuitionistic linear
 370 logic. This is done thanks to a reduction to the provability decision problem, better-studied
 371 in linear logic and known to be undecidable in both the classical and intuitionistic cases [16]:

372 **Provability:** Given a formula F , is $\vdash F$ provable?

373 Our demonstration that Provability can be encoded into Uniform interpolant is very
 374 similar to our proof in Section 4 that our counter-example has no uniform interpolant, up
 375 to additional technicalities. In particular, it is done by adapting our counter-example of

back $1^\perp = \perp$ with the lacking formula:

$$\begin{array}{c}
 \frac{}{\vdash Y^-, Y^+} \text{ (ax)} \quad \frac{\pi}{\vdash F} \text{ (!)} \\
 \hline
 \vdash Y^-, Y^+ \otimes !F \text{ (}\otimes\text{)} \\
 \hline
 \vdash Y^- \wp (Y^+ \otimes !F) \text{ (}\wp\text{)} \quad \vdash \Gamma, Y^+ \otimes (Y^- \wp ?F^\perp) \\
 \hline
 \vdash \Gamma \text{ (cut)} \\
 \hline
 \vdash \Gamma, \perp \text{ (}\perp\text{)} \\
 \vdots
 \end{array}$$

where π is a derivation of conclusion $\vdash F$, using that $\vdash F$ is assumed provable.

Then, we substitute X by 1 in ϕ_2 , obtaining a derivation ϕ_3 of conclusion sequent $\vdash \perp \wp ?(1 \otimes (\perp \wp \perp)) \wp (1 \otimes (Y^+ \otimes (Y^- \wp ?F^\perp)))$, B – remember that $X \notin \mathcal{V}(F) \cup \mathcal{V}(B)$. Now, thanks to the isomorphisms $1 \otimes E \simeq E$ and $\perp \wp E \simeq E$, this means there exists a derivation ϕ_4 of conclusion $\vdash ?\perp \wp (Y^+ \otimes (Y^- \wp ?F^\perp))$, B . Since we have the equiprovability $?\perp \dashv \vdash \perp$, and reusing the isomorphism $\perp \wp E \simeq E$, we finally obtain a derivation ϕ_5 of conclusion $\vdash Y^+ \otimes (Y^- \wp ?F^\perp)$, B , that is of $\vdash C(F)^\perp$, B .

Therefore, $C(F)$ is a uniform interpolant for $A(F)$ with respect to X , as claimed. $\blacktriangleleft$

To show there is no uniform interpolant when $\vdash F$ is not provable, we use the same infinite family of formulas $(B_n)_{n \in \mathbb{N}}$ as in Section 4, up to the substitution $Y^+ \mapsto (Y^+ \multimap (Y^+ \otimes !F))$ that links A in Equation (1) to $A(F)$ in Equation (2). That is, we will consider:

$$\begin{cases} B_0(F) & \stackrel{\text{def}}{=} (Y^+ \multimap (Y^+ \otimes !F)) \otimes (Y^+ \multimap (Y^+ \otimes !F)) \\ B_{n+1}(F) & \stackrel{\text{def}}{=} B_n(F) \otimes (Y^+ \multimap (Y^+ \otimes !F)) \end{cases} \quad (4)$$

With this viewpoint, the two following lemmas are analogous to previous results from Section 4, namely Lemma 11 and Proposition 14.

► **Lemma 22.** *Let F be a formula, and X and Y be atoms. With $A(F)$ and $(B_n(F))_{n \in \mathbb{N}}$ defined as in Equations (2) and (4), the sequent $\vdash A(F)^\perp, B_n(F)$ is provable for all $n \in \mathbb{N}$.*

Proof. It suffices to take the derivations given by Lemma 11 and to apply in each of them the substitution $Y^+ \mapsto (Y^+ \multimap (Y^+ \otimes !F))$. $\blacktriangleleft$

► **Proposition 23.** *Let F and C be formulas, and X and Y be atoms such that $X, Y \notin \mathcal{V}(F)$ and $X \notin \mathcal{V}(C)$. Assume that $\vdash F$ is not provable. With $A(F)$ and $(B_n(F))_{n \in \mathbb{N}}$ defined as in Equations (2) and (4), if $\vdash A(F)^\perp, C$ and $\vdash C^\perp, B_n(F)$ are both provable for some number $n \in \mathbb{N}$, then C contains at least $n + 2$ occurrences of Y^+ .*

Proof. Mostly similar to the proof of Proposition 14, with additional technicalities. For the sake of clarity and brevity, a demonstration is available in Appendix D. $\blacktriangleleft$

► **Proposition 24.** *Let F be a formula of linear logic. Fixing two atoms $X, Y \notin \mathcal{V}(F)$, $\vdash F$ is provable if and only if the formula $A(F)$ defined as in Equation (2) has a uniform interpolant with respect to X .*

Proof. We proceed according to the provability of F . Assume first that F is provable. Then, Proposition 21 yields a uniform interpolant for $A(F)$ with respect to X .

Assume now that F is not provable. We claim that $A(F)$ has no uniform interpolant with respect to X , using the formulas $(B_n(F))_{n \in \mathbb{N}}$ defined as in Equation (4). Observe that $X \notin \mathcal{V}(B_n)$ and that $\vdash A(F)^\perp, B_n$ is provable by Lemma 22, for all $n \in \mathbb{N}$. By Proposition 23, for any formula C such that $X \notin \mathcal{V}(C)$, if $\vdash A(F)^\perp, C$ and $\vdash C^\perp, B_n(F)$ are both provable, then C contains at least $n + 2$ occurrences of Y^+ . Hence, $A(F)$ cannot have any uniform interpolant with respect to X , concluding the proof. $\blacktriangleleft$

► **Theorem 25.** *In (classical) linear logic, the Uniform interpolant problem is undecidable.*

Proof. By Proposition 24, decidability of Uniform interpolant entails decidability of Provability. However, Provability is undecidable in propositional linear logic [16]. $\blacktriangleleft$

As was the case in Section 5 for the failure of the uniform interpolation property, it is possible to adapt our undecidability result to the *intuitionistic* case.

► **Proposition 26.** *Consider the framework of intuitionistic linear logic. Let F be a formula. Fixing two atoms $X, Y \notin \mathcal{V}(F)$, $\vdash F$ is provable if and only if the formula $A(F)$ defined as in Equation (2) has a uniform interpolant with respect to X .*

Proof. One can proceed as in the proof of Proposition 24, using conservativity and that the derivations built in the proofs of Proposition 24 and its intermediate lemmas can be framed as derivations in intuitionistic linear logic instead of classical linear logic. $\blacktriangleleft$

► **Theorem 27.** *In intuitionistic linear logic, the Uniform interpolant problem is undecidable.*

Proof. By Proposition 26, decidability of Uniform interpolant entails decidability of Provability. However, Provability is undecidable in intuitionistic propositional linear logic [16]. $\blacktriangleleft$

7 Conclusion

We presented a formula of unit-free multiplicative-exponential linear logic that has no uniform interpolant in linear logic, both classical and intuitionistic, as well as in classical and intuitionistic affine logic. This contrasts sharply with the situation of classical logic and intuitionistic logic and multiplicative-additive linear logic, that all have the uniform interpolation property. Our counter-example, *c.f.* Equation (1), sheds some light on when the contraction rule prevents uniform interpolation: it may do so when the formula A to interpolate contains both a non-contractible sub-formula (here X^+) and a contractible sub-formula (here $!(X^+ \multimap (X^+ \otimes Y^+))$), which is impossible in the three aforementioned logics but possible in linear logic and affine logic! Furthermore, please observe that our demonstration also holds in a variant of linear logic with no $?w$ -rule and/or with the $?c$ -rule replaced by the weaker selection rule $\frac{\vdash A, ?A, \Gamma}{\vdash ?A, \Gamma}$. This makes quite clear that what prevents uniform interpolation in linear logic is a contraction rule available only on some formulas (along with the constraint on the context in a $!$ -rule).

The formulas A and B_n we exhibited in Equation (1) are not overly complex, but proving no formula C can be a uniform interpolant was quite tedious as it depends heavily on the necessary structure of C . Geometry of interaction and slices were very useful tools to this end – and we do not do how to prove Proposition 14 without GOI projection graphs.

Considering the uniform interpolation property in linear logic and its fragments, since it trivially holds in the presence of quantifiers (setting $C = \exists X A$ as an interpolant for A with respect to X), and it has been proved in multiplicative-additive linear logic [1] and now been disproved for some formula of multiplicative-exponential linear logic (even allowing the

interpolant to use connectives from propositional linear logic), the principal fragment where the question stays open is the (unusual) additive-exponential linear logic. Since in this logic the contraction rule is of limited interest due to the lack of multiplicative connectives, we conjecture the method of Pitts [20] could be adapted to prove uniform interpolation in this fragment, as well as in its intuitionistic and affine variants.

A natural question following the failure of the uniform interpolation property is the subsequent decision problem: given a formula A , does it have a uniform interpolant? We showed this problem to be undecidable in both classical and intuitionistic linear logic, through a reduction to the problem of provability and by modifying our counter-example for uniform interpolation, *c.f.* Equation (2). Our reasoning might perhaps be also adapted to classical and intuitionistic affine logics, but the provability of these logics is decidable, hence we would only get a lower bound for our problem. Furthermore, this decision problem can also be considered for formulas of fragments of (intuitionistic) linear logic, *e.g.* of multiplicative-exponential (intuitionistic) linear logic. It can be easily checked that our demonstration in Section 6 also applies in the restricted setting of multiplicative-exponential linear logic, so that deciding whether a formula of multiplicative-exponential linear logic has a uniform interpolant is at least harder than provability in this fragment – which is an open problem.

References

- 1 Majid Alizadeh, Farzaneh Derakhshan, and Hiroakira Ono. Uniform interpolation in substructural logics. *The Review of Symbolic Logic*, 7(3):455–483, 2014. doi:10.1017/S175502031400015X.
- 2 James Brotherston and Rajeev Goré. Craig interpolation in displayable logics. In Kai Brännler and George Metcalfe, editors, *Proceedings of the 20th International Conference on Automated Reasoning with Analytic Tableaux and Related Methods (TABLEAUX 2011)*, volume 6793, pages 88–103. Springer, 2011. doi:10.1007/978-3-642-22119-4_9.
- 3 Vincent Danos and Laurent Regnier. Proof-nets and the Hilbert space. In Jean-Yves Girard, Yves Lafont, and Laurent Regnier, editors, *Advances in Linear Logic*, volume 222 of *London Mathematical Society Lecture Note Series*, pages 307–328. Cambridge University Press, 1995. doi:10.1017/CB09780511629150.016.
- 4 Anupam Das. Exponentials vs fixed points in linear logic, June 2024. The Proof Theory Blog. URL: <https://prooftheory.blog/2024/06/27/exponentials-vs-fixed-points-in-linear-logic/>.
- 5 Jeremy E. Dawson, James Brotherston, and Rajeev Goré. Machine-checked interpolation theorems for substructural logics using display calculi. In Nicola Olivetti and Ashish Tiwari, editors, *Proceedings of the 8th International Joint Conference on Automated Reasoning (IJCAR 2016)*, volume 9706, pages 452–468. Springer, 2016. doi:10.1007/978-3-319-40229-1_31.
- 6 Rémi Di Guardia. *Identity of Proofs and Formulas using Proof-Nets in Multiplicative-Additive Linear Logic*. Thèse de doctorat, Ecole normale supérieure de lyon - ENS LYON, September 2024. URL: <https://theses.hal.science/tel-04830060>.
- 7 Guido Fiorillo, Daniel Osorio Valencia, and Alexis Saurin. On correctness, sequentialization and interpolation. In Lionel Vaux Auclair, editor, *Ninth International Workshop on Trends in Linear Logic and Applications 2025*, 2025. URL: <https://lipn.univ-paris13.fr/TLLA/2025/abstracts/14-fiorillo-osoriovalencia-saurin.pdf>.
- 8 Wesley Fussner and Simon Santschi. Interpolation in linear logic and related systems. *ACM Transactions on Computational Logic*, 25(4), October 2024. doi:10.1145/3680284.
- 9 Dov M. Gabbay. Craig’s interpolation theorem for modal logics. In Wilfrid Hodges, editor, *Conference in Mathematical Logic — London ’70*, pages 111–127. Springer, 1972. doi:10.1007/BFb0059541.

- 526 10 Silvio Ghilardi and Marek Zawadowski. Undefinability of propositional quantifiers in the
527 modal system S4. *Studia Logica*, 55(2):259–271, June 1995. doi:10.1007/BF01061237.
- 528 11 Jean-Yves Girard. Linear logic. *Theoretical Computer Science*, 50:1–102, 1987. doi:10.1016/
529 0304-3975(87)90045-4.
- 530 12 Jean-Yves Girard. Towards a geometry of interaction. In *Categories in Computer Science*
531 *and Logic*, pages 69–108, Providence, 1989. American Mathematical Society. Proceedings of
532 Symposia in Pure Mathematics n°92. URL: [https://www.ams.org/books/conm/092/1003197/
533 conm092-1003197.pdf](https://www.ams.org/books/conm/092/1003197/conm092-1003197.pdf), doi:10.1090/conm/092.
- 534 13 Jean-Yves Girard. Proof-nets: the parallel syntax for proof-theory. In Aldo Ursini and
535 Paolo Agliano, editors, *Logic and Algebra*, volume 180 of *Lecture Notes In Pure and Applied*
536 *Mathematics*, pages 97–124, New York, 1996. Marcel Dekker. doi:10.1201/9780203748671-4.
- 537 14 Jean-Yves Girard and Yves Lafont. Linear logic and lazy computation. In Hartmut Ehrig,
538 Robert Kowalski, Giorgio Levi, and Ugo Montanari, editors, *Proceedings of the International*
539 *Joint Conference on Theory and Practice of Software Development (TAPSOFT’87)*, volume 250
540 of *Lecture Notes in Computer Science*, pages 52–66. Springer, 1987. doi:10.1007/BFB0014972.
- 541 15 Yves Lafont. The linear logic pages, 1999. URL: [https://www.i2m.univ-amu.fr/perso/yves.
542 lafont/pub/llpages.pdf](https://www.i2m.univ-amu.fr/perso/yves.lafont/pub/llpages.pdf).
- 543 16 Patrick Lincoln. Deciding provability of linear logic formulas. In Jean-Yves Girard, Yves Lafont,
544 and Laurent Regnier, editors, *Advances in Linear Logic*, volume 222 of *London Mathematical*
545 *Society Lecture Note Series*, pages 109–122. Cambridge University Press, 1995.
- 546 17 The LL Handbook Project. *Handbook of Linear Logic (draft)*. January 2022. Early draft for
547 the Linear Logic Winter School. URL: [https://ll-handbook.frama.io/ll-handbook/ll-
548 handbook-public.pdf](https://ll-handbook.frama.io/ll-handbook/ll-handbook-public.pdf).
- 549 18 L. L. Maksimova. Interpolation theorems in modal logics. Sufficient conditions. *Algebra and*
550 *Logic*, 19:120–132, 1980. doi:10.1007/BF01669837.
- 551 19 Michele Pagani and Lorenzo Tortora de Falco. Strong normalization property for second order
552 linear logic. *Theoretical Computer Science*, 411(2):410–444, 2010. doi:10.1016/j.tcs.2009.
553 07.053.
- 554 20 Andrew M. Pitts. On an interpretation of second order quantification in first order intuitionistic
555 propositional logic. *Journal of Symbolic Logic*, 57(1):33–52, 1992. doi:10.2307/2275175.
- 556 21 João Rasga, Walter Carnielli, and Cristina Sernadas. Interpolation via translations. *Mathem-*
557 *atical Logic Quarterly*, 55(5):515–534, September 2009. doi:10.1002/malq.200810013.
- 558 22 Dirk Roorda. Interpolation in fragments of classical linear logic. *Journal of Symbolic Logic*,
559 59(2):419–444, 1994. doi:10.2307/2275398.
- 560 23 Alexis Saurin. Interpolation as cut-introduction: On the computational content of Craig-
561 Lyndon interpolation. In Maribel Fernández, editor, *International Conference on Formal*
562 *Structures for Computation and Deduction (FSCD)*, volume 337 of *Leibniz International*
563 *Proceedings in Informatics (LIPIcs)*, pages 32:1–32:21. Schloss Dagstuhl – Leibniz-Zentrum
564 für Informatik, July 2025. doi:10.4230/LIPIcs.FSCD.2025.32.
- 565 24 Harold Schellinx. Some syntactical observations on linear logic. *Journal of Logic and Compu-*
566 *tation*, 1(4):537–559, 1991. doi:10.1093/LOGCOM/1.4.537.
- 567 25 Lutz Strassburger. *Linear Logic and noncommutativity in the Calculus of Structures*. Ph.D.
568 thesis, Technischen Universität Dresden, May 2003. URL: [https://www.lix.polytechnique.
569 fr/Labo/Lutz.Strassburger/papers/dissvonlutz.pdf](https://www.lix.polytechnique.fr/Labo/Lutz.Strassburger/papers/dissvonlutz.pdf).

570

A Cut-Elimination in Linear Logic

571

Cut-elimination in linear logic is defined on Tables 1 and 2. It is weakly normalising [11].

ax	$\frac{\frac{\overline{\vdash A^\perp, A} \quad (\text{ax}) \quad \pi}{\vdash A, \Gamma} \quad \frac{\pi}{\vdash A, \Gamma} \quad (\text{cut})}{\vdash A, \Gamma} \longrightarrow \frac{\pi}{\vdash A, \Gamma}$
$\wp - \otimes - 1$	$\frac{\frac{\pi}{\vdash A^\perp, B^\perp, \Gamma} \quad (\wp) \quad \frac{\phi}{\vdash A, \Delta} \quad \frac{\tau}{\vdash B, \Sigma} \quad (\otimes)}{\vdash A^\perp \wp B^\perp, \Gamma} \quad (\otimes) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, B^\perp, \Gamma} \quad \frac{\tau}{\vdash B, \Sigma} \quad (\text{cut})}{\vdash A^\perp, \Gamma, \Sigma} \quad \frac{\phi}{\vdash A, \Delta} \quad (\text{cut})}{\vdash \Gamma, \Delta, \Sigma}$
$\wp - \otimes - 2$	$\frac{\frac{\pi}{\vdash A^\perp, B^\perp, \Gamma} \quad (\wp) \quad \frac{\phi}{\vdash A, \Delta} \quad \frac{\tau}{\vdash B, \Sigma} \quad (\otimes)}{\vdash A^\perp \wp B^\perp, \Gamma} \quad (\otimes) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, B^\perp, \Gamma} \quad \frac{\phi}{\vdash A, \Delta} \quad (\text{cut})}{\vdash B^\perp, \Gamma, \Sigma} \quad \frac{\tau}{\vdash B, \Sigma} \quad (\text{cut})}{\vdash \Gamma, \Delta, \Sigma}$
$\perp - 1$	$\frac{\frac{\pi}{\vdash \Gamma} \quad (\perp) \quad \overline{\vdash 1} \quad (1)}{\vdash \Gamma} \quad (\text{cut}) \longrightarrow \frac{\pi}{\vdash \Gamma}$
$\& - \oplus_1$	$\frac{\frac{\pi}{\vdash A^\perp, \Gamma} \quad \frac{\phi}{\vdash B^\perp, \Gamma} \quad (\&) \quad \frac{\tau}{\vdash A, \Delta} \quad (\oplus_1)}{\vdash A^\perp \& B^\perp, \Gamma} \quad (\oplus_1) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, \Gamma} \quad \frac{\tau}{\vdash A, \Delta} \quad (\text{cut})}{\vdash \Gamma, \Delta}$
$\& - \oplus_2$	$\frac{\frac{\pi}{\vdash A^\perp, \Gamma} \quad \frac{\phi}{\vdash B^\perp, \Gamma} \quad (\&) \quad \frac{\tau}{\vdash B, \Delta} \quad (\oplus_2)}{\vdash A^\perp \& B^\perp, \Gamma} \quad (\oplus_2) \longrightarrow \frac{\frac{\phi}{\vdash B^\perp, \Gamma} \quad \frac{\tau}{\vdash B, \Delta} \quad (\text{cut})}{\vdash \Gamma, \Delta}$
$?d - !$	$\frac{\frac{\pi}{\vdash A^\perp, \Gamma} \quad (\text{?d}) \quad \frac{\phi}{\vdash A, ?\Delta} \quad (!)}{\vdash ?A^\perp, \Gamma} \quad (\text{?d}) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, \Gamma} \quad \frac{\phi}{\vdash A, ?\Delta} \quad (\text{cut})}{\vdash \Gamma, ?\Delta}$
$?c - !$	$\frac{\frac{\pi}{\vdash ?A^\perp, ?A^\perp, \Gamma} \quad (\text{?c}) \quad \frac{\phi}{\vdash A, ?\Delta} \quad (!)}{\vdash ?A^\perp, \Gamma} \quad (\text{?c}) \longrightarrow \frac{\frac{\pi}{\vdash ?A^\perp, ?A^\perp, \Gamma} \quad \frac{\phi}{\vdash A, ?\Delta} \quad (!)}{\vdash ?A^\perp, \Gamma, ?\Delta} \quad (\text{?c}) \longrightarrow \frac{\frac{\phi}{\vdash A, ?\Delta} \quad (!)}{\vdash \Gamma, ?\Delta, ?\Delta} \quad (\text{?c}) \longrightarrow \frac{\phi}{\vdash \Gamma, ?\Delta}$
$?w - !$	$\frac{\frac{\pi}{\vdash \Gamma} \quad (\text{?w}) \quad \frac{\phi}{\vdash A, ?\Delta} \quad (!)}{\vdash ?A^\perp, \Gamma} \quad (\text{?w}) \longrightarrow \frac{\frac{\pi}{\vdash \Gamma} \quad (\text{?w})}{\vdash \Gamma, ?\Delta}$

Table 1 Cut-elimination in Linear Logic – Key cases

$cut - cut$	$\frac{\frac{\frac{\pi}{\vdash A^\perp, B^\perp, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash B^\perp, \Gamma, \Delta} (cut) \quad \frac{\tau}{\vdash B, \Sigma}}{\vdash \Gamma, \Delta, \Sigma} (cut) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, B^\perp, \Gamma} \quad \frac{\tau}{\vdash B, \Sigma}}{\vdash A^\perp, \Gamma, \Sigma} (cut) \quad \frac{\phi}{\vdash A, \Delta}}{\vdash \Gamma, \Delta, \Sigma} (cut)$
$\wp - cut$	$\frac{\frac{\frac{\pi}{\vdash A^\perp, B, C, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash A^\perp, B \wp C, \Gamma} (\wp) \quad \frac{\tau}{\vdash A, \Delta}}{\vdash B \wp C, \Gamma, \Delta} (cut) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, B, C, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash B, C, \Gamma, \Delta} (cut) \quad \frac{\tau}{\vdash B \wp C, \Gamma, \Delta} (\wp)$
$\otimes - cut - 1$	$\frac{\frac{\frac{\pi}{\vdash A^\perp, B, \Gamma} \quad \frac{\phi}{\vdash C, \Delta}}{\vdash A^\perp, B \otimes C, \Gamma, \Delta} (\otimes) \quad \frac{\tau}{\vdash A, \Sigma}}{\vdash B \otimes C, \Gamma, \Delta, \Sigma} (cut) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, B, \Gamma} \quad \frac{\tau}{\vdash A, \Sigma}}{\vdash B, \Gamma, \Sigma} (cut) \quad \frac{\phi}{\vdash B \otimes C, \Gamma, \Delta, \Sigma} (\otimes)$
$\otimes - cut - 2$	$\frac{\frac{\frac{\pi}{\vdash B, \Gamma} \quad \frac{\phi}{\vdash A^\perp, C, \Delta}}{\vdash A^\perp, B \otimes C, \Gamma, \Delta} (\otimes) \quad \frac{\tau}{\vdash A, \Sigma}}{\vdash B \otimes C, \Gamma, \Delta, \Sigma} (cut) \longrightarrow \frac{\frac{\pi}{\vdash B, \Gamma} \quad \frac{\frac{\phi}{\vdash A^\perp, C, \Delta} \quad \frac{\tau}{\vdash A, \Sigma}}{\vdash C, \Delta, \Sigma} (cut)}{\vdash B \otimes C, \Gamma, \Delta, \Sigma} (\otimes)$
$\perp - cut$	$\frac{\frac{\frac{\pi}{\vdash A^\perp, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash A^\perp, \perp, \Gamma} (\perp) \quad \frac{\tau}{\vdash A, \Delta}}{\vdash \perp, \Gamma, \Delta} (cut) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash \Gamma, \Delta} (cut) \quad \frac{\tau}{\vdash \perp, \Gamma, \Delta} (\perp)$
$\& - cut$	$\frac{\frac{\frac{\pi}{\vdash A^\perp, B, \Gamma} \quad \frac{\phi}{\vdash A^\perp, C, \Gamma}}{\vdash A^\perp, B \& C, \Gamma} (\&) \quad \frac{\tau}{\vdash A, \Delta}}{\vdash B \& C, \Gamma, \Delta} (cut) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, B, \Gamma} \quad \frac{\tau}{\vdash A, \Delta}}{\vdash B, \Gamma, \Delta} (cut) \quad \frac{\frac{\phi}{\vdash A^\perp, C, \Gamma} \quad \frac{\tau}{\vdash A, \Delta}}{\vdash C, \Gamma, \Delta} (cut) \quad \frac{\tau}{\vdash B \& C, \Gamma, \Delta} (\&)$
$\oplus_1 - cut$	$\frac{\frac{\frac{\pi}{\vdash A^\perp, B, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash A^\perp, B \oplus C, \Gamma} (\oplus_1) \quad \frac{\tau}{\vdash A, \Delta}}{\vdash B \oplus C, \Gamma, \Delta} (cut) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, B, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash B, \Gamma, \Delta} (cut) \quad \frac{\tau}{\vdash B \oplus C, \Gamma, \Delta} (\oplus_1)$
$\oplus_2 - cut$	$\frac{\frac{\frac{\pi}{\vdash A^\perp, C, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash A^\perp, B \oplus C, \Gamma} (\oplus_2) \quad \frac{\tau}{\vdash A, \Delta}}{\vdash B \oplus C, \Gamma, \Delta} (cut) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, C, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash C, \Gamma, \Delta} (cut) \quad \frac{\tau}{\vdash B \oplus C, \Gamma, \Delta} (\oplus_2)$
$\top - cut$	$\frac{\frac{\tau}{\vdash A^\perp, \top, \Gamma} \quad \frac{\pi}{\vdash A, \Delta}}{\vdash \top, \Gamma, \Delta} (\top) \longrightarrow \frac{\tau}{\vdash \top, \Gamma, \Delta} (\top)$
$?d - cut$	$\frac{\frac{\frac{\pi}{\vdash A^\perp, B, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash A^\perp, ?B, \Gamma} (?d) \quad \frac{\tau}{\vdash A, \Delta}}{\vdash ?B, \Gamma, \Delta} (cut) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, B, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash B, \Gamma, \Delta} (cut) \quad \frac{\tau}{\vdash ?B, \Gamma, \Delta} (?d)$
$?c - cut$	$\frac{\frac{\frac{\pi}{\vdash A^\perp, ?B, ?B, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash A^\perp, ?B, \Gamma} (?c) \quad \frac{\tau}{\vdash A, \Delta}}{\vdash ?B, \Gamma, \Delta} (cut) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, ?B, ?B, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash ?B, ?B, \Gamma, \Delta} (cut) \quad \frac{\tau}{\vdash ?B, \Gamma, \Delta} (?c)$
$?w - cut$	$\frac{\frac{\frac{\pi}{\vdash A^\perp, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash A^\perp, ?B, \Gamma} (?w) \quad \frac{\tau}{\vdash A, \Delta}}{\vdash ?B, \Gamma, \Delta} (cut) \longrightarrow \frac{\frac{\pi}{\vdash A^\perp, \Gamma} \quad \frac{\phi}{\vdash A, \Delta}}{\vdash \Gamma, \Delta} (cut) \quad \frac{\tau}{\vdash ?B, \Gamma, \Delta} (?w)$
$! - cut$	$\frac{\frac{\frac{\pi}{\vdash ?A^\perp, B, ?\Gamma} \quad \frac{\phi}{\vdash A, ?\Delta}}{\vdash ?A^\perp, !B, ?\Gamma} (!) \quad \frac{\frac{\phi}{\vdash A, ?\Delta}}{\vdash !A, ?\Delta} (!)}{\vdash !B, ?\Gamma, ?\Delta} (cut) \longrightarrow \frac{\frac{\pi}{\vdash ?A^\perp, B, ?\Gamma} \quad \frac{\frac{\phi}{\vdash A, ?\Delta}}{\vdash !A, ?\Delta} (!)}{\vdash B, ?\Gamma, ?\Delta} (cut) \quad \frac{\phi}{\vdash !B, ?\Gamma, ?\Delta} (!)$

■ **Table 2** Cut-elimination in Linear Logic – Commutative cases

B Cut-Elimination in Affine Logic

Cut-elimination in affine logic is defined as in linear logic, except that the $?w - !$ key case and the $?w - cut$ commutative case are replaced by the two cases on Table 3. That cut-elimination is weakly normalising in affine logic can be proved similarly to the weak normalisation of cut-elimination in linear logic – it is *e.g.* a simple adaptation of the proof in [17].

$$\begin{array}{c}
 \hline
 w \quad \frac{\frac{\frac{\pi}{\vdash \Gamma} \quad (w)}{\vdash A, \Gamma} \quad (w) \quad \vdash A^\perp, \Delta}{\vdash \Gamma, \Delta} \quad (cut) \quad \longrightarrow \quad \frac{\pi}{\vdash \Gamma, \Delta} \quad (w) \\
 \hline
 w - cut \quad \frac{\frac{\frac{\pi}{\vdash A, \Gamma} \quad (w)}{\vdash A, B, \Gamma} \quad (w) \quad \frac{\phi}{\vdash A^\perp, \Delta}}{\vdash \Gamma, B, \Delta} \quad (cut) \quad \longrightarrow \quad \frac{\frac{\pi}{\vdash A, \Gamma} \quad \frac{\phi}{\vdash A^\perp, \Delta}}{\vdash \Gamma, \Delta} \quad (cut) \\
 \hline
 \frac{}{\vdash \Gamma, B, \Delta} \quad (w)
 \end{array}$$

Table 3 Cut-elimination in Affine Logic – Different cases with respect to Linear Logic

C Proofs by Case Study from Section 3

We provide here some cases for the easy but tedious proofs by case study of Lemmas 4 and 9 stated in Section 3.

► Lemma 4. *Let π and ϕ be derivations such that $\pi \longrightarrow \phi$. For each 0-slice ϕ' of ϕ , there exists a 0-slice π' of π such that $\pi' \longrightarrow \phi'$ or $\pi' = \phi'$.*

Proof. We distinguish cases according to the kind of the cut-elimination step. We present here only a few relevant cases, the others being trivial or similar. Precisely, we cover the $\& - \oplus_1$, $?d - !$ and $?c - !$ key cases, and the $\top - cut$ commutative case.

$\& - \oplus_1$ key case. We have

$$\frac{\frac{\frac{\pi}{\vdash A^\perp, \Gamma} \quad \frac{\phi}{\vdash B^\perp, \Gamma}}{\vdash A^\perp \& B^\perp, \Gamma} \quad (\&) \quad \frac{\frac{\tau}{\vdash A, \Delta}}{\vdash A \oplus B, \Delta} \quad (\oplus_1)}{\vdash \Gamma, \Delta} \quad (cut) \quad \longrightarrow \quad \frac{\frac{\pi}{\vdash A^\perp, \Gamma} \quad \frac{\tau}{\vdash A, \Delta}}{\vdash \Gamma, \Delta} \quad (cut)$$

and are given a 0-slice s of the reduced $\frac{\frac{\pi}{\vdash A^\perp, \Gamma} \quad \frac{\tau}{\vdash A, \Delta}}{\vdash \Gamma, \Delta} \quad (cut)$. If the *cut*-rule of this cut-elimination step is not in s , then s is also a 0-slice of the redex. Otherwise, we define the 0-slice r of the redex as making the same choices of sub-trees as in s , except for the $\&$ -rule erased during the cut-elimination step where we keep the left sub-tree $\frac{\pi}{\vdash A^\perp, \Gamma} \quad (\&_1)$ so

that $r \longrightarrow s$.

$$\begin{array}{c}
\pi = \frac{\frac{\overline{\vdash X^+, X^-} \quad (ax) \quad \vdash X^+, \Delta}{\vdash X^+, \Delta} \quad (cut) \quad \vdash X^+, \Delta}{\vdash X^+, \Delta} \quad (cut) \quad \vdash X^+, \Delta = \phi \\
\vdots \\
\vdash \Gamma \\
\vdots \\
\vdash \Gamma
\end{array}
\longrightarrow
\begin{array}{c}
\vdash X^+, \Delta \\
\vdots \\
\vdash \Gamma \\
\vdots \\
\vdash \Gamma
\end{array}
= \phi$$

■ **Figure 3** Illustration of the *ax* key case in the proof of Lemma 9

592 **?d-! key case.** Given a 0-slice s of the reduced, we define a 0-slice r of the redex as making
 593 the same choice of sub-trees as in s , except for $\&$ -rules that are above a $!$ -rule in the redex
 594 but not in the reduced, for which there is no choice to make.

595 **?c-! key case.** This case is trivial solely because a 0-slice leaves the $\&$ -rules above a $!$ -rule
 596 as they are: we make the same choices for the wanted 0-slice of the redex as the choices
 597 made in the given 0-slice of the reduced.

598 **$\top$ -cut commutative case.** We have

$$\frac{\frac{\overline{\vdash A^\perp, \top, \Gamma} \quad (\top) \quad \vdash A, \Delta}{\vdash \top, \Gamma, \Delta} \quad (cut) \quad \vdash \top, \Gamma, \Delta}{\vdash \top, \Gamma, \Delta} \quad (cut) \quad \vdash \top, \Gamma, \Delta = \phi$$


600 and are given a 0-slice s of the reduced $\frac{\overline{\vdash \top, \Gamma, \Delta} \quad (\top)}{\vdash \top, \Gamma, \Delta}$. If the $\top$ -rule of this cut-elimination
 601 step is not in s , then s is also a 0-slice of the redex. Otherwise, we define the 0-slice r of
 602 the redex as making the same choices of sub-trees as in s , except for the $\&$ -rules of π erased
 603 during the cut-elimination step where we keep an arbitrary sub-tree, say always the left one,
 604 so that $r \longrightarrow s$. ◀

605 ► **Lemma 9.** Consider a cut-elimination step $\pi \longrightarrow \phi$ between two derivations, or between
 606 two 0-slices, of conclusion $\vdash \Gamma$. Let u and v be occurrences of atoms from Γ . If there is an
 607 alternating path between u and v in $\mathcal{G}_\phi$, then there is an alternating path between them in $\mathcal{G}_\pi$.

608 **Proof.** We distinguish cases according to the kind of the cut-elimination step. We present
 609 here only the *ax* and *?c-!* key cases, the others being trivial or similar.

610 ***ax* key case.** We have

$$\pi = \frac{\frac{\overline{\vdash X^+, X^-} \quad (ax) \quad \vdash X^+, \Delta}{\vdash X^+, \Delta} \quad (cut) \quad \vdash X^+, \Delta}{\vdash X^+, \Delta} \quad (cut) \quad \vdash X^+, \Delta = \phi$$



■ **Figure 4** Illustration of the $?c - !$ key case in the proof of Lemma 9

with an alternating path p between u and v in the GOI projection graph $\mathcal{G}_\phi$. Let us call x the vertex of $\mathcal{G}_\pi$ corresponding to X^+ in the sequent $\vdash X^+, \Delta$ below the cut -rule, and x' and x'' the vertices corresponding respectively to the cut -formulas X^- and X^+ .

See Figure 3 for an illustration of $\mathcal{G}_\pi$ and $\mathcal{G}_\phi$. We observe that the vertices of $\mathcal{G}_\phi$ are those of $\mathcal{G}_\pi$ except for x' and x'' . Moreover, the edges of $\mathcal{G}_\phi$ are those of $\mathcal{G}_\pi$ except that the ax -edge $x-x'$ and the cut -edge $x'-x''$ are both removed, and that each ax -edge of endpoint x'' (corresponding to an ax -rule in τ) instead has endpoint x .

We obtain a path p' between u and v in $\mathcal{G}_\pi$ by replacing in p any ax -edge $y-x$ that does not belong to $\mathcal{G}_\pi$ by the edges $y-x''-x'-x$. Observe that p' is an alternating path since p is.

?c - ! key case. We have

$$\pi = \frac{\frac{\vdash ?A^\perp, ?A^\perp, \Delta}{\vdash ?A^\perp, \Delta} (?c) \quad \frac{\frac{\mu}{\vdash A, ?\Sigma} (!) \quad \frac{\mu}{\vdash !A, ?\Sigma} (!)}{\vdash \Delta, ?\Sigma} (cut)}{\vdash \Delta, ?\Sigma} \longrightarrow \frac{\frac{\frac{\tau}{\vdash ?A^\perp, ?A^\perp, \Delta} (?c) \quad \frac{\frac{\mu}{\vdash A, ?\Sigma} (!) \quad \frac{\mu}{\vdash !A, ?\Sigma} (!)}{\vdash \Delta, ?\Sigma, ?\Sigma} (cut)}{\vdash \Delta, ?\Sigma, ?\Sigma} (?c) \quad \frac{\mu}{\vdash A, ?\Sigma} (!) \quad \frac{\mu}{\vdash !A, ?\Sigma} (!)}{\vdash \Delta, ?\Sigma} (cut) = \phi$$

with an alternating path p between u and v in the GOI projection graph $\mathcal{G}_\phi$. Let us call S the set of vertices in $\mathcal{G}_\pi$ corresponding to the atoms occurring in the cut -formula $!A$ and $S^\perp$ the set of vertices in $\mathcal{G}_\pi$ corresponding to the atoms occurring in the cut -formula $?A^\perp$, such that there is a cut -edge between $x \in S$ and $x^\perp \in S^\perp$.

See Figure 4 for an illustration of $\mathcal{G}_\pi$ and $\mathcal{G}_\phi$. We observe that the vertices of $\mathcal{G}_\phi$ are those of $\mathcal{G}_\pi$ where S is replaced by two of its copies, that we call $S_1 = \{x_1 \mid x \in S\}$ and $S_2 = \{x_2 \mid x \in S\}$, and similarly for $S^\perp$ with $S_1^\perp = \{x_1^\perp \mid x^\perp \in S^\perp\}$ and $S_2^\perp = \{x_2^\perp \mid x^\perp \in S^\perp\}$. Furthermore, the edges of $\mathcal{G}_\phi$ are those of $\mathcal{G}_\pi$ except that:

- each cut -edge $x^\perp-x$ with endpoints $x \in S$ and $x^\perp \in S^\perp$ is replaced by two cut -edges $x_1^\perp-x_1$ and $x_2^\perp-x_2$;
- each ax -edge $y-x$ coming from the sub-derivation μ with an endpoint $x \in S$ is replaced by two ax -edges $y-x_1$ and $y-x_2$ (in case $y \in S$ we obtain the ax -edges y_1-x_1 and y_2-x_2);
- each ax -edge $y-x^\perp$ coming from the sub-derivation τ with an endpoint $x^\perp \in S^\perp$ is replaced by an ax -edge $y-x_1^\perp$ or by an ax -edge $y-x_2^\perp$, or possibly by both (in case $y \in S^\perp$ we obtain some, and at least one, edges in $\{y_1-x_1^\perp; y_1-x_2^\perp; y_2-x_1^\perp; y_2-x_2^\perp\}$).

We obtain a path p' between u and v in $\mathcal{G}_\pi$ by replacing in p each occurrence of a $x_1 \in S_1$ and of a $x_2 \in S_2$ by $x \in S$, and similarly of a $x_1^\perp \in S_1^\perp$ and of a $x_2^\perp \in S_2^\perp$ by $x^\perp \in S^\perp$. That

640 p' is an alternating path follows from the fact p is such a path. $\blacktriangleleft$

641 **D** Proof of Proposition 23 with Intermediate Results

642 We present here a proof of Proposition 23 from Section 6. It uses two intermediate results, the
643 second one being the main novelty with respect to the proof of Proposition 14 in Section 4.

644 **► Lemma 28.** *Let F and C be formulas, and X and Y be atoms such that $X \notin \mathcal{V}(F) \cup \mathcal{V}(C)$.
645 With $A(F)$ defined as in Equation (2), if $\vdash A(F)^\perp, C$ is provable, then there exists a cut-free
646 derivation of $\vdash A(F)^\perp, C$ in which no $!$ -rule has an atom X in its conclusion sequent.*

647 **Proof.** Proceed exactly as in the proof of Lemma 12. $\blacktriangleleft$

648 **► Lemma 29.** *Let F and C be formulas, and X and Y be atoms such that $X, Y \notin \mathcal{V}(F)$ and
649 $X \notin \mathcal{V}(C)$. Assume that F is not provable, and consider $A(F)$ and $(B_n(F))_{n \in \mathbb{N}}$ defined as
650 in Equations (2) and (4). Let π be a cut-free derivation of $\vdash A(F)^\perp, B_n(F)$ for some $n \in \mathbb{N}$.
651 Then in π there is no ax -rule on $\vdash Y^+, Y^-$ where both of these occurrences belong to $B_n(F)$.*

652 **Proof.** We proceed by contradiction. Such an ax -rule can only be between the occurrences
653 Y^+ and Y^- of a sub-formula $Y^+ \multimap (Y^+ \otimes !F) = Y^- \wp (Y^+ \otimes !F)$, since other occurrences
654 of Y in $B_n(F)$ are separated by a connective $\otimes$. By the sub-formula property and simple
655 (non-)provability results, this implies the following is a sub-tree of π :

$$\begin{array}{c}
 \overline{\vdash Y^-, Y^+} \quad (ax) \\
 \hline
 \overline{\vdash Y^-, Y^+, ?\Delta} \quad (?w) \quad \phi \\
 \vdash Y^-, Y^+, ?\Delta \quad \vdash !F, \Gamma \\
 \hline
 \vdash Y^-, Y^+ \otimes !F, \Gamma, ?\Delta \quad (\otimes)
 \end{array}$$

656

657 where the doubled inference rule means we apply the rule a certain number of times, and
658 where the formulas of Γ are included in
659 $\{X^-, ?(X^+ \otimes (X^- \wp (Y^+ \otimes (Y^- \wp ?F^\perp))), X^+ \otimes (X^- \wp (Y^+ \otimes (Y^- \wp ?F^\perp))), Y^+ \otimes (Y^- \wp ?F)\}$
660 with possibly some of these formulas linked by $\wp$ and formulas appearing possibly several
661 times. We substitute Y^+ by 0 in ϕ , obtaining a derivation τ of a sequent (isomorphic to)
662 $\vdash !F, \Gamma'$ where the formulas of Γ' are included in $\{X^-, ?(X^+ \otimes (X^- \wp 0)), X^+ \otimes (X^- \wp 0), 0\}$
663 with possibly some of these formulas linked by $\wp$ and formulas appearing possibly several
664 times.

665 Such a derivation τ must contain a $!$ -rule corresponding to the root connective of $!F$. By
666 the sub-formula property, this means $\vdash !F, \Gamma''$ is provable with Γ'' composed of the formula
667 $?(X^+ \otimes (X^- \wp 0))$ a certain number of times. Now, substituting X^+ by 0, we obtain a
668 derivation of conclusion $\vdash !F, \Gamma'''$ with Γ''' composed of a formula isomorphic to $?0$ a certain
669 number of times. From this last derivation, we deduce that $\vdash F$ is provable, a contradiction.
670 Indeed, if the formula $?0$ appears at least once we have the derivation

$$\begin{array}{c}
 \vdash !F, ?0, \dots, ?0 \quad \overline{\vdash \top} \quad (\top) \\
 \hline
 \vdash !F, ?0 \quad \vdash \top \quad (!) \\
 \hline
 \vdash !F \quad (cut)
 \end{array}$$

671

672 and if it does not appear we directly have a derivation of $\vdash !F$. But the last rule of a cut-free
673 derivation of $\vdash !F$ can only be $\frac{\vdash F}{\vdash !F} (!)$, so $\vdash F$ is provable and we are done. $\blacktriangleleft$

674 ► **Proposition 23.** *Let F and C be formulas, and X and Y be atoms such that $X, Y \notin \mathcal{V}(F)$
 675 and $X \notin \mathcal{V}(C)$. Assume that $\vdash F$ is not provable. With $A(F)$ and $(B_n(F))_{n \in \mathbb{N}}$ defined as in
 676 Equations (2) and (4), if $\vdash A(F)^\perp, C$ and $\vdash C^\perp, B_n(F)$ are both provable for some number
 677 $n \in \mathbb{N}$, then C contains at least $n + 2$ occurrences of Y^+ .*

678 **Proof.** We proceed as in the proof of Proposition 14. The differences with that proof are **in**
 679 **bold and colored in green.**

680 Let us call π and ϕ derivations of respectively $\vdash A(F)^\perp, C$ and $\vdash C^\perp, B_n(F)$. We assume
 681 them *cut*-free without any loss of generality. We will use the GOI projection graph of (some
 682 0-slice of) their composition by *cut* to identify Y^- 's in $C^\perp$ linked to the Y^+ 's of $B_n(F)$.
 683 Thanks to alternating paths, we will prove that these occurrences of Y^- are pairwise distinct.
 684 This yields one Y^- in $C^\perp$ for each Y^+ in $B_n(F)$, concluding the proof.

685 **Finding an alternating path in some 0-slice.** Notice that $\frac{\frac{\pi}{\vdash A(F)^\perp, C} \quad \frac{\phi}{\vdash C^\perp, B_n(F)}}{\vdash A(F)^\perp, B_n(F)} \text{ (cut)}$

686 reduces to a *cut*-free derivation of $\vdash A(F)^\perp, B_n(F)$, that we call τ . **Consider any 0-slice**
 687 **τ' of τ .** By repeated applications of Lemma 4, there is a 0-slice μ of

$$688 \quad \frac{\frac{\pi}{\vdash A(F)^\perp, C} \quad \frac{\phi}{\vdash C^\perp, B_n(F)}}{\vdash A(F)^\perp, B_n(F)} \text{ (cut)} \text{ such that } \mu \longrightarrow^* \tau'.$$

689 **By Lemma 29, there is no ax -rule on $\vdash Y^+, Y^-$ where both of these occurrences**
 690 **belong to $B_n(F)$.** Consider some occurrence Y_α^+ in $B_n(F)$, with the index α used to
 691 distinguish it from other occurrences. Since τ' is a *cut*-free 0-slice of $\vdash A(F)^\perp, B_n(F)$, there
 692 must be an ax -rule between this occurrence Y_α^+ and one of the two occurrences of Y^- in
 693 $A(F)^\perp$: call it Y_1^- . Hence, there is an alternating path (made of one ax -edge) between these
 694 two occurrences in the GOI projection graph $\mathcal{G}_\tau$ of τ . By repeated applications of Lemma 9,
 695 there also is an alternating path p between Y_1^- and Y_α^+ in $\mathcal{G}_\mu$. Thus, there is an occurrence
 696 Y_β^- and an ax -rule $\frac{}{\vdash Y_\beta^-, Y_\alpha^+} \text{ (ax)}$ in μ and $Y_\beta^- - Y_\alpha^+$ is the last edge of p . This occurrence
 697 Y_β^- can only be in $C^\perp$.

698 We use the alternating path p to show there is no sub-formula $?D$ of $C^\perp$ containing
 699 Y_β^- – say differently, there is no $?$ between the root of the formula $C^\perp$ and Y_β^- . Once
 700 this claim is proved, we are done: each occurrence of Y^+ in $B_n(F)$ is linked by an ax -edge
 701 to an occurrence of Y^- in $C^\perp$. All these occurrences of Y^- are pairwise distinct (recall
 702 Remark 8): there is no sub-formula $?D$ of $C^\perp$ containing such an occurrence, so they cannot
 703 be contracted; and there is no $\&$ -rule that can share the identified ax -rules since we are
 704 in a 0-slice, and these ax -rules cannot be above a $!$ -rule for there is no $?$ -connective nor
 705 $!$ -connective below the Y^+ of $B_n(F)$. Therefore, we have found $n + 2$ occurrences of Y^- in
 706 $C^\perp$, as wished.

707 **Exploiting the alternating path.** We prove our claim by showing a stronger result:

- 708 ■ for all occurrences of Y^- in C (respectively in $C^\perp$) belonging to p , there is no $?$ between
 709 the root of C (respectively of $C^\perp$) and these occurrences;
- 710 ■ for all occurrences of Y^+ in C (respectively in $C^\perp$) belonging to p , there is no $!$ between
 711 the root of C (respectively of $C^\perp$) and these occurrences.

712 We proceed by following the edges of p starting from its endpoint Y_1^- in $A(F)^\perp$.

Exploiting the alternating path: base case. Consider the first edge $Y_1^- - Y_2^+$ of p , with Y_2^+ necessarily in C . Using **Lemma 28**, without any loss of generality, there is no $!$ -connective between the root of C and Y_2^+ . Otherwise, since there is an ax -rule on $\vdash Y_1^-, Y_2^+$, there would be a $!$ -rule (associated to this $!$ -connective) with in its conclusion sequent a $?$ -sub-formula of $A(F)^\perp$ containing Y_1^- , meaning with $?(X^+ \otimes (X^- \wp Y^-))$: contradiction.

Exploiting the alternating path: inductive case. See Figure 2 for an illustration of our present reasoning. Assume the results holds for a prefix of p , and let us consider its last edge.

1. Suppose this prefix ends with an ax -edge $Y_k^- - Y_{k+1}^+$ with Y_{k+1}^+ in C . We know there is no $!$ between the root of C and Y_{k+1}^+ . As p is alternating, the subsequent edge is a cut -edge $Y_{k+1}^+ - Y_{k+2}^-$ with Y_{k+2}^- in $C^\perp$. Since Y_{k+1}^+ and Y_{k+2}^- are dual occurrences by definition of a cut -edge, there is no $?$ between the root of $C^\perp$ and Y_{k+2}^- .
 2. Suppose this prefix ends with a cut -edge $Y_k^+ - Y_{k+1}^-$ with Y_{k+1}^- in $C^\perp$. We know there is no $?$ between the root of $C^\perp$ and Y_{k+1}^- . As p is alternating, the subsequent edge is an ax -edge $Y_{k+1}^- - Y_{k+2}^+$ with Y_{k+2}^+ either in $B_n(F)$ or in $C^\perp$. In the first case, we are done: we have $Y_{k+2}^+ = Y_\alpha^+$ since there is no cut -edge with endpoint Y_{k+2}^+ to continue our path, and p is supposed to end in Y_α^+ . In the second case, since there is an ax -rule $\vdash Y_{k+1}^-, Y_{k+2}^+$ (ax) by definition of an ax -edge, it follows there is no $!$ between the root of $C^\perp$ and Y_{k+2}^+ – because the context condition of a corresponding $!$ -rule would prevent the presence of Y_{k+1}^- in all sequents above it.
 3. Suppose this prefix ends with an ax -edge $Y_k^- - Y_{k+1}^+$ with Y_{k+1}^+ in $C^\perp$. This case is identical to case 1 up to switching C and $C^\perp$.
 4. Suppose this prefix ends with a cut -edge $Y_k^+ - Y_{k+1}^-$ with Y_{k+1}^- in C . **This case is very similar to case 2, with Y_{k+2}^+ not in $A(F)^\perp$ since no atom of $A(F)^\perp$ is the endpoint of a cut -edge and the path cannot end in $A(F)^\perp$.**
- This proves our claim, and concludes the proof. $\blacktriangleleft$