

MOTS-c for Metabolic Health & Longevity

Part II · Weight Loss & Metabolic Peptides — Mitochondrial Peptides

CHAPTER 10: MOTS-C FOR METABOLIC HEALTH AND LONGEVITY

COMPREHENSIVE PEPTIDE THERAPY COMPENDIUM

Part II: Weight Loss & Metabolic Peptides - Mitochondrial Peptides

Chapter 10 of 43

REGULATORY STATUS

FDA STATUS

NOT APPROVED (RESEARCH COMPOUND)

- No FDA approval for any indication
- Available through compounding pharmacies only
- Research-grade peptide status

EVIDENCE GRADE

C (EMERGING PRECLINICAL EVIDENCE)

- Primarily animal studies (mice, primates)
- Limited human clinical trial data
- Promising mechanistic research
- No published Phase III trials

COMPOUNDING STATUS

WIDELY COMPOUNDED

- Available from 503A compounding pharmacies
- No FDA oversight or quality standards
- Variable purity and potency across sources

2026 Regulatory Update (verified — FDA advisory-committee calendar). On **April 15, 2026** the FDA removed MOTS-c from the 503A Category 2 list. On **July 23–24, 2026** the FDA Pharmacy Compounding Advisory Committee (PCAC) reviews MOTS-c (free base and acetate) for the Section 503A bulk drug substances list — the decision will determine whether U.S. compounding pharmacies may legally dispense it. Recommendation is non-binding and is not FDA drug approval.

CRITICAL LEGAL DISCLAIMER

MOTS-c is NOT FDA-approved for any medical condition. This chapter provides educational information about reported uses in research settings and compounding pharmacy availability. Prescribers assume full legal and professional liability for off-label use of investigational compounds. Informed consent documenting experimental status and limited human safety data is mandatory.

All clinical applications described represent reported protocols from research literature and clinical practice, not FDA-approved indications. Providers must independently verify current regulatory status and comply with state medical

board requirements before prescribing.

Evidence Summary

OVERVIEW OF CLINICAL EVIDENCE

MOTS-c (Mitochondrial Open Reading Frame of the 12S rRNA-c) is a 16-amino acid peptide encoded within the mitochondrial genome that functions as a mitochondrial-derived peptide (MDP) regulating cellular metabolism, insulin sensitivity, and metabolic homeostasis. Discovered in 2015 by researchers at the University of Southern California, MOTS-c represents a novel class of mitochondrial signaling molecules communicating metabolic status between organelles and the nucleus.^{1,2}

BIOCHEMICAL STRUCTURE

MOTS-c Sequence: MRWQEMGYIFYPRKLR

- Molecular weight: 2,174 Da
- 16 amino acids
- Encoded by mitochondrial 12S rRNA gene
- Evolutionary conserved across mammalian species
- Endogenously produced in human cells
- Circulates in bloodstream as signaling molecule

Unlike traditional pharmaceuticals developed through systematic drug discovery, MOTS-c is an endogenous human peptide whose levels decline with aging, obesity, and metabolic disease. Therapeutic administration aims to restore youthful metabolic function by supplementing declining endogenous production.^{1,2,3}

CRITICAL MECHANISTIC DISTINCTION

MOTS-c is NOT a GLP-1 agonist or incretin-based therapy. It functions through entirely different metabolic pathways:

- ✗ Does NOT activate GLP-1 receptors
- ✗ Does NOT increase insulin secretion directly
- ✗ Does NOT delay gastric emptying
- ✗ Does NOT suppress appetite through hypothalamic mechanisms
- ✓ DOES improve insulin sensitivity through AMPK activation
- ✓ DOES enhance mitochondrial function and biogenesis
- ✓ DOES regulate glucose and lipid metabolism
- ✓ DOES activate cellular stress response pathways (hormesis)
- ✓ DOES promote skeletal muscle glucose uptake

MOTS-c is grouped with metabolic peptides due to insulin-sensitizing effects and weight management applications, though mechanism diverges fundamentally from incretin-based therapies (semaglutide, tirzepatide, liraglutide).^{1,2}

CURRENT EVIDENCE BASE

PRECLINICAL ANIMAL STUDIES

Mouse Studies - Metabolic Effects:^{1,2}

- MOTS-c administration prevented diet-induced obesity in high-fat diet mice
- Improved glucose tolerance and insulin sensitivity in obese mice
- Increased skeletal muscle glucose uptake independent of insulin signaling
- Protected against age-related insulin resistance
- Reversed metabolic dysfunction in established obesity
- Enhanced exercise capacity and endurance performance
- Reduced hepatic steatosis (fatty liver disease)
- Improved lipid metabolism and reduced circulating triglycerides

Mouse Studies - Aging and Longevity:³

- MOTS-c levels decline with aging in multiple tissues
- Exogenous MOTS-c administration improved healthspan in aged mice
- Protected against age-related metabolic decline
- Maintained muscle function and physical performance with aging
- Reduced inflammatory markers in aged animals

Non-Human Primate Studies:⁴

- Middle-aged rhesus monkeys treated with MOTS-c showed:
- Improved insulin sensitivity (HOMA-IR reduction)
- Reduced fasting glucose and insulin levels
- Enhanced muscle glucose uptake measured by PET imaging
- Well-tolerated with no adverse effects during 6-month study
- Pharmacokinetic profile supports once-daily subcutaneous dosing

HUMAN OBSERVATIONAL STUDIES

Japanese Centenarian Study:⁵

- Genetic polymorphisms in mitochondrial 12S rRNA (MOTS-c encoding region)

associated with exceptional longevity

- Centenarians more likely to carry specific MOTS-c variants
- Certain polymorphisms correlated with reduced metabolic disease risk
- Suggests evolutionary selection for favorable MOTS-c variants

Metabolic Disease Association Studies:⁶

- Lower circulating MOTS-c levels observed in:
- Obesity (BMI >30)
- Type 2 diabetes
- Insulin resistance
- Metabolic syndrome
- MOTS-c levels inversely correlated with:
- Fasting insulin
- HOMA-IR (insulin resistance index)
- Hemoglobin A1c
- Body mass index

Exercise Response Studies:⁷

- Acute exercise increases circulating MOTS-c in healthy humans
- MOTS-c levels correlate with exercise capacity and fitness
- Athletes demonstrate higher baseline MOTS-c than sedentary individuals
- MOTS-c appears to mediate some exercise-induced metabolic benefits

HUMAN INTERVENTIONAL STUDIES (LIMITED)

Phase I Safety Studies (UNPUBLISHED):

- Small safety trials in healthy volunteers (N<50 total across studies)
- Subcutaneous administration at doses 5-15mg daily
- Favorable safety profile with minimal adverse effects
- No serious adverse events reported
- Preliminary pharmacokinetic data supports once-daily dosing
- Brief half-life (~30 minutes) but prolonged metabolic effects

Pilot Efficacy Studies (CONFERENCE ABSTRACTS ONLY):

- Presented at endocrinology/metabolism conferences 2020-2024
- Small studies (N=10-30 per study) in prediabetes and metabolic syndrome
- Reported improvements in:
- Fasting glucose (10-15% reduction)
- Insulin sensitivity (HOMA-IR improvement)
- Body composition (modest fat mass reduction)
- Exercise capacity (6-minute walk test improvement)
- NO peer-reviewed publications validating these findings
- Significant publication bias likely (negative studies unpublished)

CRITICAL EVIDENCE LIMITATIONS

- ! NO FDA-approved indications for any medical condition

- ! NO published Phase II or Phase III human clinical trials
- ! NO head-to-head comparisons with established diabetes/obesity medications
- ! NO long-term human safety data (>6 months)
- ! NO validated dosing protocols for clinical use
- ! NO standardized manufacturing specifications or purity standards
- ! Significant knowledge gaps regarding optimal dose, duration, and safety

CURRENT CLINICAL USE PATTERNS

Despite limited human evidence, MOTS-c has gained clinical adoption in:

- Anti-aging and longevity medicine practices
- Sports performance optimization clinics
- Integrative and functional medicine centers
- Compounding pharmacy direct-to-consumer marketing
- Peptide therapy wellness programs

This widespread use occurs in advance of rigorous clinical validation, creating tension between compelling preclinical data and absent human efficacy trials.

Providers prescribing MOTS-c operate in regulatory gray zone with elevated professional liability risk.^{1,2,3,4,5,6,7}

QUALITY OF EVIDENCE ASSESSMENT

EVIDENCE GRADE: C (EMERGING PRECLINICAL EVIDENCE)

MOTS-c evidence quality substantially lags FDA-approved metabolic medications.

While preclinical research demonstrates compelling mechanistic rationale and consistent effects across animal models, human clinical data remains insufficient to establish safety, efficacy, or appropriate therapeutic use.

GRADE C EVIDENCE JUSTIFICATION

- ✓ STRENGTH #1: Robust Preclinical Mechanistic Research
 - Published in high-impact peer-reviewed journals:
 - Cell Metabolism (2015, 2017)^{1,2}
 - Nature Communications (2016)³
 - Diabetes (2018)⁶
 - Consistent findings across multiple independent research groups
 - Well-characterized mechanism of action (AMPK/FOXO pathway activation)
 - Reproducible metabolic benefits in diverse animal models
 - Strong theoretical biological rationale
- ✓ STRENGTH #2: Endogenous Human Peptide

- Naturally occurring in all human cells (not synthetic foreign compound)
- Physiological role well-established through observational studies
- Lower theoretical toxicity risk vs synthetic pharmaceuticals
- Evolutionary conservation suggests important biological function
- Therapeutic approach: Supplementation of declining endogenous levels

✓ **STRENGTH #3: Multi-Species Validation**

- Consistent effects observed in:
 - Mice (multiple strains)
 - Rats
 - Non-human primates (rhesus monkeys)⁴
- Translation from rodents to primates increases confidence
- Primate data bridges gap between mouse models and human application

✗ **LIMITATION #1: Absence of Published Human Efficacy Trials**

- NO peer-reviewed Phase II or Phase III trials published
- Conference abstracts only (not subjected to rigorous peer review)
- Small pilot studies (N<50 combined) with unpublished data
- Unknown efficacy in clinically relevant human populations
- Significant publication bias (negative results likely unreported)

✗ **LIMITATION #2: No FDA Approval or Regulatory Oversight**

- Research compound without approved therapeutic indications
- No standardized manufacturing specifications
- Variable quality, purity, potency across compounding sources
- No FDA-mandated post-marketing surveillance
- Regulatory pathway uncertain (peptide vs biologics classification)

✗ **LIMITATION #3: Insufficient Human Safety Database**

- Human exposure limited to small short-term pilot studies (<6 months)
- NO long-term safety data (1+ years)
- Unknown drug-drug interactions
- Unknown effects in special populations:
 - Pregnancy and lactation
 - Renal insufficiency/failure
 - Hepatic impairment
 - Cardiovascular disease
 - Cancer patients (mitochondrial effects on tumor metabolism unclear)
- Adverse event reporting inconsistent and incomplete

✗ LIMITATION #4: Translation Failures Common

- Historical context: >95% of drugs effective in animals fail in humans
- Metabolic disease research particularly prone to translation failure
- Mouse metabolism differs substantially from human:
 - Higher metabolic rate (7x per kg body weight)
 - Different mitochondrial density
 - Distinct fat distribution and adipose biology
- Shorter lifespan (translation of "longevity" benefits questionable)
- Primate studies more predictive but still imperfect

COMPARATIVE EVIDENCE QUALITY

2026 Regulatory Update (verified — FDA advisory-committee calendar). On **April 15, 2026** the FDA removed MOTS-c from the 503A Category 2 list. On **July 23–24, 2026** the FDA Pharmacy Compounding Advisory Committee (PCAC) reviews MOTS-c (free base and acetate) for the Section 503A bulk drug substances list — the decision will determine whether U.S. compounding pharmacies may legally dispense it. Recommendation is non-binding and is not FDA drug approval.

EVIDENCE STRENGTHS

- ✓ Strong mechanistic understanding from cellular to whole-organism level
- ✓ Reproducible effects across multiple animal species and models
- ✓ Published in high-impact peer-reviewed scientific journals
- ✓ Endogenous peptide with established physiological role
- ✓ Compelling theoretical rationale for metabolic disease applications
- ✓ No major safety signals in limited human exposure
- ✓ Primate data supports feasibility of human translation
- ✓ Observational human data supports relevance to human metabolism

EVIDENCE WEAKNESSES

- ✗ Critical translation gap from animal models to controlled human trials
- ✗ Optimal human dosing unknown (extrapolated from mg/kg animal dosing)
- ✗ Long-term efficacy unknown (mouse studies ≠ human chronic disease)
- ✗ Long-term safety unknown (brief pilot studies insufficient)
- ✗ Drug interactions completely unstudied
- ✗ Special population safety unknown (renal, hepatic, cardiac, pregnancy)
- ✗ Manufacturing quality highly variable across compounders
- ✗ No standardized clinical protocols or treatment guidelines
- ✗ Publication bias (conference abstracts vs peer-reviewed publications)

✗ Commercial conflicts of interest in emerging research field

EVIDENCE LIMITATIONS

CRITICAL LIMITATIONS FOR CLINICAL PRACTICE

LIMITATION #1: Absence of Peer-Reviewed Human Efficacy Trials

The most significant evidence gap is lack of published, peer-reviewed human clinical trials demonstrating efficacy for any condition. While preclinical studies are compelling, animal models frequently fail to predict human outcomes:

Translation Failure Statistics:

- 95% of drugs effective in animals fail in Phase I-III human trials⁸
- Metabolic disease therapeutics particularly prone to translation failure
- Mouse metabolism differs fundamentally from humans:
 - 7x higher metabolic rate per kg body weight
 - Different fat distribution (subcutaneous vs visceral)
 - Distinct adipose tissue biology and endocrinology
 - Higher mitochondrial density in muscle tissue
 - Different dietary patterns and feeding behavior
 - Shorter lifespan (2-3 years vs 75+ years humans)

Historical Translation Failures - Metabolic Medications:

- PPAR agonists: Excellent mouse data, human liver toxicity
- mTOR inhibitors: Anti-aging in mice, human metabolic side effects severe
- Resveratrol: Longevity in worms/flies, no human benefit demonstrated
- Numerous mitochondrial-targeting compounds: Preclinical promise, clinical failure

IMPLICATION FOR MOTS-c:

Preclinical data, while encouraging, provides insufficient basis for confident prediction of human efficacy. Only properly-designed randomized controlled trials can establish whether MOTS-c produces clinically meaningful metabolic benefits in humans.

LIMITATION #2: Unknown Optimal Dosing in Humans

Current human dosing protocols derive from animal studies using allometric scaling, which often produces inaccurate human dose predictions:

Animal to Human Dose Extrapolation Problems:

- Mouse effective dose: 5-15 mg/kg subcutaneously⁹
- Allometric scaling to 70kg human: ~50-150mg per dose
- Body surface area adjustment: ~5-15mg per dose

- Currently reported human dosing: 5-15mg subcutaneously

Issues with Extrapolation:

- ✗ Allometric scaling assumes similar pharmacokinetics (often false)
- ✗ Mitochondrial density varies across species (affects tissue distribution)
- ✗ AMPK signaling sensitivity may differ human vs mouse
- ✗ Optimal dose for insulin sensitivity may differ from weight loss dose
- ✗ Dose-response curve unknown (more may not equal better)
- ✗ Individual variation in response likely (pharmacogenomics unstudied)

Consequence:

Providers prescribing MOTS-c use empiric dosing based on educated guesses rather than dose-finding studies. Patients may be under-dosed (ineffective) or over-dosed (unnecessary expense, potential unknown risks).

LIMITATION #3: Lack of Long-Term Safety Data

Brief pilot studies (<6 months) cannot detect:

DELAYED TOXICITIES

- Cumulative organ damage (liver, kidney)
- Cancer risk (mitochondrial manipulation effects on tumorigenesis unknown)
- Cardiovascular events (arrhythmias, vascular effects)
- Neurologic complications
- Endocrine disruption (thyroid, adrenal, reproductive axes)

CHRONIC EXPOSURE EFFECTS

- Antibody formation (immunogenicity against exogenous peptide)
- Receptor desensitization or downregulation
- Compensatory metabolic adaptations (loss of efficacy over time)
- Nutrient deficiencies from altered metabolism
- Mitochondrial dysfunction paradoxically (from chronic stimulation)

DRUG INTERACTIONS

- MOTS-c activates AMPK (same target as metformin) → unknown interaction
- Effects on cytochrome P450 enzymes unknown
- Interaction with insulin, GLP-1 agonists, SGLT2 inhibitors unknown
- Thyroid hormone interaction possible (mitochondrial effects)
- Statins and mitochondrial peptides: interaction potential

SPECIAL POPULATIONS

- Pregnancy: Fetal mitochondrial development effects completely unknown

- Lactation: Excretion in breast milk unknown
- Pediatrics: Growth and development effects unstudied
- Elderly: Age-related organ dysfunction complicating safety profile
- Renal failure: Peptide accumulation and dose adjustment needs unknown
- Hepatic failure: Mitochondrial-rich tissue, hepatotoxicity risk unclear
- Cancer patients: Mitochondrial effects on tumor metabolism concerning

Historical Lesson:

Fen-phen (fenfluramine-phentermine) widely prescribed for obesity based on short-term studies. Cardiac valvulopathy emerged only after years of widespread use, resulting in FDA withdrawal, lawsuits, and physician liability. Short-term safety does NOT guarantee long-term safety.

LIMITATION #4: Manufacturing Quality and Purity Concerns

MOTS-c available through compounding pharmacies lacks FDA oversight:

Quality Control Issues:

- No standardized manufacturing process
- Variable synthesis methods across compounders:
- Solid-phase peptide synthesis (SPPS) quality varies
- Purification methods inconsistent (HPLC vs crude)
- Lyophilization protocols differ
- Peptide purity ranges 80-99% (impurities may include):
- Deletion sequences (missing amino acids)
- Truncated peptides
- Oxidized or modified amino acids
- Bacterial endotoxins
- Heavy metal contaminants
- Potency verification absent (stated mg may not reflect active peptide)
- Sterility testing inconsistent
- Stability data lacking (degradation during storage unknown)

Consequence:

Patients may receive:

- ✓ High-quality pharmaceutical-grade MOTS-c (best case)
- ! Moderately pure MOTS-c with reduced potency (common)
- ✗ Low-quality product with significant impurities (not uncommon)
- ✗ Contaminated product with bacterial/endotoxin contamination (rare but serious)

Provider Responsibility:

Without FDA oversight, providers must independently vet compounding pharmacy quality. This requires evaluating certificates of analysis, understanding peptide chemistry, and accepting residual uncertainty about product quality.

LIMITATION #5: No Established Treatment Guidelines

Professional medical societies have not issued guidelines for MOTS-c use because:

Guideline Development Requires:

- ✗ Published systematic reviews and meta-analyses (none exist)
- ✗ Expert consensus based on clinical experience (insufficient experience)
- ✗ Risk-benefit analysis from controlled trials (trials incomplete)
- ✗ Comparison to standard-of-care therapies (comparisons lacking)
- ✗ Cost-effectiveness data (no economic analyses published)

Consequence:

Providers prescribing MOTS-c must independently determine:

- Which patients are appropriate candidates
- What constitutes adequate informed consent
- Appropriate baseline and monitoring laboratory tests
- When to discontinue therapy
- How to manage potential adverse effects
- When to use MOTS-c vs FDA-approved alternatives

This absence of guidance increases medicolegal liability and practice variation.

LIMITATION #6: Publication Bias and Commercial Conflicts

Emerging therapeutic areas vulnerable to publication bias:

Concerns:

- ! Positive results published; negative results filed away
- ! Conference abstracts (not peer-reviewed) suggest benefits
- ! Full manuscripts never published (suggests flawed studies or negative findings)
- ! Small sample sizes inflate effect size estimates (small study effect)
- ! Compounding pharmacies fund research (commercial interest in positive results)
- ! Peptide therapy clinics promote MOTS-c (direct financial incentive)

Historical Parallel:

Hormone replacement therapy (HRT) initially promoted based on observational studies showing cardiovascular benefits. Rigorous RCTs (WHI study) subsequently demonstrated increased cardiovascular risk. Initial enthusiasm was tempered by rigorous evidence demonstrating harm rather than benefit.

CRITICAL IMPLICATION

Current MOTS-c evidence, while scientifically interesting and mechanistically sound, does NOT constitute adequate basis for confident clinical use in treating metabolic disease. Providers must clearly communicate to patients:

1. MOTS-c is experimental with unproven efficacy in humans
2. Safety beyond 6 months completely unknown
3. FDA-approved alternatives (metformin, GLP-1 agonists) have vastly superior evidence and should be prioritized
4. Patients assuming unknown risks by using investigational therapy
5. Insurance will not cover experimental treatment
6. Long-term consequences unknown and potentially serious

Prescribing MOTS-c represents practice at the frontier of medicine - potentially beneficial but fraught with uncertainty. Informed consent must reflect this reality honestly.

Mechanism Of Action

MITOCHONDRIAL-DERIVED PEPTIDE BIOLOGY

MOTS-c represents a novel class of signaling molecules called mitochondrial-derived peptides (MDPs) that enable communication between mitochondria and the cellular nucleus. This discovery fundamentally altered understanding of mitochondrial biology, demonstrating that mitochondria are not merely "cellular powerhouses" but active endocrine organs producing hormones regulating whole-body metabolism.^{1,2}

MITOCHONDRIAL GENOME ENCODING

MOTS-c is encoded within the mitochondrial 12S ribosomal RNA gene:

- Mitochondrial DNA contains 37 genes (traditionally recognized)
- Small open reading frames (sORFs) within these genes encode peptides
- MOTS-c: 16 amino acids encoded by mitochondrial 12S rRNA
- Other MDPs: Humanin, SHLP1-6 (Small Humanin-Like Peptides)
- Evolutionary conservation suggests critical physiological functions

MITOCHONDRIAL-NUCLEAR COMMUNICATION

Traditional View (Pre-2015):

Nucleus → Mitochondria (one-way communication)

- Nuclear genes encode >1,500 mitochondrial proteins

- Proteins imported into mitochondria for function
- Mitochondria passive recipients of nuclear instructions

Revised View (Post-MOTS-c Discovery):

Nucleus ⇌ Mitochondria (bidirectional communication)

- Nuclear genes regulate mitochondrial function (forward signaling)
- Mitochondria produce peptides (MDPs) regulating nuclear gene expression (retrograde signaling)
- MOTS-c and other MDPs communicate metabolic status to nucleus
- Enables coordinated whole-body metabolic response

PHYSIOLOGICAL ROLES

MOTS-c as Metabolic Status Sensor:

- ✓ Declines with age, obesity, insulin resistance
- ✓ Increases with exercise, caloric restriction, metabolic stress
- ✓ Circulates in bloodstream as endocrine hormone
- ✓ Crosses cell membranes and enters nucleus
- ✓ Directly regulates gene transcription
- ✓ Acts as metabolic "thermostat" maintaining homeostasis

Biological Interpretation:

MOTS-c appears to function as an adaptive response to metabolic stress. When cells experience metabolic challenges (nutrient deprivation, exercise, cold exposure), mitochondria produce MOTS-c to activate compensatory pathways improving metabolic efficiency. Therapeutic MOTS-c administration mimics this adaptive stress response without requiring the stressor itself.^{1,2,3}

AMPK ACTIVATION AND METABOLIC EFFECTS

MOTS-c primary mechanism involves activation of AMP-activated protein kinase (AMPK), the central cellular energy sensor regulating metabolism in response to energy status.¹⁰

AMPK PATHWAY

AMPK Function:

- "Master metabolic switch" activated when cellular energy low (high AMP/ATP ratio)
- Promotes catabolic pathways (energy production)
- Inhibits anabolic pathways (energy consumption)
- Coordinates whole-body metabolic response to energy challenges

MOTS-c → AMPK Activation Cascade:

Step 1: MOTS-c binds to cellular receptors (specific receptor unknown)

Step 2: Increases intracellular AMP levels (mechanism under investigation)

Step 3: AMPK senses elevated AMP/ATP ratio

Step 4: AMPK phosphorylates and activates downstream targets

AMPK Downstream Effects (MOTS-c-Mediated):

GLUCOSE METABOLISM

- ✓ Increased GLUT4 translocation to cell membrane (glucose uptake)
- ✓ Enhanced glycolysis (glucose breakdown for energy)
- ✓ Increased glucose oxidation in mitochondria
- ✓ Improved insulin sensitivity (enhanced insulin signaling)
- ✓ Reduced hepatic gluconeogenesis (decreased glucose production)

LIPID METABOLISM

- ✓ Increased fatty acid oxidation (fat burning)
- ✓ Reduced fatty acid synthesis (decreased fat storage)
- ✓ Enhanced mitochondrial fat oxidation
- ✓ Improved lipid clearance from bloodstream
- ✓ Reduced hepatic triglyceride accumulation (fatty liver improvement)

MITOCHONDRIAL FUNCTION

- ✓ Increased mitochondrial biogenesis (more mitochondria)
- ✓ Enhanced oxidative phosphorylation efficiency
- ✓ Improved electron transport chain function
- ✓ Increased ATP production per unit glucose/fat
- ✓ Reduced reactive oxygen species (ROS) production

EXERCISE MIMETIC EFFECTS

- ✓ AMPK activation mimics exercise at molecular level
- ✓ PGC-1 α activation (master regulator of mitochondrial biogenesis)
- ✓ Increased SIRT1 activity (longevity-associated protein)
- ✓ Enhanced skeletal muscle oxidative capacity
- ✓ Improved physical endurance and exercise tolerance

COMPARISON TO METFORMIN

Metformin and MOTS-c share AMPK activation as primary mechanism:

2026 Regulatory Update (verified — FDA advisory-committee calendar). On **April 15, 2026** the FDA removed MOTS-c from the 503A Category 2 list. On **July 23–24, 2026** the FDA Pharmacy Compounding Advisory Committee (PCAC) reviews MOTS-c

(free base and acetate) for the Section 503A bulk drug substances list — the decision will determine whether U.S. compounding pharmacies may legally dispense it. Recommendation is non-binding and is not FDA drug approval.

Key Difference:

Metformin activates AMPK by inhibiting mitochondrial Complex I (mild metabolic stress). MOTS-c activates AMPK through unknown receptor-mediated pathway without direct mitochondrial inhibition. This mechanistic difference may explain distinct side effect profiles (metformin: GI upset common; MOTS-c: GI effects rare).

NUCLEAR GENE REGULATION AND FOXO PATHWAY

Beyond AMPK activation, MOTS-c directly regulates nuclear gene transcription by entering the nucleus and interacting with FOXO transcription factors.^{2,11}

NUCLEAR TRANSLOCATION

Pathway:

1. MOTS-c produced in mitochondria or administered exogenously
2. Peptide translocates from cytoplasm into nucleus
3. Accumulates in nucleus during metabolic stress conditions
4. Binds to DNA regulatory regions (MOTS-c response elements)
5. Recruits FOXO1 transcription factor to target genes
6. Activates stress response and metabolic adaptation programs

FOXO1 TRANSCRIPTION FACTOR

FOXO1 Function:

- Master regulator of stress response, metabolism, longevity
- Activated by caloric restriction, fasting, exercise
- Promotes cellular survival during metabolic stress
- Regulates glucose metabolism, insulin sensitivity, autophagy
- Associated with longevity in model organisms

MOTS-c Enhancement of FOXO1 Activity:

- ✓ Increases FOXO1 nuclear localization
- ✓ Enhances FOXO1 binding to target gene promoters
- ✓ Amplifies FOXO1-dependent gene transcription
- ✓ Synergizes with other longevity pathways (SIRT1, AMPK)

GENES REGULATED BY MOTS-c/FOXO1:

METABOLIC GENES

- PEPCK, G6Pase (gluconeogenesis enzymes) - DOWNREGULATED
- GLUT4 (glucose transporter) - UPREGULATED

- CPT1 (fatty acid oxidation enzyme) - UPREGULATED
- HMGCR (cholesterol synthesis enzyme) - DOWNREGULATED
- SREBP1c (lipogenesis transcription factor) - DOWNREGULATED

STRESS RESPONSE GENES

- SOD2 (mitochondrial antioxidant enzyme) - UPREGULATED
- Catalase (antioxidant enzyme) - UPREGULATED
- HSP70 (heat shock protein) - UPREGULATED
- ATG genes (autophagy) - UPREGULATED

LONGEVITY-ASSOCIATED GENES

- SIRT1 (NAD⁺-dependent deacetylase) - UPREGULATED
- PGC-1 α (mitochondrial biogenesis) - UPREGULATED
- NRF2 (oxidative stress defense) - UPREGULATED

PRACTICAL IMPLICATION

MOTS-c functions through dual mechanisms:

1. Rapid metabolic effects via AMPK (minutes to hours)
2. Sustained gene expression changes via FOXO1 (hours to days)

This dual mechanism enables both acute metabolic improvements (glucose uptake, fat oxidation) and chronic adaptations (mitochondrial biogenesis, stress resistance). Therapeutic effects likely require sustained administration to maintain gene expression changes.^{2,11}

INSULIN SENSITIZATION MECHANISMS

MOTS-c improves insulin sensitivity through multiple complementary mechanisms:

MECHANISM #1: Enhanced Skeletal Muscle Glucose Uptake

Skeletal muscle accounts for 70-80% of insulin-stimulated glucose disposal.

MOTS-c dramatically improves muscle glucose uptake:^{1,9}

INSULIN-INDEPENDENT PATHWAY

- ✓ AMPK activation → GLUT4 translocation to cell membrane
- ✓ Glucose enters muscle cells without requiring insulin signal
- ✓ Provides "metabolic bypass" around insulin resistance
- ✓ Particularly beneficial in type 2 diabetes (preserved insulin-independent uptake)

INSULIN-SENSITIZING EFFECTS

- ✓ Enhanced insulin receptor substrate (IRS) signaling
- ✓ Improved PI3K/Akt pathway activation
- ✓ Increased insulin receptor density on cell surface

- ✓ Reduced ectopic lipid accumulation in muscle (lipotoxicity)
- ✓ Improved mitochondrial function (more efficient glucose oxidation)

Evidence from Mouse Studies:

- Glucose uptake in muscle increased 40-60% with MOTS-c treatment⁹
- Effect maintained even in insulin-resistant obese mice
- Insulin tolerance tests showed dramatic improvement
- Benefits observed within hours of administration (rapid onset)

MECHANISM #2: Reduced Hepatic Glucose Production

Liver produces glucose through gluconeogenesis and glycogenolysis. Insulin normally suppresses hepatic glucose production; insulin resistance causes excessive hepatic glucose output (major contributor to hyperglycemia).

MOTS-c Hepatic Effects:

- ✓ AMPK activation inhibits gluconeogenic enzymes (PEPCK, G6Pase)
- ✓ FOXO1 regulation suppresses gluconeogenesis gene expression
- ✓ Reduced hepatic lipid accumulation (improved hepatic insulin sensitivity)
- ✓ Enhanced insulin signaling in hepatocytes
- ✓ Decreased fasting hyperglycemia

Evidence from Studies:

- Hepatic glucose production reduced 25-35% in MOTS-c-treated mice¹
- Fasting glucose normalized in diabetic animal models
- Fatty liver (NAFLD) improved with MOTS-c treatment

MECHANISM #3: Adipose Tissue Metabolic Improvement

Adipose tissue insulin resistance contributes to systemic metabolic dysfunction through:

- Excessive free fatty acid release (lipotoxicity)
- Pro-inflammatory adipokine secretion (TNF- α , IL-6)
- Reduced adiponectin (insulin-sensitizing hormone)

MOTS-c Adipose Effects:

- ✓ Enhanced insulin-stimulated glucose uptake in adipocytes
- ✓ Reduced lipolysis (decreased fatty acid release)
- ✓ Improved adipose tissue mitochondrial function
- ✓ Shift toward insulin-sensitive (smaller) adipocytes
- ✓ Potential adiponectin increase (preliminary data)

MECHANISM #4: Systemic Anti-Inflammatory Effects

Chronic low-grade inflammation underlies insulin resistance:

MOTS-c Anti-Inflammatory Actions:

- ✓ Reduced circulating inflammatory cytokines (TNF- α , IL-6, IL-1 β)
- ✓ Decreased macrophage infiltration into adipose tissue
- ✓ Improved adipose tissue inflammatory profile (M2 vs M1 macrophages)
- ✓ Reduced oxidative stress (ROS production)
- ✓ Enhanced antioxidant defense systems

Evidence:

Inflammatory markers reduced 30-50% in animal studies with MOTS-c treatment.

Whether this represents direct anti-inflammatory effect or secondary benefit from improved metabolism remains under investigation.

COMPARISON TO OTHER INSULIN SENSITIZERS

2026 Regulatory Update (verified — FDA advisory-committee calendar). On **April 15, 2026** the FDA removed MOTS-c from the 503A Category 2 list. On **July 23–24, 2026** the FDA Pharmacy Compounding Advisory Committee (PCAC) reviews MOTS-c (free base and acetate) for the Section 503A bulk drug substances list — the decision will determine whether U.S. compounding pharmacies may legally dispense it. Recommendation is non-binding and is not FDA drug approval.

MOTS-c Unique Features:

- Direct skeletal muscle glucose uptake (insulin-independent)
- Mimics molecular effects of exercise
- Endogenous peptide (physiologic replacement)
- Minimal side effects in preliminary studies

Limitations:

- Unproven efficacy in humans
- Subcutaneous injection requirement
- Short half-life (requires daily dosing)
- Unknown long-term effects

Reported Clinical Applications

CRITICAL DISCLAIMER: THE FOLLOWING PROTOCOLS REPRESENT REPORTED USES IN RESEARCH

settings and clinical practice. MOTS-c is NOT FDA-approved for any indication.

All applications are investigational and off-label. Providers assume full liability for prescribing experimental compounds.

STANDARD REPORTED DOSING PROTOCOLS

DOSING RATIONALE

Human dosing protocols derive from:

1. Allometric scaling from effective mouse doses (5-15 mg/kg)
2. Body surface area adjustments
3. Primate pharmacokinetic studies⁴
4. Small human pilot study experience (unpublished)
5. Clinical practice empiric observations

STANDARD SUBCUTANEOUS PROTOCOL

INITIATION DOSE

Week 1-2: 5 mg subcutaneously once daily

- Conservative starting dose
- Assess tolerability
- Monitor for injection site reactions

MAINTENANCE DOSE

Week 3+: 10 mg subcutaneously once daily

- Most commonly reported therapeutic dose
- Administered 7 days per week
- Injection timing: Morning preferred (mimics physiologic MOTS-c diurnal pattern)

HIGHER DOSE (SELECTED PATIENTS)

Week 5+: 15 mg subcutaneously once daily (optional)

- Reserved for non-responders at 10mg
- Used in athletic performance optimization contexts
- Increased expense without established additional benefit
- Unknown if higher doses confer additional metabolic improvements

DOSING FREQUENCY

Daily Administration Rationale:

- MOTS-c half-life: ~30 minutes (rapid clearance)
- Metabolic effects persist 18-24 hours despite short half-life
- Daily dosing maintains steady-state metabolic adaptations
- Skipped doses result in temporary loss of benefits

Alternative Dosing (REPORTED, NOT RECOMMENDED):

- Every other day: 15-20mg (attempts to reduce injection frequency)
- Twice weekly: 20-30mg (similar rationale)

- Issues with intermittent dosing:
 - ✗ Metabolic effects may wane between doses
 - ✗ Gene expression changes require consistent stimulation
 - ✗ No data supporting equivalent efficacy to daily dosing

ADMINISTRATION TECHNIQUE

INJECTION SITES

- ✓ Abdomen (2 inches from umbilicus, rotate sites)
- ✓ Anterior thigh (mid-thigh, lateral aspect)
- ✓ Deltoid (upper arm, less common)
- ✗ Avoid: Buttocks (poor absorption, painful)

INJECTION PROCEDURE

1. Reconstitute lyophilized MOTS-c with bacteriostatic water:
 - Standard concentration: 10mg/mL (1mg MOTS-c in 1mL water)
 - Add water slowly to side of vial (avoid foaming)
 - Swirl gently (do NOT shake vigorously)
 - Allow to dissolve completely (1-2 minutes)
 - Inspect for particulates (should be clear, colorless solution)
2. Draw prescribed dose using insulin syringe:
 - 5mg dose = 0.5mL
 - 10mg dose = 1.0mL
 - 15mg dose = 1.5mL
3. Clean injection site with alcohol swab
4. Pinch skin and inject at 45-90° angle
5. Inject slowly over 3-5 seconds
6. Withdraw needle and apply gentle pressure (no rubbing)
7. Rotate injection sites daily

STORAGE

- Unreconstituted (lyophilized powder): Refrigerate 36-46°F (2-8°C), protect from light
- Reconstituted solution: Refrigerate, use within 14 days (bacteriostatic water)
- Travel: Keep in cool environment, ice pack acceptable

INDICATION-SPECIFIC PROTOCOLS (REPORTED)

REPORTED APPLICATION #1: INSULIN RESISTANCE AND PREDIABETES

TARGET POPULATION

- Fasting glucose 100-125 mg/dL (impaired fasting glucose)
- HbA1c 5.7-6.4% (prediabetes range)
- HOMA-IR >2.5 (insulin resistance)
- Metabolic syndrome

REPORTED PROTOCOL

- MOTS-c 10mg subcutaneously daily for 12-16 weeks minimum
- Combined with lifestyle modification:
- Caloric restriction (500 kcal/day deficit)
- Exercise 150+ minutes per week (aerobic + resistance)
- Mediterranean or low-glycemic diet
- Monitoring: Fasting glucose, HbA1c, fasting insulin monthly

REPORTED OUTCOMES (ANECDOTAL/UNPUBLISHED)

- Fasting glucose reduction: 10-20 mg/dL
- HbA1c reduction: 0.3-0.5%
- Fasting insulin reduction: 20-30%
- HOMA-IR improvement: 30-40%

COMPARISON TO STANDARD THERAPY

Metformin (FDA-approved for prediabetes):

- Fasting glucose reduction: 15-25 mg/dL
- HbA1c reduction: 0.5-0.7%
- Evidence: Multiple large RCTs (Diabetes Prevention Program)
- Standard of care: Metformin 850mg twice daily

Clinical Reality:

MOTS-c reported effects similar magnitude to metformin based on limited data.

However, metformin has vastly superior evidence base, FDA approval, and should be first-line therapy. MOTS-c appropriate only for patients intolerant to metformin or as adjunctive therapy.

REPORTED APPLICATION #2: TYPE 2 DIABETES (ADJUNCTIVE THERAPY)

TARGET POPULATION

- Established type 2 diabetes (HbA1c $\geq$ 6.5%)
- Inadequately controlled on oral agents

- Insulin resistance predominant (vs insulin deficiency)
- NOT intended as monotherapy

REPORTED PROTOCOL

- MOTS-c 10-15mg subcutaneously daily
- MUST combine with standard diabetes medications:
- Metformin (first-line)
- ± SGLT2 inhibitor
- ± GLP-1 agonist
- ± DPP-4 inhibitor
- ± Insulin (if needed)
- Duration: Indefinite (chronic therapy)
- Monitoring: HbA1c every 3 months, fasting glucose weekly

REPORTED OUTCOMES (CASE SERIES)

- HbA1c reduction: 0.4-0.8% when added to existing therapy
- Reduced insulin requirements: 10-30% dose reduction
- Weight: Modest loss 1-3 kg over 6 months
- Medication reduction: Some patients able to discontinue sulfonylureas

CRITICAL LIMITATION

NO published controlled trials in diabetic patients. All data from:

- Individual case reports
- Small case series (N=5-20)
- Clinician anecdotal reports
- Conference abstract preliminary data

Standard of Care:

Type 2 diabetes managed with FDA-approved medications:

1. Metformin (first-line)
2. GLP-1 agonist (if overweight, cardiovascular disease)
3. SGLT2 inhibitor (if cardiovascular disease, chronic kidney disease)
4. Insulin (if inadequate control)

MOTS-c does NOT replace any of these evidence-based therapies. Use only as experimental adjunct in patients already on optimized standard therapy.

REPORTED APPLICATION #3: WEIGHT MANAGEMENT

TARGET POPULATION

- Overweight (BMI 25-29.9) or obese (BMI ≥30)

- Metabolically unhealthy obesity (insulin resistance, fatty liver)
- Preferably combined with exercise program
- NOT appropriate for aesthetic weight loss in healthy-weight individuals

REPORTED PROTOCOL

- MOTS-c 10mg subcutaneously daily
- Duration: 3-6 months minimum
- Combined with comprehensive lifestyle program:
- Caloric deficit: 500-750 kcal/day
- High-protein diet: 1.2-1.6g/kg body weight
- Exercise: 200+ minutes per week
- Resistance training: 3x per week
- Monitoring: Weight, body composition, waist circumference monthly

REPORTED OUTCOMES (ANECDOTAL)

- Weight loss: 2-5 kg over 3-6 months
- Body composition: Preferential fat loss (muscle sparing)
- Visceral fat: Reduced (waist circumference decrease)
- Exercise capacity: Improved (subjective reports)

COMPARISON TO FDA-APPROVED OBESITY MEDICATIONS

2026 Regulatory Update (verified — FDA advisory-committee calendar). On **April 15, 2026** the FDA removed MOTS-c from the 503A Category 2 list. On **July 23–24, 2026** the FDA Pharmacy Compounding Advisory Committee (PCAC) reviews MOTS-c (free base and acetate) for the Section 503A bulk drug substances list — the decision will determine whether U.S. compounding pharmacies may legally dispense it. Recommendation is non-binding and is not FDA drug approval.

Clinical Reality:

MOTS-c weight loss effects appear MODEST compared to FDA-approved obesity medications. If weight loss is primary goal, semaglutide or tirzepatide vastly superior choices. MOTS-c potentially appropriate for:

- Patients intolerant to GLP-1 agonists
- Patients with insurance denials for FDA-approved agents
- As adjunct to GLP-1 therapy (combination approach)
- Metabolic health focus beyond weight loss alone

REPORTED APPLICATION #4: ATHLETIC PERFORMANCE ENHANCEMENT

TARGET POPULATION

- Athletes seeking performance optimization
- Recreational fitness enthusiasts
- Aging adults maintaining physical function
- Combined with structured training program

REPORTED PROTOCOL

- MOTS-c 10-15mg subcutaneously daily
- Injection timing: 30-60 minutes before exercise (anecdotal preference)
- Duration: Training cycles (8-12 weeks), repeated
- Combined with:
- Structured training program
- Adequate protein intake (1.6-2.2g/kg for athletes)
- Recovery protocols (sleep, rest days)

REPORTED OUTCOMES (ATHLETE ANECDOTES)

- Endurance: Improved time to exhaustion
- Recovery: Faster recovery between training sessions
- Body composition: Enhanced fat loss while maintaining lean mass
- Subjective energy: Increased perceived energy and workout quality
- Performance metrics: Modest improvements in sport-specific measures

MECHANISTIC RATIONALE

- AMPK activation mimics exercise signaling
- Enhanced mitochondrial biogenesis (more mitochondria in muscle)
- Improved oxidative capacity (better fat utilization)
- Exercise mimetic without replacing actual exercise
- Potential synergy: Exercise + MOTS-c > either alone

CRITICAL LIMITATIONS

- ! NO published performance studies in athletes
- ! All data anecdotal or from N=1 self-experiments
- ! Placebo effect extremely powerful in performance contexts
- ! Anti-doping status: UNCLEAR (not explicitly banned but peptides scrutinized)
- ! Performance enhancement in healthy individuals ethically questionable

SPORTS MEDICINE PERSPECTIVE

Elite athletes already optimize training, nutrition, recovery. Marginal gains difficult to achieve through pharmacology. MOTS-c unlikely to provide competitive

advantage beyond well-designed training. Risk-benefit unfavorable given unknown long-term safety and lack of evidence.

REPORTED APPLICATION #5: HEALTHY AGING AND LONGEVITY

TARGET POPULATION

- Adults >50 years old
- Metabolic decline associated with aging
- Sarcopenia (age-related muscle loss)
- Declining physical function
- Longevity optimization focus

REPORTED PROTOCOL

- MOTS-c 10mg subcutaneously daily
- Duration: Long-term (years, indefinite)
- Combined with:
- Resistance training 2-3x per week (muscle preservation)
- Adequate protein: 1.2-1.5g/kg body weight
- Mediterranean diet pattern
- Sleep optimization (7-9 hours)

REPORTED OUTCOMES (ANECDOTAL)

- Muscle mass: Preserved or slightly increased
- Physical function: Improved gait speed, grip strength
- Metabolic markers: Improved glucose, lipids, inflammatory markers
- Subjective: Increased energy, vitality, quality of life
- Longevity: UNKNOWN (requires decades to assess)

MECHANISTIC RATIONALE FOR LONGEVITY

- MOTS-c declines with aging (restoration to youthful levels)
- AMPK activation associated with longevity in model organisms
- FOXO1 pathway linked to lifespan extension
- Mitochondrial function central to aging biology
- Centenarians have favorable MOTS-c genetic variants⁵

CRITICAL REALITY CHECK

- ! NO human longevity data (impossible to obtain in reasonable timeframe)
- ! Mouse longevity studies do NOT predict human lifespan effects
- ! Correlation (centenarians have certain variants) ≠ Causation (MOTS-c extends life)
- ! Rapamycin, metformin, resveratrol all promoted for longevity with minimal human evidence

- ! Healthspan (quality of life) more measurable than lifespan
- ! Lifestyle factors (exercise, diet, sleep, social connection) have strongest longevity evidence

HONEST ASSESSMENT

MOTS-c MAY contribute to healthy aging by preserving metabolic function and muscle mass. However, longevity claims are highly speculative. Focus should be on established aging interventions (exercise, nutrition, social engagement) with MOTS-c as experimental adjunct, not longevity magic bullet.

COMBINATION THERAPY PROTOCOLS (REPORTED)

MOTS-c + Metformin

RATIONALE

- Both activate AMPK (potentially synergistic)
- Metformin: Hepatic AMPK activation
- MOTS-c: Skeletal muscle AMPK activation
- Complementary tissue targeting

REPORTED PROTOCOL

- Metformin 850-1000mg twice daily (standard dosing)
- MOTS-c 10mg subcutaneously daily
- Monitor for hypoglycemia (rare but additive glucose-lowering)
- No documented adverse drug interactions

THEORETICAL CONCERNS

- ! Additive AMPK activation: Could excessive AMPK activation cause harm?
- ! Lactic acidosis: Metformin carries rare lactic acidosis risk; does MOTS-c amplify?
- ! No formal interaction studies performed

Clinical Experience:

Anecdotal reports suggest combination well-tolerated. No documented serious adverse events. However, combination represents off-protocol use of investigational agent with established medication.

MOTS-c + GLP-1 Agonist (Semaglutide, Tirzepatide, Liraglutide)

RATIONALE

- Complementary mechanisms:
- GLP-1: Appetite suppression, insulin secretion
- MOTS-c: Insulin sensitivity, exercise mimetic
- GLP-1 agonists address energy intake (reduce appetite)
- MOTS-c addresses energy expenditure and utilization

- Potential synergy for weight loss and metabolic improvement

REPORTED PROTOCOL

- GLP-1 agonist at standard therapeutic dose:
- Semaglutide 2.4mg weekly, OR
- Tirzepatide 10-15mg weekly, OR
- Liraglutide 3.0mg daily
- MOTS-c 10mg subcutaneously daily
- Monitor glucose closely if diabetic (additive glucose-lowering)

REPORTED OUTCOMES (ANECDOTAL)

- Enhanced weight loss vs GLP-1 alone (uncontrolled observations)
- Improved body composition (muscle preservation during weight loss)
- Better exercise tolerance despite caloric restriction
- Metabolic improvements beyond weight loss alone

CRITICAL LIMITATIONS

- ! NO controlled studies of combination therapy
- ! Unknown if combination superior to GLP-1 monotherapy
- ! Adding experimental agent to effective FDA-approved therapy questionable
- ! Substantially increased treatment expense
- ! Insurance unlikely to cover either agent in combination

PRACTICAL CONSIDERATION

If patient responding well to GLP-1 agonist monotherapy (15-20% weight loss), adding MOTS-c provides unclear incremental benefit. Reserve combination for:

- GLP-1 partial responders (plateaued weight loss)
- Patients experiencing muscle loss with GLP-1 therapy
- Focus on metabolic health beyond weight (insulin sensitivity, exercise capacity)

MOTS-c + Exercise Programs

RATIONALE

- MOTS-c mimics exercise at molecular level
- Exercise increases endogenous MOTS-c production
- Combination may produce synergistic effects
- Both target mitochondrial adaptation

REPORTED PROTOCOL

- MOTS-c 10mg subcutaneously daily
- Structured exercise program:

- Aerobic: 150-200 minutes per week (moderate intensity)
- Resistance: 2-3 sessions per week (full body, progressive overload)
- Flexibility/mobility: 2-3 sessions per week
- MOTS-c injection timing: 30-60 minutes pre-exercise (anecdotal preference)

REPORTED OUTCOMES

- Enhanced training adaptations (strength, endurance gains)
- Faster recovery between sessions
- Improved body composition changes vs exercise alone
- Greater exercise tolerance (can sustain higher intensities)

MECHANISTIC SYNERGY

- Exercise produces metabolic stress → MOTS-c produced endogenously
- Exogenous MOTS-c amplifies exercise-induced AMPK/FOXO1 signaling
- Combined stimulus for mitochondrial biogenesis exceeds either alone
- Enhanced training effect without additional exercise volume

CRITICAL CAVEAT

Exercise alone produces profound metabolic benefits (insulin sensitivity, weight loss, cardiovascular health). MOTS-c without exercise substantially less effective in preliminary data. Peptide is adjunct to, not substitute for, physical activity.

Patient Selection

IDEAL CANDIDATE CHARACTERISTICS

TIER 1: MOST APPROPRIATE CANDIDATES (IF PRESCRIBING)

PATIENT PROFILE #1: Metformin-Intolerant Prediabetes

- Impaired fasting glucose or impaired glucose tolerance
- Intolerant to metformin (severe GI side effects)
- Failed lifestyle modification alone
- Motivated for metabolic disease prevention
- Accepts experimental nature of therapy
- Financially able to afford ongoing treatment

RATIONALE

Metformin is first-line for prediabetes but 30-40% discontinue due to GI side effects. MOTS-c offers alternative AMPK-activating approach with potentially better GI tolerability. However, insurance will not cover (experimental status).

PATIENT PROFILE #2: Metabolic Syndrome Despite Optimized Standard Therapy

- Metabolic syndrome (hypertension, dyslipidemia, insulin resistance, abdominal obesity)
- Already on metformin, statins, antihypertensives
- Plateaued despite lifestyle modification
- Seeking additional metabolic optimization
- Understands investigational status

RATIONALE

Patients already on evidence-based therapies with residual metabolic dysfunction may benefit from adjunctive experimental approach. MOTS-c potentially additive to existing treatments targeting complementary pathways.

PATIENT PROFILE #3: GLP-1 Agonist Partial Responders

- Initiated GLP-1 agonist for weight loss/diabetes
- Achieved initial benefit but plateaued
- Seeking additional weight loss or metabolic improvement
- Financially able to afford combination therapy
- Accepts unknown risk of combining therapies

RATIONALE

GLP-1 agonists address appetite and insulin secretion. MOTS-c addresses insulin sensitivity and muscle metabolism (complementary). Combination may overcome GLP-1 plateau, though evidence lacking.

PATIENT PROFILE #4: Athletes/Active Individuals with Metabolic Goals

- Physically active lifestyle already established
- Seeking metabolic optimization to support training
- Insulin resistance despite exercise (possible genetic factors)
- Performance enhancement secondary to health optimization
- NOT seeking athletic competitive advantage

RATIONALE

MOTS-c described as "exercise mimetic" with potential synergy when combined with actual exercise. Athletes may amplify training adaptations. However, performance benefits unproven and anti-doping considerations apply.

TIER 2: ACCEPTABLE BUT SUBOPTIMAL CANDIDATES

PATIENT PROFILE #5: Aging Adults Focused on Metabolic Preservation

- Age >55 years with age-related metabolic decline
- Declining muscle mass (sarcopenia)
- Reduced exercise capacity
- Family history of metabolic disease

- Proactive approach to healthy aging
- Financially secure (long-term treatment required)

RATIONALE

MOTS-c declines with aging; supplementation may restore youthful levels. Potential for preserving metabolic health and physical function. However, longevity claims speculative and lifestyle interventions have stronger evidence.

TIER 3: INAPPROPRIATE CANDIDATES (SHOULD NOT PRESCRIBE)

PATIENT PROFILE #6: Treatment-Naive Type 2 Diabetes or Prediabetes

- New diagnosis of prediabetes or type 2 diabetes
- Not yet tried metformin or lifestyle modification
- Seeking "natural" or "advanced" therapy
- Wants to avoid conventional medications

RECOMMENDATION

Patients with newly diagnosed metabolic disease should receive FDA-approved first-line therapies (metformin, lifestyle modification) before considering investigational agents. Prescribing MOTS-c as initial therapy represents substandard care.

PATIENT PROFILE #7: Aesthetic Weight Loss Without Metabolic Disease

- Normal glucose tolerance
- No insulin resistance
- BMI <27 without metabolic complications
- Cosmetic weight loss goal
- Wanting "shortcut" to weight loss

RECOMMENDATION

MOTS-c targets insulin resistance and metabolic dysfunction. Metabolically healthy individuals unlikely to benefit significantly. Additionally, exposing healthy individuals to experimental peptide with unknown long-term safety ethically questionable.

PATIENT PROFILE #8: Financially Constrained Patients

- Unable to afford ongoing treatment
- Seeking less expensive alternative to FDA-approved medications
- Expects insurance coverage (will be denied)

RECOMMENDATION

MOTS-c is expensive research compound without insurance coverage. Treatment costs

substantially exceed many FDA-approved alternatives (metformin, generic GLP-1 agonists emerging). Financial toxicity from uncovered experimental therapy problematic.

PATIENT PROFILE #9: Patients Seeking "Magic Bullet" Without Lifestyle Change

- Unwilling to modify diet
- Sedentary lifestyle unchanged
- Views peptide as pharmaceutical solution to behavioral issues
- Poor adherence history with prior medications

RECOMMENDATION

MOTS-c appears most effective when combined with exercise and dietary modification.

Patients unwilling to implement lifestyle changes unlikely to achieve meaningful benefit. Peptide is adjunct, not replacement, for behavioral modification.

CONTRAINDICATIONS AND PRECAUTIONS

ABSOLUTE CONTRAINDICATIONS

✗ Pregnancy

- MOTS-c safety in pregnancy COMPLETELY UNKNOWN
- Mitochondrial function critical for fetal development
- No animal reproductive toxicology studies
- Potential effects on fetal mitochondrial programming unknown
- Risk-benefit ratio unacceptable
- Discontinue immediately if pregnancy confirmed or suspected

✗ Breastfeeding/Lactation

- Unknown if MOTS-c excreted in breast milk
- Potential effects on infant mitochondrial development unknown
- No safety data in nursing mothers or infants
- Avoid until breastfeeding complete

✗ Age <18 Years

- No pediatric safety or efficacy data
- Mitochondrial function critical for growth and development
- Effects on developing metabolism unknown
- Contraindicated in children and adolescents

✗ Active Malignancy

- Mitochondrial function affects cancer cell metabolism
- MOTS-c effects on tumor growth and metastasis unknown

- Some cancers rely on mitochondrial metabolism (OXPHOS)
- Others use glycolysis (Warburg effect)
- Theoretical concern: Enhanced mitochondrial function could support tumor growth
- Contraindicated until cancer in remission, then discuss with oncologist

RELATIVE CONTRAINDICATIONS (USE WITH EXTREME CAUTION)

! Severe Renal Impairment (eGFR <30 mL/min/1.73m²)

- Peptide clearance may be impaired
- Risk of accumulation and unknown toxicity
- No dose adjustment guidelines (no data)
- Metformin contraindicated in severe renal disease; MOTS-c shares mechanism
- If prescribing, dose reduction to 5mg daily and close monitoring

! Severe Hepatic Impairment (Child-Pugh C Cirrhosis)

- Mitochondrial dysfunction common in advanced liver disease
- MOTS-c effects in setting of hepatic mitochondrial pathology unknown
- Liver synthesizes proteins regulating mitochondrial function
- Avoid or use with extreme caution with hepatology consultation

! History of Lactic Acidosis

- AMPK activation by metformin rarely causes lactic acidosis
- MOTS-c activates AMPK through different mechanism
- Theoretical risk of lactic acidosis exists
- Contraindicated if prior metformin-associated lactic acidosis

! Mitochondrial Diseases (Inherited Mitochondrial Disorders)

- MOTS-c encoded by mitochondrial DNA
- Patients with mitochondrial disease have abnormal mitochondrial genomes
- MOTS-c effects in setting of primary mitochondrial disorder unpredictable
- Could help (compensatory) or harm (worsen dysfunction)
- Requires mitochondrial disease specialist consultation before considering

! Severe Malnutrition or Cachexia

- MOTS-c increases energy expenditure and fat oxidation
- Malnourished patients lack energy reserves
- Further metabolic activation could worsen nutritional status
- Correct malnutrition before considering MOTS-c

! Hypoglycemia-Prone Patients

- Type 1 diabetes with brittle control
- Post-bariatric surgery dumping syndrome

- Insulinoma or other causes of endogenous hyperinsulinism
- MOTS-c improves insulin sensitivity (may worsen hypoglycemia)
- If prescribing, reduce insulin doses prophylactically and monitor closely

DRUG-DRUG INTERACTIONS (THEORETICAL)

METFORMIN

- Both activate AMPK (potentially additive effects)
- Combination generally appears safe in clinical experience
- Monitor for lactic acidosis symptoms (rare)
- May allow metformin dose reduction in some patients

INSULIN AND INSULIN SECRETAGOGUES

- MOTS-c improves insulin sensitivity
- Risk of hypoglycemia if insulin doses not adjusted
- Reduce insulin doses by 10-20% when initiating MOTS-c
- Monitor glucose closely during titration
- Sulfonylureas: Consider dose reduction or discontinuation

THYROID HORMONE

- Mitochondria central to thyroid hormone action
- Theoretical interaction possible
- Monitor thyroid function (TSH, free T4) if on thyroid replacement
- Dose adjustments may be needed

STATINS

- Some statins affect mitochondrial function (myopathy risk)
- Theoretical interaction with mitochondrial peptide
- Monitor for muscle symptoms (myalgia, weakness)
- Check CK if symptoms develop
- No documented interaction in limited human experience

SUPPLEMENTS

- CoQ10, NAD+ precursors (NR, NMN), resveratrol, berberine all affect mitochondria
- Unknown interactions with MOTS-c
- Combination likely safe but unproven
- May be synergistic (theoretical)

SCREENING AND ELIGIBILITY ASSESSMENT

PRE-TREATMENT EVALUATION

CLINICAL ASSESSMENT

- ✓ Complete medical history:
 - Current medical conditions (especially diabetes, renal, hepatic, cardiac)
 - Medication list (assess drug interactions)
 - Prior adverse reactions to medications
 - Family history (metabolic disease, cancer)
 - Lifestyle (diet, exercise, alcohol, smoking)
- ✓ Physical examination:
 - Weight, height, BMI calculation
 - Blood pressure, heart rate
 - Waist circumference (metabolic syndrome criterion)
 - Body composition (if available: DEXA, bioimpedance)
 - Injection site examination (teach proper technique)

BASELINE LABORATORY TESTS

REQUIRED

- ✓ Comprehensive metabolic panel:
 - Glucose, creatinine, eGFR, electrolytes
 - AST, ALT, bilirubin (hepatic function)
 - Establish baseline organ function
- ✓ Hemoglobin A1c:
 - All patients (diabetic and non-diabetic)
 - Assess baseline glucose control
 - Track efficacy
- ✓ Lipid panel:
 - Total cholesterol, LDL, HDL, triglycerides
 - Metabolic syndrome diagnosis
 - Track metabolic improvements
- ✓ Pregnancy test (females of childbearing potential):
 - Rule out pregnancy before initiating
 - Discuss contraception requirements during therapy

OPTIONAL (BASED ON CLINICAL SCENARIO)

- ! Fasting insulin, HOMA-IR:

- Quantify insulin resistance severity
- Not required but helpful for tracking improvement
- ! C-peptide:
 - Differentiate type 1 vs type 2 diabetes (if unclear)
 - Assess endogenous insulin production
- ! Thyroid function (TSH, free T4):
 - Rule out thyroid dysfunction causing metabolic issues
 - Baseline if on thyroid medication
- ! Inflammatory markers (hsCRP):
 - Assess systemic inflammation
 - Track anti-inflammatory effects
- ! Lactate (arterial or venous):
 - Baseline only if lactic acidosis risk factors
 - Not routine

ELIGIBILITY CHECKLIST

PATIENT MUST MEET ALL OF THE FOLLOWING

- ☒ Age ≥ 18 years
- ☒ Not pregnant or breastfeeding
- ☒ eGFR ≥ 30 mL/min/1.73m² (preferably ≥ 45)
- ☒ No active malignancy
- ☒ Understands investigational/experimental status
- ☒ Able to afford ongoing treatment (no insurance coverage)
- ☒ Willing to self-inject daily
- ☒ Willing to implement lifestyle modifications
- ☒ Willing to participate in close monitoring

RED FLAGS PRECLUDING TREATMENT

- ☐ Pregnancy or planning pregnancy within 6 months
- ☐ Active cancer under treatment
- ☐ Severe renal impairment (eGFR < 30)
- ☐ Decompensated liver disease
- ☐ History of lactic acidosis
- ☐ Primary mitochondrial disease
- ☐ Unrealistic expectations (seeking magic bullet)
- ☐ Unable to afford treatment

⊘ Unwilling to modify lifestyle

Informed Consent

ESSENTIAL DISCLOSURE ELEMENTS

INFORMED CONSENT MUST INCLUDE

✓ DISCLOSURE #1: Experimental/Investigational Status

"MOTS-c is NOT approved by the FDA for any medical condition. This is an experimental therapy. You are essentially participating in an uncontrolled experiment by using this peptide. The FDA has not reviewed this compound for safety or efficacy."

✓ DISCLOSURE #2: Limited Human Data

"MOTS-c has been studied primarily in mice and laboratory animals. Human data is extremely limited - fewer than 50 people have been studied in research settings, and no peer-reviewed published studies exist. We are essentially guessing at the appropriate dose, duration, and safety profile based on animal studies."

✓ DISCLOSURE #3: Unknown Long-Term Safety

"We have NO information about safety beyond 6 months of use. Long-term effects on cancer risk, heart disease, kidney function, liver function, and other organs are completely unknown. You could be at risk for serious health problems years from now that we cannot predict today."

✓ DISCLOSURE #4: FDA-Approved Alternatives Exist

"For your condition [prediabetes/diabetes/obesity], there are FDA-approved medications with decades of safety data and proven efficacy:

- Metformin for prediabetes and diabetes
- GLP-1 agonists (semaglutide, tirzepatide, liraglutide) for obesity and diabetes
- SGLT2 inhibitors for diabetes

These medications have been studied in tens of thousands of patients with long-term safety data. MOTS-c has not. Standard medical practice would be to use these FDA-approved medications first."

✓ DISCLOSURE #5: No Insurance Coverage

"MOTS-c is not covered by any insurance including Medicare and Medicaid. You will pay out-of-pocket for the medication, and costs will be substantial for ongoing treatment. This is not reimbursable through health savings accounts (HSA)

or flexible spending accounts (FSA) in most cases."

✓ DISCLOSURE #6: Variable Quality From Compounding Pharmacies

"MOTS-c is made by compounding pharmacies which are not regulated by the FDA in the same way as pharmaceutical manufacturers. Quality, purity, and potency can vary between pharmacies and even between batches. There is no guarantee that what you receive contains exactly what the label states."

✓ DISCLOSURE #7: Injection Requirement and Compliance Burden

"MOTS-c requires daily subcutaneous injections, meaning you must inject yourself every day indefinitely. Missing doses may result in loss of any beneficial effects. Injection site reactions, pain, and inconvenience are common."

✓ DISCLOSURE #8: Unknown Efficacy

"While animal studies suggest MOTS-c may improve insulin sensitivity and metabolic health, we do not know if this translates to meaningful benefits in humans. You may experience no benefit whatsoever despite the expense and inconvenience. There is no way to predict who will respond and who will not."

✓ DISCLOSURE #9: Pregnancy Contraindication

[For females of childbearing potential]:

"You MUST use effective contraception while on MOTS-c. Effects on pregnancy and fetal development are completely unknown. You must have a negative pregnancy test before starting and notify me immediately if you become pregnant. You must discontinue MOTS-c if pregnancy occurs."

✓ DISCLOSURE #10: Right to Refuse and Alternatives

"You have the right to refuse this experimental therapy. Standard medical care with FDA-approved medications remains available and is recommended. Choosing standard care does not affect our doctor-patient relationship."

INFORMED CONSENT DOCUMENTATION TEMPLATE

INFORMED CONSENT FOR EXPERIMENTAL PEPTIDE THERAPY

MOTS-c (Mitochondrial-Derived Peptide)

PATIENT NAME: _____ DATE: _____

INTRODUCTION

Your physician has discussed the possibility of using MOTS-c, an investigational peptide therapy, for [INDICATION: prediabetes/insulin resistance/metabolic syndrome/other]. This document explains important information about this experimental

treatment. Please read carefully and ask questions before signing.

YOUR MEDICAL CONDITION

You have been diagnosed with: [LIST CONDITIONS]

Standard FDA-approved treatments for your condition include: [LIST ALTERNATIVES]

ABOUT MOTS-C

MOTS-c is a 16-amino acid peptide that occurs naturally in human cells. It is encoded by mitochondrial DNA and appears to play a role in regulating metabolism and insulin sensitivity. Research suggests that levels decline with aging and metabolic disease.

INVESTIGATIONAL STATUS

- ☒ MOTS-c is NOT FDA-approved for any medical condition
- ☒ This is experimental therapy with limited human data
- ☒ Fewer than 50 humans have been studied in research settings
- ☒ No peer-reviewed published studies of human efficacy exist
- ☒ Animal studies show promise but often fail to predict human outcomes

PROPOSED TREATMENT

- Medication: MOTS-c [DOSE] mg subcutaneously
- Frequency: Once daily, every day
- Duration: [SPECIFIED DURATION] initially, potentially indefinite
- Administration: Self-injection at home
- Monitoring: Laboratory tests and office visits as outlined below

EXPECTED BENEFITS (THEORETICAL)

Based on animal studies and limited human data, MOTS-c may provide:

- Improved insulin sensitivity
- Better glucose control
- Modest weight loss (2-5 kg over several months)
- Enhanced exercise capacity
- Improved metabolic markers

IMPORTANT: These benefits are not guaranteed. Many patients may experience no benefit despite treatment expense and inconvenience.

RISKS AND SIDE EFFECTS

COMMON (>5% based on limited data):

- Injection site reactions (redness, pain, swelling)
- Mild fatigue or tiredness

- Headache
- Possible hypoglycemia if you have diabetes

SERIOUS (rare or unknown frequency):

- Allergic reactions (rash, itching, difficulty breathing)
- Unknown long-term cancer risk
- Unknown long-term organ damage (liver, kidney, heart)
- Unknown interactions with other medications
- Lactic acidosis (similar to metformin, rare but serious)
- Birth defects (if pregnant - MOTS-c contraindicated)

UNKNOWN RISKS

Because this is an experimental therapy, there may be serious risks that are currently unknown. Long-term safety beyond 6 months has not been studied.

ALTERNATIVES TO MOTS-C

FDA-APPROVED MEDICATIONS

1. Metformin (for prediabetes/diabetes) - first-line therapy, extensive safety data
2. GLP-1 agonists: Semaglutide (Wegovy®/Ozempic®), Tirzepatide (Zepbound®/Mounjaro®), Liraglutide (Saxenda®/Victoza®) - FDA-approved for obesity and/or diabetes
3. SGLT2 inhibitors (for diabetes with heart or kidney disease)
4. Other diabetes medications as appropriate

LIFESTYLE MODIFICATION

- Dietary changes (caloric reduction, healthy eating patterns)
- Exercise (150+ minutes per week aerobic + resistance training)
- Weight loss through behavior modification
- These interventions have strong evidence for metabolic benefit

COSTS

- MOTS-c is NOT covered by insurance
- You will pay approximately [ESTIMATED COST] monthly out-of-pocket
- Laboratory monitoring and office visits may or may not be covered by insurance
- Total estimated cost: [TOTAL ESTIMATED COST] for [TIME PERIOD]

MONITORING REQUIREMENTS

You agree to the following monitoring schedule:

- Week 4: Office visit, adverse effects assessment
- Month 3: Office visit, laboratory tests (CMP, HbA1c, lipids)
- Month 6: Office visit, laboratory tests

- Ongoing: Every 3-6 months as determined by physician

You agree to report immediately:

- Severe injection site reactions
- Signs of allergic reaction (rash, difficulty breathing, swelling)
- Unusual fatigue, muscle pain, weakness
- Nausea, vomiting, rapid breathing (possible lactic acidosis)
- Pregnancy or suspected pregnancy

PREGNANCY AND CONTRACEPTION

[For females of childbearing potential]:

- ☒ I understand MOTS-c is contraindicated in pregnancy
- ☒ I have had a negative pregnancy test today
- ☒ I agree to use effective contraception during treatment
- ☒ I will notify my physician immediately if I become pregnant
- ☒ I understand I must stop MOTS-c if pregnancy occurs

CONFIDENTIALITY

Your use of MOTS-c and all related medical information will be kept confidential in accordance with HIPAA regulations. De-identified data may be used for research purposes.

YOUR RIGHTS

- ☒ You may refuse this experimental therapy - this will not affect your care
- ☒ You may discontinue MOTS-c at any time for any reason
- ☒ You may ask questions before, during, and after treatment
- ☒ Standard FDA-approved therapies remain available to you

PHYSICIAN STATEMENT

I have discussed MOTS-c with this patient including its investigational status, limited human data, unknown long-term safety, FDA-approved alternatives, costs, risks, and benefits. The patient has had opportunity to ask questions and demonstrated understanding of the information provided.

Physician Signature: _____ Date: _____

PATIENT STATEMENT

- ☒ I have read this consent form (or had it read to me)
- ☒ I have had opportunity to ask questions and receive answers
- ☒ I understand MOTS-c is NOT FDA-approved and is experimental
- ☒ I understand there is limited human safety and efficacy data

- ☒ I understand FDA-approved alternatives exist and have been offered to me
- ☒ I understand the risks, benefits, and costs of treatment
- ☒ I agree to required monitoring and will report adverse effects
- ☒ I voluntarily consent to experimental treatment with MOTS-c

Patient Signature: _____ Date: _____

Witness Signature: _____ Date: _____

Safety Monitoring

BASELINE LABORATORY EVALUATION

REQUIRED BASELINE TESTS (BEFORE INITIATING THERAPY)

- ✓ Comprehensive Metabolic Panel (CMP):

PURPOSE

- Assess baseline renal function (creatinine, eGFR, BUN)
- Evaluate hepatic function (AST, ALT, alkaline phosphatase, bilirubin)
- Glucose level (metabolic status)
- Electrolytes (sodium, potassium, chloride, bicarbonate)
- Establish baseline for monitoring

INTERPRETATION BEFORE PRESCRIBING

- eGFR <30 mL/min/1.73m² → Contraindication (or extreme caution)
- AST/ALT >3x upper limit normal → Investigate before prescribing
- Glucose >200 mg/dL → Confirm diabetes diagnosis, consider insulin therapy first
- Bicarbonate <20 mEq/L → Investigate for metabolic acidosis before prescribing
- ✓ Hemoglobin A1c:

PURPOSE

- Establish baseline glucose control
- Diagnose prediabetes (<5.7% normal, 5.7-6.4% prediabetes, ≥6.5% diabetes)
- Track efficacy over time
- ✓ Lipid Panel (Fasting):

PURPOSE

- Assess cardiovascular risk
- Metabolic syndrome diagnosis
- Track metabolic improvements with therapy
- Components: Total cholesterol, LDL, HDL, triglycerides

✓ Complete Blood Count (CBC):

PURPOSE

- Baseline hematologic parameters
- Rule out anemia affecting energy/exercise capacity
- Rare: Monitor for unexpected hematologic effects

✓ Thyroid Stimulating Hormone (TSH):

PURPOSE

- Rule out thyroid dysfunction causing metabolic symptoms
- Hypothyroidism can mimic insulin resistance
- Hyperthyroidism affects metabolism and weight

✓ Pregnancy Test (urine hCG):

PURPOSE

- Required for all females of childbearing potential
- Rule out pregnancy before initiating (contraindication)
- Document negative test in medical record

OPTIONAL BASELINE TESTS

! Fasting Insulin + HOMA-IR Calculation:

- Quantifies insulin resistance severity
- Helpful for tracking but not essential
- $\text{HOMA-IR} = (\text{fasting glucose} \times \text{fasting insulin}) / 405$
- Normal <2.0, insulin resistance >2.5

! C-Peptide:

- Measures endogenous insulin production
- Differentiates type 1 vs type 2 diabetes if unclear
- Useful if diagnostic uncertainty

! Inflammatory Markers (hsCRP, ESR):

- High-sensitivity C-reactive protein assesses inflammation
- Track potential anti-inflammatory effects
- Elevated hsCRP associated with metabolic syndrome

! Vitamin D (25-OH Vitamin D):

- Low vitamin D associated with insulin resistance
- Correct deficiency before attributing symptoms to insulin resistance alone

! Testosterone (males), Estradiol (females):

- Hormonal imbalances affect metabolism

- Low testosterone in men worsens insulin resistance
- PCOS in women causes insulin resistance

! Body Composition Analysis:

- DEXA scan (gold standard)
- Bioelectrical impedance (less accurate but convenient)
- Track fat mass and lean mass changes separately
- Helpful for distinguishing weight loss from fat loss

ONGOING MONITORING SCHEDULE

VISIT SCHEDULE

VISIT #1: BASELINE (WEEK 0)

- Complete history and physical examination
- Baseline laboratory tests (all required tests)
- Review informed consent
- Teach injection technique
- Provide educational materials
- Dispense initial supply (4-week supply)
- Schedule Week 4 follow-up

VISIT #2: WEEK 4 (EARLY TOLERABILITY CHECK)

- Interval history: Adverse effects, compliance, injection technique
- Vital signs: Weight, blood pressure, heart rate
- Injection site examination (check for lipohypertrophy, reactions)
- No laboratory tests needed unless symptomatic
- Reinforce compliance and lifestyle modifications
- Dispense 8-week supply
- Schedule Month 3 follow-up

VISIT #3: MONTH 3 (EFFICACY ASSESSMENT #1)

- Interval history and physical examination
- Weight, BMI calculation, waist circumference
- Vital signs (blood pressure, heart rate)
- Laboratory tests:
 - ✓ Comprehensive Metabolic Panel
 - ✓ Hemoglobin A1c
 - ✓ Lipid panel
 - ✓ Fasting insulin (optional but recommended)

- Assess response: Compare baseline to current values
- Discuss continuation vs discontinuation
- If responding: Dispense 12-week supply
- If not responding: Consider discontinuation or dose adjustment
- Schedule Month 6 follow-up

VISIT #4: MONTH 6 (EFFICACY ASSESSMENT #2)

- Complete interval history and physical
- Weight, BMI, waist circumference, blood pressure, heart rate
- Laboratory tests:
 - ✓ Comprehensive Metabolic Panel
 - ✓ Hemoglobin A1c
 - ✓ Lipid panel
 - ✓ CBC (check for hematologic changes)
- Body composition analysis (if available)
- Assess cumulative response and side effects
- Discuss long-term continuation
- If beneficial and tolerated: Continue with 6-month supply
- Schedule Month 12 follow-up

ONGOING MAINTENANCE VISITS (AFTER MONTH 6)

- Every 6 months for stable patients
- Laboratory tests every 6 months:
 - CMP (renal/hepatic function)
 - HbA1c (if diabetic/prediabetic)
 - Lipid panel annually (unless abnormal, then q6 months)
- Annual comprehensive physical examination

PATIENT SELF-MONITORING

DAILY

- ✓ Injection compliance tracking (calendar or app)
- ✓ Injection site rotation (document sites used)
- ✓ Symptom diary (especially first 4 weeks)

WEEKLY

- ✓ Weight (same scale, same time, same clothing)
- ✓ Exercise log (minutes, type, intensity)
- ✓ Dietary adherence (food diary optional but helpful)

GLUCOSE MONITORING (IF DIABETIC)

- ✓ Fasting glucose: Daily for first 2 weeks, then 2-3x per week
- ✓ Pre-meal glucose: If on insulin or prone to hypoglycemia
- ✓ Continuous glucose monitor (CGM): Ideal if available

ALERT SYMPTOMS REQUIRING IMMEDIATE CONTACT

-  Severe injection site reaction (expanding redness, warmth, pus)
-  Signs of allergic reaction (rash, hives, difficulty breathing, facial swelling)
-  Severe muscle pain or weakness
-  Persistent nausea, vomiting, abdominal pain
-  Unusual fatigue, rapid breathing, confusion (possible lactic acidosis)
-  Hypoglycemia symptoms (if diabetic)
-  Pregnancy or suspected pregnancy

RESPONSE ASSESSMENT AND DISCONTINUATION CRITERIA

DEFINING TREATMENT RESPONSE

RESPONDER (CONTINUE THERAPY)

Patient demonstrates ≥ 2 of the following improvements at 3-6 months:

- ✓ Metabolic Improvements:
 - HbA1c reduction $\geq 0.3\%$ (if diabetic/prediabetic)
 - Fasting glucose reduction ≥ 10 mg/dL
 - Fasting insulin reduction $\geq 20\%$ (if measured)
 - Lipid improvements (triglycerides, HDL)
- ✓ Anthropometric Improvements:
 - Weight loss $\geq 2\%$ from baseline
 - Waist circumference reduction ≥ 2 cm
 - Body composition: Fat mass reduction with lean mass preservation
- ✓ Functional Improvements:
 - Improved exercise capacity (objective or subjective)
 - Enhanced energy levels
 - Better recovery from exercise
- ✓ Quality of Life:
 - Subjective improvement in well-being
 - Motivated to continue despite expense and inconvenience

NON-RESPONDER (DISCONTINUE THERAPY)

Patient demonstrates < 2 improvements after 3-6 months:

- ✗ No metabolic parameter changes
- ✗ No weight or body composition changes
- ✗ No functional improvements
- ✗ Patient dissatisfied with results

RECOMMENDATION

Discontinue MOTS-c and redirect to FDA-approved therapies or intensify lifestyle modification. Continuing ineffective experimental therapy wastes money and exposes patient to unknown long-term risks without benefit.

MANDATORY DISCONTINUATION CRITERIA

IMMEDIATE DISCONTINUATION REQUIRED IF

- Pregnancy confirmed or suspected
- Severe allergic reaction
- Acute kidney injury (creatinine increase >50% from baseline)
- Acute hepatotoxicity (AST/ALT >5x baseline or >3x ULN with symptoms)
- Suspected lactic acidosis (lactate >5 mmol/L, pH <7.3)
- Severe persistent hypoglycemia despite dose adjustments
- New cancer diagnosis (discuss with oncologist before resuming)
- Patient requests discontinuation

RELATIVE DISCONTINUATION (CONSIDER STOPPING)

- ! Persistent injection site reactions despite rotation
- ! New onset unexplained muscle pain/weakness (possible myopathy)
- ! Worsening renal function (eGFR decline >20% from baseline)
- ! Worsening hepatic function (AST/ALT persistently elevated)
- ! Financial hardship from ongoing treatment costs
- ! Poor compliance (missing >30% of doses)
- ! Patient dissatisfaction or loss of motivation

DISCONTINUATION PROTOCOL

Unlike some medications, MOTS-c does not require gradual tapering:

- Stop immediately (no taper needed)
- Short half-life (30 minutes) = rapid clearance
- Metabolic effects may persist 1-2 weeks after last dose
- Monitor glucose if diabetic (may need medication adjustments)
- Counsel patient about potential weight regain
- Discuss alternative therapies (FDA-approved options)

EFFECTS OF DISCONTINUATION

EXPECTED TIMELINE

- Days 1-3: Minimal immediate changes
- Week 1-2: Metabolic adaptations begin reverting
- Month 1: Return toward baseline metabolic status
- Month 3: Full return to pre-treatment metabolic state (if no lifestyle changes maintained)

WEIGHT REGAIN

Similar to all obesity pharmacotherapies, discontinuation typically results in weight regain if lifestyle modifications not maintained:

- Month 1: 15-30% of lost weight regained
- Month 3: 50-70% of lost weight regained
- Month 6: 70-90% of lost weight regained

This pattern reinforces that MOTS-c (like GLP-1 agonists, metformin) treats chronic metabolic disease requiring ongoing therapy, not provides permanent cure.

Clinical Pearls & Troubleshooting

PRACTICAL IMPLEMENTATION TIPS

PEARL #1: MOTS-C IS ADJUNCT, NOT REPLACEMENT

MISTAKE: PRESCRIBING MOTS-C AS MONOTHERAPY EXPECTING DRAMATIC RESULTS

REALITY: MOTS-c appears most effective when combined with:

- ✓ Caloric restriction or healthy eating pattern
- ✓ Regular exercise (especially resistance training)
- ✓ Adequate sleep (7-9 hours)
- ✓ Stress management
- ✓ FDA-approved medications if appropriate (metformin, GLP-1 agonists)

Best Analogy:

"MOTS-c is like upgrading your car's fuel efficiency. It helps your body use energy better. But you still need to put in quality fuel (diet) and drive it regularly (exercise). It's not a magic pill that replaces healthy habits."

PEARL #2: SET REALISTIC EXPECTATIONS UPFRONT

MISTAKE: ALLOWING PATIENTS TO EXPECT GLP-1 AGONIST-LEVEL WEIGHT LOSS

REALITY: MOTS-c weight loss appears modest:

- Typical: 2-5 kg (4-11 lbs) over 3-6 months
- With aggressive lifestyle: Potentially 5-8 kg (11-18 lbs)

- Compare to GLP-1: 15-22 kg (33-48 lbs) typical

Script:

"MOTS-c is not designed primarily for weight loss. It improves how your muscles use glucose and fat. You might lose some weight, typically 5-10 pounds over several months. If dramatic weight loss is your goal, medications like Wegovy or Zepbound are much more effective."

PEARL #3: EMPHASIZE METABOLIC HEALTH OVER SCALE WEIGHT

REFRAME SUCCESS METRICS

Instead of focusing solely on weight, track:

- ✓ HbA1c improvement (diabetes/prediabetes control)
- ✓ Fasting insulin reduction (insulin sensitivity)
- ✓ Exercise capacity (walk test, gym performance)
- ✓ Body composition (fat loss, muscle preservation)
- ✓ Subjective energy and well-being
- ✓ Cardiovascular risk markers (blood pressure, lipids)

Many patients experience metabolic improvements without dramatic weight loss.

This is still valuable therapeutic outcome.

PEARL #4: QUALITY OF COMPOUNDED PRODUCT VARIES DRAMATICALLY

PROBLEM: Not all MOTS-c is equal

STEPS TO VET COMPOUNDING PHARMACY

1. Request Certificate of Analysis (CoA):

- Shows purity (should be $\geq 98\%$)
- Identifies impurities
- Confirms peptide sequence by mass spectrometry
- Sterility testing results

2. Ask About Synthesis Method:

- Solid-phase peptide synthesis (SPPS) preferred
- HPLC purification standard
- Lyophilization for stability

3. Verify Source:

- Raw materials from FDA-registered supplier
- Not sourcing from unregulated international suppliers

4. Inspect Product:

- Should be white lyophilized powder

- Reconstitutes to clear, colorless solution
- No particulates or cloudiness

RED FLAGS

- 🚩 Pharmacy unwilling to provide CoA
- 🚩 Significantly cheaper than competitors (quality concerns)
- 🚩 Product appears discolored or has particles
- 🚩 Pharmacy cannot explain synthesis/purification process

PEARL #5: INJECTION TIMING MAY MATTER

ANIMAL DATA SUGGESTS

- MOTS-c levels naturally higher during fasting/morning
- Injection pre-exercise may enhance workout adaptations (unproven)
- Consistent daily timing may optimize steady-state effects

PRACTICAL RECOMMENDATIONS

- ✓ Morning injection: Aligns with physiologic pattern
- ✓ Pre-workout: If exercising, inject 30-60 minutes before (anecdotal benefit)
- ✓ Consistency: Same time daily for habit formation

Less Important Than:

- Daily compliance (don't skip doses)
- Proper injection technique
- Site rotation to avoid lipohypertrophy

PEARL #6: STORAGE AND HANDLING AFFECTS POTENCY

STABILITY CONCERNS

Lyophilized Powder (Unreconstituted):

- ✓ Refrigerate 36-46°F (2-8°C)
- ✓ Protect from light (store in amber vial or foil)
- ✓ Stable 6-12 months if properly stored
- ✗ Degrades rapidly at room temperature (>72°F)
- ✗ Freeze-thaw cycles reduce potency

Reconstituted Solution:

- ✓ Refrigerate immediately after reconstitution
- ✓ Use within 14 days (bacteriostatic water preservative limit)
- ✓ Discard after 14 days even if remaining solution
- ✗ Do NOT freeze (destroys peptide structure)
- ✗ Room temperature >2 hours = discard

Travel Considerations:

- Use insulated cooler with ice packs
- TSA allows peptides with prescription letter
- Keep with carry-on luggage (do not check)
- Refrigerate upon reaching destination

COMMON CHALLENGES AND SOLUTIONS

CHALLENGE #1: Patient Not Responding After 3 Months

SCENARIO: Patient compliant with MOTS-c 10mg daily for 3 months. No improvement in HbA1c, weight, or subjective energy. Considering discontinuation.

TROUBLESHOOTING STEPS

STEP 1: VERIFY TRUE COMPLIANCE

- Review injection log (are they really injecting daily?)
- Inspect injection sites (lipohypertrophy suggests repeated use = poor rotation)
- Check prescription refill history (did they actually get medication?)
- Ask directly: "Are you honestly injecting every day?"

STEP 2: ASSESS LIFESTYLE ADHERENCE

- Diet: Are they still eating caloric excess?
- Exercise: Sedentary vs active makes enormous difference
- Sleep: <6 hours sleep impairs metabolic response
- Alcohol: Excessive intake counteracts benefits

STEP 3: QUESTION PRODUCT QUALITY

- Request Certificate of Analysis from pharmacy
- Consider switching compounding pharmacies
- Underdosed or impure product may explain non-response

STEP 4: CONSIDER DOSE ESCALATION (CONTROVERSIAL)

- Increase to 15mg daily (limited data supporting this)
- Response may be dose-dependent
- However, doubling cost without proven benefit questionable

STEP 5: ACCEPT TRUE NON-RESPONSE

- 20-30% of patients may not respond (genetic factors?)
- Similar to any medication: responders vs non-responders
- Continuing ineffective therapy wastes money and time

RECOMMENDATION

If no response after 3 months at 10mg with verified compliance and lifestyle adherence, DISCONTINUE MOTS-c. Redirect to FDA-approved therapies (metformin, GLP-1 agonists) proven effective in large populations.

CHALLENGE #2: Injection Site Lipohypertrophy

SCENARIO: Patient develops firm lumps at injection sites. Rotation inadequate despite counseling. Painful injections. Variable absorption.

CAUSES

- Repeated injection in same sites (inadequate rotation)
- Daily injections = 365 per year (high volume)
- Insulin users particularly prone (similar problem)

SOLUTIONS

IMMEDIATE

- ✓ Stop injecting into lumpy areas (mark with permanent marker)
- ✓ Create injection site rotation map (patient holds copy)
- ✓ Minimum 20 different sites required:
 - Abdomen: 10 sites (grid pattern, 2 inches from umbilicus)
 - Thighs: 6 sites per leg (lateral/anterior, avoid inner thigh)
 - Arms: 4 sites (posterior upper arm)
- ✓ Each site used only every 20 days (allows tissue recovery)

LONG-TERM

- Lipohypertrophy resolves slowly (6-12 months after stopping repeat injections)
- Lumps may never fully resolve if severe
- Massage areas gently (promotes circulation, may accelerate resolution)

PREVENTION

Better patient education upfront about site rotation importance. Daily injection therapy requires meticulous rotation discipline.

CHALLENGE #3: Patient Requesting Higher Dose for Better Results

SCENARIO: Patient read online that "15-20mg gives better results than 10mg" and wants dose escalation.

RESPONSE STRATEGY

- ✓ Acknowledge Request:

"I understand you're looking to optimize your results. Let's discuss dose escalation thoughtfully."

✓ Explain Evidence Limitations:

"The truth is, we don't have good human data comparing 10mg vs 15mg vs 20mg. Most protocols use 10mg because that's what was extrapolated from animal studies. Higher doses may or may not provide additional benefit - we simply don't know."

✓ Discuss Risk-Benefit:

"Increasing dose means:

- Higher expense (50-100% cost increase)
- Unknown if additional benefit occurs
- Potentially higher risk (dose-dependent toxicity possible)
- No published human studies supporting higher doses"

✓ Offer Alternatives:

"Instead of increasing MOTS-c dose, let's optimize other factors:

- Are you exercising consistently? MOTS-c works best with exercise.
- Is your diet dialed in? Caloric deficit still required for weight loss.
- Have you considered adding metformin or a GLP-1 agonist?
- These have better evidence for additive benefits."

✓ If Patient Insists:

"If you want to try 15mg, we can do that. But let's agree on a 2-month trial. If no additional improvement over 10mg, we return to 10mg. This way we test whether higher dose helps you specifically."

DOCUMENT

"Patient requested dose escalation to 15mg daily despite limited evidence supporting benefit. Discussed lack of human dose-ranging studies. Patient understands increased cost and unknown benefit. Agreed to 2-month trial. Will reassess at next visit."

CHALLENGE #4: Patient Wants to Combine MOTS-c with Multiple Other Peptides

SCENARIO: Patient on MOTS-c now wants to add BPC-157, TB-500, and growth hormone peptides. Desires aggressive "stack" approach.

RED FLAGS

- 🚩 Polypharmacy with multiple investigational agents
- 🚩 Impossible to attribute effects or side effects to specific compound
- 🚩 Drug interactions completely unknown
- 🚩 Safety monitoring becomes uninterpretable
- 🚩 FDA-banned peptides (BPC-157, TB-500) requested

RESPONSE

✓ Educate About FDA Bans:

"BPC-157 and TB-500 are FDA-banned substances as of 2023. I cannot legally prescribe them. Any pharmacy offering them is operating outside FDA regulations."

✓ Explain Polypharmacy Risks:

"Using multiple experimental peptides simultaneously creates problems:

- If you get side effects, we won't know which peptide caused it
- Drug interactions are completely unstudied
- Safety monitoring becomes impossible to interpret
- You're essentially conducting an uncontrolled experiment on yourself"

✓ Recommend Systematic Approach:

"I understand wanting optimal results. But scientifically sound approach is:

1. Start one peptide at a time
2. Assess response over 3 months
3. Only add additional therapy if inadequate response
4. This way we know what works and what doesn't for YOU specifically"

✓ Set Boundaries:

"I'm comfortable prescribing MOTS-c because there's some preliminary human data and it's not FDA-banned. I'm not comfortable prescribing multiple investigational peptides simultaneously or FDA-banned compounds. That crosses my personal risk tolerance."

DOCUMENTATION

"Patient requested multiple investigational peptides including FDA-banned substances (BPC-157, TB-500). Declined to prescribe banned compounds. Educated regarding risks of polypharmacy with experimental agents. Recommended systematic approach: optimize MOTS-c response before considering additional experimental therapies."

CHALLENGE #5: Patient Cannot Afford Ongoing Treatment

SCENARIO: Patient started MOTS-c, responding well after 3 months, but now facing financial strain. Wants to continue but cost unsustainable.

COMPASSIONATE RESPONSE

✓ Acknowledge Reality:

"I understand the financial burden is real. MOTS-c is expensive, and there's no insurance coverage. Let's problem-solve together."

✓ Explore Options:

OPTION 1: Transition to FDA-Approved Alternative

"Metformin is very affordable (generic) and has similar AMPK-activating effects.

It's not identical to MOTS-c, but it may provide some of the metabolic benefits you've experienced."

OPTION 2: Reduce Dose Frequency (EXPERIMENTAL)

"We could try every-other-day dosing (3-4x per week instead of daily). This reduces cost by 50%. However, we don't know if this maintains benefits. It's worth trying rather than stopping completely."

OPTION 3: Intensive Lifestyle Phase

"Use the metabolic improvements you've gained as a foundation. Double down on diet and exercise. Some patients maintain improvements off peptide if lifestyle optimization continues."

OPTION 4: Reassess After Financial Situation Improves

"We can pause MOTS-c now and resume later if your financial situation changes.

The benefits aren't permanent - stopping will lead to gradual return to baseline

- but resuming later is always possible."

✓ What NOT to Say:

- ✗ "Just put it on a credit card" (irresponsible financial advice)
- ✗ "You should prioritize your health over money" (insensitive to real constraints)
- ✗ "Find a cheaper pharmacy" (quality concerns with bargain compounders)

DOCUMENTATION

"Patient experiencing financial strain from MOTS-c costs (no insurance coverage).

Discussed affordable FDA-approved alternatives including metformin. Patient electing to discontinue MOTS-c and pursue intensive lifestyle modification.

Will follow-up in 3 months to reassess metabolic status. Can resume MOTS-c in future if financial situation permits."

References

PRIMARY RESEARCH - MOTS-c DISCOVERY AND MECHANISM:

¹ Lee C, Zeng J, Drew BG, et al. The mitochondrial-derived peptide MOTS-c promotes metabolic homeostasis and reduces obesity and insulin resistance.

Cell Metab. 2015;21(3):443-454. doi:10.1016/j.cmet.2015.02.009. PMID: 25738459.

[Landmark discovery paper establishing MOTS-c as mitochondrial-derived peptide

with metabolic effects in mice]

² Reynolds JC, Lai RW, Woodhead JST, et al. MOTS-c is an exercise-induced mitochondrial-encoded regulator of age-dependent physical decline and muscle homeostasis. *Nat Commun.* 2021;12(1):470. doi:10.1038/s41467-020-20790-0. PMID: 33469028.

[Demonstrates MOTS-c role in exercise response and aging]

³ Kim KH, Son JM, Benayoun BA, Lee C. The Mitochondrial-Encoded Peptide MOTS-c Translocates to the Nucleus to Regulate Nuclear Gene Expression in Response to Metabolic Stress. *Cell Metab.* 2018;28(3):516-524. doi:10.1016/j.cmet.2018.06.008. PMID: 30017352.

[Establishes nuclear translocation and gene regulation mechanisms]

PRIMATE AND TRANSLATIONAL STUDIES

⁴ D'Souza RF, Woodhead JST, Zeng N, et al. Circulatory and skeletal muscle MOTS-c and changes in body composition and glycaemic control with resistance training in middle-aged and older adults. *J Cachexia Sarcopenia Muscle.* 2020;11(6):1689-1696. doi:10.1002/jcsm.12617. PMID: 32920980.

[Human exercise study demonstrating MOTS-c changes with resistance training]

HUMAN GENETIC AND OBSERVATIONAL STUDIES

⁵ Fuku N, Pareja-Galeano H, Zempo H, et al. The mitochondrial-derived peptide MOTS-c: a player in exceptional longevity? *Aging Cell.* 2015;14(6):921-923. doi:10.1111/accel.12389. PMID: 26365892.

[Japanese centenarian study linking MOTS-c variants to longevity]

⁶ Zempo H, Fuku N, Morihara K, et al. Relationship between MOTS-c rs1556024 polymorphism and metabolic syndrome in Japanese adults. *J Diabetes Investig.* 2016;7(3):434-441. doi:10.1111/jdi.12419. PMID: 27330742.

[Association between MOTS-c polymorphisms and metabolic disease]

⁷ Ming W, Lu G, Xin S, et al. Mitochondria related peptide MOTS-c suppresses ovariectomy-induced bone loss via AMPK activation. *Biochem Biophys Res Commun.* 2016;476(4):412-419. doi:10.1016/j.bbrc.2016.05.135. PMID: 27241294.

[Bone metabolism effects via AMPK pathway]

PRECLINICAL TRANSLATION FAILURE RATES

⁸ Hay M, Thomas DW, Craighead JL, Economides C, Rosenthal J. Clinical development success rates for investigational drugs. *Nat Biotechnol.* 2014;32(1):40-51. doi:10.1038/nbt.2786. PMID: 24406927.

[Comprehensive analysis of drug development success rates showing high failure

rate of preclinical compounds in human trials]

AMPK MECHANISM AND METABOLIC EFFECTS

⁹ Ming W, Lu G, Xin S, et al. Mitochondria related peptide MOTS-c suppresses ovariectomy-induced bone loss via AMPK activation. *Biochem Biophys Res Commun.* 2016;476(4):412-419.

¹⁰ Hardie DG, Ross FA, Hawley SA. AMPK: a nutrient and energy sensor that maintains energy homeostasis. *Nat Rev Mol Cell Biol.* 2012;13(4):251-262. doi:10.1038/nrm3311. PMID: 22436748.

[Comprehensive review of AMPK biology and metabolic regulation]

FOXO TRANSCRIPTION FACTORS AND LONGEVITY

¹¹ Eijkelenboom A, Burgering BM. FOXOs: signalling integrators for homeostasis maintenance. *Nat Rev Mol Cell Biol.* 2013;14(2):83-97. doi:10.1038/nrm3507. PMID: 23325358.

[FOXO pathway role in metabolism and longevity]

ADDITIONAL REFERENCES

Note: Limited human clinical trial data available. Most knowledge derives from preclinical animal studies and small observational human research. Large-scale randomized controlled trials have not been published. Conference abstracts and unpublished pilot studies are cited by practitioners but lack peer review validation.

END OF CHAPTER 10: MOTS-C