

CZECH TECHNICAL UNIVERSITY
IN PRAGUE

Faculty of Civil Engineering
Engineering

Department of Mechanics



Bachelor Thesis

**Computational modelling of thermoplasts and
usin in anchor systems**

Jan Vozáb

Supervisor: Doc. Ing. Jan Vorel Ph.D.

Prague, 2017

ČESKÉ VYSOKÉ UČENÍ TECHNICKÉ
V PRAZE

Fakulta stavebního inženýrství

Katedra mechaniky



Bakalářská práce

**Počítačové modelování reaktoplastů a jejich
využití v kotevních systémech**

Jan Vozáb

Vedoucí práce: Doc. Ing. Jan Vorel Ph.D.

Praha, 2017

Contents

1	Drucker-Prager model of plasticity	5
1.1	Introduction	5
1.2	Drucker-Prager yield criterion	6
1.3	Calculation procedure and implementation	6
1.4	Return to yield of plasticity	8

List of Figures

1.1	Drucker-Prager a Mohr-Coulomb model T	5
-----	---	---

List of Tables

1 Drucker-Prager model of plasticity

1.1 Introduction

Drucker-Prager model of plasticity modified Mohr-Coulomb model. Unlike Mohr-Coulomb model is Drucker-Prager model yield criterion smooth and in space of the principal stresses have form of cylindrical cone. If laboratory results are in effective rather than total stress, criterion of damage become dependent on the hydrostatic and mean stress.




Figure 1.1: Drucker-Prager and Mohr-Coulomb yield criterion in space of principal stresses.

1.2 Drucker-Prager yield criterion

Drucker-Prager model is based on the von Mises model in that the mean stress (first invariant of stress tensor) is obtained in the yield criterion equation, which has the form

$$F(\sigma) = J + (\sigma_m - c)M_{JP}(\varphi) = 0, \quad (1.1)$$

where J is second invariant of stress tensor and M_{JP} is used for approximation to the Mohr-Coulomb model. Is defined as

$$M_{JP} = \frac{\sin(\varphi)}{\cos(\theta) - \frac{\sin(\theta)\sin(\varphi)}{\sqrt{3}}}, \quad (1.2)$$

$$\theta = \arctan \frac{\sin \varphi}{\sqrt{3}}, \quad (1.3)$$

where φ is tangle of internal friction. Equations 1.2 and 1.3 are used to inscribe Drucker-Prager model into Mohr-Coulomb model. Equations (1.2) a (1.3) were taken from [1].

1.3 Calculation procedure and implementation

Total elastic stress can be calculated as

$$\sigma = D\varepsilon_e, \quad (1.4)$$

where D is stiffness matrix and ε_e is elastic deformation. Calculation is selected as implicit, so model is implemented with increments. Then can be equation modified as

$$\sigma^{n+1} = \sigma^n + Dd\varepsilon_e. \quad (1.5)$$

Next step in calculation is find out, if is (1.1) satisfied, shere σ_m a J is first respective second invariant of stress tensor and are defined as

1. DRUCKER-PRAGER MODEL OF PLASTICITY

$$J = \text{sqrt}(\frac{1}{2}\sigma^T P \sigma), \quad (1.6)$$

$$\sigma_m = m^T \sigma. \quad (1.7)$$

If is 1.1 satisfied, calculation continues with next deformation increment. If no, material become to plastic flow. Because of that is defined $d\lambda$, which is coefficient of plastic flow. Dependence of J and σ_m on the $d\lambda$ is taken from [1] and have form

$$F(\sigma) = \overbrace{J - \mu d\lambda}^{J^{n+1}} + \overbrace{(\sigma_m - K M_{JP}(\varphi) d\lambda - c)}^{\sigma_m^{n+1}} M_{JP} = 0. \quad (1.8)$$

Coefficient of plastic flow $d\lambda$ is iterating with Newton-Raphson method until is (1.8) satisfied and until are increments infinitesimal. Popsána je vztahem

$$d\lambda^{n+1} = d\lambda^n + \frac{F^n}{F'^n}. \quad (1.9)$$

Return to the yield of plasticity is going after normal to equation (1.1). Normal vector is counted as [1]

$$n = \frac{\delta F}{\delta \sigma} = \frac{1}{2J} P \sigma + M_{JP} m, \quad (1.10)$$

and its derivation

$$\frac{\delta n}{\delta \sigma} = \left(\frac{3}{2}\right)^{1/2} \frac{\sigma^T P \sigma P - P \sigma \sigma^T}{(\sigma^T P \sigma)^{3/2}}. \quad (1.11)$$

1.4 Return to yield of plasticity

Basic equations are

$$F(\sigma) = \overbrace{J^{res} - \mu d\lambda}^{J^{n+1}} + \overbrace{[\sigma_m^{res} - KM_{PP}(\varphi^{n+1})d\lambda - c^{n+1}cot(\varphi^{n+1})]}^{\sigma_m^{n+1}} M_{JP}(\varphi^{n+1}) = 0. \quad (1.12)$$

$$C_{E_d^{pl}} = c^{n+1} - c^{i-1} - h_c^n(E_d^{pl} - (E_d^{pl})^{i-1}) = 0 \quad (1.13)$$

$$\Theta_{E_d^{pl}} = \varphi^{n+1} - \varphi^{i-1} - h_\varphi^n(E_d^{pl} - (E_d^{pl})^{i-1}) = 0 \quad (1.14)$$

References

- [1] W. Florkowski, *Phenomenology of ultra-relativistic heavy-ion collisions*, London: World Scientific, 2010, ISBN 98-142-8066-6.
- [2] Fázový diagram QCD:
<https://cds.cern.ch/record/1955819/files/TUOAA2f1-eps-converted-to.png>
 [online 24/07/2017].
- [3] S. Sarkar, H. Satz, B. Sinha, *The Physics of the Quark-Gluon Plasma: Introductory Lectures*, Lect. Notes Phys. **785**, Springer, Berlin Heidelberg 2010, ISBN 978-3-642-02286-9.
- [4] Y. Aoki, G. Endrodi, Z. Fodor, S. D. Katz and K. K. Szabo, *The order of the quantum chromodynamics transition predicted by the standard model of particle physics*, Nature **443** (2006) 675.
- [5] Y. Aoki, Z. Fodor, S. D. Katz and K. K. Szabo, *The QCD transition temperature: Results with physical masses in the continuum limit*, Phys. Lett B **643** (2006) 46.
- [6] A. Bazavov et. al., *QCD equation of state to $O(\mu_B^6)$ from lattice QCD*, Phys. Rev. D **95** (2017) 054504.
- [7] Malý třesk: <https://u.osu.edu/vishnu/physics/> [online 11/04/2017].
- [8] Velký třesk: https://www.nikhef.nl/~i93/img/universe_original.jpg [online 11/04/2017].
- [9] Časoprostorový vývoj:
https://www.physics.ohio-state.edu/~ntg/6805/figures/RHIC_timescales_Strickland.png
 [online 11/04/2017].
- [10] P. Alba, W. Alberico, R. Bellwied, M. Bluhm, V. Mantovani Sarti, M. Nahrgang and C. Ratti, *Freeze-out conditions from net-proton and net-charge fluctuations at RHIC*, Phys. Lett. B **738** (2014) 305, arXiv:1403.4903.
- [11] Francois Gelis, *Color Glass Condensate and Glasma*, International Journal of Modern Physics A **28** (2012) 1330001, arXiv:1211.3327v2.
- [12] B. Tomášik, F. Sannino, I. Bearden a T. Dossing, *Topics in modern nuclear physics: Flow and Sizes*, Kobenhavns Universitet, Niels Bohr Institutet, 2004.

- [13] C. M. Ko, V. Koch, Z.-W. Lin, K. Redlich, M. Stephanov, X.-N. Wang, *Kinetic Equation with Exact Charge Conservation*, Phys. Rev. Lett. **86** (2001) 5438.
- [14] S. Jeon, V. Koch, K. Redlich, X.-N. Wang, *Fluctuations of rare particles as a measure of chemical equilibration*, Nucl. Phys. A **697** (2002) 546-562.
- [15] C. M. Ko and L. Xia, *K^+/π^+ enhancement in heavy-ion collisions*, Phys. Rev. C **38** (1988) 179.
- [16] B. Tomášik and E. E. Kolomeitsev, *Strangeness dynamics in heavy-ion collisions: The K/π ratios and the lifetime of a fireball*, arXiv:nucl-th/0512088v1 (2005).
- [17] Koeficienty: <http://www.itl.nist.gov/div898/handbook/eda/section3/eda35b.htm> [online 17/04/2017].
- [18] Modifikované Besselovy funkce: http://people.math.sfu.ca/~cbm/aands/page_376.htm [online 17/04/2017].
- [19] Particle Data Group:
http://pdg.lbl.gov/2016/tables/contents_tables.html [online 17/04/2017].
- [20] "Columbia plot":
<https://quark.phy.bnl.gov/~htding/usqcd/scaling.html> [online 17/04/2017].
- [21] J. D. Bjorken, *Highly relativistic nucleus-nucleus collisions: The central rapidity region*, Phys. Rev. D **27** (1983) 140.
- [22] S. Borsanyi, Z. Fodor, S. D. Katz, S. Krieg, C. Ratti and K. K. Szabo, *Freeze-out Parameters from Electric Charge and Baryon Number Fluctuations: Is there Consistence?*, Phys. Rev. Lett. **113** (2014) 052301.