

# A Probabilistic Proof of the Rogers-Ramanujan Identities

By Jason Fulman

Stanford University

Department of Mathematics

Building 380, MC 2125

Stanford, CA 94305, USA

fulman@math.stanford.edu

<http://math.stanford.edu/~fulman>

Math Reviews Subject Classification: 20P05

## Abstract

The asymptotic probability theory of conjugacy classes of the finite general groups leads to a probability measure on the set of all partitions of natural numbers. A simple method of understanding these measures in terms of Markov chains is given, leading to an elementary probabilistic proof of the Rogers-Ramanujan identities. This is compared with work on the uniform measure. The main case of Bailey's lemma is interpreted as finding eigenvectors of the transition matrix of a Markov chain. It is shown that the viewpoint of Markov chains extends to quivers.

Key words: Rogers-Ramanujan, Markov chain, quiver, conjugacy class

## 1 Introduction

The Rogers-Ramanujan identities [34], [35], [37]

$$1 + \sum_{n=1}^{\infty} \frac{q^{n^2}}{(1-q)(1-q^2) \cdots (1-q^n)} = \prod_{n=1}^{\infty} \frac{1}{(1-q^{5n-1})(1-q^{5n-4})}$$
$$1 + \sum_{n=1}^{\infty} \frac{q^{n(n+1)}}{(1-q)(1-q^2) \cdots (1-q^n)} = \prod_{n=1}^{\infty} \frac{1}{(1-q^{5n-2})(1-q^{5n-3})}$$

are among the most interesting partition identities in number theory and combinatorics. Combinatorial aspects of these identities are discussed by Andrews [2],[5]. Lepowsky and Wilson [28], [29], [30] connect the Rogers-Ramanujan identities with affine Lie algebras and conformal field theory. Feigen and Frenkel [15] interpret them as a character formula for the Virasoro algebra. The author [18] applies the product form to computational group theory, obtaining a simple formula for the  $n \rightarrow \infty$  limit of the probability that an element of  $GL(n, q)$  is semisimple. Stembridge [38] proves them by adapting a method of Macdonald for calculating partial fraction expansions of symmetric formal power series. Garsia and Milne [22] offer a lengthy bijective proof, giving birth to the “involution principle” in combinatorics. Garrett, Ismail, and Stanton [21] usefully recast Roger's original proof using orthogonal polynomials. Relations with statistical mechanics appear in [8], [4], and [40]. The paper [9] gives an account of their appearance in physics.

One of the main results of this note is a first probabilistic proof of the Rogers-Ramanujan identities. In fact the  $i = 1$  and  $i = k$  cases of Theorem 1 will be proved. Theorem 1 is due to Andrews [1]. Bressoud [13] then connected it with an earlier combinatorial result of Gordon [23].

**Theorem 1** ([1]) Let  $x_n$  denote  $(1-x)\cdots(1-x^n)$ . For  $1 \leq i \leq k, k \geq 2$ ,

$$\sum_{n_1, \dots, n_{k-1} \geq 0} \frac{x^{N_1^2 + \dots + N_{k-1}^2 + N_i + \dots + N_{k-1}}}{(x)_{n_1} \cdots (x)_{n_{k-1}}} = \prod_{\substack{r=1 \\ r \neq 0, \pm i \pmod{2k+1}}}^{\infty} \frac{1}{1-x^r}$$

where  $N_j = n_j + \dots + n_{k-1}$ .

There are other elementary proofs of the Rogers-Ramanujan identities available (e.g. [11]), but the one offered here is possibly the most natural and gives insight into Bailey's Lemma.

The basic object of study in this paper is a certain one-parameter family of probability measures  $M_u$  on the set of all partitions of all natural numbers, studied in the prior article [17]. Section 2 recalls these measures and gives their group theoretic motivation. (For now we remark that they arise in the study of conjugacy classes of finite classical groups. As conjugacy classes of compact Lie groups are essentially eigenvalues up to the action of the Weyl group, their probabilistic study can be regarded as philosophically analogous to work of Dyson [14], who described the eigenvalues of random matrices of compact Lie groups in terms of Brownian motion).

Section 3 shows how to construct these measures using a non-upward-moving Markov chain on the integers. The Markov chain is diagonalizable with eigenvalues  $1, \frac{u}{q}, \frac{u^2}{q^4}, \dots$ , and a basis of eigenvectors is given. Analogous computations are done for a measure related to the uniform measure on partitions, which by work of Fristedt [16] has a Markov chain approach. It would be interesting to make a connection with the articles [36], in which a fascinating continuous space Markov chain arises in the asymptotic probability theory of the symmetric group.

With these preliminaries in place, Section 4 gives a proof of the Rogers-Ramanujan identities. The idea of the proof is simple. We compute in two ways the  $L \rightarrow \infty$  probability that the Markov chain started at  $L$  is absorbed at the point 0 after  $k$  steps. (Since the Markov chain is absorbed at 0 with probability 1 and the measure  $M_u$  corresponds to the  $L \rightarrow \infty$  limit, the time to absorption really is the most natural quantity one could consider). The sum side of the Rogers-Ramanujan identities follows from the definition of the probability measures. For the product side, the fact that the transition matrix is explicitly diagonalizable gives a sum expression. One then applies the Jacobi triple product identity (which as explained on page 21 of [2] follows easily from the  $q$ -binomial theorem) and the proof is complete.

Section 4 continues by discussing Bailey's Lemma, which is the only non-trivial step in many of the simplest proofs of the Rogers-Ramanujan identities. The most useful case of Bailey's Lemma follows immediately once a basis of eigenvectors of the transition matrix has been found. The only



$$\begin{aligned}
M_u(\lambda) &= \left[ \prod_{r=1}^{\infty} \left(1 - \frac{u}{q^r}\right) \right] \frac{u^{|\lambda|}}{q^{\sum_i (\lambda'_i)^2} \prod_i \left(\frac{1}{q}\right)_{m_i(\lambda)}} \\
&= \left[ \prod_{r=1}^{\infty} \left(1 - \frac{u}{q^r}\right) \right] \frac{u^{|\lambda|}}{q^{2[\sum_{h < i} h m_h(\lambda) m_i(\lambda) + \frac{1}{2} \sum_i (i-1) m_i(\lambda)^2]} \prod_i |GL(m_i(\lambda), q)|} \\
&= \left[ \prod_{r=1}^{\infty} \left(1 - \frac{u}{q^r}\right) \right] \frac{u^{|\lambda|} P_{\lambda}\left(\frac{1}{q}, \frac{1}{q^2}, \dots; 0, \frac{1}{q}\right)}{q^{n(\lambda)}}
\end{aligned}$$

The only fact from this section which is required for the proof of the Rogers-Ramanujan identities is the fact that  $M_u$  defines a probability measure for  $0 < u < 1$ . This can be seen using an identity from either [31] or else following Stong [39], who uses the fact that there are  $q^{n^2-n}$  unipotent elements in  $GL(n, q)$ . The first proof of Theorem 2 will use the fact that  $M_u$  is a probability measure without further comment. As there has been interest in simplifying the proofs of the Rogers-Ramanujan identities as much as possible, a second completely elementary proof of Theorem 2 will be given. From this second proof it will follow that  $M_u(\lambda)$  is a probability measure.

Although not logically necessary for this paper, we mention that for  $0 < u < 1$  and  $q$  a prime power, the measures  $M_u$  have a group theoretic description, referring the reader to the survey [20] for further discussion. Recall that the conjugacy classes of  $GL(n, q)$  are parameterized by rational canonical form. Each such matrix corresponds to the following combinatorial data. To every monic non-constant irreducible polynomial  $\phi$  over  $F_q$ , associate a partition (perhaps the trivial partition)  $\lambda_{\phi}$  of some non-negative integer  $|\lambda_{\phi}|$ . The only restrictions necessary for this data to represent a conjugacy class are that  $|\lambda_z| = 0$  and  $\sum_{\phi} |\lambda_{\phi}| \deg(\phi) = n$ . To be explicit, a representative of the conjugacy class corresponding to the data  $\lambda_{\phi}$  may be given as follows. Define the companion matrix  $C(\phi)$  of a polynomial  $\phi(z) = z^{\deg(\phi)} + \alpha_{\deg(\phi)-1} z^{\deg(\phi)-1} + \dots + \alpha_1 z + \alpha_0$  to be:

$$\begin{pmatrix}
0 & 1 & 0 & \cdots & 0 \\
0 & 0 & 1 & \cdots & 0 \\
\cdots & \cdots & \cdots & \cdots & \cdots \\
0 & 0 & 0 & \cdots & 1 \\
-\alpha_0 & -\alpha_1 & \cdots & \cdots & -\alpha_{\deg(\phi)-1}
\end{pmatrix}$$

Let  $\phi_1, \dots, \phi_k$  be the polynomials such that  $|\lambda_{\phi_i}| > 0$ . Denote the parts of  $\lambda_{\phi_i}$  by  $\lambda_{\phi_i,1} \geq \lambda_{\phi_i,2} \geq \dots$ . Then a matrix corresponding to the above conjugacy class data is:

$$\begin{pmatrix} C(\phi_i^{\lambda_{\phi_i,1}}) & 0 & 0 \\ 0 & C(\phi_i^{\lambda_{\phi_i,2}}) & 0 \\ 0 & 0 & \dots \end{pmatrix}$$

Now consider the following procedure for putting a measure on the set of all partitions of all natural numbers. Fix  $u$  such that  $0 < u < 1$ . Pick a non-negative integer such that the chance of choosing  $n$  is equal to  $(1-u)u^n$ . Then pick  $\alpha$  uniformly in  $GL(n, q)$  and take  $\lambda$  to be the partition corresponding to the polynomial  $z - 1$  in the rational canonical form of  $\alpha$ . If  $n = 0$  take  $\lambda$  to be the trivial partition. The random partition so defined obeys  $M_u$  measure. (The polynomial  $z - 1$  is considered without loss of generality. Partitions corresponding to other irreducible polynomials are probabilistically independent, and one just replaces  $q$  by raising it to the degree of the polynomial). In the limit  $u \rightarrow 1$ , one is simply studying random elements in a fixed  $GL(n, q)$  with  $n \rightarrow \infty$ . The substitutions  $u \rightarrow -u$  and  $q \rightarrow -q$  correspond to the finite unitary groups. The idea of auxilliary randomization of the dimension  $n$  is analogous to the idea of canonical ensembles in statistical mechanics.

### 3 Markov chains

#### 3.1 Group theoretical measures

The first result of this paper describes the measure  $M_u$  in terms of Markov chains. Two proofs will be given. The first proof is given in the interest of clarity and assumes that  $M_u$  is a probability measure. The second proof is more elementary.

It is convenient to set  $\lambda'_0$  (the height of an imaginary zeroth column) equal to  $\infty$ . For the entirety of this subsection,  $(x)_n$  will denote  $(1-x)(1-x/q)\cdots(1-x/q^{n-1})$ . Thus  $(x)_0 = 1$  and  $(x)_n = 0$  for  $n < 0$ . For convenience of notation, let  $P(a)$  be the  $M_u$  probability that  $\lambda_1 = a$ .  $Prob(E)$  will denote the probability of an event  $E$  under the measure  $M_u$ .

**Theorem 2** *Starting with  $\lambda'_0 = \infty$ , define in succession  $\lambda'_1, \lambda'_2, \dots$  according to the rule that if  $\lambda'_i = a$ , then  $\lambda'_{i+1} = b$  with probability*

$$K(a, b) = \frac{u^b (\frac{1}{q})_a (\frac{u}{q})_a}{q^{b^2} (\frac{1}{q})_{a-b} (\frac{1}{q})_b (\frac{u}{q})_b}.$$

*Then the resulting partition is distributed according to  $M_u$ .*

PROOF: The  $M_u$  probability of choosing a partition with  $\lambda'_i = r_i$  for all  $i$  is

$$Prob.(\lambda'_0 = \infty) \frac{Prob.(\lambda'_0 = \infty, \lambda'_1 = r_1)}{Prob.(\lambda'_0 = \infty)} \prod_{i=1}^{\infty} \frac{Prob.(\lambda'_0 = \infty, \lambda'_1 = r_1, \dots, \lambda'_{i+1} = r_{i+1})}{Prob.(\lambda'_0 = \infty, \lambda'_1 = r_1, \dots, \lambda'_i = r_i)}.$$

Thus it is enough to prove that

$$\frac{Prob.(\lambda'_0 = \infty, \lambda'_1 = r_1, \dots, \lambda'_{i-1} = r_{i-1}, \lambda'_i = a, \lambda'_{i+1} = b)}{Prob.(\lambda'_0 = \infty, \lambda'_1 = r_1, \dots, \lambda'_{i-1} = r_{i-1}, \lambda'_i = a)} = \frac{u^b(\frac{1}{q})_a(\frac{u}{q})_a}{q^{b^2}(\frac{1}{q})_{a-b}(\frac{1}{q})_b(\frac{u}{q})_b},$$

for all  $i, a, b, r_1, \dots, r_{i-1} \geq 0$ .

For the case  $i = 0$ , the equation

$$P(a) = \frac{u^a(\frac{u}{q})_{\infty}}{q^{a^2}(\frac{1}{q})_a(\frac{u}{q})_a}$$

is given a probabilistic proof in [17]. For an elementary proof of this identity, see the second proof of this theorem. For  $i > 0$  one calculates that

$$\sum_{\substack{\lambda: \lambda'_1=r_1, \dots, \lambda'_{i-1}=r_{i-1} \\ \lambda'_i=a}} M_u(\lambda) = \frac{u^{r_1+\dots+r_{i-1}}}{q^{r_1^2+\dots+r_{i-1}^2}(\frac{1}{q})_{r_1-r_2} \dots (\frac{1}{q})_{r_{i-2}-r_{i-1}}(\frac{1}{q})_{r_{i-1}-a}} P(a).$$

Similarly, observe that

$$\sum_{\substack{\lambda: \lambda'_1=r_1, \dots, \lambda'_{i-1}=r_{i-1} \\ \lambda'_i=a, \lambda'_{i+1}=b}} M_u(\lambda) = \frac{u^{r_1+\dots+r_{i-1}+a}}{q^{r_1^2+\dots+r_{i-1}^2+a^2}(\frac{1}{q})_{r_1-r_2} \dots (\frac{1}{q})_{r_{i-2}-r_{i-1}}(\frac{1}{q})_{r_{i-1}-a}(\frac{1}{q})_{a-b}} P(b).$$

Thus the ratio of these two expressions is

$$\frac{u^b(\frac{1}{q})_a(\frac{u}{q})_a}{q^{b^2}(\frac{1}{q})_{a-b}(\frac{1}{q})_b(\frac{u}{q})_b},$$

as desired. Note that the transition probabilities automatically sum to 1 because

$$\sum_{\substack{\lambda: \lambda'_1=r_1, \dots, \lambda'_{i-1}=r_{i-1} \\ \lambda'_i=a, \lambda'_{i+1}=b}} M_u(\lambda) \sum_{\substack{b \leq a \\ \lambda: \lambda'_1=r_1, \dots, \lambda'_{i-1}=r_{i-1} \\ \lambda'_i=a}} M_u(\lambda) = 1.$$

□

PROOF: (Second Proof) This proof needs only that  $M_u$  is a measure; it will emerge that  $M_u$  is a probability measure. For this proof  $P(a)$  denotes the  $M_u$  mass that  $\lambda'_1 = a$ .

One calculates that

$$\sum_{\substack{\lambda: \lambda'_1=r_1, \dots, \lambda'_{i-1}=r_{i-1} \\ \lambda'_i=a}} M_u(\lambda) = \frac{u^{r_1+\dots+r_{i-1}}}{q^{r_1^2+\dots+r_{i-1}^2}(\frac{1}{q})_{r_1-r_2} \dots (\frac{1}{q})_{r_{i-2}-r_{i-1}}(\frac{1}{q})_{r_{i-1}-a}} P(a).$$

Similarly, observe that

$$\sum_{\substack{\lambda: \lambda'_1=r_1, \dots, \lambda'_{i-1}=r_{i-1} \\ \lambda'_i=a, \lambda'_{i+1}=b}} M_u(\lambda) = \frac{u^{r_1+\dots+r_{i-1}+a}}{q^{r_1^2+\dots+r_{i-1}^2+a^2} \left(\frac{1}{q}\right)_{r_1-r_2} \cdots \left(\frac{1}{q}\right)_{r_{i-2}-r_{i-1}} \left(\frac{1}{q}\right)_{r_{i-1}-a} \left(\frac{1}{q}\right)_{a-b}} P(b).$$

Thus the ratio of these two expressions is

$$\frac{P(b)u^a}{P(a)q^{a^2} \left(\frac{1}{q}\right)_{a-b}}.$$

Since  $M_u$  is a measure, it follows that

$$\sum_{b \leq a} \frac{P(b)u^a}{P(a)q^{a^2} \left(\frac{1}{q}\right)_{a-b}} = 1.$$

From this recursion and the fact that  $P(0) = \left(\frac{u}{q}\right)_\infty$ , one solves for  $P(a)$  inductively, finding that

$$P(a) = \frac{u^a \left(\frac{u}{q}\right)_\infty}{q^{a^2} \left(\frac{1}{q}\right)_a \left(\frac{u}{q}\right)_a}.$$

Identity 2.2.8 on page 20 of [2] now implies that  $\sum_a P(a) = 1$ , so that  $M_u$  is a probability measure.

□

The algorithm of Theorem 2 is nice in that it can be implemented on a computer. An analogous Markov chain approach to the conjugacy classes of the finite symplectic and orthogonal groups has recently been worked out in [19].

Theorem 3 explicitly diagonalizes the transition matrix  $K$ , which is fundamental for understanding the Markov chain. Note that if the current distribution of the Markov chain is given by a row vector, the distribution at the next step is obtained by multiplying the row vector on the right by  $K$ .

**Theorem 3** 1. Let  $C$  be the diagonal matrix with  $(i, i)$  entry  $\left(\frac{1}{q}\right)_i \left(\frac{u}{q}\right)_i$ . Let  $M$  be the matrix  $\left(\frac{u^j}{q^{j^2} \left(\frac{1}{q}\right)_{i-j}}\right)$ . Then  $K = CMC^{-1}$ , which reduces the problem of diagonalizing  $K$  to that of diagonalizing  $M$ .

2. Let  $A$  be the matrix  $\left(\frac{1}{\left(\frac{1}{q}\right)_{i-j} \left(\frac{u}{q}\right)_{i+j}}\right)$ . Then the columns of  $A$  are eigenvectors of  $M$  for right multiplication, the  $j$ th column having eigenvalue  $\frac{u^j}{q^{j^2}}$ .

3. The inverse matrix  $A^{-1}$  is  $\left(\frac{(1-u/q^{2i})(-1)^{i-j} \left(\frac{u}{q}\right)_{i+j-1}}{q \binom{i-j}{2} \left(\frac{1}{q}\right)_{i-j}}\right)$ .



PROOF: The first part is obvious. The second part is a special case of Lemma 1 of [11]. The third part is a lemma of [3].  $\square$

The point of the proof of Theorem 3 is that once one knows (either from Mathematica or by implementing algorithms from linear algebra) what the eigenvectors are, it is a simple matter to verify the computation.

The following corollary will be useful for the proof of the Rogers-Ramanujan identities.

**Corollary 1** *Let  $E$  be the diagonal matrix with  $(i, i)$  entry  $\frac{u^i}{q^{i^2}}$ . Then  $K^r = CAE^r A^{-1}C^{-1}$ . More explicitly,*

$$K^r(L, j) = \frac{(\frac{1}{q})_L (\frac{u}{q})_L}{(\frac{1}{q})_j (\frac{u}{q})_j} \sum_{n=0}^{\infty} \frac{u^{rn} (1 - u/q^{2n}) (-1)^{n-j} (\frac{u}{q})_{n+j-1}}{q^{rn^2} (\frac{1}{q})_{L-n} (\frac{u}{q})_{L+n} q^{\binom{n-j}{2}} (\frac{1}{q})_{n-j}}.$$

PROOF: This is immediate from Theorem 3.  $\square$

### 3.2 Mixture of uniform measures

For this subsection  $q < 1$ . The measure assigns probability  $q^{|\lambda|} \prod_{i=1}^{\infty} (1 - q^i)$  to the partition  $\lambda$ . Conditioning this measure to live on partitions of a given size gives the uniform measure, an observation exploited by Fristedt [16]. As is clear from [33], this measure is very natural from the viewpoint of representation theory.

Fristedt (loc. cit.) proved that this measure has a Markov chain description. His chain affects row lengths rather than column lengths (though the algorithm would work on columns too as the measure is invariant under transposing diagrams). Nevertheless, we adhere to his notation. We use the notation that  $(x)_n = (1 - x) \cdots (1 - x^n)$ .

**Theorem 4** [16] *Starting with  $\lambda_0 = \infty$ , define in succession  $\lambda_1, \lambda_2, \dots$  according to the rule that if  $\lambda_i = a$ , then  $\lambda_{i+1} = b$  with probability*

$$K(a, b) = \frac{q^b(q)_a}{(q)_b}.$$

*Then the resulting partition is distributed according to the measure of this subsection.*

Theorem 5 diagonalizes this Markov chain, giving a basis of eigenvectors. The proof is analogous to that of Theorem 3, the second part being proved by induction.

**Theorem 5** 1. Let  $C$  be the diagonal matrix with  $(i, i)$  entry equal to  $\frac{(q)_i}{q^i}$ . Let  $M$  be the matrix with  $(i, j)$  entry  $q^i$  if  $i \geq j$  and 0 otherwise. Then  $K = CMC^{-1}$ .

2. Let  $D$  be the diagonal matrix with  $(i, i)$  entry equal to  $q^i$ . Let  $A$  be the matrix  $\left( \frac{(-1)^{i-j}}{q^{\binom{i-j}{2}} (\frac{1}{q})_{i-j}} \right)$ , so that its inverse is  $\left( \frac{1}{(\frac{1}{q})_{i-j}} \right)$  by part b of Theorem 3. Then the eigenvectors of  $M$  are the columns of  $A$ , the  $j$ th column having eigenvalue  $q^j$ .

As a corollary, one obtains a simple expression for the chance that under the measure of this subsection, the  $r$ th row has size  $j$ .

**Corollary 2**

$$K^r(L, j) = \frac{q^j q^{L(r-1)} (q)_L (\frac{1}{q})_{L-j+r-1}}{(q)_j (\frac{1}{q})_{L-j} (\frac{1}{q})_{r-1}}.$$

Letting  $L \mapsto \infty$ , the chance that the  $r$ th row has size  $j$  becomes

$$\frac{(q)_\infty q^{rj}}{(q)_j (q)_{r-1}}.$$

PROOF: To obtain the first expression, one multiplies out  $K^r = CAD^r A^{-1} C^{-1}$  and uses the  $q$ -binomial theorem

$$\sum_{m=0}^{\infty} y^m q^{(m^2+m)/2} \frac{(q)_n}{(q)_m (q)_{m-n}} = (1+yq)(1+yq^2) \cdots (1+yq^n).$$

□

The second part of the corollary can be proved directly without recourse to Markov chain theory; one simply attaches to an  $r * j$  square two partitions: one with at most  $r-1$  rows and another with at most  $j$  columns.

## 4 Rogers-Ramanujan Identities and Bailey's Lemma

The first result of this section proves the following identity of Andrews, which contains the Rogers-Ramanujan identities. In this section  $(x)_n$  denotes  $(1-x)(1-x/q) \cdots (1-x/q^{n-1})$ .

**Theorem 6** [1] For  $k \geq 2$ ,

$$\sum_{n_1, \dots, n_{k-1} \geq 0} \frac{1}{q^{N_1^2 + \dots + N_{k-1}^2} (1/q)_{n_1} \cdots (1/q)_{n_{k-1}}} = \prod_{\substack{r=1 \\ r \neq 0, \pm k \pmod{2k+1}}}^{\infty} \frac{1}{1 - (1/q)^r}$$

$$\sum_{n_1, \dots, n_{k-1} \geq 0} \frac{1}{q^{N_1^2 + \dots + N_{k-1}^2 + N_1 + \dots + N_{k-1}} (1/q)_{n_1} \dots (1/q)_{n_{k-1}}} = \prod_{\substack{r=1 \\ r \neq 0, \pm 1 \pmod{2k+1}}}^{\infty} \frac{1}{1 - (1/q)^r}$$

where  $N_j = n_j + \dots + n_{k-1}$ .

PROOF: For the first identity, we compute in two ways

$$\sum_{\lambda: \lambda' < k} M_1(\lambda).$$

One obtains the sum side by using the first definition of  $M_1$  in Section 2. For the product side, let  $u = 1$ ,  $j = 0$ ,  $r = k$ , and  $L \rightarrow \infty$  in Corollary 1. The rest is now a standard argument.

$$\begin{aligned} 1 + \sum_{n=1}^{\infty} \frac{(1 + 1/q^n)(-1)^n}{q^{n^2} q^{\binom{n}{2}}} &= 1 + \sum_{n=1}^{\infty} (-1)^n \left(\frac{1}{q}\right)^{(k+1/2)n^2 - n/2} + \left(\frac{1}{q}\right)^{(k+1/2)n^2 + n/2} \\ &= \sum_{n=-\infty}^{\infty} (-1)^n \left(\frac{1}{q}\right)^{(k+1/2)n^2} \left(\frac{1}{q}\right)^{n/2}. \end{aligned}$$

Now invoke Jacobi's triple product identity (e.g. [2])

$$\sum_{n=-\infty}^{\infty} (-1)^n v^n w^{n^2} = \prod_{n=1}^{\infty} (1 - vw^{2n-1})(1 - w^{2n-1}/v)(1 - w^{2n}).$$

The proof of the second identity is the same except that one takes  $u = \frac{1}{q}$  instead of  $u = 1$ .  $\square$

Next we discuss the most important case of Bailey's Lemma, which was alluded to in [7] and stated explicitly in [3]. A pair of sequences  $\{\alpha_L\}$  and  $\{\beta_L\}$  are called a Bailey pair if

$$\beta_L = \sum_{r=0}^L \frac{\alpha_r}{(1/q)_{L-r} (u/q)_{L+r}}.$$

Bailey's Lemma states that if  $\alpha'_L = \frac{u^L}{q^{L^2}} \alpha_L$  and  $\beta'_L = \sum_{r=0}^L \frac{u^r}{q^{r^2} (1/q)_{L-r}} \beta_r$ , then  $\{\alpha'_L\}$  and  $\{\beta'_L\}$  are a Bailey pair.

From our viewpoint, this case of Bailey's Lemma is clear. Namely let  $A, D, M$  be as in Theorem 3 (recall that  $M = ADA^{-1}$ ). Viewing  $\alpha = \vec{\alpha}_L$  and  $\beta = \vec{\beta}_L$  as column vectors, the notion of a Bailey pair means that  $\beta = A\alpha$ . This case of Bailey's Lemma follows because

$$\beta' = M\beta = ADA^{-1}\beta = AD\alpha = \alpha'.$$

It would be interesting to obtain all of Bailey's Lemma (and Theorem 1) by probabilistic arguments, and also to understand the Bailey lattice probabilistically. The reader may enjoy the survey by Bressoud [12].

## 5 Quivers

This section uses the notion of a quiver, as surveyed in Kac [27], to which the reader is referred for more detail. The basic set-up is as follows. Let  $\Gamma$  be a connected graph with  $n$  vertices labelled as  $1, \dots, n$  (where we allow loops). Let  $N, Z$  denote the natural numbers and integers respectively. Let  $f_{ij}$  be the number of edges between  $i, j$ . Associated to  $\Gamma$  is a natural bilinear form on  $Z^n$  and a root system  $\Delta \subset Z^n$ . Choose an arbitrary orientation of  $\Gamma$  so that  $\Gamma$  is a quiver. For a given dimension  $\alpha \in N^n - \{0\}$ , let  $A_\Gamma(\alpha, q)$  be the number of classes of absolutely indecomposable representations of  $\Gamma$  over the algebraic closure of a field of  $q$  elements. It is proved in [27] that  $A_\Gamma(\alpha, q)$  is a polynomial in  $q$  with integer coefficients, and that this polynomial is independent of the orientation of the graph. Kac (loc. cit.) formulated many conjectures about this polynomial. One such conjecture, which is still open, is that the constant term in  $A_\Gamma(\alpha, q)$  is the multiplicity of  $\alpha$  in the root system.

In recent work, Hua [24, 25] has given a completely combinatorial reformulation of this conjecture. To explain, let  $(1/q)_n$  denote  $(1 - \frac{1}{q}) \cdots (1 - \frac{1}{q^n})$ , and for any two partitions  $\lambda, \mu$  define  $\langle \lambda, \mu \rangle = \sum_{i \geq 1} \lambda'_i \mu'_i$ . Let  $t_j^\alpha$  be the coefficient of  $q^j$  in the polynomial  $A_\Gamma(\alpha, q)$ . Let  $b_\lambda = \prod_{i \geq 1} (\frac{1}{q})_{m_i(\lambda)}$ , where  $m_i$  is the number of parts of  $\lambda$  of size  $i$ . Let  $\lambda(1), \dots, \lambda(n)$  be an  $n$ -tuple of partitions. Set  $U_\alpha = U_1^{\alpha_1} \cdots U_n^{\alpha_n}$ . Hua's result, which reduces Kac's conjecture to a combinatorial assertion, is that

$$\sum_{\lambda(1), \dots, \lambda(n)} \frac{\prod_{1 \leq i \leq j \leq n} q^{f_{ij} \langle \lambda(i), \lambda(j) \rangle} U_1^{|\lambda(1)|} \cdots U_n^{|\lambda(n)|}}{\prod_{1 \leq i \leq n} q^{\langle \lambda(i), \lambda(i) \rangle} b_{\lambda(i)}} = \prod_{\alpha \in \Delta^+} \prod_{j=0}^{\deg(A_{\Gamma(\alpha, q)})} \prod_{i=1}^{\infty} \left( \frac{1}{1 - U_\alpha q^{j-i}} \right)^{t_j(\alpha)}.$$

A few points are in order. First, the right hand side of this equation is different from the statements in [24, 25], due to a minor slip there. Second, observe that the expression converges in the ring of formal power series in the variables  $U_1, \dots, U_n$ . This leads one to define a “probability” measure on  $n$ -tuples of partitions  $M_{\Gamma, \vec{U}}$  by assigning mass

$$\prod_{\alpha \in \Delta^+} \prod_{j=0}^{\deg(A_{\Gamma(\alpha, q)})} \prod_{i=1}^{\infty} (1 - U_\alpha q^{j-i})^{t_j(\alpha)} \frac{\prod_{1 \leq i \leq j \leq n} q^{f_{ij} \langle \lambda(i), \lambda(j) \rangle}}{\prod_{1 \leq i \leq n} q^{\langle \lambda(i), \lambda(i) \rangle} b_{\lambda(i)}}$$

to the  $n$ -tuple  $\lambda(1), \dots, \lambda(n)$ . For quivers of finite type this is a true probability measure for values of  $U_1, \dots, U_n$  sufficiently small, but in general we abuse notation by using terms from probability theory when  $U_1, \dots, U_n$  are variables.

Note that when the graph consists of a single point, this measure is simply the measure  $M_u$  from Section 2. Theorem 7 shows that the structure of a Markov chain is still present. As the idea

of the proof is the same as the second step of Theorem 2, the algebra is omitted. Let  $P(a_1, \dots, a_n)$  denote the  $M_{\Gamma, \vec{U}}$  probability that  $\lambda(1), \dots, \lambda(n)$  have  $a_1, \dots, a_n$  parts respectively.

**Theorem 7** *Let  $\lambda(1)'_1, \dots, \lambda(n)'_1$  be distributed as  $P(a_1, \dots, a_n)$ . Define  $(\lambda(1)'_2, \dots, \lambda(n)'_2)$  then  $(\lambda(1)'_3, \dots, \lambda(n)'_3)$ , etc. successively according to the rule that if  $(\lambda(1)'_i, \dots, \lambda(n)'_i)$  is equal to  $(a_1, \dots, a_n)$ , then  $(\lambda(1)'_{i+1}, \dots, \lambda(n)'_{i+1})$  is equal to  $(b_1, \dots, b_n)$  with probability*

$$K(\vec{a}, \vec{b}) = \prod_{1 \leq i \leq j \leq n} q^{f_{ij} a_i a_j} \prod_{i=1}^n \frac{U_i^{a_i}}{q^{a_i^2 (\frac{1}{q})_{a_i - b_i}}} \frac{P(b_1, \dots, b_n)}{P(a_1, \dots, a_n)}.$$

The resulting  $n$ -tuple of partitions is distributed according to  $M_{\Gamma, \vec{U}}$ .

As a final remark, observe that letting  $C$  be diagonal with entries  $\frac{1}{P(a_1, \dots, a_n)}$ , one obtains a factorization  $K = CMC^{-1}$ , where  $M$  is defined by

$$M(\vec{a}, \vec{b}) = \prod_{1 \leq i \leq j \leq n} q^{f_{ij} a_i a_j} \prod_{i=1}^n \frac{U_i^{a_i}}{q^{a_i^2 (\frac{1}{q})_{a_i - b_i}}}.$$

A very natural problem is to investigate the eigenvector matrix  $E$  of  $M$ , and  $E^{-1}$ , in order to obtain new Bailey Lemmas.

## 6 Acknowledgments

This research was supported by an NSF Postdoctoral Fellowship. The author thanks Persi Diaconis for discussions and a referee for correcting historical mistakes in the bibliography.

## References

- [1] G. Andrews, ‘An analytic generalization of the Rogers-Ramanujan identities’, *Proc. Nat. Acad. Sci.* 71 (1974), 4082-4085.
- [2] G. Andrews, *The theory of partitions. Encyclopedia of Mathematics and its Applications, Vol. 2* (Addison-Wesley, Reading, Mass.-London-Amsterdam, 1976).
- [3] G. Andrews, ‘Multiple series Rogers-Ramanujan type identities’, *Pacific J. Math.* 114 (1984), 267-283.
- [4] G. Andrews, R. Baxter, and P. Forrester, ‘Eight vertex SOS model and generalized Rogers-Ramanujan-type identities’, *J. Statist. Phys.* 35 (1984), 193-266.

- [5] G. Andrews, *q-series: their development and application in analysis, number theory, combinatorics, physics, and computer algebra*. (American Math Society, Providence, 1986).
- [6] G. Andrews, A. Schilling, and P. Forrester, ‘An  $A_2$  Bailey lemma and the Rogers Ramanujan identities’, *J. Amer. Math. Soc.* 12 (1999), 677-702.
- [7] W. Bailey, ‘Identities of the Rogers-Ramanujan type’, *Proc. London Math. Soc.* (2) 50 (1949), 1-10.
- [8] R. Baxter, *Exactly solved models in statistical mechanics*, (Academic Press, London-New York, 1982).
- [9] A. Berkovich and B. McCoy, ‘Rogers-Ramanujan identities: a century of progress from mathematics to physics’. ICM Proceedings, Volume 3 (1998).
- [10] S. Bloch and A. Okounkov, ‘The character of the infinite wedge representation’, preprint math.CO/9712009 at xxx.lanl.gov.
- [11] D. Bressoud, ‘An easy proof of the Rogers-Ramanujan identities’, *J. Number Theory* 16 (1983), 235-241.
- [12] D. Bressoud, The Bailey lattice: An introduction, in *Ramanujan Revisited*, 57-67, Academic Press, 1988.
- [13] D. Bressoud, ‘Analytic and combinatorial generalizations of the Rogers-Ramanujan identities’, *Mem. Amer. Math. Soc.* 24 (1980), No. 227.
- [14] F. Dyson, ‘A Brownian motion model for the eigenvalues of a random matrix’, *J. Mathematical Phys.* 3 (1962), 1191-1198.
- [15] B. Feigin and E. Frenkel, ‘Coinvariants of nilpotent subalgebras of the Virasoro algebra and partition identities’. I. M. Gelfand Seminar, *Adv. Soviet Math.* 16, Part 1, 139-148, Amer. Math. Soc., Providence, RI, 1993.
- [16] B. Fristedt, ‘The structure of random partitions of large integers’, *Trans. Amer. Math. Soc.* 337 (1993), 703-735.
- [17] J. Fulman, ‘A probabilistic approach to conjugacy classes in the finite general linear and unitary groups’, *J. Algebra* 212 (1999), 557-590.

- [18] J. Fulman, ‘The Rogers-Ramanujan identities, the finite general linear groups, and the Hall-Littlewood polynomials’, *Proc. Amer. Math. Soc.* 128 (2000), 17-25.
- [19] J. Fulman, ‘A probabilistic approach to conjugacy classes in the finite symplectic and orthogonal groups’, to appear in *J. Algebra*.
- [20] J. Fulman, ‘Random matrix theory over finite fields: a survey’, preprint math.GR/0003195 at xxx.lanl.gov.
- [21] K. Garrett, M. Ismail, M., and D. Stanton, ‘Variants of the Rogers-Ramanujan identities’, *Adv. in Appl. Math.* 23 (1999), 274-299..
- [22] A. Garsia and S. Milne, ‘A Rogers-Ramanujan bijection’, *J. Combin. Theory Ser. A* 31 (1981), 289-339.
- [23] B. Gordon, A combinatorial generalization of the Rogers-Ramanujan identities, *Amer. J. Math.* 83 (1961), 393-99.
- [24] J. Hua, ‘Representations of quivers over finite fields’, University of New South Wales Ph.D. Thesis, 1998.
- [25] J. Hua, ‘Representations of quivers over finite fields’, *Bull. Aust. Math. Soc.* 59 (1999), 171-2.
- [26] N. Jing, Vertex operators and Hall-Littlewood symmetric functions, *Advances in Math.* 87 (1991), 226-248.
- [27] V. Kac, Root systems, representations of quivers, and invariant theory. Lecture Notes In Mathematics 996 (Springer, Berlin,1982), 74-108.
- [28] J. Lepowsky and R.L. Wilson, ‘A new family of algebras underlying the Rogers-Ramanujan identities and generalizations’, *Proc. Nat. Acad. Sci. USA* 78 (1981), 7254-7258.
- [29] J. Lepowsky and R.L. Wilson, ‘The structure of standard modules, I: Universal algebras and the Rogers-Ramanujan identities’, *Invent. Math.* 77 (1984), 199-290.
- [30] J. Lepowsky and R.L. Wilson, ‘The structure of standard modules, II: The case  $A_1^{(1)}$  principal gradation’, *Invent. Math.* 79 (1985), 417-442.

- [31] I.G. Macdonald, *Symmetric functions and Hall polynomials*. Second Edition. (Clarendon Press, Oxford, 1995).
- [32] S. Milne and G. Lilly, ‘The  $A_l$  and  $C_l$  Bailey transform and lemma’, *Bull. Amer. Math. Soc.* 26 (1992), 258-263.
- [33] A. Okounkov, ‘Infinite wedge and measures on partitions’, preprint math.RT/9907127 at xxx.lanl.gov.
- [34] L. Rogers, ‘Second memoir on the expansion of certain infinite products’, *Proc. London Math. Soc.* 25 (1894), 318-343.
- [35] S. Ramanujan and L. Rogers, ‘Proof of certain identities in combinatory analysis’, *Proc. Cambridge Phil. Soc.* 19 (1919), 211-216.
- [36] A. Schmidt and A. Vershik, ‘Limit measures arising in the asymptotic theory of the symmetric group’, *Th. of Prob. and its Appl* 22 (1978) 72-88, 23 (1979) 42-54.
- [37] I.J. Schur, ‘Ein beitrage zur additiven zahlentheorie und zur theorie der kettenbruche’, S.B. Preuss, Akad. Wiss. Phys.-Math. KL (1917), 302-321.
- [38] J. Stembridge, ‘Hall-Littlewood functions, plane partitions, and the Rogers-Ramanujan identities’, *Trans. Amer. Math. Soc.* 319 (1990), 469-498.
- [39] R. Stong, ‘Some asymptotic results on finite vector spaces’, *Adv. in Appl. Math.* 9 (1988), 167-199.
- [40] O. Warnaar, The Andrews-Gordon identities and q-multinomial coefficients, *Commun. Math. Phys.* 184 (1997), 203-232.

Current address: Stanford University Math Department, Building 380, MC 2125, Stanford CA 94305, USA.

Current email: fulman@math.stanford.edu