



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

PHYSICA C

Physica C 388–389 (2003) 687–688

[www.elsevier.com/locate/physc](http://www.elsevier.com/locate/physc)

# Microwave absorption spectrum and reentrant phase in Bi2212 single crystal: microwave power dependence

G. Ahmad <sup>a</sup>, A. Hashizume <sup>a</sup>, S. Iwasaki <sup>a</sup>, K. Yoshii <sup>a</sup>, B.J. Reddy <sup>a</sup>,  
M. Shahabuddin <sup>b</sup>, S. Uthayakumar <sup>c</sup>, R. Jayavel <sup>c</sup>, T. Endo <sup>a,\*</sup>

<sup>a</sup> Faculty of Engineering, Mie University, Tsu, Mie 514-8507, Japan

<sup>b</sup> Department of Physics, University of Jamia Millia Islamia, Jamia Nagar, New Delhi 110025, India

<sup>c</sup> Crystal Growth Center, Anna University, Chennai 600025, India

## Abstract

Microwave power ( $P_m$ ) dependence of microwave absorption (MA) was investigated on  $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_x$  crystals at liquid nitrogen temperature. At low  $P_m$  of 0.1 mW, MA spectrum shows only a sharp first peak near zero magnetic fields ( $H_a$ ), which corresponds to Meissner phase, for  $H_a \parallel c$ -axis. MA spectrum shows a dip and a broad second peak when  $P_m$  is increased. These dip and broad peak reflect reentrant liquid phase and solid phase. The broad peak shifts to lower fields with increasing  $P_m$ . The real sample temperature is raised by 4 K with increasing  $P_m$  to 50 mW.

© 2003 Elsevier Science B.V. All rights reserved.

PACS: 74.60.Ge; 74.25.Nf; 74.22.Hs

Keywords: Bi2212 single crystal; Reentrant phase; Microwave absorption; Microwave power; Temperature rise

## 1. Introduction

Demanding for tunable high temperature superconducting microwave filters is urgent to save spaces occupied by transmission base stations. The stacked superconducting/ferromagnetic thin films are employed as the tunable microwave filters. They are operated under magnetic fields [1–5]. Then it is important to study the microwave loss properties of superconducting materials such as  $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_x$  (Bi2212) under the fields. Such a research under high microwave power is also necessary for high power components like antennas [6].

In this work, we investigated the microwave power dependence of microwave absorption (MA) in Bi2212 crystals. Combined with temperature dependence of the absorption spectrum in a solid state region of a reentrant phase [7], rises of sample temperature by the MA are estimated as a function of microwave power.

## 2. Experimental

Superconducting Bi2212 single crystals ( $T_C = 86.5$  K) were grown by self-flux method [7]. Non-resonant MA measurement was carried out on the samples employing “cavity perturbation method” [8–11]. The sample in a cryostat was put in dc magnetic field ( $H_a$ ) and a modulation field was superimposed on the dc field with an amplitude of 5 G at 100 kHz. The MA signals ( $S$ ) were measured on the samples at around liquid nitrogen temperatures ( $T$ ) as a function of  $H_a$ . The microwave power ( $P_m$ ) was varied in a range of 0.1–50 mW. The dc field  $H_a$  was applied on the sample along its  $c$ -axis ( $H_a \parallel c$ ) [12].

## 3. Results and discussion

The MA spectra of the sample measured at various  $P_m$  at low  $T$  are shown in Fig. 1. At very low  $P_m$  of 0.1 mW, the MA spectrum shows only a sharp first peak at near zero field below 50 G. When  $P_m$  is increased to 10 mW, MA spectrum evolves to show the same first peak,

\* Corresponding author. Tel.: +81-59-231-9400; fax: +81-59-231-9471.

E-mail addresses: [endo@elec.mie-u.ac.jp](mailto:endo@elec.mie-u.ac.jp), [endo@cm.elec.mie-u.ac.jp](mailto:endo@cm.elec.mie-u.ac.jp) (T. Endo).




Fig. 1. MA spectra various  $P_m$ .

followed by a dip and another new broad peak at 100–350 G. Henceforth, we call these peaks “first peak ( $P_1$ )” and “broad peak ( $P_b$ )”. Beyond this power up to 50 mW, MA spectrum is always composed of three characteristic parts; the first peak, the dip and the broad peak. However, the field positions of these structures seem to be shifted with increasing  $P_m$ .

The field position of the maximum of broad peak ( $H_{pb}$ ) is plotted in Fig. 2 as a function of  $P_m$ .  $H_{pb}$  shifts to lower fields with increasing  $P_m$ . The temperature dependence of  $H_{pb}$  ( $H_{pb}-T$ ) has been already obtained by the previous work as shown in inset of Fig. 2 [7].  $H_{pb}$  also shifts to lower fields with increasing  $T$ . From the same tendencies of  $H_{pb}-P_m$  obtained in this work and  $H_{pb}-T$ , it can be concluded that the real sample temperature  $T_{\text{samp}}$  must be raised with increasing  $P_m$  due to the larger MA, although the ambient temperature is almost the same. From these two data, we can obtain a new relation  $T_{\text{samp}}-P_m$  which is plotted in Fig. 2. The result indicates that  $T_{\text{samp}}$  rises by about 4 K when  $P_m$  is increased from 0.1 to 50 mW. This temperature rise is a considerable amount which must be taken into account when the superconducting materials are used for the high power microwave devices.

We need to explain why  $H_{pb}$  is shifted to the lower fields with increasing  $T$ . According to  $T$ -dependence of MA spectrum reported by us [7], the three structures of MA should reflect the reentrant phase diagram [13,14] as




Fig. 2.  $H_{pb}$  and estimated sample temperature  $T_{\text{samp}}$  as a function of  $P_m$ . The inset shows the previous result of  $H_{pb}-T$  [7].




Fig. 3. Model reentrant phase diagram proposed by Fisher and Lee [13], and Nelson [14].  $H_{C1}$ : lower critical field,  $H_{C2}$ : upper critical field,  $H_m$ : melting line.

shown in Fig. 3. The first peak and the broad peak reflect Meissner phase and vortex solid phase, respectively. While the dip reflects the reentrant liquid phase ( $L_1$ ) lying just above Meissner phase. Because of “nose shape” of the solid phase, the fields of  $H_a$  passing in the solid state are shifted to the lower side with increasing  $T$  as  $T_1 \rightarrow T_2 \rightarrow T_3$  as shown in Fig. 3. This is the reason why the broad peak and  $H_{pb}$  are shifted to the lower side.

#### 4. Conclusion

We investigated  $P_m$ -dependence of MA on Bi2212 at liquid nitrogen temperature. MA spectrum shows only the first peak at the low  $P_m$  while it shows the first peak, the dip and the broad peak at the higher  $P_m$ . These are reflecting Meissner phase,  $L_1$  phase and solid phase in the reentrant phase diagram.  $H_{pb}$  shifts to the lower fields with increasing  $P_m$ . With the combination of  $H_{pb}-P_m$  with  $H_{pb}-T$ , the real sample temperature is raised by 4 K with increasing  $P_m$  due to the larger MA.

#### References

- [1] M. Hein, in: J. Kuhn et al. (Eds.), High-Temperature-Superconductor Thin Film at Microwave frequencies, Springer, New York, 1999.
- [2] Z.Y. Shen et al., IEEE Trans. Appl. Supercond. 7 (1997) 2446.
- [3] A. Lauder et al., Adv. Mater. 10 (1998) 1249.
- [4] S. Hontsu et al., Supercond. Sci. Technol. 12 (1999) 836.
- [5] J. Wosik et al., Appl. Phys. Lett. 74 (1999) 750.
- [6] D.W. Face et al., Physica C 357–360 (2001) 1488.
- [7] R.D. Kale et al., Physica C 378–381 (2002) 470.
- [8] T. Endo, H. Yan, in: A. Narlikar (Ed.), Studies of High Temperature Superconductors, vol. 14, Nova Science, New York, 1995, pp. 65–106.
- [9] T. Endo, H. Yan, Jpn. J. Appl. Phys. 33 (1994) 103.
- [10] V.V. Srinivasu et al., Singapore J. Phys. 15 (1999) 9.
- [11] V.V. Srinivasu et al., in: Proc. Crystal Growth Methods and Processes, Chennai, 2002, p. 228.
- [12] K.I. Itoh et al., Physica C 357–360 (2001) 477.
- [13] M.P.A. Fisher, D.H. Lee, Phys. Rev. B 39 (1989) 2756.
- [14] D.R. Nelson, Phys. Rev. Lett. 60 (1988) 1973.