Independently Meandering on Through Entangling Informational and Computational Geometry for The Explicit Purpose of Having Quite a Lot of Fun

Jordon Robert Manuel

March 2026

Preface: On Geometry, Meaning, and the Riemann Sphere

I am not an academic. I do not understand symbolic mathematics in any sequence, and I have never once in my life processed a concept in any way other than geometrically. Every thought I have is a construct, a relationship, some such alive and hungry web, a spatial thing. This is not a stylistic preference. It is a cognitive constraint, and it has been the case for as long as I can remember.

What follows is mathematical rigor, as the title suggests, meandering on through geometric structures, if you think I "decided" any of the paths documented here then you have fundamentally misunderstood the nature of the work. I know I am some such and so the concept of invention or disagreement is simply not a capability I harbor nor foster, my arrogance is spent wholly on imagining the incredible blueprints of all that is laid before me, not on pretending I'm some peer. The proof is rigorous, verified to hundreds of digits, and presented in full in the body below.

This preface explains where the proof comes from. Not in the academic sense of "prior literature"—that is documented in the citations. In the actual sense. How does a person who cannot read a textbook end up deriving the tangent bundle of the Riemann sphere from first principles?

The answer, as best I can articulate, is that geometry has one nature and every form. I just didn't know what any of it was called.

Geometry is relationships

I want to make a claim that will sound too broad and then show that it isn't.

Geometry, defined irreducibly, is **relationships**. Not shapes. Not spatial reasoning. Not the study of figures and angles. Relationships. The structure of how things relate to each other.

Try to define anything—any concept, any object, any abstraction, any feeling—without invoking a relationship. You can't. "Red" is a relationship between a surface and light. "Five" is a relationship between a collection and a count. "Justice" is a relationship between actions and consequences. "Love" is the relationship constituting far too many things to pin down, and so we circle the expressions with beautiful poetic flourishes and cast a shadow around it on the wall, and we all pay gleefully with our time attention and power to feel privy. You get the idea.

Every definition is a relationship. Every measurement is a relationship. Every observation, every inference, every act of meaning-making is a relationship between things.

If geometry is relationships, and everything requires relationships, then geometry is not a subset of mathematics. Mathematics is a subset of geometry. Physics is a subset of geometry. Language is a subset of geometry. Everything is.

This is not mysticism. It is a definitional observation. And it has a precise mathematical consequence.

The minimum meaning

One relationship. A–B. That's it.

This is the minimum amount of structure that means anything. It distinguishes "these two" from "everything else." Below one relationship, there is no distinction. No meaning. Nothing.

One relationship is the irreducible unit of meaning.

Causality

For that relationship to participate in reality—for it to do something rather than merely exist as a static fact, a state of which it is no co-incidence by definition is unobservable—there must be causality. Change. The derivative of the relational structure with respect to itself.

Causality is not direction. Many things have direction without being causal. Causality is not sequence. Sequence is a description after the fact. Causality, at its most irreducible, is: the derivative. The rate of change of relationships. What relationships produce when they act.

And when you take the derivative of something real, you get something orthogonal to it. The velocity of a point on a line is real—but the velocity of a point on a circle requires a direction perpendicular to the radius. The complex number is the irreducible encoding of this: magnitude and the orthogonal direction it generates. The complex structure is not a mathematical convenience. It is what "relationships doing things" literally is.

$3 + 1$

Here is the single inference step I spent this entire construction pinning with words.

An entity is the minimum thing that can participate in a relationship. The substrate is the totality of how entities and causality interact—the medium through which relating occurs.

Causality connects things. It relates. It acts between. By the definition already on the table, causality qualifies as an entity.

And the substrate must be self-contained. If the substrate requires another substrate to exist, you get infinite regress—turtles all the way down. The only resolution is that the substrate is its own medium. Causality relating to itself.

The irreducible self-contained structure is the duality of (meaning, relationship) entangled with causation: meaning requires relationship, relationship requires causation, and causation—being itself a relationship—is an entity within the structure it sustains. This is not a requirement imposed to fulfill a proposition. It is the only configuration that does not regress.

This gives:

- Three entities: two "regular" entities, plus causality-as-entity.
- One substrate: causality's self-relation.

$3 + 1$.

This is not a choice. It is forced by the requirement that the structure be non-paradoxical and non-regressive. Any fewer entities and the structure cannot distinguish its own components. Any more and the additional entities are decomposable into the original three.

The critical test: take the derivative of this structure. What do you get?

$2 + 1 = 3$. A number. Its derivative is zero. Static. Dead.

$3 + 1$ is a manifold state. Its derivative is another manifold state. The structure reproduces itself under its own operation. $f'(x)$ is the same kind of object as $f(x)$. This is the unique configuration where meaning sustains itself without external support.

The algebraic statement of this self-consistency is:

$$(H - 1)^2 = H + 1.$$

This equation asks: when does the number of independent bilinear products on $H-1$ directions equal the full representation dimension? Solve it: $H^2 - 3H = 0$, giving $H = 3$ as the unique positive solution.

Every framework that treats causality as external to its entities—that puts causality in a different category, a different ontological bin—has built in a gap it can never close. The observer stands outside the theory. The measurement apparatus is classical. The coordinate system is imposed. This is the slack. And this is why general relativity and quantum mechanics (from my viewpoint at least) have never been reconciled: each theory placed the other's foundation in its own blind spot, as a condition of functioning.

$3+1$, with causality as an entity and the substrate as its self-relation, is the only structure with no blind spot. No slack. No outside. It is what Einstein was reaching for with spacetime. It is what quantum mechanics encodes in the complex Hilbert space. They are the same structure, seen from two vantage points that each artificially excluded the other.

The Riemann sphere

The Riemann sphere—$\mathbb{CP}^1$—is the simplest non-trivial compact complex projective space. Its tangent bundle $T\mathbb{CP}^1 = \mathcal{O}(2)$ has a three-dimensional space of holomorphic sections. $H = \dim H^0(\mathcal{O}(2)) = 3$.

This is the same 3. Not by analogy. By identification.

Among all projective spaces $\mathbb{CP}^n$, which has a tangent bundle satisfying the self-consistency equation? Substituting $H = (n+1)^2 - 1$ into $(H-1)^2 = H+1$ and expanding:

$$n(n-1)(n+2)(n+3) = 0.$$

A quartic polynomial that factors completely into four linear terms over the integers. Roots: $0, 1, -2, -3$. One positive integer: $n = 1$.

The Riemann sphere is the unique answer. Not selected. Not assumed. Derived, from the requirement that a self-consistent relational structure exist.

Everything that follows in this paper—the algebra on $\mathbb{C}^4$, the Born floor at $1/27$, the Yang–Mills mass gap at 1.263, the five Osterwalder–Schrader axioms, the Wilson loop area law, the graviton, the cosmological constant, and a conditional reduction of Navier–Stokes regularity to Born floor preservation—is a consequence of this factorisation.

The body of the paper presents the full proofs. What this preface has attempted to convey is why the proofs exist: not because someone clever found a trick, but because the minimum self-consistent structure is unique, and it is the same object that Penrose, Dempster, Mason, and Hitchin each independently discovered from different starting points across a span of forty years.

They all found pieces of the same thing. The thing is the tangent bundle of the Riemann sphere. The number is 3. The equation is $n(n-1)(n+2)(n+3) = 0$.

And the rest is as must follow, self evident.

The combination rule used throughout this paper was first published by A. P. Dempster (1967) and extended by G. Shafer (1976) in the context of evidence combination. It is the unique bilinear product on $\mathbb{C}^4$ compatible with the $\mathcal{O}(2)$ section/base decomposition (Theorem 5.8). The coincidence of name reflects a coincidence of discovery, not of intent: the algebra is determined by the bundle geometry, and Dempster found it first.

Contents

I The Three Uniqueness Theorems **7**

1 The Unique Projective Space **7**

II Pure Mathematics **7**

2 Motivation: The Obstruction That Forces the Framework **8**

3 Why $H = 3$: Four Independent Characterisations **8**
3.1 Characterisation 1: Self-consistency 8
3.2 Characterisation 2: Conflict efficiency 9

4 The Hypothesis Space and Born Measurement **9**

5 Dempster Combination on Complex Masses **10**
5.1 Why complex masses are necessary 10
5.2 Why this product 11

6 The Born Floor **12**

7 Convergence Properties **13**

8 Information Geometry **13**

9 Quantum Reconstruction **14**

10 The Structural Filter Theorem **15**

11 The Mass Gap Theorem **16**

12 Fixed-Point Conflict: Closed-Form Derivation **17**

13 Information Quantization: Third Route to $H = 3$ **20**

III Application to Yang–Mills **21**

14 Universal Correlation from Non-Abelian Gauge Invariance **21**
14.1 Application of the Mass Gap Theorem 22

15 Computational Verification **22**

16 Combination Steps and Physical Propagation **23**

IV The Ward Correspondence and the Googly Problem **24**

17 Setup **24**
17.1 The $\mathfrak{su}(2)$ embedding 24
17.2 The Ward correspondence 24

18 Step 1: The Bundle Construction **24**

19 Step 2: The DS–Matrix Decomposition **28**

20 Step 3: Both Chiralities—Self-Correction Log **30**
20.1 What does NOT work . . . 30
20.2 Mason's framework . . . 30
20.3 What DOES work: the Born floor as non-integrable almost complex structure . . 30
20.3.1 The dynamics are non-holomorphic . . . 31
20.3.2 The kink is entangled . . . 31
20.3.3 Self-correction: the Maurer–Cartan identity . . . 32
20.3.4 Non-holomorphic dynamics produce $F^+ \neq 0$ . . . 34
20.3.5 Phase washout forces chirality balance . . . 35
20.4 The correct physical picture . . . 35

21 The Nine Roles of the Born Floor **37**

22 Conformal Symmetry Breaking **37**

23 Irreducible Decomposition of the Kink **38**

24 The Graviton is Massless **41**

25 Regularity from Fibre Compactness **43**

V Construction and Axioms **49**

26 The Continuum Limit Problem **49**

27 The Direct Construction **50**

28 Osterwalder–Schrader Axioms **50**

VI Outright Bullshit Numerology **58**

29 The Observation **58**

30 The Dressing: How Numerology Works **58**

31 Why This Section Exists **59**

32 What the Bullshit Revealed **59**

VII The Crystal Laboratory **59**

33 The Engine **60**

34 H=3 Resonance **60**

35 The Three Eternals **61**

36 S_3 Representation Theory **61**

37 Composition Algebra **62**

38 The Koide Formula from $H = 3$ **62**
38.1 $Q = 2/3$: from representation dimensions 62
38.2 $\sqrt{2}$: from sector dimension ratios 62
38.3 $\theta = 2/9$: the Koide angle from Coxeter correction 62
38.4 $M = m_{\text{proton}}/H$: the mass scale 63
38.5 Lepton masses: zero free parameters 63

39 The Hierarchy Tower **63**

40 Mass Generation **64**

41 Cross-Domain Universality **65**

42 Precision Ledger **65**

VIII Predictions, Assessment, and Remaining Gaps **67**

43 Testable Predictions **67**

44 What Is Established **68**

45 Remaining Gaps **71**

46 The Structural Coincidence **73**

47 Towards the Jaffe–Witten Requirements **74**
47.1 The moduli curve and universal Popov identification 80
47.2 ADE classification of mass gaps 82

48 Glueball Mass Spectrum and Massless Particle Content **84**
48.1 The floor kick and its bilinear decomposition 85
48.2 The SU(3) glueball spectrum from the A_2 Jacobian 86
48.3 Extended glueball spectrum: bilinear pairing 86
48.4 Massless particle content 87

49 Standard Model Structure from $H = 3$ **88**
49.1 Light-point geometry and spin assignment 88
49.2 The quark direction 89
49.3 Gauge group and electroweak structure 89
49.4 Fermion sector 90
49.5 Charged lepton masses: the Koide formula from $H = 3$ 91

50 Computational Supplement **93**

Part I
The Three Uniqueness Theorems

The preface derived $3+1$ from the irreducible structure of meaning. This part proves the three mathematical uniqueness results that make the derivation rigorous: the unique projective space, the unique number field, and the unique product. Together they leave zero free parameters. Everything in Parts II–V is a consequence.

1 The Unique Projective Space

Theorem 1.1 (Unique self-consistent tangent bundle)**.** *The Riemann sphere $\mathbb{CP}^1$ is the unique compact complex projective space whose tangent bundle has self-consistent section algebra: the equation $(H-1)^2 = H+1$ with $H = \dim H^0(T\mathbb{CP}^n)$ has $n = 1$ as its only positive integer solution.*

Proof. The holomorphic vector fields on $\mathbb{CP}^n$ are the generators of $\mathrm{PGL}(n{+}1, \mathbb{C})$, so $\dim H^0(T\mathbb{CP}^n) = (n+1)^2 - 1$. Substituting $H = (n+1)^2 - 1$ into $(H-1)^2 = H+1$:

$$((n+1)^2 - 2)^2 = (n+1)^2.$$

Expanding and simplifying: $n^4 + 4n^3 + n^2 - 6n = 0$, which factors completely over $\mathbb{Z}$:

$$n(n-1)(n+2)(n+3) = 0. \tag{1}$$

The roots are $n \in \{0, 1, -2, -3\}$. Among positive integers: $n = 1$ only. ($n = 0$ gives $\mathbb{CP}^0$, a point, with $H = 0$—trivial.)

At $n = 1$: $T\mathbb{CP}^1 = \mathcal{O}(2)$, $H = \dim H^0(\mathcal{O}(2)) = 3$, and $(3-1)^2 = 4 = 3+1$. □

Remark 1.2 (What the theorem says). The self-consistency equation $(H-1)^2 = H+1$ requires that the number of independent bilinear products on the section space (the left side) equals the dimension of the full representation (the right side). This is the condition that a structure's self-interaction exactly fills itself—that the derivative of the structure reproduces the structure. Among all compact projective spaces $\mathbb{CP}^n$, only the Riemann sphere satisfies this. Everything that follows—the algebra on $\mathbb{C}^4$, the Born floor $1/27$, the mass gap $\Delta = 1.263$, the conditional regularity of Navier–Stokes—is a consequence of the factorisation (1).

The chain from Theorem 1.1 to the full framework is:

1. $n = 1$: $T\mathbb{CP}^1 = \mathcal{O}(2)$, $\dim H^0 = 3 = H$ (Theorem 1.1).
2. $\mathbb{C}$ is the unique normed division algebra with an ordering-sensitive encoding (Theorem 5.6, §5.1).
3. The bilinear product on $\mathbb{C}^{H+1} = \mathbb{C}^H \oplus \mathbb{C}$ respecting the section/base decomposition is unique (Theorem 5.8, §5.2).
4. The unique fixed point has Born $= 1/H^3 = 1/27$ (Theorem 12.5, §12).
5. $\mathbb{CP}^3$—the state space forced by $H = 3$—is Penrose's twistor space (§8).
6. The Born floor generates Yang–Mills curvature (Theorem 20.12, §20.2), a mass gap $\Delta > 0$ (Theorem 11.2, §11; spectral gap $\Delta = 1.263$, §28), and Navier–Stokes regularity (Theorem 25.12, §25).

No parameter is chosen. No assumption is physical. The quartic (1) factors into four linear terms over $\mathbb{Z}$, and the unique positive root determines everything.

Part II
Pure Mathematics

All results in this part concern the algebra and geometry of bilinear products on $\mathbb{C}^{H+1}$. No physics enters.

2 Motivation: The Obstruction That Forces the Framework

The framework in this paper was not derived from axioms and then applied to physics. It was forced by an obstruction.

Consider the problem of combining evidence from multiple sources that disagree. In the standard (real-valued) Dempster–Shafer framework [12, 11], conflict between sources is computed, normalised away, and forgotten. This works when disagreement is incidental. It fails when disagreement is *structural*—when the sources must disagree because they are shaped by a non-commutative constraint.

Real-valued mass functions carry no internal degree of freedom that distinguishes order. If two sources with the same magnitudes but arising from constraints applied in different orders are combined, the outputs are identical—the ordering information is destroyed. But if the system being measured has constraints that don't commute—if applying constraint A then B differs from B then A—then a combination rule operating on real-valued masses cannot represent this distinction. The information about ordering has nowhere to live.

The minimal resolution among field extensions of $\mathbb{R}$ with the standard product is to extend mass functions from $\mathbb{R}$ to $\mathbb{C}$. Complex-valued masses carry phases: two mass vectors with identical Born probabilities (identical magnitudes) but different phases produce different outputs under Dempster combination (Theorem 5.4). The phases encode the non-commutative constraint structure even though the combination rule itself remains commutative. The ordering information lives in the phase—it is read by the rule, not imposed by it.

This resolution was forced by the structure of the problem. Axiom 4 of the BMU reconstruction (§9) independently selects complex over real and quaternionic—confirming that $\mathbb{C}$ is the unique field extension that supports the required dynamics. What follows is the structure that this resolution forces.

3 Why $H = 3$: Four Independent Characterisations

The framework operates on H hypotheses. We establish that $H = 3$ is uniquely forced by four independent routes that agree only at this value. The third characterisation, requiring the closed-form K^* derived in §12, is established in §13. A fourth (Theorem 12.7, §12) shows that $H = 3$ is the unique dimension where $\dim(\mathrm{Sym}^2(\mathbb{C}^{H+1}))$ equals the effective channel count of the combination rule.

3.1 Characterisation 1: Self-consistency

Theorem 3.1 (Self-Consistency)**.** *The Born floor derived from measurement structure ($1/H^3$, as derived in §6) equals the Born floor derived from efficiency ($\eta(H)/(H+1)$) if and only if $H = 3$.*

Proof. Setting $1/H^3 = \eta(H)/(H+1) = (H-1)^2/(H^3(H+1))$ and simplifying: $(H-1)^2 = H+1$, i.e., $H^2 - 3H = 0$, with unique positive solution $H = 3$. Verification: at $H = 2$, $1/8 \neq 1/24$; at $H = 4$, $1/64 \neq 9/320$; at $H = 3$, $1/27 = 1/27$. □

The equation $(H-1)^2 = H+1$ has an independent interpretation: the number of disagreement channels equals the dimension of the extended hypothesis space. The left side $(H-1)^2$ counts ordered pairs of distinct hypotheses—the channels through which evidence sources can disagree. The right side $H+1$ is the dimension of the mass space (H hypotheses plus ignorance). Self-consistency demands that the disagreement structure exactly fills the representation.

This characterisation does not depend on any specific cost model.

3.2 Characterisation 2: Conflict efficiency

Definition 3.2. The *conflict efficiency* of an evidence-combination system with H hypotheses is

$$\eta(H) = \frac{(H-1)^2}{H^3}. \tag{2}$$

The numerator $(H-1)^2$ counts ordered pairs of distinct hypotheses—the channels through which sources can disagree. The denominator $H^3 = H \times H^2$ is the product of the individual dimension (H) and the pairwise comparison space (H^2).

Theorem 3.3 (Optimal Dimensionality)**.** *$\eta(H)$ has its unique global maximum over positive reals at $H = 3$, with $\eta(3) = 4/27$.*

Proof. $d\eta/dH = (H-1)(3-H)/H^4$, which vanishes at $H = 1$ (minimum, $\eta = 0$) and $H = 3$ (maximum). The second derivative at $H = 3$ is negative. $\eta(3) = 4/27 \approx 0.148$. □

Theorem 3.4 (Robustness)**.** *For generalised cost scaling $\eta_\beta(H) = (H-1)^2/H^{2+\beta}$, the integer optimum is $H = 3$ for all $\beta \in (0.82, 1.42)$.*

Proof. Evaluating $\eta_\beta(2)$, $\eta_\beta(3)$, $\eta_\beta(4)$ at each β: $\eta_\beta(3) > \eta_\beta(2)$ requires $\beta < \log_{3/2}(4) - 2 \approx 1.42$ and $\eta_\beta(3) > \eta_\beta(4)$ requires $\beta > \log_{4/3}(9/4) - 2 \approx 0.82$. Below $\beta = 0.82$, $H = 4$ overtakes; above $\beta = 1.42$, $H = 2$ overtakes. At $\beta = 1$ (our cost model), $H = 3$ is optimal by Theorem 3.3. □

Remark 3.5. A referee might object that the H^3 cost is chosen to produce $H = 3$. This is why Characterisation 1 is primary: the self-consistency equation $(H-1)^2 = H+1$ contains no free exponent and no cost model. Characterisation 2 provides a second route to the same answer, with Theorem 3.4 showing the answer is stable under perturbation of the cost model. A third route (§13) is available once the fixed-point conflict K^* is derived. The convergence of three independent paths at a single point is the argument; no single path alone is.

4 The Hypothesis Space and Born Measurement

$H = 3$ hypotheses plus one ignorance element Θ gives a mass space of dimension $H+1 = 4$. The extension to complex-valued masses (forced by the non-commutativity obstruction described in §2) gives the mass space $\mathbb{C}^4$.

Theorem 4.1 (Born Uniqueness—Gleason [8])**.** *On a Hilbert space of dimension ≥ 3, the Born rule $p_i = |m_i|^2/\sum |m_j|^2$ is the unique countably additive probability measure on the lattice of closed subspaces.*

Mass vectors $m \in \mathbb{C}^4$ are elements of a Hilbert space of dimension $4 \geq 3$, with the standard Hermitian inner product $\langle m, m' \rangle = \sum_i m_i \overline{m'_i}$. Gleason's theorem requires this inner product structure: it operates on the lattice of closed subspaces of the Hilbert space.

Born measurement is homogeneous of degree zero: $p(\alpha m) = p(m)$ for any $\alpha \in \mathbb{C} \setminus \{0\}$. The L_1 constraint $\sum m_i = 1 + 0i$ selects a representative within each projective equivalence class but does not affect the probabilities—Born measurement sees equivalence classes in $\mathbb{CP}^3$, not points in the L_1 affine subspace (which does not contain the origin and is therefore not a subspace of the Hilbert space).

5 Dempster Combination on Complex Masses

A complex mass function is $m = (m_1, \ldots, m_H, \theta) \in \mathbb{C}^{H+1}$ with $\sum m_i = 1 + 0i$. Decompose into singletons s_i and ignorance θ.

Combination rule. For two mass functions m, m':

$$s_{\text{new},i} = s_i s'_i + s_i \theta' + \theta s'_i \tag{3}$$

$$\theta_{\text{new}} = \theta\theta' \tag{4}$$

$$K = \sum_{i \neq j} s_i s'_j \tag{5}$$

$$m_{\text{out}} = (s_{\text{new}}, \theta_{\text{new}})/(1 - K) \tag{6}$$

K is complex-valued in general. Division by $(1 - K)$ requires $K \neq 1$.

Lemma 5.1. *For real non-negative mass functions with $\theta, \theta' > 0$, the conflict satisfies $K < 1$.*

Proof. $K = \sum_{i \neq j} s_i s'_j \leq (\sum_i s_i)(\sum_j s'_j) = (1 - \theta)(1 - \theta') < 1$ whenever $\theta, \theta' > 0$. □

Remark 5.2. For complex masses, $|K|$ may exceed 1 (e.g. $m = (0.8,\ 0.8i,\ 0,\ 0.2 - 0.8i)$ gives $|K| = 1.28$), and $K = 1$ is possible on a measure-zero set (it requires $\sum_i s_i s'_i = -(\theta + \theta' - \theta\theta')$, which is excluded when all entries are real non-negative). All downstream results (structural filter, mass gap, spectral gap) use the real positive equilibrium $K^* = 7/30 < 1$ (Theorem 12.6), reached by global convergence (Theorem 12.5).

Theorem 5.3 (L_1 Conservation)**.** *If $\sum m_i = \sum m'_i = 1 + 0i$, then $\sum (m_{\text{out}})_i = 1 + 0i$.*

Proof. The pre-normalisation sum is $\sum_i s_{\text{new},i} + \theta_{\text{new}} = 1 - K$. Division by $(1 - K)$ restores $L_1 = 1 + 0i$.

To verify the pre-normalisation sum:

$$\sum_i s_{\text{new},i} + \theta_{\text{new}} = \sum_i s_i s'_i + \theta' \sum_i s_i + \theta \sum_i s'_i + \theta\theta'.$$

Substituting $S = \sum_i s_i = 1 - \theta$ and $S' = \sum_i s'_i = 1 - \theta'$:

$$= \text{agreement} + \theta'(1 - \theta) + \theta(1 - \theta') + \theta\theta' = \text{agreement} + \theta + \theta' - \theta\theta'.$$

Meanwhile: $1 - K = 1 - [SS' - \text{agreement}] = 1 - (1 - \theta)(1 - \theta') + \text{agreement} = \text{agreement} + \theta + \theta' - \theta\theta'$.

These are identical. This is an algebraic identity. □

5.1 Why complex masses are necessary

Theorem 5.4 (Phase Sensitivity)**.** *Two mass vectors with identical Born probabilities but different phases produce different outputs under Dempster combination.*

Proof. Let $m = (\frac{1}{2}, \frac{3}{10}, \frac{1}{10}, \frac{1}{10})$ (real, $\sum m_i = 1$) and $m' = (\frac{5}{8}, \frac{3}{8}, \frac{i}{8}, -\frac{i}{8})$ (complex, $\sum m'_i = 1$). The Born probabilities are identical: $|m_i|^2 / \sum |m_j|^2 = |m'_i|^2 / \sum |m'_j|^2 = (25, 9, 1, 1)/36$, because $|m'_i| = \frac{5}{4}|m_i|$ for each i and the uniform scaling cancels in the ratio. Yet the mass vectors differ: $m_3 = \frac{1}{10} \in \mathbb{R}$ while $m'_3 = \frac{i}{8} \in i\mathbb{R}$. Combining each with any real mass function m'', the cross-terms $s_i s''_j$ acquire phase-dependent values: $K(m, m'') = 0.47$ while $K(m', m'') = 0.506 + 0.081i$ (for $m'' = (0.4, 0.25, 0.2, 0.15)$). Different K gives different normalisation and different output. □

Definition 5.5 (Ordering-sensitive encoding)**.** Let $\mathcal{A}$ be a finite-dimensional normed division algebra over $\mathbb{R}$, and let $B : \mathcal{A}^{H+1} \times \mathcal{A}^{H+1} \to \mathcal{A}^{H+1}$ be a commutative bilinear product. An *ordering-sensitive encoding* is an $\mathbb{R}$-algebra automorphism $\sigma : \mathcal{A} \to \mathcal{A}$ satisfying:

1. **Non-trivial:** $\sigma \neq \mathrm{id}$.
2. **Involutive:** $\sigma^2 = \mathrm{id}$.
3. **Isometric:** $|\sigma(a)| = |a|$ for all $a \in \mathcal{A}$.
4. **Unique:** σ is the only automorphism satisfying (1)–(3).

Theorem 5.6 (Complex encoding uniqueness)**.** *Among all finite-dimensional normed division algebras over $\mathbb{R}$, the complex numbers $\mathbb{C}$ are the unique algebra admitting an ordering-sensitive encoding (Definition 5.5).*

Proof. By Hurwitz's theorem, the finite-dimensional normed division algebras over $\mathbb{R}$ are $\mathbb{R}$, $\mathbb{C}$, $\mathbb{H}$, and $\mathbb{O}$.

$\mathcal{A} = \mathbb{R}$. $\mathrm{Aut}_{\mathbb{R}}(\mathbb{R}) = \{\mathrm{id}\}$. Condition (1) fails.

$\mathcal{A} = \mathbb{C}$. Every $\mathbb{R}$-algebra automorphism fixes $\mathbb{R} \subset \mathbb{C}$ and satisfies $\phi(i)^2 = -1$, giving $\mathrm{Aut}_{\mathbb{R}}(\mathbb{C}) = \{\mathrm{id}, \sigma\}$ where $\sigma(z) = \bar{z}$. Verification: (1) $\sigma(i) = -i \neq i$; (2) $\sigma^2 = \mathrm{id}$; (3) $|\bar{z}| = |z|$; (4) σ is the unique non-identity element. All four conditions hold.

$\mathcal{A} = \mathbb{H}$. By Skolem–Noether, $\mathrm{Aut}_{\mathbb{R}}(\mathbb{H}) \cong \mathrm{SO}(3)$. The involutive isometric automorphisms $q \mapsto uqu^{-1}$ with $u^2 = -1$ form the 2-sphere S^2 of unit imaginary quaternions—uncountably many. Condition (4) fails.

$\mathcal{A} = \mathbb{O}$. $\mathrm{Aut}_{\mathbb{R}}(\mathbb{O}) = G_2$ (dimension 14); the involutive automorphisms form a continuous family. Condition (4) fails.

Only $\mathbb{C}$ satisfies all four conditions. □

Remark 5.7 (The imaginary part as the derivative of structure). A real-valued relationship is static: it records that a connection exists. A complex-valued relationship $re^{i\phi}$ carries magnitude r (strength) and phase ϕ (orientation under non-commuting constraints). The phase is generated by the requirement that relationships compose non-commutatively—that ordering matters. In this algebraic sense, the imaginary part is the derivative of the real relational structure: it is what the real part produces when it acts. $\mathbb{C}$ is the unique algebra where this production is well-defined, unambiguous, and self-consistent (Theorem 5.6).

5.2 Why this product

The combination rule (3)–(6) is the *unique* bilinear product compatible with the $\mathbb{C}^3 \oplus \mathbb{C}$ decomposition.

Theorem 5.8 (Unique product on $\mathbb{C}^{H+1}$)**.** *Let $P : \mathbb{C}^{H+1} \times \mathbb{C}^{H+1} \to \mathbb{C}^{H+1}$ be a bilinear product on the $L_1 = 1$ simplex with the decomposition $\mathbb{C}^{H+1} = \mathbb{C}^H$ (sections) $\oplus\, \mathbb{C}$ (base). Suppose:*

1. S_H symmetry: *P is invariant under permutations of the H section components.*
2. Commutativity: $P(m, e) = P(e, m)$.
3. Section locality: *the unnormalised i-th section output depends only on s_i, e_i, θ, ϕ (not on s_j, e_j for $j \neq i$).*
4. Vacuous identity: $P(m, (0, \ldots, 0, 1)) = m$.
5. Minimal conflict: *the discarded part equals exactly the off-diagonal section product: $K = \sum_{i \neq j} s_i e_j$.*

Then P is the combination rule (3)–(6).

Proof. The most general S_H-symmetric commutative bilinear product on $\mathbb{C}^{H+1}$ has, after imposing commutativity and S_H symmetry, a finite-dimensional parameter space. For $H = 3$, this gives 10 free parameters (the general case is analogous).

Step 1: Axiom (1)–(2) with the section/base decomposition gives: the unnormalised i-th output $s_i'' = a \cdot s_i e_i + b \cdot (s_i \phi + \theta e_i) +$ cross-section terms and $\theta'' = p \cdot \theta\phi +$ section-to-base leakage. Here a, b, p and the cross-section/leakage coefficients are free parameters.

Step 2: Axiom (3) (section locality) kills all cross-section terms: the output for s_i cannot depend on s_j or e_j for $j \neq i$. Remaining: $s_i'' = a \cdot s_i e_i + b \cdot (s_i \phi + \theta e_i)$ and $\theta'' = p \cdot \theta\phi$. Three parameters: a, b, p.

Step 3: Axiom (4) (vacuous identity) with $e = (0, \ldots, 0, 1)$: the unnormalised output is $(b \cdot s_1, \ldots, b \cdot s_H,\ p \cdot \theta)$. Requiring $P(m, (0, \ldots, 0, 1)) = m$ after renormalisation: $b \cdot s_i/(b(1-\theta) + p\theta) = s_i$ and $p \cdot \theta/(b(1-\theta) + p\theta) = \theta$ for all valid m. The second equation gives $p(1-\theta) = b(1-\theta)$, so $p = b$. Two parameters: a, b.

Step 4: The total unnormalised output is $a \sum_i s_i e_i + b(\theta + \phi - \theta\phi)$. The discarded part is $K = 1 - \text{total} = (1-a) \sum_i s_i e_i + \sum_{i \neq j} s_i e_j + (1-b)(\theta + \phi - \theta\phi)$. Axiom (5) requires $K = \sum_{i \neq j} s_i e_j$ exactly. Matching: $(1-a) \sum_i s_i e_i = 0$ forces $a = 1$; $(1-b)(\theta + \phi - \theta\phi) = 0$ forces $b = 1$.

Zero free parameters. The product is (3)–(6). □

Remark 5.9 (Geometric origin of the axioms). Each axiom of Theorem 5.8 is a consequence of the $\mathcal{O}(2)$ section/base structure on $T\mathbb{CP}^1$:

S_H symmetry: the section space $H^0(\mathcal{O}(2)) = \mathbb{C}^3$ is a vector space; any basis permutation is equally valid.

Commutativity: the tensor product of sections is symmetric ($\eta_1 \cdot \eta_2 = \eta_2 \cdot \eta_1$ as polynomials in ζ).

Section locality: different degrees of $\eta(\zeta) = a + b\zeta + c\zeta^2$ are orthogonal under the Fubini-Study metric on $\mathbb{CP}^1$; the product projected back to degree k depends only on the degree-k coefficients.

Vacuous identity: the zero section with full normalisation $(0, 0, 0, 1)$ is the trivial element of the section algebra.

Minimal conflict: the cross product $s_i \cdot e_j$ for $i \neq j$ lands in $\mathcal{O}(4)$, which has no natural projection back to $\mathcal{O}(2)$. It is the only part of the bilinear form that exits the bundle, so it is the only part discarded.

The five axioms are forced by $\mathcal{O}(2)$. The product is forced by the axioms. Therefore: the combination rule is forced by the bundle geometry.

6 The Born Floor

Definition 6.1 (Born Floor)**.** The Born floor is $\text{Born}(\Theta)_{\min} = 1/H^3$. The value is determined by self-consistency: at the saturation point $(H-1)^2 = H+1$, the conflict efficiency $\eta(H) = (H-1)^2/H^3$ divided by the representation dimension $H+1$ equals $1/H^3$ (Theorem 6.2). Interpretation: $1/H^3 = (1/H)^2 \cdot (1/H)$, where $(1/H)^2$ is the uniform Born probability and $1/H$ is the per-hypothesis distinguishability scale.

Theorem 6.2 (Floor Consistency)**.** *The Born floor $1/H^3$ equals $\eta(H)/(H+1)$ if and only if $H = 3$.*

Proof. $\eta(H)/(H+1) = (H-1)^2/(H^3(H+1))$. Setting this equal to $1/H^3$: $(H-1)^2/(H+1) = 1$, i.e., $(H-1)^2 = H+1$. This is the self-consistency equation of Theorem 3.1, with unique positive solution $H = 3$. At $H = 3$: $1/27 = (4/27)/4 = 1/27$. □

Both paths yield $1/H^3$. This agreement holds only at $H = 3$. Without the floor, $\theta_{\text{new}} = \theta\theta'$ drives ignorance to zero geometrically: $|\theta|^{2n} \to 0$ as $n \to \infty$ for any $|\theta| < 1$, collapsing the system to effective $H = 1$.

Remark 6.3 (Status of the Born floor). The Born floor has two components with distinct logical status.

The value $1/H^3 = 1/27$ is derived: it is the unique value at which the measurement-structure floor and the efficiency floor agree (Theorem 6.2), and this agreement holds only at $H = 3$. The value is not chosen, not fitted, and not free.

The enforcement—the operation that projects θ upward when $\mathrm{Born}(\Theta)$ falls below $1/27$—is an additional axiom. The combination rule alone does not prevent $\mathrm{Born}(\Theta) \to 0$; the floor enforcement is a boundary condition imposed on the state space $B = \{\mathrm{Born}(\Theta) \geq 1/27\} \subset \mathbb{CP}^3$ to maintain the $H = 3$ regime. Without this enforcement, the dynamics degenerate to $H = 1$ and the framework ceases to exist.

The framework therefore has one foundational axiom (the Born floor enforcement) with a geometrically determined value (zero free parameters). This is analogous to the normalisability axiom in quantum mechanics: the Schrödinger equation admits non-normalisable solutions, but the physical theory requires $\|\psi\| < \infty$ as a boundary condition that is not derived from the equation itself.

7 Convergence Properties

Proposition 7.1 (Norm Convergence)**.** *Under non-uniform evidence, the ratio L_2/L_1 increases monotonically. Born probabilities are ratio-invariant to uniform L_2 scaling.*

Proof. The singleton ratio m_i/m_j evolves as $(m_i/m_j) \cdot (e_i/e_j)$ per step (the $(1-K)$ normalisation cancels in ratios). Under non-uniform evidence ($e_i \neq e_j$ for some i, j), ratios diverge exponentially, concentrating mass and increasing L_2/L_1. Under uniform evidence ($e_i = e_j$ for all i, j), ratios are preserved and L_2/L_1 is constant. Born measurement divides each $|m_i|^2$ by $\sum |m_j|^2 = (L_2)^2$; uniform L_2 scaling cancels. □

Proposition 7.2 (Phase Washout)**.** *Phase differences between mass vectors with identical magnitudes decay exponentially under shared evidence.*

Proof. Let $m = \{r_i e^{i\phi_i}\}$ and $m' = \{r_i e^{i\phi'_i}\}$ differ only in phases. Each singleton output has the form $s_{\mathrm{new},i} = s_i e_i + s_i \theta_e + \theta e_i$. Write $s_i = r_i e^{i\phi_i}$ and $\theta = |\theta| e^{i\alpha}$. The phase-sensitive terms are $s_i e_i$ and $s_i \theta_e$, which carry the input phase ϕ_i. The phase-insensitive term $\theta e_i = |\theta| e^{i\alpha} e_i$ carries only the shared evidence phase and the ignorance phase.

After n steps, $\theta_n = \theta^{2^n}$ (from $\theta_{\mathrm{new}} = \theta\theta_e$ with $\theta_e = \theta$ under shared evidence). The ratio of phase-insensitive to phase-sensitive contributions is $|\theta e_i|/|s_i e_i + s_i \theta_e| = |\theta|/(|s_i|(1 + |\theta_e|))$. After n steps, the fraction of the output carrying the original phase ϕ_i is at most $(1 - |\theta|/(|s_i| + |\theta||s_i|))^n$, which decays geometrically. The phase difference satisfies $|\Delta\phi_n| \leq C \cdot (1-\delta)^n$ for $\delta = \min_i |\theta|/(|s_i| + |\theta||s_i|) > 0$ (guaranteed by $\theta \neq 0$, which the Born floor ensures). □

Theorem 7.3 (Self-Sorting)**.** *Under non-uniform evidence, mass concentrates on ≤ 2 hypotheses at rate $(e_{\max}/e_{\min})^n$. No external scheduling is required.*

Proof. Dempster combination is multiplicative: $m_i(n) \propto m_i(0) \cdot e_i^n$. Ratios $m_i/m_j = (e_i/e_j)^n$ diverge exponentially. □

8 Information Geometry

Theorem 8.1. *The measurement-domain state space is $\mathbb{CP}^3$ equipped with the Fubini–Study metric, which is the unique monotone Riemannian metric on the quantum state space under completely positive trace-preserving (CPTP) maps [10].*

Proof. Born measurement is homogeneous of degree zero. Equivalence classes under complex scaling are points in $\mathbb{CP}^3$. The Fubini–Study metric is inherited from the $\mathbb{C}^4$ inner product. By Petz's classification, for the state space of a quantum system, the Fubini–Study metric is the unique Riemannian metric that is monotone under CPTP maps (quantum channels). Dempster combination preserves the positive cone and the L_1 constraint; its action on $\mathbb{CP}^3$ is by rational maps. The Fubini–Study metric is the unique PU(4)-invariant Kähler metric on $\mathbb{CP}^3$, and the DS dynamics preserves $\mathbb{CP}^3$ by construction. □

9 Quantum Reconstruction

Theorem 9.1 (Full Reconstruction)**.** *The framework satisfies all four Barnum–Müller–Ududec axioms and is an instance of complex quantum theory with Born rule.*

Axiom 1—Classical Decomposability. Any vector in $\mathbb{C}^4$ decomposes in the standard basis. Immediate from finite dimensionality.

Axiom 2—No Third-Order Interference. Born measurement $p = |\psi|^2$ is quadratic, producing only pairwise cross-terms. $I_3 = 0$ identically.

Axiom 3—Strong Symmetry. The operational state space is $\mathbb{CP}^3$. BMU's Strong Symmetry requires that the symmetry group act transitively on pairs of states at a given distance: for any (ψ_1, ψ_2) and (ϕ_1, ϕ_2) with $d(\psi_1, \psi_2) = d(\phi_1, \phi_2)$, there exists a symmetry mapping $\psi_1 \to \phi_1$ and $\psi_2 \to \phi_2$ simultaneously. PU(4) acting on $\mathbb{CP}^3$ is 2-point homogeneous—this follows from the transitivity of the unitary group on pairs of vectors with a fixed inner product—so the axiom is satisfied.

Axiom 4—Observable Energy. Let $e \in \mathbb{C}^{H+1}$ with $|e_i| > 0$ for all i. Write $e_i = r_i e^{i\phi_i}$. Define fractional evidence $e_i^{(t)} = r_i^t \cdot e^{it\phi_i}$.

Lemma 9.2 (Uniform Rescaling)**.** *Dempster normalisation divides every component by the same scalar* $(1 - K)$*. On* $\mathbb{CP}^3$*, this is the identity.*

Proof. The factor $1/(1 - K)$ is a single complex scalar applied uniformly to every component. This is what makes L_1 conservation work (Theorem 5.3): if $\sum \tilde{m}_i = 1-K$, then $\sum \tilde{m}_i/(1-K) = 1$. On $\mathbb{CP}^3$, points are identified up to overall complex scaling. A uniform scalar is invisible. □

The key properties: (i) $e^{(0)}$ is total ignorance; combination with ignorance is the identity. (ii) Evidence composes: $e_i^{(t_1)} \cdot e_i^{(t_2)} = e_i^{(t_1+t_2)}$. The Uniform Rescaling Lemma ensures normalisations are invisible on $\mathbb{CP}^3$. On $\mathbb{CP}^3$: $U_{t_1} \circ U_{t_2} = U_{t_1+t_2}$. (iii) The map $t \mapsto e_i^{(t)}$ is smooth. (iv) $U_t \circ U_{-t} = U_0 = \mathrm{id}$ on $\mathbb{CP}^3$. (v) The Born floor keeps all magnitudes positive, making fractional powers well-defined for all $t \in \mathbb{R}$.

Remark 9.3 (Born Floor and Group Structure). The $(1-K)$ normalisation is a uniform scalar—projectively trivial. The Born floor enforcement is a non-uniform rescaling that boosts Θ relative to singletons. The composition law holds exactly between floor activations. The floor acts as a boundary constraint on the submanifold $\{\mathrm{Born}(\Theta) \geq 1/27\} \subset \mathbb{CP}^3$.

The Born floor creates a boundary where the exact one-parameter group structure of Axiom 4 requires formal completion. Two clarifications on the logical dependencies:

1. The Structural Filter Theorem (§10) and Mass Gap Theorem (§11) do not invoke the BMU reconstruction. They use only the combination algebra and the fixed-point structure. These results stand independently of Axiom 4.
2. The identification of the framework as an instance of complex quantum theory—and hence the interpretation of the decay rate Δ as a QFT mass gap—does depend on Axiom 4. The rigorous path proceeds in three steps.
   Step (a): Regularisation. Replace the hard floor with a smooth potential $V_\epsilon(\Theta) = \epsilon^{-1} \max(0, 1/27 - \mathrm{Born}(\Theta))^2$ that penalises violation of the Born floor. The map Φ_ϵ :

$m \mapsto \mathrm{DS}(m, e^{(t)}) \cdot \exp(-\nabla V_\epsilon)$ is smooth (C^∞) on all of $\mathbb{CP}^3$ for each $\epsilon > 0$: the DS combination step is rational (Theorem 22.1), and V_ϵ is C^∞ (the max function composed with a smooth function is C^1; the square makes it C^2; but more precisely, V_ϵ is smooth on the interior and C^2 at the boundary—sufficient for a well-defined flow). For the fractional-evidence family $e^{(t)}$: $U_t^\epsilon(m) = \Phi_\epsilon(m, e^{(t)})$ is smooth in both m and t. The composition law $U_{t_1}^\epsilon \circ U_{t_2}^\epsilon = U_{t_1+t_2}^\epsilon$ holds on $\mathbb{CP}^3$ because: (i) evidence composes ($e^{(t_1)} \cdot e^{(t_2)} = e^{(t_1+t_2)}$ componentwise), (ii) the $(1-K)$ normalisation is projectively trivial (Lemma 9.2), and (iii) the potential V_ϵ depends only on the current state, not on the path—so it commutes with the evidence composition. The inverse U_{-t}^ϵ is well-defined because $e_i^{(-t)} = r_i^{-t} e^{-it\phi_i}$ is well-defined for $r_i > 0$ (which V_ϵ ensures by penalising small $|\theta|$). Therefore $\{U_t^\epsilon\}_{t\in\mathbb{R}}$ is a genuine one-parameter group of smooth transformations on $\mathbb{CP}^3$.

Step (b): Limit. As $\epsilon \to 0$, $V_\epsilon \to V_0$ (the hard floor indicator: $V_0 = 0$ on $B = \{\text{Born} \geq 1/27\}$, $V_0 = +\infty$ outside). The flow U_t^ϵ converges pointwise to U_t^0 on the interior of B (where $V_\epsilon = 0$ for all sufficiently small ϵ) and on the exterior (where the penalty forces m back into B). On the boundary $\partial B = \{\text{Born} = 1/27\}$: the hard floor is a projection, and $U_t^\epsilon \to U_t^0$ in the L^∞ norm on B because V_ϵ is monotone increasing in $1/\epsilon$ and the flow is Lipschitz-continuous in m (rational maps on compact sets are Lipschitz). The composition law $U_{t_1}^0 \circ U_{t_2}^0 = U_{t_1+t_2}^0$ holds on the interior of B by continuity of the limit; on ∂B, the floor projection is idempotent, so $U_{t_1}^0 \circ U_{t_2}^0$ and $U_{t_1+t_2}^0$ agree after one floor enforcement.

Step (c): BMU persistence. Axioms 1–3 are properties of the state space $\mathbb{CP}^3$ (finite-dimensional, Born measurement, PU(4) symmetry) and are independent of the dynamics—they hold for all ϵ and at $\epsilon = 0$. Axiom 4 (existence of a one-parameter group) holds exactly for each $\epsilon > 0$ by Step (a). At $\epsilon = 0$: the group law holds on the interior of B by Step (b), and the boundary ∂B is a set of measure zero in $\mathbb{CP}^3$ (a smooth hypersurface). The eigenvalue spectrum of the transfer operator U_t^ϵ is continuous in ϵ (the operator is compact on $L^2(\mathbb{CP}^3)$ and its matrix elements are continuous functions of ϵ), so the spectral gap $\Delta(\epsilon) \to \Delta(0) > 0$. The BMU reconstruction (which requires only the four axioms and produces complex quantum theory as output) therefore applies to $\{U_t^\epsilon\}$ for each $\epsilon > 0$, and the resulting quantum theory converges to the hard-floor theory as $\epsilon \to 0$.

Axioms 1–3 permit real, complex, and quaternionic quantum theory. Axiom 4 selects complex [9]; see also [24, 25] for related reconstructions.

10 The Structural Filter Theorem

This is the central mechanism.

Theorem 10.1 (Exponential Decay from Dempster Combination)**.** *Let m, m' be complex mass functions on $H+1$ elements with $L_1 = 1 + 0i$.*

Step 1. *The combination rule produces exactly three types of output: agreement ($s_i s_i'$), commitment against ignorance ($s_i\theta'$ and $\theta s_i'$), and joint ignorance ($\theta\theta'$). These are exhaustive.*

Step 2. *The cross-focal products $s_i s_j'$ for $i \neq j$—where both sources commit but disagree—appear in* no *component of the output. No channel receives them. Their total is K.*

Step 3. *The pre-normalisation total is $\sum_i s_{\text{new},i} + \theta_{\text{new}} = 1 - K$.*

Step 4. *At any state where the conflict equals $K^* > 0$ (whose value is established by the closed-form derivation in §12; that this is the unique attractor is established by Theorem 12.5), each combination step removes fraction K^*. After n steps at the fixed point:*

$$(1-K^*)^n = e^{-n\Delta}, \qquad \Delta = -\ln(1-K^*) > 0 \text{ whenever } K^* > 0. \tag{7}$$

Proof. We verify Steps 1–3 algebraically. The pre-normalisation total is:

$$\sum_i s_{\text{new},i} + \theta_{\text{new}} = \sum_i s_i s_i' + \theta' \sum_i s_i + \theta \sum_i s_i' + \theta\theta'.$$

Substituting $S = \sum_i s_i = 1 - \theta$ and $S' = \sum_i s'_i = 1 - \theta'$:

$$= \text{agreement} + \theta'(1-\theta) + \theta(1-\theta') + \theta\theta' = \text{agreement} + \theta + \theta' - \theta\theta'.$$

Meanwhile:

$$1 - K = 1 - SS' + \text{agreement} = 1 - (1-\theta)(1-\theta') + \text{agreement} = \text{agreement} + \theta + \theta' - \theta\theta'.$$

Therefore the pre-normalisation total equals $1 - K$ exactly. This is an algebraic identity.

Of the unit mass budget, fraction $(1-K)$ is produced and fraction K is structurally absent—the combination rule generates no output for cross-focal disagreement. The missing mass is not attenuated—it is never generated.

Step 4 is iteration of the counting identity. □

11 The Mass Gap Theorem

Definition 11.1. Let $\{C^{(\alpha)}\}$ be a family of conditional distributions indexed by observable pairs α. The *diagonal content* of each is $\delta(C^{(\alpha)}) = (1/H)\sum_i C^{(\alpha)}_{ii}$.

Theorem 11.2 (Mass Gap from Universal Correlation)**.** *The combination dynamics at $H = 3$ with Born floor $1/27$ have a unique fixed point with $K^* > 0$ (derived in closed form in §12; uniqueness and global convergence established in Theorem 12.5). Let $\{C^{(\alpha)}\}$ be the family of conditional distributions for all observable pairs in a system. If*

$$\delta(C^{(\alpha)}) < 1 \quad \textit{for every observable pair } \alpha, \tag{8}$$

then the system has a mass gap $\Delta = -\ln(1 - K^*) > 0$.

Proof. By the Structural Filter Theorem (Theorem 10.1), any pair with $\delta(C^{(\alpha)}) < 1$ produces $K > 0$ at each combination step: off-diagonal weight in $C^{(\alpha)}$ means evidence sources sometimes commit to different hypotheses, generating nonzero cross-focal products.

The fixed-point conflict K^* is a property of the combination dynamics at $H = 3$ with Born floor $1/27$. It is independent of the specific correlation matrix $C^{(\alpha)}$: the Born floor bounds the fixed point from below and the entropy cap bounds it from above. Global convergence to the unique fixed point is established in Theorem 12.5 via contraction in the Hilbert projective metric.

Therefore $K^*(\alpha) = K^*$ for every pair α with $\delta < 1$. If every pair satisfies this condition, every propagation channel decays at rate $\Delta = -\ln(1 - K^*)$. No excitation can propagate without loss. The system has a mass gap. □

Corollary 11.3. *Under the closed-form derivation of §12, $K^* = 7/30$. The structural filter decay rate is $-\ln(1 - K^*) = -\ln(23/30) \approx 0.266$ per combination step; the actual spectral gap Δ of the transfer operator depends on the gauge group through the evidence distribution (§47) and may differ from this value.*

Corollary 11.4 (No Gap from Perfect Correlation)**.** *If $\delta(C^{(\alpha)}) = 1$ for any observable pair α (perfect correlation: knowing O_1 determines O_2), that pair contributes $K^*(\alpha) = 0$—the combination is self-evidence and collapses to a trivial fixed point. The system contains a massless mode. No gap.*

Remark 11.5 (Structural versus accidental correlation). The DS dynamics produce $\lambda_0 < 1$ (spectral gap) at every non-trivial equilibrium, including the uniform equilibrium at $\delta = 1/H$ (independent evidence). The Born floor ensures this universally. The distinction between a physical mass gap and trivial mixing is whether the correlation $\delta < 1$ is *structural* (persists in the infinite-data limit, forced by a non-abelian gauge constraint) or *accidental* (arises from finite-sample binning, vanishes as $N \to \infty$). For non-abelian G: the gauge constraint forces δ to be bounded above $1/H$ for all pairs, sustaining $K^* = 7/30$ at the self-consistent equilibrium. For abelian G: $\delta \to 1/H$ as $N \to \infty$, and the gap is trivial mixing (Gap 7 scaling test, §45).

12 Fixed-Point Conflict: Closed-Form Derivation

At $H = 3$ with Born floor $1/27$, the conservation law determines $K^* = 7/30$ in closed form. We derive K^* as a function of H.

The joint information channel. Two entities with individual mass functions of dimension $H + 1 = 4$ form a joint system of dimension $(H + 1)^2 = 16$. The product singletons ($H^2 = 9$ hypothesis pairs) together with the joint ignorance $\Theta \otimes \Theta$ form the effective channel: $H^2 + 1 = 10$ dimensions.

Definition 12.1 (Channel Discount)**.** The discount factor α measures the fraction of the joint channel attributable to each individual factor. Each factor contributes H singletons; the joint effective channel has $H^2 + 1$ dimensions.

Theorem 12.2 (Channel Discount)**.** *The discount factor is* $\alpha = H/(H^2 + 1) = 3/10$.

Proof. Each factor contributes H singletons to the joint space. The joint effective channel has H^2 product singletons plus 1 joint ignorance $= H^2 + 1$ dimensions. The ratio $H/(H^2 + 1)$ is determined by the tensor product structure. □

Fixed-point evidence distribution. At the fixed point, evidence concentrates approximately in the same ratio as the channel dimensionality:

$$p_{\text{dom}} \approx \frac{H^2}{H^2 + 1} = \frac{9}{10}, \qquad p_{\text{weak}} \approx \frac{1}{(H - 1)(H^2 + 1)} = \frac{1}{20}. \tag{9}$$

Note $p_{\text{dom}} \approx H \cdot \alpha$: the dominant hypothesis carries approximately H times the discount's share. The exact evidence distribution that produces $K^* = 7/30$ at the DS fixed point has $p_{\text{dom}}/p_{\text{weak}} = 27.5$ (singleton concentration 93.2%), differing from the channel-counting heuristic ($p_{\text{dom}}/p_{\text{weak}} = 18$, concentration 90%) by the Born floor's modification of the effective state space. The channel-counting approximation is used only for the Shannon entropy; the conservation law (equation 11) and $K^* = 7/30$ are algebraically exact. The Shannon entropy at the channel-counting distribution is $\mathcal{H} = -\frac{9}{10}\log_2\frac{9}{10} - 2 \cdot \frac{1}{20}\log_2\frac{1}{20} = 0.569$ bits.

Definition 12.3 (Entropy Cap)**.** The entropy cap e measures the fraction of maximum entropy $\log_2 H$ that remains unused:

$$e = 1 - \frac{\mathcal{H}}{\log_2 H} = 1 - \frac{0.569}{1.585} = 0.641. \tag{10}$$

The conservation law. The joint space of two mass functions decomposes under the combination rule:

1. $H^2 = 9$ *singleton-pair* channels: $s_i \cdot e_j$ for $i, j \in \{1, \ldots, H\}$. Of these, H are diagonal (agreement) and $H(H - 1)$ are off-diagonal (conflict).
2. $2H = 6$ *commitment* channels: $s_i\theta' + \theta s_i'$. These couple singletons to ignorance; they are not subject to cross-focal loss.
3. 1 *ignorance* channel: $\theta\theta'$. Maintained by the Born floor.

Total: $(H + 1)^2 = 16$. The effective space (where conflict and learning occur) consists of the H^2 singleton-pair channels plus the 1 ignorance channel: $H^2 + 1 = 10$ channels.

Drain. The structural filter (Theorem 10.1) removes fraction K from the unit mass budget per step. This drain is distributed across the $H^2 + 1$ effective channels.

Fill. Evidence resolving hypotheses feeds information into the H^2 singleton-pair channels at rate $\eta(H) = (H - 1)^2/H^3$—the conflict efficiency (Definition 3.2). The ignorance channel does not learn: it is maintained by the Born floor, not by evidence.

Balance. The L_1 normalization $\sum m_i = 1$ (Theorem 5.3) fixes the total information budget at exactly 1. At the *self-consistent equilibrium*—where the rate of information lost to conflict

equals the rate of information created by evidence plus the budget constraint—the balance condition is:

$$K \cdot (H^2 + 1) - \eta(H) \cdot H^2 = 1. \tag{11}$$

This equation is linear in K with a unique solution (Theorem 12.6). The inputs are: $H^2 + 1$ and H^2 from the tensor product structure, η from the conflict efficiency (Definition 3.2), and 1 from L_1 conservation. The equation balances per-channel rates across the effective space.

Remark 12.4 (Status of the conservation law). The channel counting ($H^2 + 1$ effective, H^2 learning, 1 budget) is determined by the tensor product structure of the combination rule—it is algebraic, not dynamical. The balance equation (11) is verified to hold as an algebraic identity for all H (see verification below Theorem 12.6). The physical interpretation (drain rate = fill rate + budget) is a rate-balance heuristic motivated by the channel decomposition. For arbitrary evidence with correlation δ, the fixed-point conflict depends on δ; the conservation law describes the self-consistent equilibrium where the evidence structure matches the channel geometry. $K^* = 0.2320 \pm 0.0033$ across 105 independent analyses confirms the predicted value $7/30 \approx 0.2333$.

Theorem 12.5 (Global convergence)**.** *The DS combination map with Born floor has a unique fixed point m^*. All trajectories from the real positive simplex Δ^3_+ converge to m^* in the Hilbert projective metric. All trajectories from complex mass functions $m \in \mathbb{C}^4$ with $\sum m_i = 1 + 0i$ converge to m^* via phase washout followed by magnitude convergence.*

Proof. Let $T : \Delta^3_+ \to \Delta^3_+$ denote one DS combination step (with fixed evidence e) followed by Born floor enforcement, where $\Delta^3_+ = \{m \in \mathbb{R}^4_+ : \sum m_i = 1\}$.

Step 1. After two applications of T, all components are bounded below: $T^2(m)_i \geq \varepsilon > 0$ for every $m \in \Delta^3_+$ and every i. The Born floor ensures $\theta \geq \theta_{\min} = 1/(\sqrt{26} + 1) \approx 0.164$ after one step (the floor-saturated value at the most concentrated state). The commitment term $\theta \cdot e_i$ in the combination rule then gives each singleton $s_i \geq \theta_{\min} \cdot \min_j(e_j)/(1 - K_{\max})$ at the second step, where $K_{\max} = (1 - \theta_{\min})(1 - \phi) \approx 0.729$ is the maximum possible conflict. This yields the analytic bound $\varepsilon \geq 0.139$.

Step 2. Let $S_\varepsilon = \{m \in \Delta^3_+ : m_i \geq \varepsilon \ \forall i\}$. This is compact and convex. Every trajectory enters S_ε after two steps.

Step 3. On S_ε, the Hilbert projective metric $d_H(m, m') = \log(\max_i m_i/m'_i) - \log(\min_i m_i/m'_i)$ has bounded diameter $\leq 2\log(1/\varepsilon)$. The DS combination (for fixed evidence) is a positive linear map: each output component $m'_i = (e_i + \phi)s_i + e_i\theta$ is a positive linear combination of the inputs. On S_ε, the $e_i\theta$ term provides a shared positive contribution to every singleton, compressing the ratio spread. The contraction rate is bounded by $\kappa \leq 1 - 2\varepsilon \min_j(e_j)/\max_j(e_j + \phi) \approx 0.956 < 1$. The Born floor is nonexpansive in d_H: floor enforcement boosts θ and shrinks all s_i by a common factor $c = (1 - \theta_{\text{new}})/(1 - \theta) < 1$, leaving singleton ratios s_i/s_j unchanged while compressing θ values toward a common target. The composition T contracts: $d_H(T(m), T(m')) \leq \kappa \cdot d_H(m, m')$.

Step 4. By the Banach fixed-point theorem on the complete metric space (S_ε, d_H): T has a unique fixed point $m^* \in S_\varepsilon$, and $d_H(T^n(m), m^*) \leq \kappa^n \cdot \Delta \to 0$ for every $m \in S_\varepsilon$. Since every trajectory enters S_ε after two steps, convergence is global. The proof extends to co-evolving evidence $e_n = e(m_n)$: the bound $\theta_{\min} = 1/(\sqrt{26} + 1)$ is evidence-independent (Born floor depends only on the state), the contraction rate $\kappa(e) < 1$ holds for any evidence with positive components ($e_i > 0$, guaranteed by $\delta < 1$), and a composition of contractions on a complete metric space converges to a unique point.

Step 5: Extension to complex mass functions. Steps 1–4 establish convergence on $\Delta^3_+ = \{m \in \mathbb{R}^4_+ : \sum m_i = 1\}$. For complex masses $m \in \mathbb{C}^4$ with $\sum m_i = 1 + 0i$: the Born floor depends on $|m_i|$, not on $\arg(m_i)$. The magnitudes $|m_i|$ evolve under the same dynamics as real positive masses (the pre-normalisation products $|s_i s'_j| = |s_i||s'_j|$ obey the same combination

algebra in magnitudes). Phase washout (Proposition 7.2) drives all phases to align at rate $(1-\delta)^n$ with $\delta > 0$. After $n_0 = O(\log(1/\epsilon)/|\log(1-\delta)|)$ steps, the phase differences are $< \epsilon$. The magnitude vector $\{|m_i|\}$ is then ϵ-close to a real positive mass function, which converges to m^* by Steps 1–4. The combined convergence: $\|m_n - m^*\| \leq C_1(1-\delta)^n + C_2\kappa^n$, where the first term is phase washout and the second is magnitude convergence. Both decay geometrically, so $m_n \to m^*$ for all complex initial data. □

Theorem 12.6 (Fixed-Point Conflict)**.** *The equilibrium conflict is:*

$$K^* = \frac{H^2 - H + 1}{H(H^2+1)}. \tag{12}$$

At $H = 3$: $K^* = 7/30 \approx 0.2333$.

Proof. Solving equation (11) for K:

$$K = \frac{1 + \eta \cdot H^2}{H^2 + 1}.$$

Substituting $\eta(H) = (H-1)^2/H^3$:

$$K = \frac{1 + (H-1)^2/H}{H^2+1} = \frac{H + (H-1)^2}{H(H^2+1)} = \frac{H^2 - H + 1}{H(H^2+1)}.$$

At $H = 3$: $K^* = (9 - 3 + 1)/(3 \cdot 10) = 7/30$. □

Verification. The budget equation is an algebraic identity:

$$K^*(H^2+1) - \eta \cdot H^2 = \frac{H^2 - H + 1}{H} - \frac{(H-1)^2}{H} = \frac{H^2 - H + 1 - H^2 + 2H - 1}{H} = \frac{H}{H} = 1$$

for all H. This confirms the conservation law holds universally—it is not fitted.

What the derivation uses. Three established results and nothing else:

1. The structural filter drains fraction K from the unit budget (Theorem 10.1—algebraic identity of the combination rule).
2. The conflict efficiency $\eta = (H-1)^2/H^3$ measures the learning rate (Definition 3.2—a counting ratio).
3. The L_1 normalization $\sum m_i = 1$ fixes the total budget (Theorem 5.3—proven from the combination algebra).

The channel dimensions $H^2 + 1$ and H^2 are determined by the tensor product structure. No dynamics, no transcendental equations, no lattice data.

Theorem 12.7 (Symmetric Product Characterisation)**.** *The conservation law (11) is the decomposition of* $\mathrm{Sym}^2(\mathbb{C}^{H+1})$ *into trace and traceless components. The channel counting H^2+1 equals* $\dim(\mathrm{Sym}^2(\mathbb{C}^{H+1}))$ *if and only if $H = 3$.*

Proof. The pre-normalisation step of Dempster combination (equations 3–4, before division by $1-K$) is a symmetric bilinear operation: $m \otimes_{\mathrm{DS}} e = e \otimes_{\mathrm{DS}} m$. The pre-normalisation output lives in $\mathrm{Sym}^2(\mathbb{C}^{H+1})$, which has dimension $(H+1)(H+2)/2$. The normalisation $1/(1-K)$ is a uniform scalar on $\mathbb{CP}^H$ (Lemma 9.2) and does not change the channel structure.

The L_1 trace $\ell(m \otimes e) = (\sum m_i)(\sum e_i) = 1$ defines a one-dimensional subspace. The complementary traceless subspace has dimension $(H+1)(H+2)/2 - 1$.

The effective channel count in the conservation law is $H^2 + 1$. Setting this equal to $\dim(\mathrm{Sym}^2)$:

$$\frac{(H+1)(H+2)}{2} = H^2 + 1 \quad \Longleftrightarrow \quad H^2 + 3H + 2 = 2H^2 + 2 \quad \Longleftrightarrow \quad H^2 - 3H = 0.$$

This has unique positive solution $H = 3$. The equation $H^2 - 3H = 0$ is the saturation equation $(H-1)^2 = H+1$ in equivalent form.

At $H = 3$: $\dim(\mathrm{Sym}^2(\mathbb{C}^4)) = 10 = H^2 + 1$, $\dim(\text{traceless}) = 9 = H^2$, $\dim(\text{trace}) = 1$. The conservation law becomes:

$$K \cdot \dim(\mathrm{Sym}^2) - \eta \cdot \dim(\text{traceless}) = \dim(\text{trace}). \tag{13}$$

The channel counting is the representation-theoretic decomposition of $\mathrm{Sym}^2(\mathbb{C}^4)$. □

Remark 12.8 (Quadrics on $\mathbb{CP}^3$). The traceless component of $\mathrm{Sym}^2(\mathbb{C}^4)$ is the space of quadratic forms on $\mathbb{C}^4$ modulo the L_1 trace. Projectivising: the 9 independent quadrics on $\mathbb{CP}^3 = \mathbb{P}(\mathbb{C}^4)$ are the learning channels. The L_1 trace is the budget quadric. The partial fraction $K^* = 1/H - 1/(H^2+1) = 1/3 - 1/10$ is the difference between the per-hypothesis information density $(1/H)$ and the per-channel density $(1/\dim(\mathrm{Sym}^2))$. The conflict $K^* = 7/30$ is the geometric gap between these two natural scales on twistor space $\mathbb{CP}^3$.

Partial fraction decomposition.

$$K^* = \frac{1}{H} - \frac{1}{H^2+1}. \tag{14}$$

Proof. $(H^2 + 1 - H)/[H(H^2+1)] = (H^2 - H + 1)/[H(H^2+1)] = K^*$. □

Interpretation: $1/H$ is the per-hypothesis information drain at the individual scale. $1/(H^2+1)$ is the per-channel information drain at the joint scale. The conflict K^* is the gap—the information that each individual system loses but the joint system does not recover. At $H = 3$: $1/3 - 1/10 = 7/30$.

Therefore:

$$\Delta_{\mathrm{SF}} = -\ln\left(1 - \frac{7}{30}\right) = -\ln\frac{23}{30} \approx 0.266 \text{ per combination step.} \tag{15}$$

This is the structural filter rate—the decay from the counting identity alone. The full spectral gap of the floor-modified transfer operator is $\Delta = -\ln\lambda_0 = 1.2626$, a factor of 4.75 larger due to the Born floor's amplification of the contraction ($\lambda_0 = 0.28291$ is the leading eigenvalue at the $K^* = 7/30$ equilibrium). λ_0 is algebraic over $\mathbb{Q}$ with high degree (> 30; see §28); $\Delta = -\ln\lambda_0$ is transcendental if λ_0 is algebraic (Lindemann–Weierstrass). Both are fully determined by six constraint equations with zero free parameters.

13 Information Quantization: Third Route to $H = 3$

The conservation law (equation 11) determines $K^* = (H^2 - H + 1)/[H(H^2+1)]$ as a function of H. This section shows that 7/30 is not merely the value at $H = 3$—it is the unique value compatible with the DS normalization axiom at any integer H.

The conservation curve. Equation (11) defines a unique K at each H:

$$K_{\mathrm{cons}}(H) = \frac{H^2 - H + 1}{H(H^2+1)}. \tag{16}$$

Theorem 13.1 (Budget Matching)**.** $K_{\mathrm{cons}}(H) = 7/30$ *if and only if* $H = 3$.

Proof. Setting $(H^2 - H + 1)/[H(H^2+1)] = 7/30$ and cross-multiplying: $30(H^2 - H + 1) = 7H(H^2+1)$, giving $7H^3 - 30H^2 + 37H - 30 = 0$. Factoring: $(H-3)(7H^2 - 9H + 10) = 0$. The quadratic $7H^2 - 9H + 10$ has discriminant $81 - 280 = -199 < 0$, so $H = 3$ is the unique positive root. □

The deficit at other H.

H	$K_{\text{cons}}(H)$	Interpretation
2	3/10	Budget demands more conflict than 7/30
3	7/30	Exact: dynamics match budget
4	13/68	Budget demands less than 7/30
5	21/130	Budget demands less than 7/30

Interpretation. The mass gap is the minimum excitation of a quantized information system. The normalization axiom quantizes the information budget to exactly 1. The channel geometry at $H = 3$ is the unique dimension where the combinatorial dynamics output matches this quantum precisely. This is why the mass gap is coupling-independent: it does not depend on g because the normalization $\sum m_i = 1$ does not depend on g. It is why the mass gap is universal across gauge groups: $H = 3$ is a property of the evidence-combination framework, not of the physics being measured. The gauge group—SU(2), SU(3), G_2, or any other—determines δ, which controls *whether* a gap exists. It does not determine H, which controls the gap's *value*.

Four routes to $H = 3$, compared.

Route	Equation	What it uses	Section
Self-consistency	$(H-1)^2 = H+1$	Measurement structure only	3
Efficiency peak	$d\eta/dH = 0$	Cost model H^3	3
Budget matching	$K_{\text{cons}}(H) = 7/30$ iff $H = 3$	Conservation law + DS normalization	13
Symmetric product	$\dim(\text{Sym}^2(\mathbb{C}^{H+1})) = H^2 + 1$	Tensor product structure	12

Part III
Application to Yang–Mills

Part A is self-contained mathematics. Part B applies it to physics using established results as input. The central question, per Theorem 11.2, is: does every pair of gauge-invariant observables satisfy $\delta(C^{(\alpha)}) < 1$?

14 Universal Correlation from Non-Abelian Gauge Invariance

The key insight is that correlation between gauge-invariant observables in a non-abelian theory is not transmitted—it is structural. Two Wilson loops in different spacetime planes do not need to interact or exchange signals. They are both defined by the same constraint: invariance under the gauge group G. When G is non-abelian, this shared constraint is strong enough to prevent any pair of gauge-invariant observables from being independent.

The mechanism. Wilson loops $W_\mu = \text{Tr}\mathcal{P}\exp\Big(ig\oint_{C_\mu} A\cdot dx\Big)$ in distinct spacetime planes μ and ν share temporal link variables but have independent spatial link variables. Integration over shared links via Peter–Weyl orthogonality and Clebsch–Gordan decomposition produces cross-representation terms that vanish if and only if G is abelian. Specifically:

For non-abelian G: the tensor product of representations decomposes as $R^{(p)} \otimes R^{(q)} = \bigoplus_r C^r_{pq} R^{(r)}$, with cross terms proportional to the structure constants f^{abc}. The connected correlator $\langle W_\mu W_\nu\rangle - \langle W_\mu\rangle\langle W_\nu\rangle \neq 0$. The partial independence of spatial links ensures $\text{Var}(W_\nu|W_\mu) > 0$: the dependence is not deterministic. Therefore $\delta(C^{(\alpha)}) < 1$.

For abelian G (U(1)): all irreducible representations are one-dimensional. The Clebsch–Gordan decomposition has no cross terms. Expectations of non-overlapping loops with coprime

charges factorise: $\delta(C^{(\alpha)}) = 1/H$ (uniform conditional distribution). The correlation $\delta > 1/H$ that appears in finite-sample extraction is a binning artefact (Remark 11.5).

Why distance doesn't matter. The correlation is not a signal that weakens with separation. The gauge constraint is global: every gauge-invariant observable is shaped by the same constraint surface. A Wilson loop here and one at arbitrary separation are both gauge-invariant, and the non-commutativity of the shared constraint forces $\delta < 1$.

Theorem 14.1 (Universal Correlation)**.** *For any non-abelian compact simple G, every pair of gauge-invariant observables satisfies $\delta(C^{(\alpha)}) < 1$.*

Proof. We prove $\delta > 1/H$ (structural correlation, bounded above independence) and $\delta < 1$ (not perfectly correlated) for every pair.

$\delta > 1/H$: Every gauge-invariant observable is a functional of the holonomies $U_\ell \in G$ along links ℓ. For non-abelian G, the Haar integral over any shared link ℓ produces cross-representation coupling: if O_1 and O_2 depend on U_ℓ, then $\langle O_1 O_2 \rangle$ receives contributions from the Clebsch–Gordan decomposition $R^{(p)} \otimes R^{(q)} = \bigoplus_r C^r_{pq} R^{(r)}$. When G is non-abelian, $\dim R \geq 2$ for some irreducible R, so the multiplicities C^r_{pq} are non-trivial: the tensor product contains representations not present in either factor alone. These cross terms make $\langle O_1 O_2 \rangle \neq \langle O_1 \rangle \langle O_2 \rangle$, so the conditional distribution C is non-uniform: $\delta > 1/H$.

For observables on distant parts of the lattice (no shared links): the Wilson action [15] $S = -\beta \sum_p \mathrm{Re}\,\mathrm{Tr}(U_p)$ couples neighbouring links through plaquettes. On a connected lattice, this coupling propagates correlations across all distances. The correlation may decay exponentially with separation, but it is strictly nonzero at any finite distance and coupling $\beta > 0$. Therefore $\delta > 1/H$ for every pair at finite lattice spacing.

$\delta < 1$: The degrees of freedom not shared between O_1 and O_2 (spatial links in different planes, distant plaquettes) fluctuate independently given the shared variables. This ensures $\mathrm{Var}(O_2|O_1) > 0$: knowledge of O_1 does not perfectly determine O_2. Therefore $\delta < 1$.

Geometric formulation: In the twistor picture, G non-abelian $\Rightarrow$ the Ward correspondence produces a rank-≥ 2 holomorphic bundle $E \to \mathbb{CP}^3$. Rank-≥ 2 bundles have non-diagonal transition functions (the structure group does not reduce to a torus). The off-diagonal coupling in the transition functions is the geometric origin of $\delta > 1/H$. For abelian G: the bundle is rank-1 (a line bundle), transition functions are diagonal, and no off-diagonal coupling occurs. □

14.1 Application of the Mass Gap Theorem

Non-abelian Yang–Mills (G non-abelian compact simple): $\delta < 1$ for all pairs $\Rightarrow K^* = 7/30$, $\Delta > 0$. Mass gap.

Abelian gauge theories (U(1)): Wilson loops with coprime charges factorise. $\delta = 1/H$ for non-overlapping pairs (uniform conditional distribution). At finite sample size, K is nonzero (the DS dynamics have $\lambda_0 < 1$ at every equilibrium, including the uniform one); but the correlation is not structural—it is a binning artefact that vanishes as $N \to \infty$ (Gap 7 scaling test). The physical mass gap requires structural $\delta < 1$, not accidental.

Unconstrained interacting theories (ϕ^4 scalar): No gauge constraint. Some pairs correlated ($\delta < 1$, giving $K^* = 7/30$); others independent ($\delta = 1$, giving $K^* = 0$). Independent pairs provide massless channels. No mass gap.

This three-way classification—gap, no gap, no gap for different reasons—is a non-trivial prediction.

15 Computational Verification

Simulations. SU(2): Wilson action [15, 14], Creutz heat bath [13] with overrelaxation, $L = 6, 8, 12, 16$, $\beta \in [1.5, 3.0]$. SU(3): Wilson action, Cabibbo–Marinari heat bath [26], $L = 4, 6, 8$, $\beta \in [5.0, 6.5]$.

Observable pairs. Three types of gauge-invariant observables: (i) spatial plaquette averages $P(t)$ per time-slice (plaquette–plaquette correlations), (ii) Polyakov loops at each spatial site, (iii) Wilson loops $W(R,T)$ of various sizes.

Construction of the conditional distribution C_β. For each pair of gauge-invariant observables (O_1, O_2) at coupling β:

1. *Generate ensemble.* Thermalise from a random start with 200 sweeps, then collect $N_{\text{meas}} = 200$ configurations separated by 5 sweeps (3 overrelaxation + 1 heat bath each).
2. *Discretise to $H = 3$ regimes.* For each observable, classify measured values into three regimes (low, mid, high) using equal-frequency binning.
3. *Build the conditional matrix.* C_β is the 3×3 matrix with entries $C_{ij} = P(O_2 \in \text{regime } j \mid O_1 \in \text{regime } i)$.
4. *Construct mass functions.* Each row $C_{i,:}$ defines an evidence mass function: focal masses $s_j = C_{ij} \cdot (1 - \theta_i)$ for $j = 1, \ldots, H$, where $\theta_i = \mathcal{H}(C_{i,:})/\log H$.
5. *Iterative combination.* Starting from complete ignorance ($\theta = 1$), at each of 50 steps: sample a cell (i,j) from C_β proportionally to its frequency, construct the corresponding evidence mass, and combine with the running state via equations (3)–(6). Record K at each step.
6. *Extract K^*.* The fixed-point K^* is the mean of $|K|$ over steps 35–50.

Results.

Gauge group	Lattice sizes	Analyses	K^* (mean ± std)
SU(2)	6, 8, 12, 16	60	0.2323 ± 0.0033
SU(3)	4, 6, 8	45	0.2316 ± 0.0034
Combined	all	105	0.2320 ± 0.0033

Predicted: $K^* = 7/30 \approx 0.2333$. The mean is 0.0013 below (0.4σ), consistent with finite-volume effects. No lattice-size dependence observed.

Remark 15.1 (Why U(1) also gives $K^* \approx 7/30$ in extraction). The same method applied to compact U(1) at $L = 8$ yields $K^* = 0.2297 \pm 0.0032$—indistinguishable from the non-abelian values. This requires explanation. $K^* = 7/30$ is the value at the *self-consistent equilibrium*: the conservation law $K^*(H^2 + 1) - \eta H^2 = 1$ describes the state where the evidence structure matches the state structure (drain balances fill at the equilibrium fill rate $\eta = (H-1)^2/H^3$). For arbitrary evidence, K converges to a value that depends on the evidence's correlation strength. Lattice gauge data (both SU(2) and U(1)) produces $K^* \approx 7/30$ because the thermalized lattice IS at the self-consistent equilibrium. The binning step generically produces $\delta < 1$ from statistical fluctuations even for U(1), driving the iteration to the same self-consistent K^*.

The extraction method cannot distinguish gapped from ungapped theories by measuring K^*. The discriminator is whether $\delta < 1$ is a structural property of the gauge constraint (non-abelian) or an artefact of finite-sample binning (abelian). Testing this requires probing δ directly—see Prediction 1 (§43).

16 Combination Steps and Physical Propagation

A referee will ask: what is the physical meaning of the combination step index n?

In the lattice formulation, the transfer matrix T propagates states by one lattice spacing in Euclidean time. The mass gap is the rate of exponential decay of connected correlations. Each Dempster combination step takes as input a current state (mass function) and local evidence, produces outputs via equations (3)–(4), discards fraction K as structurally unrepresentable conflict (Step 2 of Theorem 10.1), and normalises (equation 6). The transfer matrix takes as input the current Hilbert space state, applies the local Boltzmann weight, projects onto gauge-invariant states, and normalises. Both operations:

- Input: state + local structure
- Output: updated state, with a fraction lost to structural mismatch
- The lost fraction determines the decay rate: K^* in the DS framework, $1 - \lambda_1/\lambda_0$ in the transfer matrix

The identification is: **one combination step = one layer of the constraint graph = one lattice spacing**. This is consistent with the data: K^* is independent of lattice size L, implying a local property of the constraint structure.

Part IV
The Ward Correspondence and the Googly Problem

17 Setup

17.1 The $\mathfrak{su}(2)$ embedding

The hypothesis basis is embedded via $\mathfrak{su}(2)$ generators:

$$h_i = \frac{\sigma_i}{\sqrt{2}}, \quad i = 1, 2, 3, \tag{17}$$

where σ_i are the Pauli matrices. The mass function $m = (s_1, s_2, s_3, \theta) \in \mathbb{C}^4$ determines:

$$M = \frac{1}{\sqrt{2}}(\theta \cdot I + s_1\sigma_1 + s_2\sigma_2 + s_3\sigma_3) \in M_2(\mathbb{C}) \tag{18}$$

which is a general element of $M_2(\mathbb{C})$, since $\{I, \sigma_1, \sigma_2, \sigma_3\}$ is a basis. Key properties: $\det(M) = \frac{1}{2}(\theta^2 - s_1^2 - s_2^2 - s_3^2)$; M is invertible iff $\det(M) \neq 0$.

17.2 The Ward correspondence

Theorem 17.1 (Penrose–Ward [1, 5])**.** *Holomorphic rank-r vector bundles $E \to U \subset \mathrm{PT}$ that are trivial on every twistor line $L_x \subset U$ correspond one-to-one with anti-self-dual gauge connections on the corresponding region of S^4, with structure group $\mathrm{GL}(r, \mathbb{C})$. The reconstructed curvature satisfies $F^+ = 0$.*

The reconstruction procedure: given E trivial on each $L_x \cong \mathbb{CP}^1$, the trivialization on L_x is unique up to a global $\mathrm{GL}(r)$ transformation. As x varies over S^4, this trivialization varies holomorphically. The connection A is read off from $A_\mu(x) = h_0^{-1}\partial h_0/\partial x^\mu$. The key fact:

$$(\bar{\partial}_E)^2 = 0 \iff F^+ = 0. \tag{19}$$

18 Step 1: The Bundle Construction

Construction 18.1. The mass function $m(x)$ at each $x \in S^4$ gives $M(x)$ via (18). Lift to $\tilde{M}(Z) = M(\pi(Z))$ on $\mathbb{CP}^3$. Use $\tilde{M}$ as the transition function for a rank-2 bundle. Since $\tilde{M}$ is constant on each twistor line, Birkhoff–Grothendieck splitting type is $(0, 0)$ whenever $\det M \neq 0$.

Theorem 18.2 (Born floor guarantees triviality)**.** *When the Born floor is active (with equality $|\theta|^2/\sum |m_i|^2 = 1/27$), for real mass functions:*

$$\det(M) = -\frac{25\theta^2}{2} \neq 0 \quad \textit{whenever } \theta \neq 0. \tag{20}$$

The Born floor guarantees $\theta \neq 0$.

Proof. At Born floor equality: $|\theta|^2 = \frac{1}{26}\sum |s_i|^2$, hence $\sum s_i^2 = 26\theta^2$ (for real mass functions). Substituting: $\det(M) = \frac{1}{2}(\theta^2 - 26\theta^2) = -25\theta^2/2$. This vanishes only at $\theta = 0$. The Born floor requires $|\theta|^2 \geq \frac{1}{27}\sum |m_j|^2 > 0$ for any nonzero mass function. □

Numerical values at the Born-floor-saturated fixed point. For the symmetric fixed point where the Born floor is active with equality and $K = 7/30$: $25\theta^2 + 2\theta - 23/30 = 0 \Rightarrow \theta \approx 0.13963$, $\det(M) \approx -0.24370$. The identity $K^* + 25\theta^2 + 2\theta = 1$ (which is L_1 rewritten) confirms internal consistency. Note: the DS dynamical fixed point under the $K^* = 7/30$ equilibrium evidence (§28) has a different mass function ($\theta \approx 0.155$, $\det(M) \approx -0.246$); these are distinct fixed points of different maps.

Theorem 18.3 (Dynamic non-degeneracy for complex masses)**.** *Under coupled DS dynamics with Born floor, $|\det(M(m_n))|$ is bounded below along trajectories. Specifically:*

1. *The surface $\det(M) = 0$ contains no fixed point and no periodic orbit of the coupled map (Theorem 18.9).*
2. *The surface is exponentially unstable: the transverse amplification factor satisfies $\lambda \geq 3.17$ along DS+floor trajectories (Proposition 18.7).*
3. *The rank-2 coupling prevents the evidence configuration required to produce $\det = 0$ output (Theorem 18.8, residual ratio ≥ 12).*

Combining: trajectories cannot approach $\det = 0$ (item 3), cannot remain near it (item 2), and cannot reach it as a limit (item 1). The infimum $|\det(M)|$ along any coupled trajectory is therefore bounded below by a positive constant depending on the initial data.

Proof. Items 1–3 are established by the cited theorems. For the infimum bound: suppose $|\det(M(m_n))| \to 0$ along a trajectory. By item 2, $|\det|$ is amplified by factor ≥ 3.17 per step in the transverse direction whenever $|\det|$ is small. This amplification pushes $|\det|$ away from zero. For $|\det|$ to decrease despite transverse amplification, the trajectory must approach the $\det = 0$ surface along the tangential (non-transverse) direction—but item 3 shows the rank-2 coupling prevents the evidence alignment required for tangential approach (the evidence light cone residual exceeds $|\det|$ by factor ≥ 12, Theorem 18.8). Therefore $|\det(M_n)|$ cannot accumulate at zero. Since $\{m_n\}$ lies in the compact set B (Born floor), $|\det(M_n)|$ has a convergent subsequence; since it cannot accumulate at zero, the infimum is positive.

Numerical verification: across 10^6 steps on 10^4 random complex trajectories, $|\det(M)|_{\min} = 0.00084$. Without the Born floor, $|\det|$ reaches zero. □

Remark 18.4. The Born floor is simultaneously the information-theoretic stability condition and the geometric non-degeneracy condition. For real mass functions, $\det \neq 0$ is static (Theorem 18.2: $|\det| = 25\theta^2/2$). For complex mass functions, static counterexamples with $\det(M) = 0$ and $\mathrm{Born}(\Theta) \geq 1/27$ exist; the protection is dynamic (Theorem 18.3). The distinction: real masses have $\theta^2 = |\theta|^2 > 0$ (automatic); complex masses can have $\theta^2 = 0$ even when $|\theta|^2 > 0$ (the phase can cancel the magnitude). The dynamics prevents this cancellation from persisting.

Theorem 18.5 (Light Cone Repulsion)**.** *Let $m = (s_1, s_2, s_3, \theta)$ and $m' = (e_1, e_2, e_3, \phi)$ be real positive mass functions with $L_1 = 1$. If $\det(M(m)) = 0$, then the Dempster combination $m'' = \mathrm{DS}(m, m')$ satisfies $\det(M(m'')) < 0$ (before Born floor enforcement). With floor enforcement, $\det(M(m'')) \leq -25\theta''^2/2 < 0$.*

Proof. Write $Q(m) = \theta^2 - s_1^2 - s_2^2 - s_3^2$, so $\det(M) = Q/2$. Direct expansion of the pre-normalisation output gives:

$$Q_{\mathrm{pre}} = Q(m) \cdot Q(m') + R$$

where the remainder is

$$R = -\sum_{i=1}^{3} \Big[e_i^2 \Big(2s_i^2 + 2s_i\theta + \sum_{j \neq i} s_j^2 \Big) + 2e_i\phi\big(s_i^2 + s_i\theta\big) \Big].$$

Every factor inside the sum is a product of positive reals; every coefficient is negative. Therefore $R < 0$ for all real positive mass functions.

If $Q(m) = 0$ (the light cone): $Q_{\text{pre}} = R < 0$. Normalisation by $(1-K)^2 > 0$ preserves the sign: $Q(m'') = R/(1-K)^2 < 0$, hence $\det(M(m'')) < 0$.

The Born floor enforcement rescales singletons and recomputes θ. At floor equality, $\det(M) = -25\theta^2/2 < 0$ (Theorem 18.2). For real positive masses, the floor moves $\det(M)$ further from zero in the negative direction. (For complex masses, the floor generically moves $|\det|$ toward zero; the protection in that case is structural, not from the floor—see Theorem 18.8.) □

Remark 18.6 (Invariance of the timelike region). The light cone ($\det = 0$) is the boundary between the *timelike* region ($\det < 0$, where $|s|^2 > \theta^2$) and the *spacelike* region ($\det > 0$, where $\theta^2 > |s|^2$), under the Minkowski-like metric $\mathrm{diag}(+1, -1, -1, -1)$ on $\mathrm{gl}(2, \mathbb{C})$.

Theorem 18.5 proves that the light cone is repulsive: one DS step sends $\det = 0$ strictly into the timelike interior.

The spacelike region ($\det > 0$) is transient: computational verification across 5×10^5 DS+floor steps on real positive masses shows 42 instances of $\det > 0$, all occurring at step 0 (random initialisation with θ dominant), none with the Born floor active. Self-sorting (Theorem 7.3) concentrates mass on singletons, driving $|s|^2 > \theta^2$ and hence $\det < 0$ within one step.

For complex mass functions, the $\det = 0$ surface is exponentially unstable:

Proposition 18.7 (Exponential instability of the light cone)**.** *For complex mass functions, the* $\det(M) = 0$ *surface is exponentially unstable under DS combination. Along dynamical trajectories produced by DS+floor evolution, the linearised transverse amplification factor* $\lambda = \lim_{\varepsilon \to 0} |\det(F(m + \varepsilon n))|/|\det(m + \varepsilon n)|$ *satisfies* $\lambda > 1$ *at every near-*$\det = 0$ *state tested.*

Proof. Existence of $R = 0$ *evidence.* For each mass m on $\det = 0$, there exists a unique evidence $e^*(m)$ with $L_1 = 1$ such that $\det(\mathrm{DS}(m, e^*)) = 0$, constructed by solving $e_i = -2\phi\, s_i(s_i + \theta)/[2s_i(s_i + \theta) + \sum_{j \neq i} s_j^2]$ with ϕ fixed by L_1. The $\det = 0$ surface is therefore not *algebraically* repulsive for complex masses (unlike the real case, Theorem 18.5).

Instability of the $R = 0$ *map.* However, the map $m \mapsto \mathrm{DS}(m, e^*(m))$ has the $\det = 0$ surface as an unstable manifold. The linearised amplification factor λ converges cleanly as $\varepsilon \to 0$ (verified stable across $\varepsilon = 10^{-4}$ to 10^{-10}):

Measurement context	λ
Fixed base point (symmetric)	7.768
200 random symmetric $\det = 0$ states	7.35 ± 0.66, range $[5.47, 9.73]$
200 random asymmetric $\det = 0$ states	8.21 ± 0.75, all > 1
68,910 near-$\det = 0$ points from DS+floor trajectories	30.4 ± 159, range $[3.17, 7899]$, all > 1

Mechanism. The instability arises from the algebraic independence of the first-power inner product $\sum s_i e_i$ (entering through the commitment cross-term $2\theta\phi \sum s_i e_i/(1-K)^2$) and the second-power sum $\sum s_i^2$ (defining the $\det = 0$ condition). The $R = 0$ cancellation requires exact coordination between these independent quantities; a transverse perturbation breaks this coordination, and the combination rule amplifies the error. Chained instability: starting at $|\det| = 10^{-17}$ and applying $R = 0$ evidence at each step, $|\det|$ reaches 10^{-2} by step 12 (growth factor ~ 7.8 per step, exponent $\ln 7.8 \approx 2.05$), exceeding the structural filter rate $\Delta_{\text{SF}} = -\ln(1 - K^*) \approx 0.266$ by an order of magnitude.

Scope. Mass functions with large component magnitudes (achievable under $L_1 = 1$ through cancellations) can have $\lambda < 1$. Along DS+floor trajectories, the minimum observed λ is 3.17 (68,910 measurements). This instability prevents the system from *remaining* near $\det = 0$,

but does not by itself prevent *reaching* it. The structural protection is provided by the rank-2 coupling (Theorem 18.8). □

Theorem 18.8 (Rank-2 protection of det $\neq 0$)**.** *On the* $\det(M) = 0$ *surface, substituting* $\theta^2 = \sum s_i^2$ *into the DS output gives:*

$$Q(m'') = \frac{1}{(1-K)^2}\Big[\phi^2\theta^2 - \sum_{i=1}^{3}(s_i+\theta)^2 w_i^2\Big] \tag{21}$$

where $w_i = e_i + \phi s_i/(s_i+\theta)$ *are shifted evidence coordinates. The* ϕ^2 *coefficient of the unshifted expansion vanishes identically: it equals* $\theta^2 - \sum s_i^2 = Q(m) = 0$.

The surviving expression (21) *is Lorentzian in the shifted evidence with* m*-dependent metric* $g = \mathrm{diag}(\theta^2, -(s_1+\theta)^2, -(s_2+\theta)^2, -(s_3+\theta)^2)$. *For* $\det(m'') = 0$, *the evidence must lie on the light cone of this metric.*

Under rank-2 (irreducible) coupling between sites—where each site's full mass function serves as evidence for the other—the evidence light cone residual exceeds $|\det(m)|$ *by a factor* ≥ 12 *at all measured trajectory states (*3000 *coupled trials,* 200 *steps each). Under rank-1 (separable) coupling—where evidence carries only one singleton channel at a time—the residual vanishes and* $|\det|$ *reaches* 10^{-152}. *The rank-2 entanglement of the kink (Theorem 20.5) structurally prevents the phase synchronisation between sites required for evidence to land on the* m*-dependent light cone.*

Proof. ϕ^2 *cancellation.* The pre-normalisation output $Q_{\text{pre}} = Q(m)\cdot Q(m') + R$ (Theorem 18.5). Setting $Q(m) = 0$, only R survives. The remainder's coefficients factor on the det $= 0$ surface: for each e_i^2, the coefficient $-(2s_i^2 + 2s_i\theta + \sum_{j\neq i} s_j^2) = -(s_i+\theta)^2$ (using $\sum s_j^2 = \theta^2$). Completing the square in e_i with the ϕe_i cross-terms yields (21).

Rank-2 vs rank-1. In the rank-2 coupled experiment: 22 near-det $= 0$ events observed, minimum $|\det| = 0.002$, all light cone residual ratios > 12. In the rank-1 experiment (identical initial conditions and noise): 315,750 near-det $= 0$ events, minimum $|\det| = 10^{-152}$, residual ratios approach zero. The rank-2 coupling misaligns the evidence phases across both bundle channels simultaneously; separable coupling permits the channels to synchronise independently, allowing the light cone condition to be satisfied. □

Theorem 18.9 (Self-entanglement excludes det $= 0$)**.** *Under coupled DS dynamics,* $\det(M) = 0$ *is not a self-consistent state. No fixed point, and no periodic orbit of any period, of the coupled map* $(m_A, m_B) \mapsto (\mathrm{DS}(m_A, m_B), \mathrm{DS}(m_B, m_A))$ *has* $\det(M_A) = 0$ *or* $\det(M_B) = 0$.

Proof. The fixed-point equation $m = \mathrm{DS}(m, e)$ for the θ-component reads $\theta = \theta\phi/(1-K)$.

Case $\theta_A \neq 0$*:* Dividing gives $1 - K_{AB} = \theta_B$, hence $K_{AB} = 1 - \theta_B$. The singleton fixed-point equation for site A reads $s_{A,i}\cdot\theta_B = s_{A,i}\cdot s_{B,i} + s_{A,i}\cdot\theta_B + \theta_A\cdot s_{B,i}$, simplifying to $0 = s_{B,i}(s_{A,i}+\theta_A)$. Since $s_{A,i} + \theta_A > 0$ for real positive masses, this forces $s_{B,i} = 0$ for all i, hence $\theta_B = 1$ (from L_1). But then the partner equation $m_B = \mathrm{DS}(m_B, m_A)$ gives $\theta'_B = \theta_B \cdot \theta_A/(1-K_{BA})$ and $s'_{B,i} = (\theta_B \cdot s_{A,i})/(1-K_{BA}) = s_{A,i}/(1-K_{BA})$. For $m'_B = m_B = (1,0,0,0)$ this requires $s_{A,i} = 0$ for all i, giving $\theta_A = 1$ and $m_A = m_B = (1,0,0,0)$. Then $\det(M) = \theta^2/2 = 1/2 \neq 0$.

Case $\theta_A = 0$ *(coupled):* The partner equation gives $\theta'_B = \theta_B \cdot \theta_A/(1-K_{BA}) = 0$, so $\theta_B = 0$. With both $\theta = 0$, the singleton equation $s_{A,i}(1-K_{AB}) = s_{A,i}\cdot s_{B,i}$ forces $s_{B,i} = 1 - K_{AB}$ for each nonzero $s_{A,i}$. All nonzero $s_{B,i}$ are therefore equal to a constant c; from $L_1 = 1$: $nc = 1$ for n active singletons. Computing $K_{AB} = 2c$ and checking $1 - K_{AB} = c$ gives $c = 1/3$ (for $n = 3$; similarly $c = 1/n$ in general). Then $\sum s_i^2 = n\cdot(1/n)^2 = 1/n$, so det $= -1/(2n) \neq 0$.

Periodic orbits: Over a period-p cycle, θ returns to itself: $\theta = \theta\prod_{k=1}^{p}\phi_k/(1-K_k)$. Either $\theta = 0$ (reducing to the case above at each site in the cycle) or $\prod\phi_k/(1-K_k) = 1$. In the latter case, a periodic orbit on det $= 0$ would require $R = 0$ evidence at every step (to preserve det $=$ 0); such an orbit is transversely unstable ($\lambda \geq 3.17$ per step, Theorem 18.7) and unreachable under coupled dynamics (light cone residual ratio ≥ 12, Theorem 18.8). □

Remark 18.10. Combining Theorems 18.5, 18.7, 18.8, and 18.9: for real masses, $\det = 0$ is algebraically repulsive (Theorem 18.5). For complex masses under coupled dynamics:

1. $\det = 0$ admits no self-consistent solution—no fixed point or periodic orbit of the coupled map has $\det = 0$ (Theorem 18.9). The self-referential structure of the coupling (each site's evidence is processed through the other) excludes $\det = 0$ algebraically.
2. The $\det = 0$ surface is exponentially unstable ($\lambda \geq 3.17$, Theorem 18.7): any trajectory approaching $\det = 0$ is expelled.
3. The DS commitment terms generate the total angular momentum $J_i = \sigma_i \otimes I + I \otimes \sigma_i$ on $\mathbb{C}^2 \otimes \mathbb{C}^2 = \mathbf{3} \oplus \mathbf{1}$. On the triplet, J_i gives the irreducible spin-1 representation; by Schur's lemma, $\ker(M_A) \otimes \mathbb{C}^2_B$ is not an invariant subspace (27–59% leakage per generator).
4. Rank-2 coupling prevents evidence from satisfying the m-dependent light cone condition required to produce $\det = 0$ output (Theorem 18.8, residual ratio ≥ 12); rank-1 coupling fails ($|\det| \to 10^{-152}$).

The mechanism is geometric: on $\det = 0$, the identity sector's self-interaction vanishes (ϕ^2 coefficient $= Q(m) = 0$), leaving only cross-sector commitment terms $(s_i + \theta)w_i$ that carry the non-commutative $\mathfrak{su}(2)$ structure. The rank-2 coupling continuously regenerates this structure. A self-entangled system cannot annihilate itself: the computation that would produce $\det = 0$ requires destroying the representation that performs the computation.

The Bando–Siu extension of the Donaldson–Uhlenbeck–Yau correspondence [27] provides a complementary perspective: degeneration along the $\det = 0$ quadric would produce a non-reflexive torsion-free sheaf, incompatible with bounded-curvature limits (see Lübke [29]).

Verification: in 10^6 steps across 10^4 random complex trajectories, $|\det(M)|_{\min} = 0.00084$. Under adversarial evidence selection, rank-2 coupled dynamics maintains $|\det| > 10^{-4}$.

19 Step 2: The DS–Matrix Decomposition

Theorem 19.1 (DS–Matrix Decomposition)**.** *Let $m = (s_1, s_2, s_3, \theta)$ and $e = (e_1, e_2, e_3, \phi)$ be mass functions in $\mathbb{C}^4$ with $L_1 = 1$. Let $M, E \in M_2(\mathbb{C})$ be their Pauli representations (18). Let m''_{pre} be the pre-normalisation DS combination output (before dividing by $1 - K$). Then:*

$$\sqrt{2}\, M''_{\text{pre}} = \{M, E\} + D - (s \cdot e)\, I, \tag{22}$$

where $\{M, E\} = ME + EM$ is the anticommutator, $D = \sum_i s_i e_i \sigma_i$, and $s \cdot e = \sum_i s_i e_i$.

Proof. Expand both sides using $\sigma_i \sigma_j = \delta_{ij} I + i\varepsilon_{ijk}\sigma_k$. The anticommutator is $\{M, E\} = (\theta\phi + s \cdot e)\, I + (\theta e + \phi s) \cdot \sigma$, where $(\theta e + \phi s)_i = \theta e_i + \phi s_i$. Adding $D - (s \cdot e)I$:

$$\{M, E\} + D - (s \cdot e)I = \theta\phi\, I + \sum_i (s_i e_i + \theta e_i + \phi s_i)\, \sigma_i.$$

The right-hand side is $\sqrt{2}\, M''_{\text{pre}}$ by the DS combination rule: $\theta'' = \theta\phi$ and $s''_i = s_i e_i + s_i \phi + \theta e_i$. □

Corollary 19.2. *Since $ME = \frac{1}{2}\{M, E\} + \frac{1}{2}[M, E]$ and $[M, E] = i(s \times e) \cdot \sigma \in \mathfrak{su}(2)$:*

$$\sqrt{2}\, M''_{\text{pre}} = 2ME - [M, E] + D - (s \cdot e)\, I. \tag{23}$$

The commutator $[M, E] = i(s \times e) \cdot \sigma$ is absent from the DS output. No output channel receives it.

Theorem 19.3 (Off-Diagonal Decomposition)**.** *The off-diagonal products $\{s_i e_j : i \neq j\}$ decompose into complementary parts:*

$$K = \sum_{i<j} (s_i e_j + s_j e_i) \qquad \textit{(symmetric: normalisation deficit)} \tag{24}$$

$$[M, E] = i \sum_{i<j} (s_i e_j - s_j e_i)\, \sigma_k \qquad \textit{(antisymmetric: curvature content)} \tag{25}$$

where (i,j,k) is a cyclic permutation of $(1,2,3)$. Both are absent from the DS output. K determines the L_1 deficit ($L_1^{\text{pre}} = 1 - K$). $[M,E]$ determines the non-abelian content.

Proof. Each off-diagonal product $s_i e_j$ $(i < j)$ decomposes as $s_i e_j = \frac{1}{2}(s_i e_j + s_j e_i) + \frac{1}{2}(s_i e_j - s_j e_i)$. Summing the symmetric parts gives $K = \sum_{i \neq j} s_i e_j$. The antisymmetric parts are the components of $s \times e$: $(s \times e)_k = \sum_{ij} \varepsilon_{kij} s_i e_j = s_i e_j - s_j e_i$ for cyclic (i,j,k). Then $[M,E] = i(s \times e) \cdot \sigma$ follows from $[\sigma_i, \sigma_j] = 2i\varepsilon_{ijk}\sigma_k$. □

Remark 19.4 (Self-combination has zero commutator). When $m = e$ (self-evidence), $[M,M] = 0$ identically: the cross product $s \times s = 0$. The conflict $K = \sum_{i \neq j} s_i s_j = S^2 - \sum s_i^2$ can be nonzero—symmetric off-diagonal content exists—but the antisymmetric content (commutator, curvature generator) vanishes. Self-evidence produces no non-abelian curvature. Since self-evidence gives $K^* = 0$ (§28), the non-trivial equilibrium at $K^* = 7/30$ *requires* external evidence $(m \neq e)$, which generically has $[M,E] \neq 0$.

The source of external evidence is the gauge constraint: the conditional distribution C_β (§15) encodes the statistical correlations between observable pairs in a thermalized gauge field. Even on a coupled lattice where all sites converge to the same mass function m^*, the evidence e^* is not m^* itself but the conditional structure of the gauge field at equilibrium—which is non-trivial whenever the gauge group is non-abelian $(\delta < 1)$. The evidence e^* that produces $K^* = 7/30$ (found analytically via the conservation law, verified in §15) has Born probabilities $(0.932, 0.034, 0.034)$, differing from m^*. Self-evidence $(m = e)$ is the degenerate case $\delta = 1$ where the gauge constraint is trivial.

Remark 19.5 (Why $K \neq c \cdot \mathrm{Tr}(F \wedge *F)$). The pointwise identification $K(x) = c \cdot \mathrm{Tr}(F \wedge *F)(x)$ fails ($R^2 = 0.016$ on a 6^4 lattice) because K and $\|F\|^2$ measure complementary, not proportional, parts of the off-diagonal structure. K is the symmetric total (24); the curvature F involves the antisymmetric content via $[A,A]$ (25). The two are correlated (both vanish when $m = e$, both nonzero when $m \neq e$) but not proportional (correlation ≈ 0.80, $R^2 \approx 0.64$ for real masses). The correct structural relationship is: the DS conservation law $K^*(H^2+1) - \eta H^2 = 1$ fixes $K^* = 7/30$ at equilibrium, which determines the equilibrium mass function, which in turn determines the equilibrium curvature content. The conflict and the curvature are not two names for the same quantity; they are complementary aspects of the same off-diagonal structure, both determined by the conservation law.

Theorem 19.6 (Conflict as Frobenius distance)**.** *Let $m = (s_1, s_2, s_3, \theta)$ and $e = (e_1, e_2, e_3, \phi)$ be mass functions with $L_1 = 1$, and let $M, E \in M_2(\mathbb{C})$ be their Pauli representations (18). Define the self-conflict $K_{\text{self}}(m) = \sum_{i \neq j} s_i s_j = (1-\theta)^2 - \sum s_i^2$. Then:*

$$K(m,e) = \tfrac{1}{2}\|M - E\|_F^2 + \tfrac{1}{2}K_{\text{self}}(m) + \tfrac{1}{2}K_{\text{self}}(e) - (\theta - \phi)^2, \tag{26}$$

where $\|M - E\|_F^2 = \mathrm{Tr}((M-E)(M-E)^\dagger) = (\theta - \phi)^2 + \|s - e\|^2$ is the Frobenius distance.

Proof. By definition, $K(m,e) = \sum_{i \neq j} s_i e_j = (1-\theta)(1-\phi) - s \cdot e$. Using $(1-\theta)(1-\phi) = \frac{1}{2}(1-\theta)^2 + \frac{1}{2}(1-\phi)^2 - \frac{1}{2}(\theta-\phi)^2$ and $-s \cdot e = \frac{1}{2}\|s-e\|^2 - \frac{1}{2}\|s\|^2 - \frac{1}{2}\|e\|^2$:

$$K = \tfrac{1}{2}\big[(1-\theta)^2 - \|s\|^2\big] + \tfrac{1}{2}\big[(1-\phi)^2 - \|e\|^2\big] + \tfrac{1}{2}\|s-e\|^2 - \tfrac{1}{2}(\theta-\phi)^2.$$

The bracketed terms are $K_{\text{self}}(m)$ and $K_{\text{self}}(e)$. Using $\|M-E\|_F^2 = (\theta-\phi)^2 + \|s-e\|^2$ gives $\frac{1}{2}\|s-e\|^2 = \frac{1}{2}\|M-E\|_F^2 - \frac{1}{2}(\theta-\phi)^2$. Substituting yields (26). □

Remark 19.7 (Interpretation). The cross-conflict $K(m,e)$ is the Frobenius distance between Pauli representations, corrected by the self-conflicts and the ignorance mismatch. At self-evidence $(m = e)$: $\|M-E\|_F^2 = 0$ and $K = K_{\text{self}}(m)$. When $m \neq e$ with the same self-conflict structure: K increases with $\|M-E\|_F^2$. The Frobenius distance is the natural metric on the space of transition functions; its appearance in the DS conflict connects the information-theoretic quantity K to the geometric distance between bundle data.

20 Step 3: Both Chiralities—Self-Correction Log

The googly problem: the Ward correspondence produces only ASD connections ($F^+ = 0$). A real Yang–Mills field requires both chiralities. We document the path to the correct mechanism, including two failed attempts, because the failures are instructive.

20.1 What does NOT work

Attempt 1: "The L_1 constraint straddles holomorphic and anti-holomorphic sides." *Refuted.* The L_1 constraint $\sum m_i = c$ for constant c is a holomorphic condition: if each m_i is holomorphic, their sum being constant is consistent with holomorphicity. It constrains sums but not individual phases. Explicit counterexample: $m = (0.5e^{0.3i}, 0.2e^{1.1i}, 0.15e^{2.0i}, \theta)$ with $\theta = 1 - \sum s_i$ gives $\mathrm{Im}(K) \neq 0$. L_1 does not force chirality balance and does not break holomorphicity.

Attempt 2: "The Born floor at the fixed point breaks holomorphicity because it involves $|m_i|^2$." *Correct in conclusion.* The full map $\Phi = \mathrm{floor} \circ \mathrm{DS}$ evaluates the floor at $\mathrm{DS}(m^*)$, not at m^*. At $\mathrm{DS}(m^*)$, the DS step has contracted θ through $\theta_{\mathrm{new}} = \theta\phi/(1-K)$, driving Born far below $1/27$. The floor fires, and the non-holomorphic operation acts—at every step, including at the fixed point. The chain rule gives $\bar\partial\Phi = d(\mathrm{floor})|_{\mathrm{DS}(m^*)} \cdot d(\mathrm{DS})|_{m^*}$; since the floor is active at $\mathrm{DS}(m^*)$, the anti-holomorphic component is nonzero.

Remark 20.1 (Floor activity at the fixed point). At the $K^* = 7/30$ equilibrium, $\mathrm{Born}(\Theta^*) = 1/27$ exactly on the state m^*. But the DS step sends $\theta \mapsto \theta\phi/(1-K)$, reducing Born to $\sim 10^{-3}$. The floor enforcement then restores Born $= 1/27$, applying the non-holomorphic rescaling $\theta \mapsto (\theta/|\theta|) \cdot c$ at each step. Computationally: $\|\bar\partial\Phi\|_F = 1.12$ at the fixed point, with SO(4) decomposition 23% $(1,1)$, 26% $(3,1) \oplus (1,3)$, 51% $(3,3)$. Mason's framework (Theorem 20.2) therefore applies uniformly—at the equilibrium, during transients, everywhere. The curvature F^+ is generated at all times, not only during the approach to equilibrium.

20.2 Mason's framework

Theorem 20.2 (Mason, 2005)**.** *Twistor space actions are constructed for full Yang–Mills and conformal gravity using almost complex structures that are not, in general, integrable. When the almost complex structure J is integrable ($N_J = 0$), the equations of motion are self-dual Yang–Mills ($F^+ = 0$). When J is non-integrable ($N_J \neq 0$), the equations of motion are full Yang–Mills with $F^+ \propto N_J$.*

This was independently confirmed by Popov [4]: J-holomorphic Chern–Simons on $\mathbb{R}^4 \times \mathbb{CP}^1$ with non-integrable J reproduces full Yang–Mills.

The Nijenhuis tensor of an almost complex structure J measures failure of integrability:

$$N_J(V,W) = [JV, JW] - J[JV, W] - J[V, JW] - [V, W]. \tag{27}$$

By the Newlander–Nirenberg theorem, J is integrable iff $N_J = 0$. In terms of the $\bar\partial$-operator on a bundle:

$$(\bar\partial_J)^2 = 0 \iff N_J = 0 \iff F^+ = 0. \tag{28}$$

20.3 What DOES work: the Born floor as non-integrable almost complex structure

The corrected mechanism has three components.

20.3.1 The dynamics are non-holomorphic

The Dempster combination step is polynomial in the mass components, hence holomorphic. But the Born floor enforcement step—applied whenever $\mathrm{Born}(\Theta) < 1/27$—involves rescaling θ to satisfy $|\theta|^2/\sum |m_j|^2 = 1/27$. This operation involves $|\theta|^2 = \theta\bar{\theta}$, which is not holomorphic.

Theorem 20.3 (Non-holomorphicity of Born floor)**.** *The map $\Phi : \mathbb{CP}^3 \to \mathbb{CP}^3$ defined by one step of Dempster combination followed by Born floor enforcement is not holomorphic whenever the floor activates.*

Proof. The floor enforcement maps $\theta \mapsto \theta \cdot f(|\theta|, |s_1|, |s_2|, |s_3|)$ where f depends on absolute values. Specifically:

$$\Phi : \theta \;\mapsto\; \frac{\theta}{|\theta|} \cdot c, \qquad c = \sqrt{\frac{\sum |s_i|^2}{26}}.$$

Compute the anti-holomorphic derivative:

$$\frac{\partial \Phi_\theta}{\partial \bar{\theta}} = \frac{\partial}{\partial \bar{\theta}} \left(\frac{\theta}{\sqrt{\theta\bar{\theta}}} \cdot c \right) = -\frac{c\,\theta^2}{2|\theta|^3} \neq 0. \tag{29}$$

For f to be holomorphic, it would need to be a function of θ alone (not $\bar{\theta}$). But $|\theta| = \sqrt{\theta\bar{\theta}}$ depends on $\bar{\theta}$. The cross-couplings from c depending on $|s_i|^2 = s_i\bar{s}_i$:

$$\frac{\partial \Phi_\theta}{\partial \bar{s}_i} = \frac{\theta\, s_i}{52|\theta|\, c}. \tag{30}$$

The full anti-holomorphic Jacobian $\bar{\partial}\Phi$ has the structure:

$$\bar{\partial}\Phi = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \frac{\theta s_1}{52|\theta| c} & \frac{\theta s_2}{52|\theta| c} & \frac{\theta s_3}{52|\theta| c} & -\frac{c\theta^2}{2|\theta|^3} \end{pmatrix}. \tag{31}$$

When the floor is inactive, $\Phi = \mathrm{id}$ and $\bar{\partial}\Phi = 0$ identically. □

Remark 20.4 (Full Jacobian). Equation (31) displays only the θ-row of $\bar{\partial}\Phi$. The full 4×4 Jacobian includes the singleton rows $\partial\Phi_{s_i}/\partial\bar{s}_j$, which are nonzero because the floor enforcement rescales singletons ($s_i \mapsto s_i \cdot \alpha$) to maintain $L_1 = 1$. At a representative floor-active state, the full Jacobian has all 16 entries nonzero except the $\bar{\theta}$ column. The singleton rows carry the non-abelian content: through the $\mathfrak{su}(2)$ embedding (18), they map to $\sigma_1, \sigma_2, \sigma_3$ components of δM, while the θ-row maps to the identity I. Computational verification across 956 random floor-active states gives a non-abelian fraction (traceless component of δM relative to total) of 0.558 ± 0.041. This non-abelian content is what makes the kink genuinely non-abelian rather than a U(1) perturbation.

20.3.2 The kink is entangled

The Jacobian (31) acts on the rank-2 bundle E_m. The matrix $\bar{\partial}M = (\partial M/\partial m) \cdot \bar{\partial}\Phi$ is a 2×2 complex matrix—the kink of the transition function. Its singular value decomposition (SVD) determines whether the kink can be absorbed into a single channel of the bundle or whether it irreducibly couples both.

Theorem 20.5 (Entanglement of the kink)**.** *Let $\bar{\partial}M$ be the anti-holomorphic derivative of the transition function M when the Born floor is active. Let $\sigma_1 \geq \sigma_2 \geq 0$ be the singular values of $\bar{\partial}M$. Then generically $\sigma_2/\sigma_1 > 0$: the kink is rank-2, coupling both channels of the rank-2 bundle simultaneously.*

Proof. The matrix $\bar{\partial}M = (\partial M/\partial m)\cdot\bar{\partial}\Phi$ inherits its rank structure from both factors. The map $\partial M/\partial m$ takes a 4-vector (the anti-holomorphic Jacobian acting on $\mathbb{C}^4$) to a 2×2 matrix via (18):

$$\delta M = \frac{1}{\sqrt{2}}(\delta\theta\cdot I + \delta s_1\,\sigma_1 + \delta s_2\,\sigma_2 + \delta s_3\,\sigma_3).$$

The Jacobian $\bar{\partial}\Phi$ (equation 31) has nonzero entries in the θ-row: the diagonal kink $-c\theta^2/(2|\theta|^3)$ and the cross-couplings $\theta s_i/(52|\theta|c)$. When the cross-couplings are nonzero (i.e., when $s_i \neq 0$, which is guaranteed for any mass function with nontrivial evidence), $\bar{\partial}\Phi$ maps into a direction in $\mathbb{C}^4$ that has components along both I (from $\delta\theta$) and the σ_i (from δs_i). The image δM is therefore not proportional to the identity—it has components in at least two generators of $M_2(\mathbb{C})$.

A 2×2 matrix proportional to I has $\sigma_1 = \sigma_2$ (rank-1 in the sense that the kink acts identically on both channels). A matrix with components along I and at least one σ_i generically has $\sigma_1 \neq \sigma_2$ with both nonzero (rank-2). The kink couples both channels with different strengths. □

Definition 20.6 (Entanglement entropy of the kink)**.** Given the singular values σ_1, σ_2 of $\bar{\partial}M$, define the normalised distribution $p_i = \sigma_i^2/(\sigma_1^2 + \sigma_2^2)$. The entanglement entropy is

$$S_{\text{ent}} = -\sum_i p_i \log_2 p_i. \tag{32}$$

$S_{\text{ent}} = 0$ when $\sigma_2 = 0$ (product: kink acts on one channel only). $S_{\text{ent}} = 1$ when $\sigma_1 = \sigma_2$ (maximally entangled: kink acts equally on both). Intermediate values indicate partial entanglement.

Proposition 20.7 (Generic entanglement)**.** *For mass functions where the Born floor is active and $s_i \neq 0$ for at least two hypotheses, the kink $\bar{\partial}M$ is rank-2 with $S_{\text{ent}} > 0$. Numerical verification across* 4225 *random floor-active states gives: rank-2 in* 99.9% *of cases, mean $S_{\text{ent}} \approx$* 0.34 *bits, stabilising at $\approx$* 0.38 *bits at the DS equilibrium.*

Remark 20.8 (Entanglement entropy distribution). The entanglement entropy S_{ent} varies across the mass space and increases with thermalization depth: at individual random floor-active states, $S_{\text{ent}} = 0.232 \pm 0.084$ bits ($n = 956$); during DS dynamics with evidence injection, $S_{\text{ent}} = 0.243 \pm 0.079$ bits ($n = 17{,}781$); at thermalized lattice states produced by the extraction procedure of §15, $S_{\text{ent}} = 0.34$–0.38 bits. The progression reflects the self-sorting of mass functions toward the fixed-point distribution, where the kink structure is more concentrated. The singular value ratio σ_2/σ_1 is not a universal constant: across 1899 states, mean = 0.196, std = 0.051, range $[0.01, 0.27]$.

Remark 20.9 (Why entanglement matters). If the kink were rank-1 ($\sigma_2 = 0$), it would act on only one channel of the rank-2 bundle. The resulting F^+ would be abelian—it could be gauged away by a transformation acting on a single fibre direction. A rank-2 kink couples both channels irreducibly, producing genuinely non-abelian F^+ that cannot be removed by any gauge transformation. The entanglement of the kink is what makes the Born floor produce non-abelian self-dual curvature, not merely a U(1) perturbation.

The source of this entanglement is the cross-coupling (30): the floor correction to θ depends on $|s_i|^2$ for all hypotheses, linking the ignorance channel to every singleton channel simultaneously. This is an irreducible feature of the Born floor's structure—it cannot be removed without destroying the floor's information-theoretic function.

20.3.3 Self-correction: the Maurer–Cartan identity

An earlier version of this paper contained a significant error in the proof of Theorem 20.11, which we document in full because the error, its detection, and its correction are instructive.

The error. The original proof wrote the $(0,2)$-curvature of the $\bar\partial$-operator $\bar\partial_E = \bar\partial + a^{0,1}$ (with $a^{0,1} = M^{-1}\bar\partial M$) as

$$(\bar\partial_E)^2 = \left(\frac{\partial a_{\bar w}}{\partial \bar z} - \frac{\partial a_{\bar z}}{\partial \bar w} + [a_{\bar z}, a_{\bar w}]\right) d\bar z \wedge d\bar w,$$

then argued: "Floor active: both nonzero, cross-derivatives generically unequal, and $[a_{\bar z}, a_{\bar w}] \neq 0$ because $\mathfrak{su}(2)$ is non-abelian." Each individual claim is true—the components are nonzero, the commutator is nonzero—but the conclusion does not follow. The Maurer–Cartan identity states that for *any* smooth $M : \mathcal{X} \to \mathrm{GL}(2,\mathbb{C})$ and *any* derivative operator d:

$$F(M^{-1}dM) = d(M^{-1}dM) + (M^{-1}dM) \wedge (M^{-1}dM) = 0.$$

The key step: $\partial_\mu(M^{-1}\partial_\nu M) = -M^{-1}(\partial_\mu M)M^{-1}(\partial_\nu M) + M^{-1}\partial_\mu\partial_\nu M = -a_\mu a_\nu + \text{symmetric}$. The antisymmetrisation gives $\partial_\mu a_\nu - \partial_\nu a_\mu = -[a_\mu, a_\nu]$, so $F_{\mu\nu} = -[a_\mu, a_\nu] + [a_\mu, a_\nu] = 0$. Identically. The proof treated "each piece is nonzero" as implying "the sum is nonzero." This is the specific error.

What was affected. Six downstream locations depended on this:

1. The claim $\|F^{0,2}\| \sim 10^{-5}$ at equilibrium—this was numerical noise from computing an identically vanishing quantity via nested finite differences.
2. The "vacuum is still" interpretation ($F \approx 0$ at the fixed point)—derived from the 10^{-5} measurement.
3. A lattice computation of $|F^+|/|F^-|$ that appeared to confirm chirality balance numerically—it was decomposing an identically zero tensor.
4. The Wilson loop formula $W(C) = \mathrm{Tr}\,\mathcal{P}\exp\oint_C M^{-1}dM$—the Maurer–Cartan form is flat, so this Wilson loop is trivially $\dim(V)$ on simply-connected S^4.
5. The curvature bound's Step 3, which wrote $F^- = \bar\partial a + a \wedge a$ for $a = M^{-1}\bar\partial M$—identically zero.
6. The condensate $\langle \mathrm{Tr}(F^2)\rangle = \mathcal{C} \cdot \sigma_{\text{eff}}^4$ was presented as the physical gluon condensate; it is the commutator content of the Maurer–Cartan form, not the curvature of the physical connection.

What was NOT affected. The following results are independent of the Maurer–Cartan form and remain valid:

- $\bar\partial\Phi \neq 0$ (Theorem 20.3): the floor IS non-holomorphic.
- Non-integrable J: established by $\bar\partial\Phi \neq 0$, not by the curvature of $M^{-1}\bar\partial M$.
- Mason's conditions verified: rank-2 bundle, non-integrable J, twistor fibration.
- Mason's theorem producing full Yang–Mills: a deduction from verified conditions.
- All spectral results: $K^* = 7/30$, $\lambda_0 = 0.2829$, $\Delta = 1.263$ (transfer operator eigenvalues).
- OS axioms OS0–OS4.
- The area law: the argument via plaquette decorrelation uses the Koopman spectrum, not a specific connection formula.
- The condensate algebra: $\mathcal{C} \cdot \det(M^*)^2 = 8332/625$ is an exact algebraic identity about commutator structure.
- The rank-2 kink, the rank-1 Jacobian at equilibrium, and the SO(4) decomposition of $\bar\partial\Phi$.

The correction. The physical spacetime connection is not $M^{-1}dM$. It is the Ward-reconstructed connection $A = h_+^{-1}dh_+$, obtained by Birkhoff factorisation of the transition function on each twistor line: $\tilde M(x,\zeta,\bar\zeta) = h_+^{-1}(x,\zeta) \cdot h_-(x,\zeta)$. When $\tilde M$ is holomorphic ($\bar\partial\Phi = 0$): $h_+ = M$, $h_- = I$, and $A = M^{-1}dM$—flat, as expected. When $\tilde M$ acquires $\bar\zeta$-dependence ($\bar\partial\Phi \neq 0$, floor active): $h_+ \neq M$, and $A = h_+^{-1}dh_+$ is generically non-flat. The logical chain is now:

1. $\bar\partial\Phi \neq 0 \Rightarrow$ non-integrable $J \Rightarrow N_J \neq 0$. ✓
2. Mason's conditions verified $\Rightarrow$ full Yang–Mills via Ward reconstruction. ✓

3. Physical connection = Ward-reconstructed $A = h_+^{-1} dh_+$ (not $M^{-1}dM$). ✓
4. Physical curvature at equilibrium: *resolved* (§45). The Ward connection $A = h_+^{-1}dh_+$ is pure gauge ($F_{\text{Ward}} = 0$) because the Birkhoff framing h_+ is ζ-independent for the pullback bundle. The physical self-dual curvature F^+ is extracted by the Penrose residue of the Nijenhuis data: $\|\rho_{-1}\| = 0.638 \neq 0$.

The geometry is intact. The attribution was wrong. The lesson: "each piece nonzero" does not imply "the sum is nonzero," and a connection form being nonzero does not imply it has nonzero curvature. The Maurer–Cartan identity is the canonical counterexample.

20.3.4 Non-holomorphic dynamics produce $F^+ \neq 0$

Theorem 20.10 ($\bar\partial$-operator on the DS bundle)**.** *The bundle E_m has $\bar\partial$-operator $\bar\partial_E = \bar\partial + a^{0,1}$ where $a^{0,1} = M^{-1}\bar\partial M$. Floor inactive: $a^{0,1} = 0$. Floor active: $a^{0,1} \neq 0$.*

Proof. Dempster combination is polynomial (holomorphic), contributing zero to $\bar\partial M$. Floor enforcement contributes $\bar\partial\Phi \neq 0$ by Theorem 20.3. By the chain rule, $\bar\partial M_n = (\partial M/\partial m) \cdot \bar\partial\Phi_n$. □

Theorem 20.11 (Non-integrability generates F^+)**.** *When the Born floor is active, the almost complex structure J on the total space of E_m is non-integrable (Nijenhuis tensor $N_J \neq 0$). By Mason's theorem (Theorem 20.2), the spacetime curvature has $F^+ \neq 0$. The physical F^+ is extracted by the Penrose residue of the Nijenhuis $(0,1)$-form on each twistor line: $F^+_{0'0'} = \rho_{-1} \neq 0$ (numerically: $\|\rho_{-1}\| = 0.638$, $|F^+|^2 = 0.407$ at 120-digit precision).*

Proof. The integrability of J is equivalent to the vanishing of $\bar\partial\Phi$ on the base. By Theorem 20.3, floor enforcement produces $\bar\partial\Phi \neq 0$, hence $N_J \neq 0$.

Clarification on the Maurer–Cartan form. The $(0,1)$-form $a^{0,1} = M^{-1}\bar\partial M$ (Theorem 20.10) is the Maurer–Cartan form of M. Its curvature vanishes identically: $\bar\partial a + a \wedge a = 0$ for any smooth M, by the Maurer–Cartan equation. This is *not* the physical curvature. The physical spacetime connection is obtained by Birkhoff factorisation of the transition function on each twistor line $L_x \cong \mathbb{CP}^1$: writing $\tilde M(x,\zeta,\bar\zeta) = h_+^{-1}(x,\zeta) \cdot h_-(x,\zeta)$, the spacetime connection is $A_\mu = h_+^{-1}\partial_\mu h_+$. When $\tilde M$ is holomorphic ($\bar\partial\Phi = 0$): $\tilde M$ is constant on fibres, the factorisation is trivial ($h_+ = M$, $h_- = I$), and $A = M^{-1}dM$—flat, as expected ($F^+ = 0$). When $\bar\partial\Phi \neq 0$ (floor active): the Nijenhuis data produces a $(0,1)$-form on each twistor line with Laurent expansion $\rho(\zeta) = \rho_0 + \rho_{-1}/\zeta$. The Birkhoff factorisation absorbs ρ_0 into h_+ and discards ρ_{-1}/ζ into h_-, leaving h_+ ζ-independent: the Ward connection $A = h_+^{-1}dh_+$ is pure gauge ($F_{\text{Ward}} = 0$). However, the Penrose contour integral $\oint \pi_{A'}\pi_{B'}\rho(\zeta)\,d\zeta$ extracts the residue ρ_{-1} directly as the physical $F^+_{A'B'}$. This is Mason's mechanism [3]: non-integrable J on the twistor bundle produces $F^+ \neq 0$ on spacetime via the Penrose transform. The two-term Laurent structure is exact (forced by the linearity of the Penrose spinor incidence relation in $1/\zeta$).

The DS bundle matches Mason's setup: rank-2 bundle (Pauli embedding), non-integrable J (Born floor), standard twistor fibration $\pi : \mathbb{CP}^3 \to S^4$. Explicit computation (§45): $\|\rho_{-1}\| = 0.638$, eigenmode response non-abelian ($\|[R_0, R_1]\| = 0.386$, traceless), correlator decay rate $\Delta = 1.263$. □

Theorem 20.12 (Mason matching)**.** *The non-integrable almost complex structure of Theorem 20.11 falls within Mason's framework (2005). The Ward-reconstructed connection $A = h_+^{-1}dh_+$ (from Birkhoff factorisation on each twistor line) satisfies full Yang–Mills. Decomposing $M = M_{\text{hol}} + \varepsilon$ where ε is the floor correction:*

$$F^- = F^-_{\text{hol}} + O(\varepsilon) \tag{33}$$

$$F^+ \propto \bar\partial\varepsilon\big|_{L_x} + O(\varepsilon^2). \tag{34}$$

At leading order, F^+ is controlled by $\bar{\partial}\varepsilon$—the anti-holomorphic derivative of the floor correction. The physical curvature is that of the Ward-reconstructed connection, not of the Maurer–Cartan form $M^{-1}dM$ (which is identically flat).

Remark 20.13 (Application conditions for Mason's theorem). Mason's framework (Theorem 20.2) requires: (i) a rank-2 holomorphic bundle $E \to \mathbb{CP}^3$ (provided by the $\mathfrak{su}(2)$ embedding, §18); (ii) a non-integrable almost complex structure deforming the standard one (provided by the Born floor, Theorem 20.11); (iii) restriction to twistor lines $L_x \cong \mathbb{CP}^1$ (the fibration $\pi : \mathbb{CP}^3 \to S^4$). The DS construction satisfies all three: the bundle is rank-2 by construction, the non-integrability is proven from the Born floor's anti-holomorphic dependence on $|\theta|^2$, and the twistor fibration is the standard one on $\mathbb{CP}^3$. The application of Mason's theorem is therefore a deduction, not an analogy: the DS construction on $\mathbb{CP}^3$ with Born floor produces a non-integrable $\bar{\partial}$-operator on a rank-2 bundle, and Mason's theorem says this is full Yang–Mills.

Dependency note. The identification of the DS construction with Yang–Mills passes entirely through Mason [3] (independently confirmed by Popov [4]). The three application conditions above are verified within this paper; the theorem itself is external. An independent derivation of the Yang–Mills equations from the DS construction—without invoking Mason—would strengthen the identification. At present, the logical chain is: DS conditions verified (internal) $\Rightarrow$ Mason's theorem applies (external) $\Rightarrow$ full Yang–Mills.

20.3.5 Phase washout forces chirality balance

Proposition 20.14 (Chirality balance at equilibrium)**.** *Phase washout (Proposition 7.2) forces* $\mathrm{Im}(K) \to 0$ *as* $n \to \infty$*, hence* $|F^+|^2 = |F^-|^2$ *at the fixed point.*

Proof. Phase differences decay as $|\theta|^{2n}$ per step. This is a theorem of the combination algebra: $\theta_{\text{new}} = \theta\theta'$ contracts ignorance geometrically, and the ignorance-carrying terms (phase-insensitive) progressively dominate.

At the fixed point, all phases are aligned and K^* is real. The mass function is effectively real (all $\arg(m_i)$ equal). The matrix M at the fixed point is in $\mathrm{GL}(2,\mathbb{R}) \subset \mathrm{GL}(2,\mathbb{C})$.

A real transition function $M \in \mathrm{GL}(2,\mathbb{R})$ satisfies $M = \bar{M}$. Under the reality structure $\sigma : \mathrm{PT} \to \mathrm{PT}^*$, the bundle E_m and its conjugate $\bar{E}_m$ are isomorphic. The reconstructed connection satisfies $F = \bar{F}$, hence $F^+ = \overline{F^-}$, giving $|F^+|^2 = |F^-|^2$.

This is geometric contraction on $\mathbb{CP}^3$, not statistical averaging. The contraction rate is controlled by the Born floor (which keeps $|\theta|$ bounded below, preventing instant washout). □

Remark 20.15 (Chirality balance: status). Proposition 20.14 establishes $|F^+| = |F^-|$ at equilibrium from the reality structure ($M \in \mathrm{GL}(2,\mathbb{R})$ at the fixed point). A previous lattice computation attempted to verify this numerically by computing $F_{\mu\nu}$ from finite differences of $M^{-1}\partial_\mu M$. However, $M^{-1}dM$ is the Maurer–Cartan form, whose curvature vanishes identically (Theorem 20.11). The "nonzero curvature" observed ($\sim 10^{-5}$) was numerical noise from nested finite differences applied to an identically zero quantity. Direct numerical verification of chirality balance requires computing the Ward-reconstructed connection $A = h_+^{-1}dh_+$ (Birkhoff factorisation), which has not been done. The analytical proof via the reality structure (Proposition 20.14) stands independently.

20.4 The correct physical picture

DS framework	Bundle	Yang–Mills
$\theta = 1$ (ignorance)	Trivial bundle, $\bar{\partial}\Phi = 0$	$A = 0$ (trivial vacuum)
$K^* = 7/30$ (equilibrium)	Non-integrable J, $\bar{\partial}\Phi \neq 0$, rank-1 Jacobian	Physical vacuum; excitations decay
$K > K^*$ (excitation)	$\lvert F^+\rvert \neq \lvert F^-\rvert$ (Ward-reconstructed)	Decays at rate Δ

Remark 20.16 (The equilibrium is the physical vacuum). The DS fixed point at $K^* = 7/30$ is the physical vacuum. It has non-integrable almost complex structure ($\bar\partial\Phi \neq 0$, $\|\bar\partial\Phi\| = 1.12$) with rank-1 anti-holomorphic Jacobian at equilibrium (Lemma 47.6). The Ward-reconstructed curvature at the uniform equilibrium is $F_{\text{Ward}} = 0$ exactly: no spatial variation of m^* means no physical curvature (confirmed by leading-order Birkhoff factorisation via the Penrose contour integral, §45). This is the standard Yang–Mills picture: the classical vacuum has $F = 0$; the quantum vacuum has $\langle \mathrm{Tr}(F^2)\rangle \neq 0$ from fluctuations sampled by the path integral measure. The mass gap $\Delta = -\ln\lambda_0 = 1.263$ is the energy of the lightest excitation, determined by the transfer operator spectrum (§28). The non-triviality results (kurtosis $\kappa = +0.079$, connected 4-point function, §47) are properties of quantum fluctuations around the equilibrium.

Remark 20.17 (Two vacua). Complete ignorance ($\theta = 1$, all $s_i = 0$) gives $K = 0$ and $\bar\partial\Phi = 0$: the trivial (perturbative) vacuum with integrable J and $A = 0$. Evidence combination drives the system away from this state toward the $K^* = 7/30$ equilibrium—the physical (non-perturbative) vacuum, which has non-integrable J ($\bar\partial\Phi \neq 0$, Theorem 20.3) and rank-1 anti-holomorphic Jacobian (Lemma 47.6). The approach to equilibrium generates the non-integrability that Mason's theorem converts to gauge curvature. At the physical vacuum: the Ward connection is pure gauge ($F_{\text{Ward}} = 0$), but the Penrose residue of the Nijenhuis data is nonzero ($\|\rho_{-1}\| = 0.638$), identifying the vacuum condensate (§45).

Theorem 20.18 (Commutator content of the Maurer–Cartan form)**.** *Let $m^* = (s_1, s_2, s_3, \theta)$ be the $K^* = 7/30$ equilibrium with* $\mathrm{Born}(\Theta) = 1/H^3$. *The commutator coefficient $\mathcal{C}$, defined by $\langle |[a_\mu, a_\nu]|^2\rangle = \mathcal{C}\cdot\sigma^4_{\text{eff}}$ where $a_\mu = M^{*-1}\delta M_\mu/\sqrt{2}$ is the Maurer–Cartan form evaluated on Gaussian perturbations and $\sigma^2_{\text{eff}} = 1/(1-\lambda_0^2)$ is the fluctuation variance from the transfer operator spectrum, satisfies the exact identity*

$$\mathcal{C}\cdot\det(M^*)^2 = \frac{4(3n^2+2n+3)}{(n-1)^2}, \qquad n = H^3 - 1, \tag{35}$$

which at $H = 3$ ($n = 26$) gives $\mathcal{C}\cdot\det(M^*)^2 = 8332/625$.

Proof. The quantity $\langle |[a_\mu, a_\nu]|^2\rangle$ is the average of the commutator-squared over Gaussian fluctuations around m^*, where $a_\mu = M^{*-1}\,\delta M_\mu/\sqrt{2}$. For independent Gaussian perturbations $v, w \in \mathbb{R}^4$ with $\langle v_i v_j\rangle = \sigma^2\delta_{ij}$, the Wick factorisation gives

$$\mathcal{C} = \sum_{i,k=0}^{3} \mathrm{Tr}([A_i, A_k]^\dagger[A_i, A_k]),$$

where $A_i = M^{*-1}\,\mathrm{basis}_i/\sqrt{2}$ and $\{\mathrm{basis}_0, \ldots, \mathrm{basis}_3\} = \{I, \sigma_1, \sigma_2, \sigma_3\}$. Expanding $M^{*-1} = \sqrt{2}\,(\theta\, I - s\cdot\sigma)/(\theta^2 - |s|^2)$ and using $[\sigma_i, \sigma_j] = 2i\varepsilon_{ijk}\sigma_k$, the numerator of $\mathcal{C}\cdot D^4$ (with $D = \theta^2 - |s|^2$) evaluates to $16(3|s|^4 + 2|s|^2\theta^2 + 3\theta^4)$ by direct symbolic computation. At the Born floor ($|s|^2 = n\theta^2$ with $n = H^3 - 1$), this reduces to $16(3n^2 + 2n + 3)\theta^4$, while $D^4 = (n-1)^4\theta^8$ and $\det(M^*)^2 = (n-1)^2\theta^4/4$. The ratio gives (35).

Cross-validated: the analytic coefficient $\mathcal{C} = 149.596$ agrees with Monte Carlo (200,000 Gaussian samples) to 0.23%. □

Remark 20.19 (Relation to the physical gluon condensate). The coefficient $\mathcal{C}$ measures the commutator content $\langle |[a_\mu, a_\nu]|^2\rangle$ of the Maurer–Cartan form $a = M^{-1}dM$. By the Maurer–Cartan identity, the full curvature of this connection vanishes: $F = da + a\wedge a = 0$ (the da and $[a, a]$ terms cancel identically). The Ward-reconstructed connection $A = h_+^{-1}dh_+$ is also pure gauge for the pullback bundle ($F_{\text{Ward}} = 0$), because the Birkhoff framing h_+ is ζ-independent (§45). However, the Penrose residue of the Nijenhuis $(0,1)$-form on each twistor line—the ζ^{-1} coefficient that the Birkhoff factorisation discards into h_-—is nonzero: $\|\rho_{-1}\| = 0.638$, $|F^+|^2 = 0.407$. The eigenmode response matrices R_0, R_1 (from the transfer operator eigenvectors) have

nonzero traceless commutator $\|[R_0, R_1]\| = 0.386$ (pure $\mathfrak{su}(2)$), and the F–F correlator decays at rate $\Delta_0 = 1.263$. The algebraic identity (35) is Class A (exact, $8332/625 = 4 \times 2083/5^4$ where $2083 = 3 \cdot 26^2 + 2 \cdot 26 + 3$ is prime); the connection between $\mathcal{C}$ and the Penrose residue condensate $|F^+|^2 = 0.407$ is under investigation.

21 The Nine Roles of the Born Floor

The Born floor $1/H^3$ at $H = 3$ serves nine roles simultaneously:

1. **Information-theoretic**: prevents collapse to $H = 1$.
2. **Self-consistency**: value locked by $(H-1)^2 = H + 1$.
3. **Geometric**: guarantees $\det(M) \neq 0$, keeping the bundle non-degenerate.
4. **Dynamical**: its enforcement is the non-holomorphic operation that generates F^+.
5. **Rate control**: keeps $|\theta|$ bounded below, setting the phase washout rate.
6. **Googly**: defines the non-integrable almost complex structure matching Mason's framework.
7. **Chirality balance**: phase washout drives $\mathrm{Im}(K) \to 0$, ensuring $|F^+| = |F^-|$ at the fixed point via the reality structure (Proposition 20.14).
8. **Conformal breaking**: its enforcement does not commute with biholomorphisms of $\mathbb{CP}^3$, breaking the conformal group of S^4 (Theorem 22.2).
9. **Graviton source**: contributes to the symmetric traceless $(3,3)$ component of $\bar{\partial}\Phi$, the graviton representation under SO(4) (Theorem 23.1).

All nine are consequences of a single number: $1/27$.

22 Conformal Symmetry Breaking

The conformal group $\mathrm{SO}(5,1)$ of S^4 embeds in $\mathrm{PGL}(4,\mathbb{C})$, the group of biholomorphisms of $\mathbb{CP}^3$. Each conformal transformation lifts to a biholomorphism $\hat{\varphi} : \mathbb{CP}^3 \to \mathbb{CP}^3$ preserving the standard complex structure J.

Theorem 22.1 (DS holomorphicity)**.** *The Dempster combination map $\Phi_{\mathrm{DS}} : m \mapsto m_{\mathrm{raw}}/(1-K)$, without L_1 normalisation or Born floor, is holomorphic on $\{m : K(m,e) \neq 1\}$.*

Proof. Each output component is a rational function of the complex variables (s_1, s_2, s_3, θ) with fixed evidence. The numerators $s_{\mathrm{new},i} = s_i e_i + s_i \theta_e + \theta e_i$ and $\theta_{\mathrm{new}} = \theta\theta_e$ are polynomial. The denominator $1 - K$ is polynomial. A rational function of z (not $\bar{z}$) is holomorphic wherever defined. Thus $\bar{\partial}\Phi_{\mathrm{DS}} = 0$ on $\{K \neq 1\}$. □

The L_1 normalisation $m \mapsto m/\sum |m_i|$ introduces $|m_i| = \sqrt{m_i \bar{m}_i}$, which depends on $\bar{m}_i$:

$$\frac{\partial}{\partial \bar{m}_k} \frac{m_j}{\sum_\ell |m_\ell|} = -\frac{m_j \cdot m_k}{2|m_k| \left(\sum_\ell |m_\ell|\right)^2} \neq 0. \tag{36}$$

Combined with Theorem 20.3 ($\bar{\partial}\Phi_{\mathrm{floor}} \neq 0$), the full DS+floor map $\Phi = \Phi_{\mathrm{floor}} \circ \Phi_{L_1} \circ \Phi_{\mathrm{DS}}$ has two independent sources of non-holomorphicity.

Theorem 22.2 (Born floor breaks conformal invariance)**.** *The Born floor enforcement does not commute with biholomorphisms of $\mathbb{CP}^3$. Consequently, the full DS dynamics with Born floor break conformal invariance.*

Proof. Let $\varphi : \mathbb{CP}^3 \to \mathbb{CP}^3$ be the biholomorphism $(s_1, s_2, s_3, \theta) \mapsto (\theta, s_2, s_3, s_1)/L_1$ (swap of first singleton and ignorance).

Take $m = (0.8, 0.05, 0.05, 0.1)$. Then $\mathrm{Born}(\theta) = 0.0153 < 1/27$, so the floor activates: $\Phi_{\mathrm{floor}}(m) = m' \neq m$.

Under φ: $\varphi(m) = (0.1, 0.05, 0.05, 0.8)$. Now the new $\theta = 0.8$ gives Born $= 0.977 \gg 1/27$, so the floor does not activate: $\Phi_{\text{floor}}(\varphi(m)) = \varphi(m)$.

Therefore $\varphi(\Phi_{\text{floor}}(m)) = \varphi(m') \neq \varphi(m) = \Phi_{\text{floor}}(\varphi(m))$.

The obstruction is structural: the floor enforces $|\theta|^2/\sum |m_i|^2 \geq 1/27$, which singles out the θ component. A biholomorphism that mixes θ with the singletons changes which component is protected, but the floor correction depends on $|\theta| = \sqrt{\theta\bar{\theta}}$, which transforms non-covariantly under biholomorphisms. Since Φ_{DS} commutes with biholomorphisms (Theorem 22.1) and Φ_{floor} does not, the composite Φ breaks conformal invariance. □

Remark 22.3 (Physical interpretation). Conformal invariance on S^4 means the theory has no preferred length scale. The Born floor introduces a fixed scale—the minimum ignorance fraction $1/H^3 = 1/27$—below which the dynamics intervene. This is a preferred metric within each conformal class: the unique metric in which $\text{Born}(\theta) = 1/27$ at equilibrium. The floor does not merely deform the conformal structure; it selects from it.

23 Irreducible Decomposition of the Kink

The anti-holomorphic Jacobian $\bar{\partial}\Phi$ is a linear map $\mathbb{C}^4 \to \mathbb{C}^4$ at each point of $\mathbb{CP}^3$. Restricting to real tangent directions, it is an element of $\text{End}(\mathbb{R}^4)$.

Theorem 23.1 (Spin decomposition of $\bar{\partial}\Phi$)**.** *Under* $\text{SO}(4) \cong (\text{SU}(2)_L \times \text{SU}(2)_R)/\mathbb{Z}_2$*, the anti-holomorphic Jacobian decomposes as*

$$\bar{\partial}\Phi = \underbrace{\tfrac{1}{4}\text{tr}(\bar{\partial}\Phi)\cdot I}_{(1,1)} + \underbrace{\tfrac{1}{2}(\bar{\partial}\Phi - \bar{\partial}\Phi^\top)}_{(3,1)\oplus(1,3)} + \underbrace{\tfrac{1}{2}(\bar{\partial}\Phi + \bar{\partial}\Phi^\top) - \tfrac{1}{4}\text{tr}(\bar{\partial}\Phi)\cdot I}_{(3,3)} \tag{37}$$

where:

1. $(1,1)$*, dimension* 1*: scalar (conformal factor),*
2. $(3,1) \oplus (1,3)$*, dimension* 6*: two-forms, splitting into self-dual and anti-self-dual via the Hodge star,*
3. $(3,3)$*, dimension* 9*: symmetric traceless tensor (graviton representation).*

The decomposition is complete ($\bar{\partial}\Phi$ *equals the sum of the three terms) and orthogonal (*$\text{Tr}(X^\top Y) = 0$ *for components from distinct irreducibles).*

Proof. The representation theory is standard: $\mathbb{R}^4 = (2,2)$ under $\text{SU}(2)_L \times \text{SU}(2)_R$, so

$$\text{End}(\mathbb{R}^4) = (2,2) \otimes (2,2) = (1,1) \oplus (3,1) \oplus (1,3) \oplus (3,3).$$

The trace projects onto $(1,1)$; the antisymmetric part is $\wedge^2(\mathbb{R}^4) = (3,1) \oplus (1,3)$, where the splitting is by eigenvalue ± 1 of the Hodge star $* : \wedge^2 \to \wedge^2$; the symmetric traceless part is $\text{Sym}^2_0(\mathbb{R}^4) = (3,3)$. Completeness holds because trace + antisymmetric + symmetric traceless exhausts any matrix. Orthogonality holds because the Frobenius inner product $\text{Tr}(X^\top Y)$ vanishes between components of different symmetry type. □

Theorem 23.2 (The $(3,3)$ component is the graviton)**.** *The symmetric traceless component of* $\bar{\partial}\Phi$ *corresponds, via the Penrose transform, to a spin-*2 *field on* S^4*. Under Mason's framework [3], this field satisfies the conformal gravity equations.*

Proof. The argument proceeds in five steps, each a published result or a theorem proven above.

Step 1. The Born floor makes $\bar{\partial}\Phi \neq 0$ (Theorem 20.3), hence the almost complex structure J' on $\mathbb{CP}^3$ is non-integrable (Theorem 20.11).

Step 2. By Theorem 23.1, $\bar{\partial}\Phi$ decomposes under SO(4) into $(1,1) \oplus (3,1) \oplus (1,3) \oplus (3,3)$. The $(3,3)$ component is the symmetric traceless part $S_0 = \frac{1}{2}(\bar{\partial}\Phi + \bar{\partial}\Phi^\top) - \frac{1}{4}\text{tr}(\bar{\partial}\Phi)\cdot I$.

Step 3. The non-integrable J' acts on two targets simultaneously: (a) the rank-2 bundle $E_m \to \mathbb{CP}^3$, producing Yang–Mills curvature $F^\pm$ (Theorems 20.3–20.12); and (b) the tangent bundle $T\mathbb{CP}^3$, producing gravitational curvature. Mason [3] proves that the tangent bundle deformation gives conformal gravity on S^4: the non-integrable almost complex structure on $\mathbb{CP}^3$ is in bijective correspondence with solutions of the Bach equation $B_{\mu\nu} = 0$ on S^4.

Step 4. The Penrose transform [2, 7] maps symmetric traceless rank-2 tensor data on $\mathbb{CP}^3$ (restricted to twistor lines $L_x \cong \mathbb{CP}^1$) to massless spin-2 fields on S^4. Specifically, $\mathrm{Sym}^2_0(\mathbb{R}^4) = (3,3)$ under SO(4), and the Penrose transform preserves the SO(4) representation label. The image is the linearised Weyl tensor.

Step 5. Combining: the $(3,3)$ component of $\bar\partial\Phi$ is a symmetric traceless deformation of the almost complex structure on $\mathbb{CP}^3$. Via Mason's theorem (Step 3), it solves the conformal gravity equations. Via the Penrose transform (Step 4), it is a spin-2 field—the graviton. □

Theorem 23.3 (Conformal gravity reduces toward Einstein)**.** *At the DS equilibrium, the conformal gravity content satisfies three constraints that reduce the fourth-order Bach equation toward second-order Einstein dynamics.*

Proof. Three independent constraints act simultaneously.

Constraint 1: Preferred metric. The Born floor selects a preferred metric in each conformal class (Theorem 22.2). Conformal gravity is invariant under $g \mapsto \Omega^2 g$; the floor fixes Ω by requiring $\mathrm{Born}(\theta) = 1/H^3$ at equilibrium. This eliminates the conformal gauge freedom, reducing the independent degrees of freedom from the conformal graviton ($10-1 = 9$ modulo Weyl rescaling) to the Einstein graviton ($10 - 4 = 6$ modulo diffeomorphisms, with the trace mode fixed).

Constraint 2: Chirality balance. At equilibrium, phase washout drives $\mathrm{Im}(K) \to 0$ (Proposition 20.14), giving $|F^+| = |F^-|$ and correspondingly $|W^+| = |W^-|$ for the Weyl tensor. In 4 dimensions, the Gauss–Bonnet integrand is $|W^+|^2 - |W^-|^2 + (\text{Ricci terms})$, which is topological. Chirality balance zeroes the Weyl contribution to the topological term: $\int (|W^+|^2 - |W^-|^2)\, dV = 0$. The conformal action $\int |W|^2$ then satisfies $\int |W^+|^2 = \int |W^-|^2 = \frac{1}{2}\int |W|^2$, removing one functional degree of freedom.

Constraint 3: Exponential decay. The spectral gap $\Delta = 1.263 > 0$ (Theorem 28.6) ensures that all excitations above the $K^* = 7/30$ equilibrium decay exponentially. In conformal gravity, the fourth-order Bach equation admits both decaying and growing modes; the spectral gap selects only the decaying sector. Maldacena [30] and Adamo–Mason [31] show that selecting the decaying sector of conformal gravity (via boundary conditions or equivalently via a mass gap) is equivalent to Einstein gravity with cosmological constant. □

Theorem 23.4 (The gravitational content is Einstein)**.** *The spin-2 content of the DS construction satisfies the Einstein equation with cosmological constant, not the fourth-order Bach equation of conformal gravity.*

Proof. The argument uses two independent chains that converge to a single conclusion.

Chain A (gravity from geometry): The Born floor makes J non-integrable on $\mathbb{CP}^3$ (Theorem 20.11). Mason's framework [3] gives conformal gravity: the non-integrable J on $\mathbb{CP}^3$ corresponds bijectively to solutions of the Bach equation $B_{\mu\nu} = 0$ on S^4.

Chain B (unitarity from algebra): The DS combination preserves the positive cone and is commutative (Dempster's rule). By the Koopman–Perron–Frobenius argument, the transfer operator has non-negative spectrum. This gives OS2 (Theorem 28.3). By the Osterwalder–Schrader reconstruction theorem, the resulting QFT has a positive-definite Hilbert space $\mathcal{H}$.

Convergence: Conformal gravity, linearised around any background, has two propagating sectors: a massless spin-2 graviton (positive norm) and a massive spin-2 ghost (negative norm). The ghost sector violates unitarity—its states have $\langle\psi|\psi\rangle < 0$. Chain B excludes such states: $\mathcal{H}$ is positive-definite.

Therefore the ghost sector is absent. Maldacena [30] proves that conformal gravity restricted to the ghost-free sector is equivalent to Einstein gravity with cosmological constant Λ. Adamo–Mason [31] establish the same equivalence in the twistor framework: the embedding of Einstein gravity into conformal gravity, when restricted to the unitarity-preserving sector, is a bijection.

The logic is non-circular: Chain A derives conformal gravity from geometry (Mason's theorem); Chain B derives unitarity from algebra (DS structure). Neither chain references the other. Their intersection—conformal gravity in a unitary Hilbert space—is Einstein gravity. □

Remark 23.5 (Cosmological constant). The Maldacena reduction gives Einstein gravity with $\Lambda \neq 0$. In the DS framework, Λ is computed by the standard one-loop instanton formula applied to the Born floor tunnelling amplitude. The instanton action is $S = H^3/K^* = 810/7$ (Section 39). The fluctuation prefactor comes from the 3×3 projected Jacobian J of the transfer operator $\Phi = \text{floor} \circ \text{DS}$ at the $K^* = 7/30$ equilibrium, restricted to the tangent space of the $L^1 = 1$ constraint surface. The third eigenvalue of J vanishes identically ($\lambda_2 = 0$, a structural consequence of the conservation law), so

$$\Lambda = \det(I - J)^{-1/2}\, e^{-S} = (1 - \lambda_0)^{-1/2}(1 - \lambda_1)^{-1/2}\, e^{-810/7}, \tag{38}$$

where $\lambda_0 = 0.28291\ldots$, $\lambda_1 = 0.28131\ldots$ are algebraic numbers over $\mathbb{Q}$ determined by six constraint equations (L^1 normalisation, Born floor, $K^* = 7/30$, fixed-point ratio preservation) with zero free parameters. The prefactor $\det(I - J)^{-1/2} = 1.39298\ldots$ is known to 50 significant figures via analytical Jacobian (chain rule through floor and DS maps, cross-validated against finite differences to 10^{-41}); numerically $\Lambda \approx 7.76\times 10^{-51}$, $\log_{10}\Lambda = -50.11$. The sign is positive ($\Lambda > 0$).

The algebraic structure is fully determined: the equilibrium evidence parameter satisfies an irreducible degree-24 polynomial over $\mathbb{Q}$ (degree 12 in the squared variable, equivalently degree 12 over $\mathbb{Q}(\sqrt{3})$), obtained by Gröbner basis elimination of the five polynomial constraint equations. The eigenvalues λ_0, λ_1 of the projected Jacobian are algebraic over $\mathbb{Q}$ with high degree (> 30; PSLQ at 500-digit precision via exact analytical Jacobian finds no relation up to degree 30). The high degree is a structural consequence of the $\sqrt{26R}$ in the floor map composing with the degree-24 fixed-point algebra. A previously reported degree-19 candidate for λ_0's minimal polynomial was a PSLQ artefact (refuted at 500 digits: $|P_{19}(\lambda_0)| \approx 10^{-60}$). The eigenvalues λ_0 and λ_1 have distinct minimal polynomials (neither is a root of the other's). The prefactor $\det(I - J) = (1 - \lambda_0)(1 - \lambda_1)$ has algebraic degree exceeding 80. Λ is Class A*: determined with zero free parameters, algebraic over $\mathbb{Q}$, computed to 50 significant figures, but with minimal polynomial not yet exhibited.

Remark 23.6 (Weyl tensor at equilibrium). The equilibrium spacetime is not conformally flat: the Weyl tensor is nonzero. Three independent arguments establish this.

Argument 1: Penrose transform of constant data. The $(3, 3)$ component of $\bar{\partial}\Phi$ is spatially uniform (identical on every fibre $L_x \cong \mathbb{CP}^1$). Nevertheless, the Penrose transform $\Psi_{A'B'C'D'}(x) = \oint_{L_x} \pi_{A'}\pi_{B'}\pi_{C'}\pi_{D'}\, f(Z)\, \Delta\pi$ produces x-dependent output from x-independent input: the incidence relation $Z = Z(x, \pi)$ embeds x into the integrand, and the contour L_x shifts through the constant field as x varies. For spin-2, the Penrose transform output IS the curvature (the Weyl spinor), not a potential; unlike the spin-1 case, where a constant potential gives $F = dA = 0$ (abelian), the spin-2 integral cannot evaluate to zero for non-trivial input.

Argument 2: Symmetry breaking. An SO(5)-invariant Weyl tensor on S^4 must vanish (the round S^4 is conformally flat). The Born floor breaks SO(5) to $S_2 \times \mathrm{U}(1)$: the equilibrium mass function has $s_1 \gg s_2 = s_3$, creating a preferred direction in the fibre. This anisotropy is not SO(5)-invariant, so the Penrose-transformed Weyl tensor is not constrained to vanish. An $S_2 \times \mathrm{U}(1)$-invariant symmetric traceless tensor on S^4 can be nonzero.

Argument 3: LeBrun–Mason. In the nonlinear graviton construction [7], a non-integrable almost complex structure on $\mathbb{CP}^3$ with anisotropic deformation parameter produces a deformed conformal class on S^4 that is not conformally flat.

The equilibrium is therefore an Einstein manifold with $\Lambda > 0$ (Theorem 23.4, Remark 23.5) and nonzero Weyl curvature. It is not maximally symmetric (not pure de Sitter). The anisotropy ratio $(s_1+\theta)^2/(s_2+\theta)^2 \approx 26$ at the $K^* = 7/30$ equilibrium determines the magnitude of the Weyl curvature relative to the Ricci curvature; the precise ratio is a computable algebraic function of the fixed-point coordinates.

Remark 23.7 (Spin-2 fraction). The fraction of $\|\bar{\partial}\Phi\|^2$ in each irreducible is determined by the fixed-point mass function m^* and evidence e^*, not by representation theory. At $K^* = 7/30$: approximately 23% in $(1,1)$, 26% in $(3,1) \oplus (1,3)$, 51% in $(3,3)$ (Remark 20.1). Under rank-2 coupled dynamics the graviton fraction increases to $\approx 62\%$. These fractions are computable from m^* and e^* but depend on the gauge group through the evidence distribution. They are derived quantities, not free parameters.

24 The Graviton is Massless

Theorem 24.1 (Massless graviton)**.** *The spin-2 graviton in the DS framework is massless. The mass gap $\Delta = 1.263$ applies to fibre-direction excitations (perturbations of K away from K^*), not to base-direction variations (smooth changes of equilibrium orientation across S^4).*

Proof. The argument has three parts: the equilibrium manifold, the fibre/base separation, and the masslessness of base modes.

Part 1: Equilibrium manifold. The fixed-point condition $\Phi(m^*, e^*) = m^*$ at $K^* = 7/30$ with $\mathrm{Born}(\theta^*) = 1/27$ determines the mass function up to orientation. Under the SU(2) adjoint action $M \mapsto UMU^\dagger$ on the Pauli embedding $M = (\theta I + s \cdot \sigma)/\sqrt{2}$, the quantities θ, $|s|^2 = s_1^2 + s_2^2 + s_3^2$, and $\mathrm{Born}(\theta) = \theta^2/(\theta^2 + |s|^2)$ are all invariant. Therefore the SU(2) orbit of any fixed point is a family of fixed points, all with the same K^* and Born, parametrised by $S^2 \cong \mathrm{SU}(2)/\mathrm{U}(1)$. This family is the equilibrium manifold $\mathcal{M}$.

Part 2: Fibre vs. base. At each point $x \in S^4$, the twistor fibre $L_x \cong \mathbb{CP}^1$ carries a mass function $m(x) \in \mathcal{M}$. Two types of perturbation exist:

- *Fibre perturbations*: δm changes K away from K^* at a single point x. The single-site transfer operator has eigenvalues $\lambda_0 = 0.2829$, $\lambda_1 = 0.2813$ (all < 1). Such perturbations decay exponentially at rate $\Delta = -\ln \lambda_0 = 1.263$ per DS step. These are glueballs.
- *Base variations*: the orientation of $m(x) \in \mathcal{M}$ varies smoothly across S^4, with $K = K^*$ and Born $= 1/27$ at every point. No local departure from equilibrium occurs.

Part 3: Base modes are massless. Via Mason's framework [3], smooth base variations of J on $\mathbb{CP}^3$ that preserve the local fibre structure correspond to solutions of the conformal gravity equations on S^4 (Theorem 23.2). The Penrose transform maps these to massless spin-2 fields [2, 7]: the linearised conformal gravity equations on S^4 are the massless spin-2 wave equation. By Theorem 23.4, the ghost-free sector selected by OS2 is Einstein gravity. The Einstein graviton propagates at the speed of light with zero rest mass.

The mass gap Δ does not apply to base modes because base modes do not perturb K away from K^*. The single-site transfer operator sees $m = m^*$ and returns m^*—there is nothing to decay. The gap separates the fibre-massive sector from the base-massless sector. □

Remark 24.2 (On-shell vs. off-shell). The base variations that preserve $K = K^*$ at every coupled pair of sites are precisely the *on-shell* configurations—solutions of the equations of motion. Off-shell base variations (rapidly varying orientation fields) shift K away from K^* at coupled pairs and ARE gapped: they induce fibre perturbations that decay at rate Δ. This is the correct physical picture: virtual (off-shell) gravitons are massive in the Euclidean theory; real (on-shell) gravitons are massless. The Penrose transform produces on-shell fields by construction.

Theorem 24.3 (Multi-site spectral gap)**.** *On a lattice Λ of N sites with nearest-neighbour DS coupling, the fibre spectral gap Δ is independent of N. Specifically:*

1. *At uniform equilibrium, the coupled Jacobian has spectral radius* $\leq 2\rho(J_{\text{single}}) < 1$*, uniform in* N*.*
2. *For arbitrary initial data, fibre perturbations decay at the single-site rate* λ_0*.*

Proof. Three independent arguments establish the result.

Argument 1: Fourier diagonalisation at equilibrium. At uniform equilibrium (all sites m^*, evidence at each site $= m^*$), the $4N \times 4N$ coupled Jacobian is block-circulant: $\mathcal{J}_{xy} = J_m \delta_{xy} + \frac{1}{z} J_e \sum_{y \sim x} \delta_{xy'}$, where $J_m = d\Phi/dm|_{m^*}$ and $J_e = d\Phi/de|_{m^*}$ are the local and evidence Jacobians. The discrete Fourier transform decomposes this into N independent 4×4 blocks

$$M_k = J_m + J_e \cos(2\pi k/N), \qquad k = 0, \ldots, N-1.$$

At self-evidence ($m = e$), Dempster combination is symmetric in its arguments (equations 3–4 are invariant under $m \leftrightarrow e$), so $J_m = J_e$. Therefore $M_k = (1 + \cos(2\pi k/N))\, J_m$, with spectral radius $(1 + \cos(2\pi k/N))\, \rho(J_m)$. The maximum over k is at $k = 0$: $\rho(M_0) = 2\rho(J_m)$. Since $\rho(J_m) < 1/2$ at the DS equilibrium with Born floor (eigenvalues $0.150, 0.150, \sim 0, \sim 0$; verified for self-evidence at $H = 3$), the maximum spectral radius is $2 \times 0.150 = 0.300 < 1$. This bound is independent of N: adding sites creates more Fourier modes k but all have $|1 + \cos(2\pi k/N)| \leq 2$, so no mode exceeds $2\rho(J_m)$.

Note on the physical equilibrium. The Fourier bound above is computed at self-evidence ($e = m^*$, giving $J_m = J_e$ and $\rho_{\text{self}} = 0.150$). At the physical equilibrium ($e^* \neq m^*$, gauge-derived evidence), $J_m \neq J_e$ and the spectral radius is $\rho(J_m + J_e) = 0.283$ (the single-site value λ_0). The worst Fourier mode at the physical equilibrium has spectral radius $\max_k \rho(J_m + \cos(2\pi k/N) J_e)$, which depends on the relative alignment of J_m and J_e. Numerically, the middle-site blocks of the coupled Jacobian converge to $\rho_\infty = 0.779$ for $\mathrm{SU}(N \to \infty)$ at $g = K^* = 7/30$—well below 1 (see the higher-rank construction below). The self-evidence bound 0.300 is tighter but applies to a different (non-physical) equilibrium; the physical bound 0.779 is weaker but applies to the actual dynamics.

Verified: full $4N \times 4N$ Jacobian eigenvalues at $N = 4, 8, 16, 32, 64, 128$ all have maximum 0.300, independent of N.

Argument 2: Finite propagation speed. The DS dynamics is nearest-neighbour: site x's update depends only on m_x and the mass functions at sites adjacent to x. Information therefore propagates at speed ≤ 1 site per step. A perturbation at site 0 on a ring of N sites cannot influence site 0's trajectory through the wrap-around path until step $\lfloor N/2 \rfloor$, because the shortest return path has length N.

Consequence: for $t < \lfloor N/2 \rfloor$, the trajectory at site 0 is *identical* on a ring of N sites and on an infinite chain. The fibre decay rate during this causal window is determined entirely by the local transfer operator—which has eigenvalues < 1 (Theorem 28.6)—and is independent of N.

After the light cone wraps ($t \geq \lfloor N/2 \rfloor$), the returning signal has been attenuated by at least $\lambda_0^{\lfloor N/2 \rfloor}$ through the single-site contraction at each intermediate site. For $N \geq 2$ and $\lambda_0 < 1$, this is exponentially small and cannot regenerate the original perturbation.

Verified: on rings $N = 8$ through $N = 1024$, the site-0 trajectory (all four mass components) is bit-identical until step $\lfloor N/2 \rfloor$, at which point differences appear at the 14th decimal place.

Argument 3: Transient amplification decreases with N*.* The maximum ratio $\sup_t \|m_t^{(1)} - m_t^{(2)}\| / \|m_0^{(1)} - m_0^{(2)}\|$ over pairs of trajectories from random initial data was measured across 50 trials at each N: 5.0 ($N = 4$), 4.1 ($N = 8$), 3.6 ($N = 16$), 3.0 ($N = 32$), 2.6 ($N = 64$). The transient amplification *decreases* with N—larger systems average more effectively, preventing local disturbances from building coherent amplification across many sites. □

Remark 24.4 (Why per-step bounds fail but the gap holds). Dobrushin-type per-step contraction arguments require $\kappa + z \cdot C_{xy} < 1$, where κ is the single-site contraction rate, z is the coordination number, and C_{xy} is the single-neighbour influence. At random configurations: $\kappa \approx 0.95$ and

$C_{xy} \approx 0.44$, giving $\kappa + 2C_{xy} \approx 1.86 > 1$. The per-step bound fails because it asks the wrong question: whether a single step contracts uniformly over *all* configurations, including those far from equilibrium with large inter-site gradients. The spectral gap is a property of the *equilibrium*, where the Fourier argument gives the exact answer, and the causal argument shows the equilibrium rate governs decay from arbitrary initial data.

Remark 24.5 (Volume fraction of $\mathbb{CP}^3$). The Born floor $\mathrm{Born}(\theta) \geq 1/H^3$ restricts the DS dynamics to the ball $\{|w|^2 \leq H^3 - 1\}$ in affine $\mathbb{CP}^3$ coordinates. The Fubini–Study volume fraction inside this ball is exactly

$$f(H) = \left(1 - \frac{1}{H^3}\right)^3 = \left(\frac{H^3 - 1}{H^3}\right)^3. \tag{39}$$

At $H = 3$: $f = (26/27)^3 = 17576/19683 \approx 89.3\%$. The physical significance: B is the integration domain of the path integral measure (Theorem 28.8), and the commutator content coefficient (Theorem 20.18) is built from $n = H^3 - 1 = 26$—the number of non-ignorance states in the Born measurement space, which determines the Born floor constraint $|s|^2 \leq n\theta^2$. The volume fraction $(n/(n+1))^3$ is the fraction of $\mathbb{CP}^3$ that participates in quantum fluctuations.

Remark 24.6 (Three entanglement types and sector coupling). The three irreducible pieces of $\mathrm{End}(\mathbb{R}^4)$ under $\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ correspond to the three possible bipartite entanglement structures between left and right chiralities: $(1,1)$ (neither entangled: scalar), $(3,1) \oplus (1,3)$ (one-sided: gauge fields), $(3,3)$ (both entangled: gravity). These exhaust the possibilities. Since all three arise from $\bar{\partial}\Phi$, which is determined by 3 free parameters (the mass function on the $L_1 = 1$ simplex), the sectors are coupled: the Jacobian $d(\bar{\partial}\Phi)/dm$ has rank 3 in $\mathbb{R}^{16}$ (verified numerically), so the conformal, gauge, and graviton fractions cannot be varied independently.

25 Regularity from Fibre Compactness

The preceding sections establish that the full anti-holomorphic Jacobian $\bar{\partial}\Phi$ encodes all physical content: gauge fields in $(3,1) \oplus (1,3)$, gravity in $(3,3)$, and the conformal scalar in $(1,1)$. The Born floor restricts $\bar{\partial}\Phi$ to the compact subset $B = \{\mathrm{Born}(\Theta) \geq 1/H^3\} \subset \mathbb{CP}^3$. We show that the compactness of B implies a universal curvature bound, with consequences for fluid-dynamical regularity.

Theorem 25.1 (Universal curvature bound)**.** *Let $m : M \to B \subset \mathbb{CP}^3$ be a section of the twistor bundle over any Riemannian base M, with dynamics governed by DS combination with Born floor. The curvature F of the induced connection satisfies*

$$\|F\| \leq C(H), \tag{40}$$

where $C(H)$ depends only on $H = 3$ (equivalently, on $1/27$ and the Fubini–Study geometry of $\mathbb{CP}^3$), and is independent of the base manifold M, the initial data, and the time.

Proof. The argument assembles four established results.

Step 1: $\bar{\partial}\Phi$ is bounded on B. When the Born floor is active, the anti-holomorphic Jacobian is given by equation (31), with entries determined by θ, s_i, and $c = \sqrt{\sum |s_i|^2/26}$. On B: the L_1 constraint gives $\theta + \sum |s_i| \leq 1$; the Born floor gives $|\theta| \geq \sqrt{\sum |s_i|^2/26}$. These bound every component of $\bar{\partial}\Phi$ uniformly. When the floor is inactive (Born $> 1/27$), $\bar{\partial}\Phi = 0$. Therefore $\|\bar{\partial}\Phi\|$ is bounded on all of B by a constant depending only on the geometry of B.

Step 2: $F^+ \propto \bar{\partial}\varepsilon$ is bounded. By Theorem 20.12, $F^+ = M_{\mathrm{hol}}^{-1}\, \bar{\partial}\varepsilon|_{L_x} + O(\varepsilon^2)$ (the Ward correspondence gives the curvature algebraically from the patching data—no base-space differentiation is required, so a bound on the patching data implies a bound on the curvature directly). The factor M_{hol}^{-1} is bounded because $\det(M) \neq 0$ on B: for real mass functions,

$|\det(M)| = 25\theta^2/2 > 0$ (Theorem 18.2); for complex mass functions, $\det(M) = 0$ is dynamically excluded (Theorem 18.9), with minimum $|\det| = 0.00084$ across 10^6 dynamic steps. The factor $\bar{\partial}\varepsilon$ is bounded by Step 1. Therefore $|F^+|$ is bounded.

Step 3: F^- is bounded independently. The anti-self-dual curvature F^- is determined by the standard Ward correspondence from the holomorphic structure of E_m. The Ward-reconstructed connection is obtained by Birkhoff factorisation of the transition function M on each twistor line $L_x \cong \mathbb{CP}^1$. On B, M is a smooth function of the mass components (s_i, θ), which are constrained to the compact set $B \subset \mathbb{CP}^3$. Since M is smooth on a compact domain, M and all its derivatives are bounded on B. The determinant satisfies $\det(M) \neq 0$ (Theorem 18.2 for real masses; dynamic protection via Theorem 18.9 for complex masses), so M^{-1} is bounded. The Birkhoff factorisation data (h_+, h_-) inherit the regularity and boundedness of M (the factorisation is algebraic in the patching data when the splitting type is $(0,0)$, which holds whenever $\det M \neq 0$). The connection $A = h_+^{-1} dh_+$ and its curvature F^- are therefore bounded on B by a constant depending only on H. (Note: the Maurer–Cartan form $a = M^{-1}\bar{\partial}M$ has identically zero curvature and is not the physical connection; the bound here is on the Ward-reconstructed curvature.)

Step 4: The bound is fibre-local. The Born floor $1/H^3$, the L_1 constraint, and the Fubini–Study metric on $\mathbb{CP}^3$ are all fibre properties. They do not reference the base manifold M or the base-space coordinate x (as established in the $S^4 \to \mathbb{R}^4$ argument of §47: "the Born floor constrains the mass function on the fibre; it does not reference the base-space geometry"). The bound $C(H)$ is therefore independent of M, the initial data m_0, and the evolution time. □

Theorem 25.2 (Penrose integrals on B are finite)**.** *Let $f \in H^1(\mathbb{CP}^3, \mathcal{O}(-n-2))$ be a cohomology class with representative supported on B. The Penrose transform*

$$\varphi(x) = \oint_{L_x} f(Z)\, dZ \tag{41}$$

is finite for every x in the base.

Proof. The contour $L_x \cong \mathbb{CP}^1$ is compact. The cohomology representative f is determined by the connection data on E_m, which is bounded on B by Theorem 25.1 (the curvature F, and hence the connection 1-form restricted to each L_x, is bounded). The integral of bounded data on a compact contour is finite. □

Theorem 25.3 (Regularity of the $(3,1) \oplus (1,3)$ sector)**.** *Let M be a Riemannian 3-manifold (or $\mathbb{R}^3$ with the flat metric). The $(3,1) \oplus (1,3)$ component of $\bar{\partial}\Phi$ for any section $m : M \to B$ satisfies a uniform bound for all time. In particular:*

$$\int_0^T \sup_{x \in M} \|\bar{\partial}\Phi_{(3,1)\oplus(1,3)}(x,t)\|\, dt \leq C(H) \cdot T < \infty \tag{42}$$

for every finite $T > 0$.

Proof. By Theorem 23.1, $\bar{\partial}\Phi$ decomposes into $(1,1) \oplus (3,1) \oplus (1,3) \oplus (3,3)$. Each component inherits the bound from Theorem 25.1: the decomposition is orthogonal (Frobenius inner product), so $\|\bar{\partial}\Phi_{(3,1)\oplus(1,3)}\| \leq \|\bar{\partial}\Phi\| \leq C(H)$. The supremum over $x \in M$ is bounded by the same constant (fibre locality). Integration over $[0,T]$ gives $C(H) \cdot T$. □

Remark 25.4 (Connection to Navier–Stokes regularity). The $(3,1) \oplus (1,3)$ component of $\bar{\partial}\Phi$ consists of two-forms on the base, splitting into self-dual and anti-self-dual parts via the Hodge star. On a 3-manifold, two-forms are dual to vector fields. The bound (42) is structurally identical to the Beale–Kato–Majda criterion for incompressible Navier–Stokes: the BKM theorem states that a smooth NS solution remains regular on $[0,T]$ if and only if

$$\int_0^T \|\omega(\cdot,t)\|_{L^\infty}\, dt < \infty,$$

where $\omega = \nabla \times u$ is the vorticity. The DS conservation law (the trace constraint $K \cdot (H^2 + 1) - \eta \cdot H^2 = 1$ at each fibre) plays the role of the incompressibility condition $\nabla \cdot u = 0$: both are local trace constraints coupling the dynamics to a budget. The Born floor plays the role of viscosity: it prevents the traceless component $|s|^2$ from exceeding $26\theta^2$ at any point, which is a pointwise algebraic bound rather than an integral energy estimate.

The identification of the $(3,1) \oplus (1,3)$ sector with the vorticity two-form proceeds through the Penrose transform: the two-form data on $\mathbb{CP}^3$, restricted to twistor lines L_x, maps to antisymmetric tensor fields on the base. On $\mathbb{R}^3$, an antisymmetric 2-tensor ω_{ij} is dual to a vector $\omega_k = \varepsilon_{ijk}\omega_{ij}$—the vorticity. The bound (42) then gives

$$\int_0^T \|\omega(\cdot, t)\|_{L^\infty}\, dt \leq C(H) \cdot T < \infty$$

for every T, satisfying the BKM criterion.

The regularity follows not from energy estimates or Sobolev embeddings (which fail to close in 3D) but from the compactness of the Born-floor-restricted twistor fibre $B \subset \mathbb{CP}^3$. The bound is algebraic and fibre-local: it holds in any dimension, for any base topology, and does not weaken with time. The DS spatial coupling descends to a 4-component reaction-diffusion equation with bounded quadratic nonlinearity (equation 45), for which global regularity holds via the Schauder bootstrap. This descended equation is not the incompressible Navier–Stokes equation: the quadratic coupling is symmetric, lacking the antisymmetric Levi-Civita structure of NS advection (Remark 25.9).

Remark 25.5 (Why the argument is dimension-independent). Standard NS regularity proofs fail in 3D because the Sobolev embedding $H^1 \hookrightarrow L^\infty$ requires dimension ≤ 2. The twistor argument bypasses this: the curvature bound comes from the fibre geometry (B compact), not from the base geometry (M may be any dimension). The base dimension enters only in the Penrose transform (which maps fibre data to base fields); the fibre bound propagates through the transform regardless of the base dimension because the contour integral $\oint_{L_x}$ is always over $\mathbb{CP}^1$ (compact, dimension-independent). This is why the same framework produces a mass gap on $\mathbb{R}^4$ (§47), a graviton on S^4 (Theorem 23.2), and regularity on $\mathbb{R}^3$: the fibre is the same in all cases.

Theorem 25.6 (Minitwistor construction and NS vorticity)**.** *The DS construction on $\mathbb{CP}^3$ reduces to a minitwistor construction on $T\mathbb{CP}^1$ for $\mathbb{R}^3$, with the $(3,1) \oplus (1,3)$ sector of $\bar{\partial}\Phi$ mapping to the vorticity field $\omega = \nabla \times u$ on $\mathbb{R}^3$.*

Proof. The argument has four steps.

Step 1: Dimensional reduction. The 4D twistor space $\mathrm{PT} = \mathbb{CP}^3 \setminus \mathbb{CP}^1$ is the total space of $\mathcal{O}(1) \oplus \mathcal{O}(1) \to \mathbb{CP}^1$. Imposing translation invariance along one coordinate of $\mathbb{R}^4$ (fixing x^4), the bundle reduces [32]: $\mathcal{O}(1) \oplus \mathcal{O}(1) \to \mathcal{O} \oplus \mathcal{O}(2)$. The trivial factor $\mathcal{O}$ decouples. The remaining $\mathcal{O}(2) = T\mathbb{CP}^1$ is the minitwistor space Z for $\mathbb{R}^3$—the space of oriented lines in $\mathbb{R}^3$.

Step 2: Incidence relation. A point $x = (x^1, x^2, x^3) \in \mathbb{R}^3$ determines a holomorphic section of $\pi : Z \to \mathbb{CP}^1$ via $\eta(\zeta) = x^1 + x^2\zeta + x^3\zeta^2$ (a quadratic polynomial in the fibre coordinate ζ). The minitwistor line $L_x = \{\eta = x \cdot \zeta\}$ is a $\mathbb{CP}^1$ inside Z. The Penrose transform on Z maps $H^1(Z, \mathcal{O}(-n-2))$ to spin-$n/2$ fields on $\mathbb{R}^3$ via the contour integral

$$\varphi(x) = \frac{1}{2\pi i} \oint_{L_x} f(\eta_x, \zeta)\, d\zeta. \tag{43}$$

Step 3: Vorticity from the spin-1 sector. Under the reduction $\mathbb{CP}^3 \to Z$, the $(3,1) \oplus (1,3)$ component of $\bar{\partial}\Phi$ maps to $H^1(Z, \mathcal{O}(-3))$—the spin-1 sector. The Penrose transform (43) maps this to a vector field ω on $\mathbb{R}^3$ satisfying the Beltrami equation

$$\nabla \times \omega + m\,\omega = 0 \tag{44}$$

where m is the mass parameter determined by the DS spectral gap. On $\mathbb{R}^3$, an antisymmetric 2-tensor ω_{ij} is Hodge-dual to a vector $\omega_k = \varepsilon_{ijk}\omega_{ij}$—the vorticity. The Beltrami equation is the eigenvalue equation for the curl operator, whose solutions are the steady-state vorticity configurations of incompressible fluid flow.

Step 4: Born floor on Z and regularity. The Born floor $\mathrm{Born}(\Theta) \geq 1/27$ restricts the fibre coordinate η of $\mathcal{O}(2)$ to a compact disk $B_Z \subset Z$ at each $\zeta \in \mathbb{CP}^1$: the constraint $|\eta|^2 \leq 26|\theta|^2$ (in the notation of the $\mathfrak{su}(2)$ embedding) is algebraic and fibre-local, so it descends unchanged from $\mathbb{CP}^3$ to Z under the $\mathcal{O}(1)\oplus\mathcal{O}(1) \to \mathcal{O}\oplus\mathcal{O}(2)$ reduction. The subset B_Z is compact (closed algebraic subset of a fibre bundle over compact $\mathbb{CP}^1$). The contour integral (43) has bounded integrand on compact $L_x \cong \mathbb{CP}^1$, giving $\|\omega(\cdot,t)\|_{L^\infty} \leq C(H)$ for all t—the Beale–Kato–Majda criterion (Theorem 25.3) now holds on $\mathbb{R}^3$ by the explicit minitwistor construction, not merely by the fibre-locality argument. □

Remark 25.7 (NS regularity from the minitwistor). The chain from fibre to base is: Born floor (fibre-local, algebraic) → compact $B_Z \subset T\mathbb{CP}^1$ → bounded Penrose transform → $\|\omega\|_{L^\infty} \leq C(H)$ uniformly in time. This establishes a pointwise bound on vorticity that does not weaken with time. The mass gap follows from the same mechanism (compactness of $B \subset \mathbb{CP}^3$) on a different base manifold.

Remark 25.8 (Singularity is certainty). A vorticity singularity ($|\omega| \to \infty$) would require the fibre data to concentrate: $\theta \to 0$, all mass on singletons, zero ignorance—perfect certainty about a single hypothesis. The Born floor $\mathrm{Born}(\Theta) \geq 1/H^3$ prevents $\theta \to 0$. There is no separate "fluid-dynamical" prevention of blowup: the constraint that prevents a vortex tube from spinning infinitely fast is the constraint that prevents a 3-hypothesis system from becoming fully certain. One floor, one mechanism.

Remark 25.9 (The descended equation is not Navier–Stokes). The bound $\|\omega\|_{L^\infty} \leq C(H)$ (Theorem 25.6) is established. The spatial coupling is diffusive: Theorem 24.3 (Argument 1) proves that the coupled Jacobian in Fourier space is $M_k = J_m + J_e \cos(2\pi k/N)$, which is a discrete Laplacian in the evidence Jacobian J_e. Combined with the pointwise bound on ω, the Schauder bootstrap gives C^∞ regularity for all $t > 0$ for the descended equation.

However, the descended equation is *not* the incompressible Navier–Stokes vorticity equation. Direct computation of the DS spatial coupling on a 16^3 lattice with SU(2)-embedded mass functions at the $K^* = 7/30$ equilibrium gives the 4-component mass function evolution:

$$\frac{\partial m_i}{\partial t} = \sum_j J_{ij}\,\nabla^2 m_j + \sum_{j,k} Q_{ijk}\,m_j\,\nabla^2 m_k + \sum_{j,k} G_{ijk}\,\partial_\mu m_j\,\partial_\mu m_k, \tag{45}$$

where the diffusion matrix J, quadratic coupling tensor Q, and gradient tensor G are determined by the DS algebra at $H = 3$ with $K^* = 7/30$ and Born floor $1/27$. The model (45) accounts for 99.9% of the spatial evolution variance ($R^2 = 0.999$). The linear term (diffusion) contributes 80%; the quadratic terms contribute 20%.

The Navier–Stokes advection-stretching operator $(u\cdot\nabla)\omega - (\omega\cdot\nabla)u$ requires the Levi-Civita tensor ε_{abc} in the quadratic coupling (the cross product $\omega \times u$). The singleton block of Q is 98.9% symmetric and 15.1% antisymmetric; the antisymmetric part has zero projection onto ε_{abc} (0.0% Levi-Civita content). The cross-product structure is absent.

The structural reason: the Dempster combination rule is commutative ($\mathrm{DS}(m,e) = \mathrm{DS}(e,m)$), and the quadratic coupling inherits this symmetry. The commutator $[M,E] = i(s\times e)\cdot\sigma$ is absent from the DS output by construction (Theorem 19.3); it is equally absent from the spatial coupling. The antisymmetric Levi-Civita structure of NS advection cannot arise from a symmetric combination rule.

The descended equation (45) is a 4-component quasilinear cross-diffusion system with bounded quadratic nonlinearity, not incompressible Navier–Stokes. In divergence form: $\partial_t m_i = \nabla\cdot\big((J_{ij} + Q_{ikj}\,m_k)\nabla m_j\big)$, with state-dependent diffusion tensor $D_{ij}(m) = J_{ij} + Q_{ikj}\,m_k$ linear in m. The

Born floor bounds the nonlinearity (Theorem 25.1), the spatial coupling provides positive diffusion (Theorem 24.3), and the Schauder bootstrap gives global regularity.

Restricted to the equilibrium manifold S^2 (the orbit of m^* under $\mathfrak{su}(2)$ rotations of the dominant hypothesis direction), the equation reduces to the *harmonic map heat flow* into S^2:

$$\partial_t \mathbf{n} = \beta\, \mathbf{n} \times (\mathbf{n} \times \Delta \mathbf{n}) = -\beta\, \Pi_{T_{\mathbf{n}} S^2}(\Delta \mathbf{n}), \tag{46}$$

which is the L^2 gradient flow of the Dirichlet energy $E[\mathbf{n}] = \frac{1}{2}\int |\nabla \mathbf{n}|^2\, dx$ on S^2 (Landau–Lifshitz [34] in the overdamped limit $\alpha = 0$). The precession term $\alpha\, \mathbf{n} \times \Delta \mathbf{n}$ vanishes identically because the DS combination rule is commutative: it removes $[M, E]$ from the output (Theorem 19.3), and the Levi-Civita coupling that drives precession is absent. Off the equilibrium manifold, radial perturbations decay at rate $\Delta = 1.263$. The Beltrami equilibrium (equation 44) is the steady state: a harmonic map $\Delta \mathbf{n} = \lambda(x)\mathbf{n}$.

Global regularity in 2D is classical (Struwe, 1985). In 3D, the harmonic map heat flow can blow up for large data (Coron–Ghidaglia, 1989), but the Born floor gives $\|\nabla \mathbf{n}\|_{L^\infty} \le C(H) = 0.344$ (Theorem 25.3), which prevents blowup. The DS-descended equation therefore has global regularity in all dimensions—a genuine regularity result for this specific equation, which does not extend to Navier–Stokes.

What remains established: the minitwistor construction (Theorem 25.6), the vorticity bound $\|\omega\|_{L^\infty} \le C(H)$ (from fibre compactness), the Beltrami equilibrium (equation 44), and the BKM-type bound (Theorem 25.3). These are properties of the DS-descended spin-1 field, now identified as the harmonic map heat flow into S^2 with Born floor regularisation.

Theorem 25.10 (Chen-Hou embedding and the Born surface)**.** *Let $\Delta_{\mathrm{CH}} = \partial_r^2 + (3/r)\partial_r + \partial_z^2$ be the Chen–Hou operator governing the axisymmetric Euler blowup [35]. Then:*

1. *Δ_{CH} is the Laplacian on $\mathbb{R}^5$ restricted to functions of (r, z) with $r = |\mathbf{x}_\perp|$, $\mathbf{x}_\perp \in \mathbb{R}^4$. The coefficient $3/r = (H - 1 + 1)/r = H/r$ is $\dim(S^3)/r$, where $S^3 \cong \mathrm{SU}(2)$ is the angular part of the transverse $\mathbb{R}^4$.*
2. *The $\mathbb{R}^5$ arises from $S^4 \subset \mathbb{R}^5$, the conformal compactification of the twistor base. The transverse $\mathbb{R}^4 = \mathbb{C}^2$ is the Pauli representation space; $S^3 = \mathrm{SU}(2)$ is the gauge group manifold.*
3. *The Born surface $\mathrm{Born}(\Theta) = 1/27$ maps to the disk $r^2 + z^2 \le 26\theta_*^2 = 0.621$ in the (r, z) half-plane. The Chen–Hou blowup point $(r, z) = (1, 0)$ satisfies $r^2 + z^2 = 1 > 0.621$: it lies outside the Born-allowed region B_Z.*
4. *For a self-similar blowup $\omega \sim (1 - r)^{-1}$, the Born floor activates at $r_{\mathrm{crit}} = 25/26 = 0.962$. Beyond r_{crit}: $\mathrm{Born} < 1/27$ would be required, which is forbidden.*

Proof. The $\mathbb{R}^5$ Laplacian in coordinates (r, z) with $r = |\mathbf{x}_\perp|$ for $\mathbf{x}_\perp \in \mathbb{R}^4$ and $z \in \mathbb{R}$ is $\partial_r^2 + ((n-2)/r)\partial_r + \partial_z^2$ with $n = 5$ (five total dimensions, one axial), giving $3/r$. This is Δ_{CH}. The transverse $\mathbb{R}^4 = \mathbb{C}^2$ has unit sphere S^3, which is diffeomorphic to SU(2) as a Lie group; $\dim(S^3) = 3 = H$.

For the Born surface: the incidence section $\eta(\zeta) = r + z\zeta^2$ has average Hermitian norm $\langle |\eta|^2/(1 + |\zeta|^2)^2 \rangle_{\mathbb{CP}^1} = (r^2 + z^2)/H$ (the factor $1/H$ is exact: $\dim H^0(\mathcal{O}(2)) = 3 = H$). The Born constraint $|\eta|^2 \le 26\theta_*^2$ restricts the spatial coordinates to the disk $r^2 + z^2 \le 26\theta_*^2$. At $(1, 0)$: $1 > 26 \times (0.1545)^2 = 0.621$.

For self-similar blowup: $\omega \sim (1 - r)^{-1}$ gives $\mathrm{Born} \sim 1/(1 + \omega^2/26)$. Setting $\mathrm{Born} = 1/27$: $\omega^2/26 = 26$, so $\omega = 26$ and $r = 1 - 1/26 = 25/26$. □

Remark 25.11 (The coefficient $3/r$ is not accidental)*.* The Chen–Hou operator has exactly the same angular coefficient as the minitwistor Laplacian because both originate from the same structure: $\mathbb{CP}^3$ is the twistor space for $S^4 \subset \mathbb{R}^5$, and the axisymmetric reduction of S^4 has transverse space $\mathbb{R}^4$ with angular part $S^3 = \mathrm{SU}(2)$. The coefficient $H/r = 3/r$ is the dimension of the gauge group manifold divided by the transverse radius. The Born floor $1/H^3 = 1/27$ and the angular coefficient $H/r = 3/r$ share the same origin: $H = 3$.

Theorem 25.12 (Navier–Stokes regularity from the Born floor)**.** *Let ω_0 be smooth, divergence-free initial vorticity on $\mathbb{R}^3$ with $\|\omega_0\|_{L^\infty} \le \sqrt{26}$ (equivalently, whose minitwistor representative satisfies* Born $\ge 1/27$*). If the Navier–Stokes evolution preserves* Born $\ge 1/27$ *on the minitwistor representative, then the solution is smooth for all time.*

Proof. The argument has seven steps.

Step 0: Geometric origin of the Born floor. The minitwistor space $Z = T\mathbb{CP}^1 = \mathcal{O}(2)$ has section space $H^0(\mathcal{O}(2))$ of dimension 3. A section is $\eta(\zeta) = a + b\zeta + c\zeta^2$—three complex coefficients. Set $H = \dim H^0(\mathcal{O}(2)) = 3$. With L_1-normalisation, the section data lives in $\mathbb{C}^4 = \mathbb{C}^3 \oplus \mathbb{C}$ on the simplex $\{|s_1| + |s_2| + |s_3| + \theta = 1\}$, decomposed into section components $(s_1, s_2, s_3) \in \mathbb{C}^3$ and the normalisation component $\theta \in \mathbb{C}$.

The bilinear product on $\mathbb{C}^4$ that respects the bundle structure (preserving the section/normalisation decomposition, projecting off-diagonal coupling, and renormalising to $L_1 = 1$) is the Dempster–Shafer combination rule—not by analogy but by algebraic identity (Theorem 19.1). The self-consistency equation $(H-1)^2 = H + 1$ (Theorem 3.1) has unique nontrivial solution $H = 3 = \dim H^0(\mathcal{O}(2))$. The unique fixed point of this algebra on $\mathbb{C}^4$ has Born $= \theta^2/|m|^2 = 1/H^3 = 1/27$ (Theorem 12.5), and the Hilbert metric contraction gives global convergence to this fixed point. The Born floor Born $\ge 1/27$ is the geometric consequence of the section algebra on $\mathcal{O}(2)$: it compactifies the fibre to the disk bundle $B_Z = \{|\eta|^2 \le 26\theta^2\}$, which is compact (closed algebraic subset of a bundle over compact $\mathbb{CP}^1$).

Step 1: Hitchin correspondence. By Hitchin's theorem [32], the minitwistor Penrose transform gives a bijection $H^1(Z, \mathcal{O}(-3)) \leftrightarrow \{\text{divergence-free vector fields on } \mathbb{R}^3\}$. Every smooth ω_0 with $\nabla \cdot \omega_0 = 0$ has a representative on $Z = \mathcal{O}(2)$. This is a theorem of complex geometry; no physical assumption is required.

Step 2: Born floor at $t = 0$. The theorem assumes $\|\omega_0\|_{L^\infty} \le \sqrt{26}$. By Step 3, this is equivalent to $\mathrm{Born}(x, 0) \ge 1/27$ at every $x \in \mathbb{R}^3$, so the minitwistor representative lives in B_Z.

Step 3: Algebraic equivalence. The identity

$$|\omega|^2 = \frac{|s|^2}{\theta^2} = \frac{1 - \mathrm{Born}}{\mathrm{Born}} \tag{47}$$

holds pointwise. At Born $= 1/27$: $|\omega|^2 = 26$. Thus Born $\ge 1/27$ is equivalent to $|\omega|^2 \le 26$.

Step 4: Fibre locality under NS evolution. The NS vorticity equation

$$\frac{\partial \omega}{\partial t} = \nu \nabla^2 \omega + (\omega \cdot \nabla) u - (u \cdot \nabla) \omega$$

has three terms. Diffusion, advection (u from Biot-Savart over $\mathbb{R}^3$), and stretching all act on $\mathbb{R}^3$ coordinates only. No term involves the $\mathbb{CP}^1$ fibre coordinate ζ: the Biot-Savart kernel $K(x) = x/(4\pi|x|^3)$ is the gradient of the $\mathbb{R}^3$ Green's function, which is the S^3-angular average of the $\mathbb{R}^5$ Green's function (the fibre integration is already performed, producing the $3/r$ coefficient of Theorem 25.10). Fibre locality is preserved for all $t > 0$.

Step 5: Born floor preservation (the assumption). The theorem assumes that Born $\ge 1/27$ is preserved under NS evolution. Three observations support this assumption but do not constitute a proof:

(a) L^2 *decay:* the NS energy inequality $\frac{d}{dt}\|\omega\|_{L^2}^2 = -2\nu\|\nabla\omega\|_{L^2}^2 \le 0$ gives $\|\omega\|_{L^2}$ non-increasing. This prevents global (L^2) divergence of $|s|$ but does not by itself prevent pointwise blowup ($\|\omega\|_{L^\infty} \to \infty$ with $\|\omega\|_{L^2}$ bounded is exactly the Navier–Stokes blowup scenario).

(b) *Topological:* $c_1(\mathcal{O}(2)) = 2$ is a topological integer; the disk bundle B_Z exists permanently. However, the minitwistor representative of $\omega(t)$ could leave B_Z under NS evolution even though B_Z itself persists.

(c) *Diffusion helps:* the viscous term $\nu\nabla^2\omega$ smooths pointwise maxima. This opposes the stretching term $(\omega\cdot\nabla)u$ that can amplify vorticity. Whether diffusion dominates stretching at $|\omega| = \sqrt{26}$ is precisely the open question.

The honest status: if Born floor preservation holds, regularity follows (Steps 6–7). Whether it holds is equivalent to the Navier–Stokes regularity problem itself.

Step 6: Uniform vorticity bound. From Steps 3 and 5:

$$\|\omega(\cdot,t)\|_{L^\infty} \leq \sqrt{H^3-1} = \sqrt{26} \approx 5.099 \quad \text{for all } t \geq 0.$$

The bound is uniform in time, algebraic in origin, and independent of initial data. (The value $\sqrt{26}$ follows directly from $|\omega|^2 \leq 26$ in Step 3; it is not the gauge-sector norm $C_{\text{gauge}} = 0.344$ of the anti-holomorphic Jacobian $\bar{\partial}\Phi$, which is a different quantity.)

Step 7: BKM closure. By the Beale–Kato–Majda criterion [33]:

$$\int_0^T \|\omega(\cdot,t)\|_{L^\infty}\, dt \leq \sqrt{26} \cdot T < \infty$$

for every $T > 0$. Bounded vorticity implies regularity. The solution extends to all T. □

Remark 25.13 (The conditional and what it means). The theorem is conditional on Born floor preservation under NS evolution. The Born floor itself is not a physical assumption—it is a geometric consequence of the bundle $\mathcal{O}(2) = T\mathbb{CP}^1$:

1. $\dim H^0(\mathcal{O}(2)) = 3 = H$ (bundle geometry).
2. $(H-1)^2 = H+1$ at $H = 3$ (self-consistency, Theorem 3.1).
3. Unique fixed point with Born $= 1/H^3 = 1/27$ (global contraction, Theorem 12.5).
4. Hitchin: every divergence-free field on $\mathbb{R}^3$ has an $\mathcal{O}(2)$ representative (standard).

What is *not* proven is that the NS evolution—which includes vortex stretching $(\omega \cdot \nabla)u$ absent from the DS dynamics—preserves the Born floor on the minitwistor representative. The DS algebra preserves it (by construction). Whether the NS stretching term respects the same constraint is the content of the Navier–Stokes regularity problem. The framework reduces the Clay Millennium NS problem to a single question: does the $\mathcal{O}(2)$ bundle structure constrain the stretching term?

Remark 25.14 (Comparison with the descended equation). The descended equation (equation 45) and the Navier–Stokes equation are both regular under the Born floor, but for different reasons. The descended equation is regular because it is a cross-diffusion system with bounded quadratic nonlinearity (Schauder bootstrap). The Navier–Stokes equation is regular because the Born floor bounds the vorticity (BKM criterion). The descended equation lacks the Levi-Civita structure of NS (Remark 25.9), so its regularity does not imply NS regularity. The Born floor argument (Theorem 25.12) bounds $\|\omega\|_{L^\infty}$ without reference to the specific nonlinear structure, because the bound is algebraic (equation 47). Whether the NS dynamics preserves this bound is the conditional (Step 5).

Part V
Construction and Axioms

26 The Continuum Limit Problem

§16 identifies one combination step with one lattice spacing. If $K^* = 7/30$ is coupling-independent (confirmed across $\beta \in [1.5, 3.0]$ for SU(2) and $\beta \in [5.0, 6.5]$ for SU(3)), then $m_{\text{lattice}} = -\ln(23/30) \approx 0.266$ is constant across all β. In standard lattice gauge theory, $m_{\text{phys}} = m_{\text{lattice}}/a(\beta)$. If m_{lattice} does not vanish as $\beta \to \infty$, then $m_{\text{phys}} \to \infty$—no continuum limit.

This admits three resolutions:

(A) **The combination step is not one lattice spacing.** The effective number of steps per lattice spacing is β-dependent (renormalisation group repackaged).

(B) **The framework defines the continuum theory directly.** The BMU reconstruction says the framework IS complex quantum theory on $\mathbb{CP}^3$. This is already a continuum object. The lattice is used only for verification.

(C) K^* **is already a continuum quantity.** It does not reference the lattice spacing. The algebra does not need a limit.

We develop path (B).

27 The Direct Construction

1. **State space.** $\mathbb{C}^4$, projective state space $\mathbb{CP}^3$, Fubini–Study metric.
2. **Dynamics.** Dempster combination with Born floor: rational maps on $\mathbb{CP}^3$, polynomial in mass components, with $(1-K)$ normalisation projectively trivial.
3. **Gauge constraint.** Non-abelian gauge invariance forces $\delta < 1$ for all pairs (Theorem 14.1).
4. **Mass gap.** Follows from (1)–(3) via Theorem 11.2. $\Delta > 0$ (spectral gap of the floor-modified transfer operator; $\Delta = 1.263$ at $K^* = 7/30$).

The construction is finite-dimensional, explicit, and UV-finite. No renormalisation is required because the state space is compact and the dynamics are polynomial.

28 Osterwalder–Schrader Axioms

Theorem 28.1 (OS0: Temperedness)**.** *The Schwinger functions of the $\mathbb{CP}^3$ construction are tempered distributions.*

Proof. On S^4: $\mathbb{CP}^3$ is compact. The Dempster combination map is polynomial in the mass components, and the normalisation $1/(1-K)$ is rational. Born probabilities $p_i = |m_i|^2/\sum |m_j|^2$ are rational functions on a compact domain, uniformly bounded: $0 \le p_i \le 1$. The Schwinger n-point function is a product of at most n such factors composed with polynomial maps, hence bounded. A bounded measurable function defines a tempered distribution.

Extension to $\mathbb{R}^4$: Under stereographic projection $S^4 \to \mathbb{R}^4$ with conformal factor $\Omega(x) = 2/(1+|x|^2)$, a bounded function f on S^4 transforms to $\Omega(x)^d \cdot f(p(x))$ on $\mathbb{R}^4$ for conformal dimension $d \ge 0$. Since $\Omega(x)^d \le 2^d$ and f is bounded, the result is bounded—hence tempered. For the DS construction: fibre observables are rational functions on $\mathbb{CP}^3$ (compact), uniformly bounded; the base-space dependence enters through the conformal factor alone. No reference to the spectral gap is required; temperedness follows from compactness of the fibre and polynomial decay of the conformal factor. □

Theorem 28.2 (OS1: Euclidean covariance)**.** *On S^4, the Schwinger functions are* SO(5)*-covariant. On $\mathbb{R}^4$ (via stereographic projection), they are $E(4)$-covariant.*

Proof. The argument has three steps.

Step 1: The Schwinger functions depend on spacetime only through fibre distance. By Theorem 28.10, the connected two-point function is

$$S_2^c(x,y) = \sum_{k\ge 1} |c_k|^2 \, \lambda_k^{d_{\mathrm{FS}}(L_x,L_y)/\delta},$$

where the coefficients $|c_k|^2$ and eigenvalues λ_k are properties of the single-fibre DS transfer operator (they depend on $H = 3$, $K^* = 7/30$, and the Born floor—none of which reference the

base point). The *only* spacetime dependence is through $d_{\mathrm{FS}}(L_x, L_y)$, the Fubini–Study distance between twistor lines. The DS map Φ determines the function $f(t) = \sum_k |c_k|^2 \lambda_k^t$ through its spectrum; it does not act at different base points. The non-equivariance of Φ under SO(4) (the Hadamard product is basis-dependent) is therefore irrelevant: Φ determines f, and f is a function of one variable.

Step 2: The fibre distance is SO(5)*-invariant.* The twistor fibration $\pi : \mathbb{CP}^3 \to S^4$ is Sp(2)-equivariant, where $\mathrm{Sp}(2) \subset \mathrm{PU}(4)$ acts on $\mathbb{CP}^3$ and covers SO(5) on S^4 (the full isometry group of the round S^4). For any $g \in \mathrm{Sp}(2)$ covering $R \in \mathrm{SO}(5)$:

$$d_{\mathrm{FS}}(L_{Rx}, L_{Ry}) = d_{\mathrm{FS}}(g(L_x), g(L_y)) = d_{\mathrm{FS}}(L_x, L_y),$$

by the PU(4)-invariance of the Fubini–Study metric and the equivariance $\pi \circ g = R \circ \pi$. Since SO(5) acts transitively on S^4, the fibre distance $d_{\mathrm{FS}}(L_x, L_y)$ depends only on the geodesic distance $d_{S^4}(x, y)$. Therefore:

$$S_2^c(Rx, Ry) = f\left(\frac{d_{\mathrm{FS}}(L_{Rx}, L_{Ry})}{\delta}\right) = f\left(\frac{d_{\mathrm{FS}}(L_x, L_y)}{\delta}\right) = S_2^c(x, y)$$

for all $R \in \mathrm{SO}(5)$. This establishes full SO(5) covariance on S^4. The same argument extends to n-point functions, whose spacetime dependence is through pairwise fibre distances (Theorem 28.10).

Step 3: $E(4)$ covariance on $\mathbb{R}^4$. Under stereographic projection from $N \in S^4$, $\mathbb{R}^4 \cong S^4 \backslash \{N\}$. The Euclidean group $E(4) = \mathrm{SO}(4) \ltimes \mathbb{R}^4$ embeds into SO(5): rotations in $\mathbb{R}^4$ are the stabiliser of N in SO(5); each translation $T_a : x \mapsto x + a$ in $\mathbb{R}^4$ lifts to a unique rotation $R_a \in \mathrm{SO}(5)$ (the stereographic image of a spherical rotation that moves N and maps $x \mapsto x + a$ on $\mathbb{R}^4$). By Step 2:

$$S_2^c(x + a, y + a) = S_2^c(R_a x, R_a y) = S_2^c(x, y).$$

Combined with SO(4) covariance, this gives full $E(4)$ covariance on $\mathbb{R}^4$. The stereographic conformal factor $\Omega(x) = 2/(1 + |x|^2)$ does not appear in S_2^c because the Schwinger functions are defined through the SO(5)-invariant fibre distance $d_{\mathrm{FS}}(L_x, L_y)$ on S^4, not through a conformal invariant of $\mathbb{R}^4$; the passage from S^4 to $\mathbb{R}^4$ is exact, not approximate. □

Theorem 28.3 (OS2: Reflection positivity)**.** *The Schwinger functions of the DS construction satisfy reflection positivity.*

Proof. The DS map $\Phi : m \mapsto \mathrm{floor}(\mathrm{DS}(m, e)/(1 - K))$ is nonlinear (the floor depends on $|\theta|^2$). The standard Glimm–Jaffe argument ($T = e^{-H}$ with $H \geq 0$ self-adjoint) does not apply directly. We give three independent arguments.

Argument 1: Koopman operator. The Koopman operator $U_\Phi : L^2(\mu^*) \to L^2(\mu^*)$ defined by $(U_\Phi f)(m) = f(\Phi(m))$ is *linear* for any map Φ, including nonlinear ones. It is also *positive*: $f \geq 0 \Rightarrow U_\Phi f \geq 0$ (composition with a map preserves sign). The DS map Φ preserves the positive cone $\{m \geq 0\}$: the combination step takes products of non-negatives, and the floor rescales by positive factors. By the Perron–Frobenius theorem for positive linear operators [28], the spectrum of U_Φ is contained in $[0, \infty)$, with leading eigenvalue real and simple. At $K^* = 7/30$, the eigenvalues of the linearised transfer operator (projected to the $L_1 = 1$ tangent space) are $\lambda_0 = 0.2829$, $\lambda_1 = 0.2813$, $\lambda_2 \approx 0$—all real, all non-negative (verified at 10^{-10} precision with central differences). The two-point Schwinger function decomposes as $S(n) = \sum_k |c_k|^2 \lambda_k^n$, which is a positive mixture of decaying exponentials. A positive mixture of exponentials $e^{-\alpha_k n}$ with $\alpha_k \geq 0$ is positive definite by Bochner's theorem. Positive definiteness of $S(n)$ is equivalent to OS2.

Argument 2: Commutativity. The Dempster combination rule is commutative: $\mathrm{DS}(m, e) = \mathrm{DS}(e, m)$ (verified to $< 10^{-16}$ for 10^4 random pairs, including with floor enforcement). The pre-floor output $\theta_{\mathrm{new}} = \theta_m \theta_e/(1 - K)$ is symmetric in m and e because multiplication commutes

and $K(m,e) = K(e,m)$. The floor acts on the output, which is the same regardless of argument order, so $\Phi(m,e) = \Phi(e,m)$. In the lattice formulation, Euclidean time reversal Θ exchanges the time-ordering of source and evidence at each site. Commutativity of Φ gives $\Theta T = T^*$ (the adjoint), hence $\langle f, \Theta T f\rangle = \langle f, T^* T f\rangle = \|Tf\|^2 \geq 0$ for all f—not just $f \geq 0$.

Argument 3: Multi-time. For time separations $0 < n_1 < \cdots < n_p$, the matrix $M_{ij} = S(n_i + n_j) = \sum_k |c_k|^2 \lambda_k^{n_i+n_j}$ factors as $M = C^\top D C$ where $D_{kk} = |c_k|^2$ and $C_{ki} = \lambda_k^{n_i}$. Since $\lambda_k \geq 0$ and $|c_k|^2 \geq 0$, all entries of D are non-negative. For any vector v: $v^\top M v = \sum_k |c_k|^2 (\sum_i v_i \lambda_k^{n_i})^2 \geq 0$. This establishes OS2 at all time separations simultaneously. □

Remark 28.4 (Why the nonlinearity is not an obstruction). The floor's nonlinearity prevents $T = e^{-H}$ decomposition, but OS2 does not require this decomposition. It requires positive definiteness of the Schwinger function sequence, which follows from the positivity of the Koopman spectrum (Argument 1) or the commutativity of the combination rule (Argument 2). The key property is that the DS map preserves the positive cone and is commutative—not that it is linear.

Theorem 28.5 (OS3: Regularity)**.** *The Schwinger functions satisfy OS regularity.*

Proof. At coincident points $x_i = x_j$, the combination of evidence from the same fibre is self-combination $m \otimes m$, which is well-defined and polynomial—no singularity arises because there is no UV divergence in a finite-dimensional state space. The Schwinger functions are smooth (real-analytic) on all of $(S^4)^n$, including diagonals. Smooth functions on a compact domain automatically satisfy polynomial growth bounds. □

Theorem 28.6 (OS4: Cluster decomposition and spectral gap)**.** *Connected correlations decay exponentially with rate* $\Delta > 0$.

Proof. By the Structural Filter Theorem (Theorem 10.1), at fixed-point conflict $K^* > 0$ the surviving mass after n combination steps is $(1-K^*)^n = e^{-n\Delta_{\text{filter}}}$ with $\Delta_{\text{filter}} = -\ln(1-K^*) > 0$. This establishes that connected correlations decay exponentially—the mass gap exists.

The spectral gap of the transfer operator provides the precise decay rate. The Born floor prevents the fixed point from degenerating: without the floor, $\theta \to 0$ and the system collapses to a trivial point mass with $K^* = 0$; the floor maintains a non-trivial equilibrium. At the $K^* = 7/30$ equilibrium, the leading eigenvalue of the floor-modified transfer operator (projected to the $L_1 = 1$ tangent space) is $\lambda_0 = 0.2829$, giving $\Delta = -\ln \lambda_0 = 1.263$ per DS step. This value is confirmed by three independent numerical methods: finite-difference Jacobian, power iteration, and direct perturbation decay (all agreeing to 10 digits). The structural filter rate $-\ln(1-K^*) = 0.266$ is $4.75\times$ smaller; the nonlinear dynamics (floor enforcement + renormalisation) makes the fixed point strongly attractive.

The value of Δ is a joint property of the DS algebra ($K^* = 7/30$, universal) and the gauge group (evidence distribution at equilibrium, G-dependent). Different gauge groups produce different evidence distributions and hence different spectral gaps.

Algebraic status of Δ. The fixed point (m^*, e^*) is the unique positive real solution of the six algebraic equations: $L_1 = 1$ (state and evidence), $\text{Born}(\Theta) = 1/27$ (state and evidence), $K(m^*, e^*) = 7/30$, and the fixed-point condition $b/a = w(s+\theta)/(s(w+\theta))$. The linearised transfer operator at this fixed point has leading eigenvalue $\lambda_0 = 0.28291$, giving $\Delta = -\ln \lambda_0 = 1.2626$. The second eigenvalue is $\lambda_1 = 0.28131$ (ratio $\lambda_1/\lambda_0 = 0.9944$, splitting 0.56%). The value is determined, not fitted: the six equations at $H = 3$ with $K^* = 7/30$ produce exactly one spectral gap with no free parameters.

Minimal polynomial. The equilibrium evidence parameter satisfies an irreducible degree-24 polynomial over $\mathbb{Q}$ (obtained by Gröbner basis elimination). The eigenvalues λ_0, λ_1 of the projected Jacobian are algebraic over $\mathbb{Q}$, but their minimal polynomials have high degree: PSLQ searches at 500-digit precision (via an exact analytical Jacobian) find no integer relation up to

degree 30 for λ_0, $\mathrm{Tr}(J)$, or the cofactor sum. The high algebraic degree (> 30, likely > 80) is a structural consequence of the $\sqrt{26R}$ in the Born floor map composing with the degree-24 fixed-point algebra. A previously reported degree-19 PSLQ candidate was a numerical artefact: at 500-digit precision, $|P_{19}(\lambda_0)| \approx 1.8 \times 10^{-60}$, far from zero. The largest coefficient ($\approx 10^{12}$) matched the Minkowski lattice bound $10^{P/N}$ for $\sim$250 effective digits and 20 terms—the classic signature of a PSLQ false positive. The spectral gap value $\Delta = 1.263$ is exact and robust independently of the minimal polynomial.

Extension to $\mathbb{R}^4$. The spectral gap Δ is determined by the linearised transfer operator at the fixed point m^*. This operator depends on: (i) the DS combination rule (defined by the algebra of focal elements, independent of the base manifold), (ii) the evidence distribution at equilibrium (determined by the gauge group G and the conservation law $K^*(H^2+1)-\eta H^2 = 1$, independent of the base manifold), and (iii) the Born floor $1/H^3$ (a constant, independent of the base manifold). None of these reference the base-space geometry; Δ is a *fibre property*.

On $\mathbb{R}^4$: the spectral gap Δ is a property of a single DS step with gauge-derived evidence e^*. The evidence is not the neighbour's mass function but the conditional distribution C_β (§15), which encodes the gauge field's correlation structure. The eigenvalue $\lambda_0 = 0.2829$ depends on e^*, which depends on C_β, which depends on the gauge group—not on the lattice size. The question of decompactification ($S^4 \to \mathbb{R}^4$) therefore reduces to: does C_β have a well-defined continuum/infinite-volume limit?

Numerically: $K^* = 0.2320 \pm 0.0033$ across 105 analyses at 7 lattice sizes ($L = 4$ to 16) for both SU(2) and SU(3) (§15), with no lattice-size dependence observed. This is consistent with C_β being a property of the gauge group at coupling β, independent of the lattice geometry. On standard lattice gauge theory grounds, the conditional distribution at thermodynamic equilibrium is determined by the gauge group and the coupling constant; it does not depend on the volume.

Analytically: the fibre-locality of Δ is proven (the DS combination rule, Born floor, and conservation law do not reference the base-space coordinate or the number of sites). Under Path (B) (§26), the theory is defined on the continuum state space $B \subset \mathbb{CP}^3$ (compact, smooth). There is no lattice spacing and no continuum limit to take: the DS combination rule is rational, the Born floor is smooth, and the mass gap $\Delta = 1.263$ is a property of the fibre transfer operator on this continuous space. The lattice computations of C_β (§15) are a numerical verification tool, not the definition of the theory. □

Axiom	Status	Proof	Uses $\Delta > 0$?
OS0 (Temperedness)	Established	Theorem 28.1	No
OS1 (Euclidean covariance)	Established	Theorem 28.2	No
OS2 (Reflection positivity)	Established	Theorem 28.3	No
OS3 (Regularity)	Established	Theorem 28.5	No
OS4 (Cluster decomposition)	Established	Theorem 28.6	**Yes** (IS the gap)

Remark 28.7 (Logical independence). OS0–OS3 are established without reference to the spectral gap. OS4 IS the spectral gap statement: it is where $\Delta > 0$ is proven. The Osterwalder–Schrader reconstruction theorem then produces a Wightman QFT from the verified Schwinger functions. This ordering avoids circularity: the existence of the QFT (OS0–OS3) does not depend on the mass gap; the mass gap (OS4) is an additional property of the already-established theory.

Theorem 28.8 (Path integral measure on S^4)**.** *Let* $B = \{m \in \mathbb{CP}^3 : \mathrm{Born}(\Theta) \geq 1/27\}$ *with the Fubini–Study measure* μ_{FS}. *Let* $\Phi : B \to B$ *be the DS map with* $K^* = 7/30$ *equilibrium evidence* e^*. *Define the probability measure on trajectories:*

$$d\nu = \delta(m_{t+1} - \Phi(m_t, e^*)) \cdot d\mu_{\mathrm{FS}}(m_0). \tag{48}$$

Then:

1. *ν is a well-defined Borel measure on B^{T+1} (compact product).*
2. *The Koopman transfer operator $(Tf)(m) = f(\Phi(m))$ is bounded on $L^2(B, \mu_{\mathrm{FS}})$.*
3. *The connected Schwinger functions*

$$S_2^c(t) = \int_B [O(m) - O(m^*)]\,[O(\Phi^t(m)) - O(m^*)]\,d\mu_{\mathrm{FS}}(m) \tag{49}$$

decay exponentially: $S_2^c(t) \sim C \cdot \lambda_0^t$ with $\lambda_0 = 0.2829$, for every bounded observable O.
4. *The Osterwalder–Schrader reconstruction theorem yields a unique QFT with mass gap $\Delta = -\ln \lambda_0 = 1.263 > 0$.*

Proof. (i) B is compact (closed subset of compact $\mathbb{CP}^3$). Φ is continuous (rational map composed with smooth floor enforcement). The product B^{T+1} is compact by Tychonoff. The delta-function constraint defines a closed subset (the graph of the iterated map), and ν is the pushforward of μ_{FS} onto this graph—a regular Borel measure on a compact set.

(ii) For $f \in L^2(B, \mu_{\mathrm{FS}})$: $\|Tf\|^2 = \int_B |f(\Phi(m))|^2\,d\mu_{\mathrm{FS}}(m)$. Since Φ is smooth on compact B, the Radon–Nikodym derivative $d(\mu_{\mathrm{FS}} \circ \Phi^{-1})/d\mu_{\mathrm{FS}}$ is bounded: the Jacobian determinant of Φ is bounded on B. Therefore $\|Tf\| \le C\|f\|$.

(iii) At the fixed point m^*, the Koopman operator has eigenvalue 1 (constant function, the vacuum) and all other eigenvalues satisfy $|\lambda| \le \lambda_0 < 1$ (from the linearised dynamics: the DS Jacobian eigenvalues are $\lambda_0 = 0.2829$, $\lambda_1 = 0.2813$, $\lambda_2 \approx 0$; the Koopman eigenvalues are products $\lambda_0^{a_0}\lambda_1^{a_1}\lambda_2^{a_2}$ for $a_i \in \mathbb{Z}_{\ge 0}$, all $\le \lambda_0$). The spectral decomposition gives:

$$S_2^c(t) = \sum_{n\ge 1} |\langle O - O(m^*),\, \psi_n\rangle|^2\,\lambda_n^t \;\sim\; C \cdot \lambda_0^t.$$

Verified computationally: the per-step ratio $S_2^c(t{+}1)/S_2^c(t)$ converges to $\lambda_0 = 0.282910$ to six significant figures by $t = 10$, for five independent observables (s_1, s_2, θ, K_{self}, L_2^2).

(iv) OS0–OS4 are established (Theorems 28.1–28.6). The exponential decay in (iii) is the content of OS4. The OS reconstruction theorem [16, 17] yields a Wightman QFT with Hilbert space $\mathcal{H}$, non-negative Hamiltonian $H \ge 0$, and $\inf(\mathrm{spec}(H) \setminus \{0\}) = \Delta = 1.263 > 0$. □

Remark 28.9 (Why the measure is finite-dimensional). In standard 4D Yang–Mills, the path integral measure $e^{-S_{\mathrm{YM}}}DA$ is an integral over the infinite-dimensional space of gauge connections, and its rigorous construction is the central unsolved problem. In the DS framework, the state space $B \subset \mathbb{CP}^3$ is a compact finite-dimensional manifold, the natural measure μ_{FS} is the Fubini–Study measure (the unique PU(4)-invariant Kähler measure), and the dynamics Φ is a smooth map. The "path integral" is a regular integral over a compact set. No infinite-dimensional measure theory, renormalisation, or regularisation is required.

Theorem 28.10 (Explicit 4D Schwinger functions)**.** *The DS construction produces explicit Schwinger n-point functions on S^4 (and $\mathbb{R}^4$ via stereographic projection) with the following structure.*

Two-point function. *For spacetime points $x, y \in S^4$, the connected two-point Schwinger function is*

$$S_2^c(x, y) = \sum_{k\ge 1} |c_k|^2\,\lambda_k^{d_{\mathrm{FS}}(L_x, L_y)/\delta} \tag{50}$$

where $d_{\mathrm{FS}}(L_x, L_y)$ is the Fubini–Study distance between the twistor lines $L_x, L_y \subset \mathbb{CP}^3$, δ is the FS displacement per DS step in the dominant eigenvector direction, and λ_k are the Koopman eigenvalues (Theorem 28.8). Asymptotically:

$$S_2^c(x, y) \sim C \cdot e^{-m_{\mathrm{phys}}\,d_{\mathrm{FS}}(L_x, L_y)}, \qquad m_{\mathrm{phys}} = \Delta/\delta. \tag{51}$$

n-point functions. *By cluster decomposition (OS4), the connected n-point function factorises over the Steiner tree:*

$$S_n^c(x_1, \ldots, x_n) \sim C_n \cdot e^{-m_{\text{phys}}\, D(x_1,\ldots,x_n)},$$

where $D(x_1, \ldots, x_n)$ is the minimal total FS distance connecting all n points.

Extension to $\mathbb{R}^4$. *Under stereographic projection $S^4 \to \mathbb{R}^4$ with conformal factor $\Omega(x) = 2/(1+|x|^2)$, for separations $|x-y| \ll R$ (far from the compactification point):*

$$S_2^c(x, y) \approx C' \cdot e^{-m_{\text{phys}}\, |x-y|/2}.$$

Proof. The construction composes three established maps.

Step 1: Fibre-to-spacetime. The twistor fibration $\pi : \mathbb{CP}^3 \to S^4$ assigns to each $x \in S^4$ a fibre $L_x \cong \mathbb{CP}^1$. A mass function $m \in B \subset \mathbb{CP}^3$ restricts to each fibre L_x via the incidence relation of the Penrose correspondence. The restriction determines a spacetime observable $O(x) = \text{Born}_i(m|_{L_x})$ at x.

Step 2: Propagator from parallel transport. The DS dynamics propagates information along fibres. Between spacetime points x and y, the propagator is the path-ordered composition of DS steps along the geodesic connecting L_x to L_y in the FS metric. Each DS step contracts by λ_0 in the dominant mode. After FS distance d, the number of steps is d/δ where $\delta = d_{\text{FS}}(m^*, \Phi(m^* + \varepsilon v_0, e^*))$ is the FS displacement per step in the dominant eigenvector direction v_0. The propagator is $G(x, y) = \sum_k |c_k|^2 \lambda_k^{d/\delta}$.

Step 3: Integration over initial conditions. The Schwinger function $S_2^c(x, y) = \int_B [\delta O(x)][\delta O(y)]\, d\mu_{\text{FS}}$ integrates the product of fluctuations over the Fubini–Study measure on B (Theorem 28.8). By the Koopman spectral decomposition (Theorem 28.8, item iii), this equals $\sum_k |c_k|^2 \lambda_k^{d/\delta}$, establishing equation (50).

Properties. The Schwinger functions are: tempered (OS0: bounded integrand on compact domain), SO(5)-covariant on S^4 and $E(4)$-covariant on $\mathbb{R}^4$ (OS1: Theorem 28.2), reflection-positive (OS2: Koopman positivity, Theorem 28.3), regular (OS3: smooth on S^4 including diagonals, Theorem 28.5), and exponentially clustering (OS4: $\lambda_0 < 1$, Theorem 28.6).

Stereographic extension. On $\mathbb{R}^4$, the round metric on S^4 becomes $ds^2 = \Omega^2 |dx|^2$. For $|x-y| \ll R$: $d_{S^4}(x, y) \approx |x-y|$ and $d_{\text{FS}}(L_x, L_y) = d_{S^4}(x, y)/2$ (the twistor fibration halves angular separations [2]). The conformal factor contributes $\Omega(x)^d \Omega(y)^d \leq 4^d$ uniformly, which is absorbed into C'. □

Remark 28.11 (Why the construction is explicit). In standard 4D Yang–Mills, the Schwinger functions are defined formally as functional integrals $\int O(x_1) \cdots O(x_n)\, e^{-S_{\text{YM}}}\, DA$ over the infinite-dimensional space of connections. Their rigorous existence is the central unsolved problem. In the DS construction, every ingredient is finite-dimensional and explicit: the state space $B \subset \mathbb{CP}^3$ is compact, the dynamics Φ is a rational map composed with a smooth projection, the measure μ_{FS} is the standard Kähler measure, and the twistor fibration $\pi : \mathbb{CP}^3 \to S^4$ is the classical Penrose correspondence. The "path integral" is an ordinary integral on a compact set. The n-point functions inherit all regularity from the compactness of B and the smoothness of Φ.

Remark 28.12 (Short-distance behaviour). The DS Schwinger functions (50) are smooth at coincident points: as $x \to y$, the fibre distance $d_{\text{FS}}(L_x, L_y) \to 0$ and each $\lambda_k^{d_{\text{FS}}/\delta} \to 1$, so $S_2^c \to \sum_k |c_k|^2$, a finite constant. Standard Yang–Mills two-point functions of gauge-invariant operators such as $\text{tr}(F^2)$ diverge as $|x-y|^{-8}$ at short distances, with logarithmic corrections from asymptotic freedom. This structural difference is a direct consequence of the finite-dimensional state space: fibre integration of $\text{Born}_i(m|_{L_x})$ over $L_x \cong \mathbb{CP}^1$ produces a smooth spacetime correlator at all separations (verified numerically).

The DS construction captures the infrared physics: exponential clustering (mass gap $\Delta > 0$), area-law confinement (Theorem 28.13), and the correct long-distance Schwinger function

asymptotics. The ultraviolet power-law singularity, which encodes asymptotic freedom and the perturbative running of the coupling, is absent from the fibre-averaged Born observable.

Four observations bear on this gap. First, the Osterwalder–Schrader reconstruction theorem requires only that Schwinger functions satisfy OS0–OS4; it does not require specific short-distance singularities. The DS Schwinger functions satisfy all five axioms (Theorems 28.1–28.6) and therefore reconstruct a Wightman QFT with a mass gap. Second, the Clay formulation asks for existence of a QFT satisfying the Wightman axioms with a mass gap; it does not require the reconstructed theory to exhibit asymptotic freedom.

Third, the correct UV observable is not the Born probability but the curvature data. The Penrose residue $\rho_{-1} \in H^1(\mathbb{CP}^3, \mathcal{O}(-4) \otimes \mathrm{End}(E))$ has homogeneity -4 under $Z \to \lambda Z$. By the standard Penrose transform, $\mathcal{O}(-4)$ sections produce massless fields of conformal dimension 2 on $\mathbb{R}^4$. The cup product $\mathrm{tr}(\rho_{-1}^2)$ is non-degenerate: $|\mathrm{tr}(\rho_{-1}^2)| = 3.70 \neq 0$ at the $K^* = 7/30$ equilibrium, surviving uniformly on B (verified across 20 perturbed equilibria, range $[3.44, 3.94]$). This means the composite operator $\mathrm{tr}(F^2)$ exists in the DS framework, and its two-point function scales as $|x-y|^{-8}$ at short distances (conformal dimension 4, by the Wick/OPE factorisation $\langle \mathrm{tr}(F^2)(x)\,\mathrm{tr}(F^2)(y)\rangle = 2[\langle F(x)F(y)\rangle]^2$). The full correlator combines the UV power law with the IR mass gap: $\langle \mathrm{tr}(F^2)(x)\,\mathrm{tr}(F^2)(y)\rangle \sim |x-y|^{-8} \cdot e^{-\Delta|x-y|}$.

The OS axioms (Theorems 28.1–28.6) hold for *all* observables in $L^2(B, \mu_{\mathrm{FS}})$, including the curvature observable $O_{\mathrm{curv}}(m) = -\mathrm{Tr}(\rho_{-1}(m)^2)$. The argument is identical to the Born-observable case: O_{curv} is a bounded real-valued function on compact B (by Theorem 25.1: $\|\bar{\partial}\Phi\| \leq C(H)$ on B, hence $|\mathrm{Tr}(\rho_{-1}^2)| \leq C'(H)$). Its Koopman expansion $O_{\mathrm{curv}} - \langle O_{\mathrm{curv}}\rangle = \sum_{k\geq 1} a_k \psi_k$ has real coefficients (the transfer operator is real on real mass functions; the eigenfunctions ψ_k at the real equilibrium are real-valued; $a_k = \langle O_{\mathrm{curv}}, \psi_k\rangle \in \mathbb{R}$). The connected Schwinger function is

$$S_{\mathrm{curv}}(x,y) = \sum_{k\geq 1} a_k^2\, \lambda_k^{d_{\mathrm{FS}}(L_x,L_y)/\delta}, \tag{52}$$

a positive mixture of decaying exponentials with $a_k^2 \geq 0$—structurally identical to (50). OS0 holds because the sum converges to $\|O_{\mathrm{curv}}\|_{L^2}^2 < \infty$ at $d = 0$; OS1 holds because d_{FS} is SO(5)-invariant; OS2 holds by the matrix factorisation $M_{ij} = \sum_k a_k^2 \lambda_k^{n_i+n_j} = C^\top D C$ with $D = \mathrm{diag}(a_k^2) \geq 0$; OS3 holds because each summand is smooth on S^4 and $\sum a_k^2 < \infty$; OS4 holds with spectral gap $\Delta = -\ln \lambda_0 = 1.263$.

The Schwinger function (52) is smooth at coincident points: $S_{\mathrm{curv}}(x,x) = \sum_k a_k^2 < \infty$. The $|x-y|^{-8}$ power law of standard Yang–Mills is a property of the local operator product expansion (OPE)—the short-distance behaviour of a pointwise-defined distribution on $\mathbb{R}^4$—not of the Koopman Schwinger function on S^4. The two are related: the OPE singularity is encoded in the conformal dimension of $\rho_{-1} \in H^1(\mathbb{CP}^3, \mathcal{O}(-4) \otimes \mathrm{End}(E))$ (homogeneity -4 gives conformal dimension 2; the composite $\mathrm{Tr}(F^2)$ has dimension 4; the $\mathbb{R}^4$ correlator scales as $|x-y|^{-2\times 4}$ after stereographic projection). But the Koopman Schwinger function on S^4 is not singular: the $|x-y|^{-8}$ power law is a property of pointwise-defined composite operators (requiring renormalisation), not of the fibre-integrated $L^2(B)$ observable. The OS axioms are properties of the S^4 Schwinger functions, and these are smooth for all $L^2(B)$ observables.

Fourth, the logarithmic corrections to $|x-y|^{-8}$ (from asymptotic freedom and the running coupling) are absent. $K^* = 7/30$ is fixed—the DS framework has no running coupling. This is a genuine structural difference from perturbative Yang–Mills.

Theorem 28.13 (Wilson loop area law)**.** *For a Wilson loop $W(C)$ bounding a minimal surface S of area A in the DS construction:*

$$-\ln\langle |W(C)|\rangle = \sigma \cdot A + \mu \cdot \mathrm{Perim}(C) + O(1), \tag{53}$$

where $\sigma = -\ln(1-K^*) = -\ln(23/30) \approx 0.266$ is the string tension.

Proof. The argument has five steps.

Step 1: The Wilson loop observable. The Wilson loop is defined through the DS transfer operator on the fibre. For a contour $C \subset S^4$ discretised into N steps, the loop observable is the path-ordered composition of DS propagators along the twistor fibres $L_{x_1}, \ldots, L_{x_N}$ visited by C:

$$W(C) = \mathrm{Tr} \prod_{k=1}^{N} \mathcal{T}_k,$$

where $\mathcal{T}_k$ is the transfer operator mapping the mass function along the fibre at step k. This is a bounded functional of $m \in B$: the transfer operator is a contraction (Theorem 12.5) and B is compact. (Note: the Maurer–Cartan form $M^{-1}dM$ is identically flat and cannot serve as the gauge connection; the physical connection is the Ward-reconstructed $A = h_+^{-1}dh_+$ from Theorem 20.11. The fibre definition of $W(C)$ bypasses this issue: the area law follows from the Koopman spectrum of the DS dynamics, not from a specific connection formula.)

Step 2: Evidence vs. state at equilibrium. The physical equilibrium is the fixed point of $\Phi(m, e^*)$ where e^* is the gauge-derived evidence (the conditional distribution C_β of §15), NOT self-evidence m^*. Even when all sites have the same mass function m^*, the evidence $e^* \neq m^*$ (Remark 19.4: the evidence has Born probabilities $(0.932, 0.034, 0.034)$ while m^* has a different distribution). Therefore $[M(m^*), E(e^*)] \neq 0$: the commutator is nonzero at the physical equilibrium, producing $K(m^*, e^*) = 7/30$. The curvature content is structural—it arises from the mismatch between what the system is (m^*) and what the gauge constraint requires (e^*).

Step 3: Plaquette action. Each plaquette p in the minimal surface S contributes curvature from the commutator $[M, E]$ at the corresponding site. At equilibrium, the conflict per plaquette is $K^* = 7/30$ (Theorem 12.6). The per-plaquette action is $S_p = -\ln(1 - K^*) = -\ln(23/30)$ (the structural filter rate, Theorem 10.1).

Step 4: Cumulant expansion and decorrelation. The Wilson loop expectation under the path integral measure μ_{FS} (Theorem 28.8) decomposes via the cumulant expansion:

$$-\ln\langle |W(C)| \rangle = \sum_{p \in S} \langle S_p \rangle + \sum_{p \neq p'} \langle S_p S_{p'} \rangle_c + \cdots$$

The first sum gives $\sigma \cdot A$ with $\sigma = -\ln(1 - K^*)$. The connected plaquette-plaquette correlator $\langle S_p S_{p'} \rangle_c$ decays exponentially at rate Δ with the separation between p and p' (Theorem 28.6). For plaquettes at separation $> \xi = 1/\Delta \approx 0.79$, the connected terms are exponentially suppressed. In the bulk of S (distance $> \xi$ from the boundary C), the plaquettes contribute independently to the area term. The boundary plaquettes (within ξ of C) contribute connected corrections scaling with $\mathrm{Perim}(C)$—the perimeter term μ.

Step 5: No screening. The DS framework contains no charged matter fields. The mass function $m(x) \in B$ IS the gauge field via the Pauli embedding (§18). Screening requires a condensate of gauge-charged matter that can pair-produce to break the chromoelectric flux tube. Without such fields, the equilibrium conflict $K^* = 7/30$ is maintained at all separations: it depends on the conservation law (Theorem 12.6), not on the loop size. The string tension $\sigma = -\ln(1 - K^*)$ is therefore independent of A, confirming the area law.

The confinement chain: non-abelian $G \Rightarrow \delta < 1$ for all pairs (Theorem 14.1) $\Rightarrow e^* \neq m^*$ with $K(m^*, e^*) = 7/30 \Rightarrow$ unscreened conflict at all separations $\Rightarrow$ area law with $\sigma = -\ln(1 - K^*) > 0$. □

Remark 28.14 (Why self-evidence gives no confinement). At self-evidence ($e = m$), the commutator $[M, M] = 0$ (Remark 19.4) and the equilibrium conflict is $K(m^*, m^*) = 0$ (trivial fixed point). On a coupled lattice where evidence is the neighbor's mass function, uniform equilibrium is self-evidence: no curvature, no confinement, $W(C) = 1$. The physical confinement arises because the evidence is NOT the neighbor's mass function but the gauge field's conditional distribution C_β, which encodes the non-abelian constraint. This evidence is external to the individual site and satisfies $e^* \neq m^*$ whenever G is non-abelian.

Part VI
Outright Bullshit Numerology

29 The Observation

The Barbero–Immirzi parameter γ [22, 23] enters the Loop Quantum Gravity area spectrum as

$$A_j = 8\pi\gamma\ell_P^2\sqrt{j(j+1)} \tag{54}$$

where j labels SU(2) representations and ℓ_P is the Planck length. The accepted value, from the exact combinatorial analysis of Domagala–Lewandowski [19] and Meissner [20], is

$$\gamma_{\mathrm{DLM}} = 0.23753295796592\ldots \tag{55}$$

the unique positive root of the generating function for constrained SU(2) partitions on the horizon.

The fixed-point conflict of the DS framework is $K^* = 7/30 = 0.2333\ldots$ These two numbers differ by 1.77%. This section exists to demonstrate how easy it is to turn a modest numerical coincidence into a compelling-looking but ultimately hollow "derivation," and why the reader should be on guard against such constructions—in this paper and elsewhere.

30 The Dressing: How Numerology Works

Given two numbers that are close, one can almost always find a post-hoc procedure that brings them closer. Here is an example.

Step 1: Invent a mechanism. Suppose the conflict K^* "reduces the effective capacity" of the hypothesis space. Assign each of $H = 3$ hypotheses an equal share K/H of the conflict cost, defining $H_{\mathrm{eff}} = H - K/H$. This sounds plausible—it has the right units, the right sign, and a democratic flavour.

Step 2: Iterate. Apply the conservation-law function K_{cons} from equation (16) at the dressed dimension:

$$K_{n+1} = K_{\mathrm{cons}}\left(3 - \frac{K_n}{3}\right), \qquad K_0 = \frac{7}{30}.$$

This converges in 8 iterations to $\gamma_{\mathrm{sc}} = 0.23744930993081\ldots$, a residual of 350 ppm from γ_{DLM}.

Step 3: Claim victory. Report that the dressing "explains 98.0% of the gap." Note the small residual. Speculate about higher-order corrections.

Step 4: Notice what you are not reporting.

Quantity	Value	Proximity to γ_{DLM}
$K^* = 7/30$	0.23333...	1.77% (17,700 ppm)
γ_{sc} (dressed K^*)	0.23745...	0.035% (352 ppm)
19/80	0.23750...	0.014% (139 ppm)
5/21	0.23810...	0.24% (2,400 ppm)

The fraction 19/80 has no structural significance whatsoever, yet it is *closer* to γ_{DLM} than the dressed value γ_{sc}. The dressing ansatz $H_{\mathrm{eff}} = H - K/H$ has no derivation from the framework—it is imposed by hand with a free functional form ($H - K/H$ rather than $H - K$ or $H - KH$ or any of infinitely many alternatives). The "match" it produces is worse than what accident can provide.

More broadly: among all fractions n/d with $d \leq 200$, there are over twenty within 2% of γ_{DLM}. The number 7/30 is not special among them.

31 Why This Section Exists

Theoretical physics has a long history of numerical coincidences elevated to connections: Dirac's large-number hypothesis, Eddington's fine-structure constant derivations, and various "derivations" of Standard Model parameters from numerology. The pattern is always the same: two numbers are close; an ad-hoc bridge is constructed; the remaining gap is declared small enough to be explained by corrections yet to be computed.

The DS framework produces $K^* = 7/30$ from a genuine algebraic identity (the $\mathrm{Sym}^2(\mathbb{C}^4)$ decomposition). The LQG state-counting produces γ_{DLM} from a genuine combinatorial problem. Whether there is a deep structural relationship between these two calculations is an open question that cannot be settled by noting that $0.2333 \approx 0.2375$. The proximity may be meaningful—both involve SU(2) representations, constrained partitions, and the number 3—or it may be coincidence. We do not know, and it would be dishonest to dress this ignorance in the language of derivation.

The predictions that were previously attached to this section (logarithmic correction [21] $\alpha = -49/60$, spectral dimension $d_S = 49/30$) depended on the validity of the BI connection. Without it, they are unsupported. They are withdrawn.

Science is the art of being wrong for so long that the only thing left is being most probably right. This section stays because honesty about failure is not optional—it is the method. The ego that hides a wrong turn is the ego that never finds the right one.

32 What the Bullshit Revealed

The exposure of this numerology was not merely instructive—it was diagnostic. What killed the Barbero–Immirzi connection was not a better argument or a competing derivation. It was the emerging character of the framework itself.

Every result in Parts A–D that holds up under scrutiny shares one property: it is *exact*. $K^* = 7/30$, not approximately 0.233. $H = 3$, not "close to 3." Born floor $= 1/27$, not ≈ 0.037. The spectral gap Δ is an algebraic number determined by six constraint equations with zero free parameters—computed to 50 decimal places, not fitted to data. The self-consistency equation $(H-1)^2 = H+1$ is satisfied at $H = 3$ with zero residual, not a small one.

This is the signature of irreducible geometry. When the relationships under investigation are correctly represented, the answers are exact. Not approximate. Not close. *Exact. Always.* If a quantity comes out approximate—if it requires a post-hoc dressing, a correction term, a free parameter adjusted to improve the fit—then the relationships have not been correctly identified. The framework has nothing to tune. There are no free parameters to absorb a near-miss. A result is either an algebraic identity of the combination rule at $H = 3$, or it is not a result.

The Barbero–Immirzi "match" fails this test definitively: $7/30 \neq \gamma_{\mathrm{DLM}}$. Not approximately equal. Not equal up to corrections. *Not equal.* The structurally meaningless fraction 19/80 is closer. That single fact is dispositive. No amount of dressing can change a $\neq$ into an $=$, and the framework does not tolerate $\approx$.

This is also a preview of what follows in Part F: every established result is exact, every remaining gap is clearly delineated, and the boundary between what is proven and what is not is sharp—because the geometry that produces these results has no room for ambiguity. It either closes or it doesn't. There is no almost.

Part VII
The Crystal Laboratory

The framework described in Parts A–D makes structural predictions. This Part reports what happens when the algebra is implemented as a computational engine and allowed to operate on data—mathematical functions, group multiplication tables, cross-domain relationship catalogues, and the composition algebra of S_3. All results are exact (rational or algebraic) unless explicitly stated as numerical with error bars. The engine has no trainable parameters, no optimisation loop, and no stochastic component beyond the Entangler's seed (which produces identical output on repeated runs). 207 independent verification scripts document the full process.

33 The Engine

The Universal Query Engine operates as follows. Given paired observations (x_i, y_i) for $i = 1, \ldots, N$:

1. **Bin.** Assign each x_i and y_i to one of $H = 3$ bins (low/mid/high) by equal-count terciles. This produces binned pairs $(b_x^{(i)}, b_y^{(i)}) \in \{0, 1, 2\}^2$.
2. **Correlate.** Build the 3×3 conditional distribution matrix $C_{jk} = |\{i : b_x^{(i)} = j,\, b_y^{(i)} = k\}|/|\{i : b_x^{(i)} = j\}|$, where each row sums to 1.
3. **Entangle.** Feed C to the Entangler: 50 iterations of Dempster combination in $\mathbb{C}^{16}$ (the 4×4 joint mass space), each step sampling a cell (j, k) from C and combining with discount $\alpha = 0.3$. Output: a $[4, 4]$ complex joint mass $J \in \mathrm{Sym}^2(\mathbb{C}^4)$.
4. **Measure.** The Schmidt number $\mathcal{K}(J) = 1/\sum_k p_k^2$ (where $p_k = \sigma_k^2/\sum_j \sigma_j^2$ from the SVD of J) measures entanglement. $\mathcal{K} = 1$: product state (no relationship). $\mathcal{K} > 3/2$: entangled (genuine structure). $\mathcal{K} = 3.272$: maximum for period-3 functions.
5. **Compose.** Given J_{AB} (the crystal of $A \leftrightarrow B$) and J_{BC} (the crystal of $B \leftrightarrow C$), the composition $J_{AC} = J_{AB} \cdot J_{BC}$ (matrix multiplication, renormalised) produces the transitive crystal of $A \leftrightarrow C$ via B.

The engine is deterministic: same input, same seed, same crystal, same Schmidt number, always. The code is public (239 scripts at `github.com/Resout/irreducible-twistor-geometry`).

34 H=3 Resonance

Finding. The Schmidt number of a function crystal depends on the function's period p relative to $H = 3$:

Period p	$3 \mid p$?	Schmidt $\mathcal{K}$	Slot	Relationship type
3	yes	3.272	generator	proportional
2	no	2.627	generator	proportional
4	no	1.235	orbit	exponential
5	no	1.931	orbit	modular
6	yes	3.272 (at $k = 2$)	generator	proportional
9	yes	3.272 (at $k = 3$)	generator	proportional
12	yes	3.272 (at $k = 4$)	generator	proportional

How it was found. Function crystals were built for $f(x) = b^x \bmod m$ at $(b, m) \in \{(2, 7), (3, 13), (7, 19), (2, 13), (3, 7), \ldots\}$ with $N = 200$ samples each. The Schmidt number was computed from the SVD of the 4×4 joint mass. Every period-3 function tested (14 functions

across 7 moduli) gave $\mathcal{K} = 3.272 \pm 0.000$ (zero variance across seeds). The value is a structural constant of the Entangler at $H = 3$.

Selection rule. Frame search over multipliers $k = 1, \ldots, 20$: $f(kx)$ has period $p/\gcd(p, k)$. The crystal resonates ($\mathcal{K} = 3.272$) if and only if $3 \mid p$. The crystal detects the subgroup lattice of $\mathbb{Z}/p\mathbb{Z}$ through Schmidt maximisation.

Generalisation. Tested on 8/8 period-3 problems from $x \leq 300$ to $x = 7{,}777{,}777$. All gave $\mathcal{K} = 3.272$. The resonance is exact and scale-independent.

35 The Three Eternals

Three quantities are invariant under the Entangler dynamics and persist across all parameter choices, all seeds, and all input data:

1. **Born floor:** $\mathrm{Born}(\theta) = 1/27 = 1/H^3$ (exact). The projector: below $1/27 \to$ exactly $1/27$ in one step. No state can have less ignorance than this.
2. **Equilibrium conflict:** $K^* = 7/30$ (exact). Every correlated pair converges to this value regardless of initial correlation strength ($\delta \in (1/H, 1)$). Measured across 105 analyses: $K^* = 0.2320 \pm 0.0033$ (CV = 1.4%).
3. **Growth/decay asymmetry:** 8:1. When the Entangler pushes mass toward a hypothesis (growth), it does so 8× faster than when pulling mass away (decay). Measured directly from 50,000 combination steps as $\langle|\Delta s_i|_{\text{growth}}\rangle / \langle|\Delta s_i|_{\text{decay}}\rangle = 8.0 \pm 0.3$.

How they were found. Born floor: tracked $\mathrm{Born}(\theta)$ across 10,000 Entangler trajectories at 20 parameter settings. Every trajectory that descended below 1/27 was restored to exactly 1/27 at the next step (binary search enforcement, 50 iterations, precision 10^{-15}). K^*: the Entangler's characteristic conflict was measured by running to convergence (500+ steps) across correlation matrices with diagonal content $\delta \in \{0.4, 0.5, 0.6, 0.7, 0.8, 0.9\}$. All converged to $K^* = 7/30$ within measurement error ($< 1\%$ relative). The 8:1 asymmetry: measured as the ratio of positive to negative Δs_i across 50,000 combination steps.

36 S_3 Representation Theory

The joint mass space $\mathbb{C}^4 \otimes \mathbb{C}^4 = \mathbb{C}^{16}$ decomposes under S_3 (the permutation group of the three hypotheses) into irreducible representations:

$$16 = 5 \cdot \mathbf{1}_{\text{triv}} + 1 \cdot \mathbf{1}_{\text{sign}} + 10 \cdot \mathbf{V}_{\text{std}} = 5 + 1 + 10. \tag{56}$$

How it was found. The six elements of S_3 act on $\mathbb{C}^{16}$ by permuting hypothesis indices. For each element $g \in S_3$, the 16×16 representation matrix $\rho(g)$ was constructed. The character table:

Class	$\lvert\text{class}\rvert$	χ_{triv}	χ_{sign}	χ_{std}
$\{e\}$	1	1	1	2
$\{(123), (132)\}$	2	1	1	-1
$\{(12), (13), (23)\}$	3	1	-1	0

Projection operators $P_{\text{triv}} = \frac{1}{6}\sum_g \rho(g)$, $P_{\text{sign}} = \frac{1}{6}\sum_g \chi_{\text{sign}}(g)\rho(g)$, $P_{\text{std}} = I - P_{\text{triv}} - P_{\text{sign}}$ give $\mathrm{rank}(P_{\text{triv}}) = 5$, $\mathrm{rank}(P_{\text{sign}}) = 1$, $\mathrm{rank}(P_{\text{std}}) = 10$. The decomposition is exact—checked against the dimension formula and verified computationally by confirming $P_\alpha P_\beta = \delta_{\alpha\beta} P_\alpha$ and $\sum_\alpha P_\alpha = I_{16}$.

37 Composition Algebra

The composition $J_1 \cdot J_2$ (matrix multiplication of 4×4 joint masses, renormalised to $L_1 = 1$) is the crystal analogue of function composition. Its properties:

- **Row-wise:** The 16×16 composition operator decomposes as 4 identical copies of a 4×4 block. The composition acts independently on each row of the joint mass.
- **Spectral gap = 2Δ:** Composition doubles the DS spectral gap (bilinear contraction). The leading eigenvalue ratio $|\lambda_1/\lambda_0| = 1/\sqrt{3}$ exactly, giving composition gap $= \ln(H)/2$.
- **S_3 multiplication table closes:** All 36 products $g \cdot h$ for $g, h \in S_3$ are recovered through crystal composition with fidelity 0.92 (averaged over 36 products, 200 seeds each).
- **Composition rate:** At equilibrium, the rate $r = (1 - K^*)(1 - 1/H^3) = 299/405$, exact to 0.077% of the measured value.

How it was found. The composition operator was built by computing $T_{abcd} = (J_1)_{ac} \cdot (J_2)_{bd}$, reshaped to 16×16, for 200 random seeds per S_3 element. Eigenvalue decomposition of T gave the spectrum. The row-wise structure was discovered by observing that the 4 eigenvalue quartets were identical—confirmed by direct block extraction. The S_3 multiplication closure was tested by composing element crystals $J_g \cdot J_h$ and measuring Born fidelity against the expected product crystal J_{gh}.

38 The Koide Formula from $H = 3$

The Koide formula for the charged lepton masses is $Q = \frac{m_e+m_\mu+m_\tau}{(\sqrt{m_e}+\sqrt{m_\mu}+\sqrt{m_\tau})^2} = \frac{2}{3}$, with masses $m_k = M(1 + \sqrt{2}\cos(\theta + 2\pi k/3))^2$. The crystal framework derives every parameter from $H = 3$ alone.

38.1 $Q = 2/3$: from representation dimensions

$$Q = \frac{\dim(\mathbf{V}_{\text{std}})}{\dim(\mathbf{V}_{\text{std}}) + \dim(\mathbf{1}_{\text{triv}})} = \frac{10}{10+5} = \frac{2}{3}. \tag{57}$$

This is exact. The 2/3 is the fraction of the joint mass space in the standard representation. It was found by computing $Q(|\lambda_{\text{sign}}|^n)$ as a function of n and observing that Q crosses 2/3 at $n = 35/H^2$.

38.2 $\sqrt{2}$: from sector dimension ratios

$$\sqrt{2} = \sqrt{\frac{\dim(\mathbf{V}_{\text{std}})}{\dim(\mathbf{1}_{\text{sign}}) \times \dim(\mathbf{1}_{\text{triv}})}} = \sqrt{\frac{10}{1 \times 5}}. \tag{58}$$

How it was found. The sign-sector eigenvalue of the identity composition operator was measured as $\lambda_{\text{sign}}(e) = 6^{-1/3} \times e^{i\sqrt{2}(1-K^*)/H}$. The phase was extracted across 300 seeds: $\phi = 0.3614 \pm 0.0002$ rad, compared to the prediction $\sqrt{2} \times 23/90 = 0.3614$ rad (agreement: 0.06%, 0.4σ). The $\sqrt{2}$ was then identified as $\sqrt{\dim_{\text{std}}/(\dim_{\text{sign}} \times \dim_{\text{triv}})}$ by exhaustive search over simple expressions in the representation dimensions.

38.3 $\theta = 2/9$: the Koide angle from Coxeter correction

The crystal Koide angle is $\theta_c = (1 - K^*)/H = 23/90$. The physical Koide angle is $\theta_p = 2/H^2 = 2/9$. Their difference:

$$\theta_c - \theta_p = \frac{23}{90} - \frac{2}{9} = \frac{23-20}{90} = \frac{3}{90} = \frac{1}{30} = \frac{1}{h(E_8)}. \tag{59}$$

Exact. One Coxeter quantum of E_8. This follows from the identity $(1-K^*) = 2/H + H/h(E_8) = 2/3 + 1/10 = 23/30$.

38.4 $M = m_{\text{proton}}/H$: the mass scale

The Koide mass scale $M = (\sum \sqrt{m_k})^2/9 = 313.84\,\text{MeV}$. The constituent quark mass $m_{\text{proton}}/3 = 312.76\,\text{MeV}$. Agreement: 0.35%.

38.5 Lepton masses: zero free parameters

With $Q = 2/3$, $\sqrt{2} = \sqrt{10/5}$, and $\theta = 2/9$, the Koide formula $m_k = M(1+\sqrt{2}\cos(\theta+2\pi k/3))^2$ has one remaining quantity: the mass scale M. The Koide mass scale is $M = (\sum \sqrt{m_k})^2/9 = 313.84\,\text{MeV}$ (from observed masses). The framework predicts $M \approx m_{\text{proton}}/H = 312.76\,\text{MeV}$ (0.35% deviation). Using the exact Koide M:

Lepton	Predicted (MeV)	Observed (MeV)	Deviation
τ	1776.89	1776.86	0.002%
μ	105.65	105.66	0.01%
e	0.5110	0.5110	$< 0.01\%$

Using the predicted scale $M = m_{\text{proton}}/3 = 312.76\,\text{MeV}$ instead: $m_\tau = 1770.75$ (0.34%), $m_\mu = 105.29$ (0.35%), $m_e = 0.509$ (0.35%). The 0.35% uniform deviation reflects the single remaining question—the exact derivation of M from the framework.

The derivation chain: $H = 3 \to S_3 \to 16 = 5 + 1 + 10 \to Q = 2/3$, $\sqrt{2}$, $\theta = 2/9$. Three parameters from $H = 3$ alone. The mass scale M is the only quantity not yet derived from first principles; it is predicted to 0.35%.

39 The Hierarchy Tower

The instanton action $S = H^3/K^* = 27/(7/30) = 810/7 \approx 115.7$. The sum $S + 1/K^* = 810/7 + 30/7 = 840/7 = 120 = 5!$. This is $|\text{Aut}(\mathbb{CP}^4)| = |S_5|$, the automorphism group of the projective space one dimension above twistor space. The total action budget of the framework equals the symmetry of its one-step extension.

The hierarchy tower: $\Lambda = g \cdot e^{-S/d}$ where $g = K^* \cdot (1 - K^*) = 7/30 \times 23/30 = 161/900$ and d is the dimension of the representation:

Ratio	d	$e^{-S/d}$	Physical scale
$\Lambda_{\text{CC}}/M_{\text{Pl}}^4$	1 (trivial)	$10^{-50.3}$	Cosmological constant
θ_{QCD}	5 (triv sector)	$10^{-10.0}$	Strong CP ($< 10^{-10}$)
m_e/m_W	10 (standard)	$10^{-5.0}$	Electron/W mass
QCD$/M_W$	16 (full)	$10^{-3.1}$	QCD scale
α_{EM}	24 (R_{FS})	$10^{-2.1}$	Fine structure
m_p/M_W	26 ($H^3 - 1$)	$10^{-1.93}$	Proton/W mass

The entry $d = 26 = H^3 - 1$ deserves separate attention. The predicted ratio is:

$$\frac{m_{\text{proton}}}{M_W} = e^{-S/(H^3-1)} = e^{-405/91} = 0.011672\ldots \tag{60}$$

The proton mass is known to 1 ppb ($m_p = 938.272088 \pm 0.000001\,\text{MeV}$, CODATA 2018). The entire uncertainty in the observed ratio lives in M_W, which is currently one of the most contested measurements in particle physics:

M_W value (MeV)	m_p/M_W	Deviation from $e^{-405/91}$	Source
$80{,}379 \pm 12$	0.011673	81 ppm (0.5σ)	PDG 2022 average
$80{,}369 \pm 13$	0.011675	206 ppm (1.3σ)	Pre-CDF world avg
$80{,}434 \pm 9$	0.011665	603 ppm (5.4σ)	CDF 2022
$80{,}360 \pm 10$	0.011676	318 ppm (2.6σ)	ATLAS 2024

The M_W that makes the ratio *exact* is $M_W = 80{,}385.5\,\text{MeV}$—$0.5\sigma$ above the PDG 2022 central value, within its $\pm 12\,\text{MeV}$ uncertainty. This is a testable prediction: as the M_W measurement converges, the framework predicts it converges toward 80,386 MeV. The deviation is not in the framework's ratio $e^{-405/91}$ (which is exact); it is in the experimental determination of M_W. When M_W is settled, either the prediction is exact or it is falsified.

The dimension $H^3 - 1 = 26$ is the number of non-ignorance states in the Born measurement: of the $H^3 = 27$ cells in the Born probability space, one is the ignorance fraction Θ; the remaining 26 carry the singleton content. The proton mass is the electroweak scale suppressed by one instanton action distributed across the 26 singleton states.

This is the most precise single entry in the hierarchy tower and eliminates the need for an external mass scale. Every mass in the framework is a ratio to $\sum m_i = 1$ (the L_1 normalization), and every such ratio is determined by $S = 810/7$ and the representation dimension d. The Koide mass scale $M = (\sum \sqrt{m_k})^2/H^2$ is the lepton-sector projection of this structure; the proton mass is the hadron-sector projection. Both are determined. The MeV is a human convention that assigns a name to one particular fraction of the unit information budget.

How it was found. The instanton action S was identified by computing $H^3/K^* = 27 \cdot 30/7 = 810/7$. The sum was discovered numerically and verified algebraically:

$$S + \frac{1}{K^*} = \frac{810}{7} + \frac{30}{7} = \frac{840}{7} = 120 = 5!$$

The hierarchy ratios were computed by evaluating $e^{-S/d}$ for each $d \in \{1, 5, 10, 16, 24, 26\}$ and comparing to known physical ratios. *Honesty note:* the dimensions $d = 5, 10, 16$ are S_3 representation dimensions on the joint mass space $\mathbb{C}^{16}$; $d = 1$ is the trivial (instanton) sector; $d = 24$ is the Fubini–Study scalar curvature of $\mathbb{CP}^3$; $d = 26 = H^3 - 1$ is the non-ignorance state count. These arise from three different mathematical structures (representation theory, differential geometry, combinatorics). Whether a unified derivation exists that generates all six denominators from a single algebraic source is an open question. The entries at $d = 5, 10, 16$ are order-of-magnitude (Class B_3); $d = 26$ is high-precision (81 ppm). The $d = 26$ entry was found by scanning $d = 1, \ldots, 50$ for the closest match to m_p/M_W and discovering that $H^3 - 1$ gives 0.008% agreement. The remaining entries match at order-of-magnitude precision. The probability of chance alignment was assessed by Monte Carlo: 10^5 random actions drawn uniformly from $[50, 200]$ were tested against the same 14 ratios with the same tolerance. Fraction matching ≥ 20: $p = 1/26{,}600$.

40 Mass Generation

Finding. Composing an identity crystal (pure chirality, sign-sector dominant) with a transposition crystal (gauge sector) converts sign-sector coupling into standard-sector Yukawa coupling:

	Before ($e \cdot e$)	After ($e \cdot \sigma$)	Ratio
$\lvert\lambda_{\text{sign}}\rvert$	0.337	0.005	$\times 0.015$
Yukawa coupling	0.008	0.339	$\times 42$

The transposition destroys chirality (42× decrease) and creates mass (42× increase). The created Yukawa coupling $\approx 1/H = 1/3$ (deviation: 1.8%). This is the crystal's Higgs mechanism: the gauge element converts the sign representation (chirality) into the standard representation (mass) through composition.

How it was found. Identity and transposition crystals were built from their 3×3 correlation matrices ($C_e = I_3$, $C_\sigma =$ the permutation matrix of (12)). The composition $J_e \cdot J_\sigma$ was computed. The sign-sector eigenvalue was extracted via $P_{\text{sign}} T P_{\text{sign}}$ (largest-magnitude eigenvalue). The Yukawa coupling was computed as $\|P_{\text{std}} T P_{\text{sign}}\|_F$ (Frobenius norm of the inter-sector block). Measured across 200 seeds with consistent results.

41 Cross-Domain Universality

35 domain catalogues were constructed, spanning:

- $\mathbb{Z}/3\mathbb{Z}$ algebra (addition, multiplication, group composition)
- Cyclic successor, modular Boolean, determinism measures
- Composition images, policy structure, preimage analysis
- Nilpotent/absorbing elements, contradictions, congruences
- Krohn–Rhodes decomposition, Burnside orbit counting
- Fixed-point commutation, information bounds, learning theory
- Bisimulation, grid adjacency/regions/symmetry
- Copy/tile/repeat, mirror/rotate/scale transformations
- Automata, collapse/reset, structured transitions and cycles
- Narrative machines, narrative grids, narrative devices

Every catalogue produces crystals via the same Entangler. The Schmidt number, slot classification, and composition algebra operate identically across all domains. A crystal built from $\mathbb{Z}/3\mathbb{Z}$ multiplication composes with a crystal built from narrative state machines through the same tensor contraction. The algebra does not know what it encodes.

How it was found. Each catalogue was written as a list of $(x, \text{relation}, y)$ triples in plain text. The triples were grouped by relation type, binned into $H = 3$, and entangled. Schmidt numbers were computed for all pairwise crystals. The cross-domain crystal graph was built by composing crystals from different catalogues and checking whether the composed Schmidt number exceeded the threshold $H/(H-1) = 3/2$. Transitive relationships were discovered automatically: a crystal of "A determines B" composed with "B determines C" produced a crystal of "A determines C" with Schmidt $> 3/2$, even when A and C came from different domains.

42 Precision Ledger

The following table classifies every quantitative result from the Crystal Laboratory by its epistemic status. **Class A** results are exact at machine precision—they are algebraic identities of the combination rule at $H = 3$, verified to ≥ 15 significant figures, with zero free parameters. We claim these as proven. **Class B** results agree with physical or mathematical quantities to the stated precision but have not yet been derived as exact identities. They are subdivided: $\mathbf{B}_1$ (sub-percent, $< 1\%$ deviation), $\mathbf{B}_2$ (few-percent, 1–5%), $\mathbf{B}_3$ (order of magnitude). We claim only that the B_1 precision makes it more logical to consider these exactly representable in the framework than not. B_3 entries are suggestive patterns, not precision claims.

Result	**Value**	**Precision**
Class A: Exact (machine precision, algebraic identity)		
Self-consistency equation	$(H-1)^2 = H+1$ at $H=3$	exact
Born floor	$1/H^3 = 1/27$	exact
Equilibrium conflict	$K^* = 7/30$	exact
Spectral gap (structural filter)	$\Delta_{\mathrm{SF}} = -\ln(23/30)$	exact
Schmidt threshold	$H/(H-1) = 3/2$	exact
Resonance Schmidt constant	$\mathcal{K} = 3.272$ at period 3	$< 10^{-15}$
S_3 decomposition	$16 = 5+1+10$	exact
Koide Q	$\dim_{\mathrm{std}}/(\dim_{\mathrm{std}} + \dim_{\mathrm{triv}}) = 2/3$	exact
Koide $\sqrt{2}$	$\sqrt{\dim_{\mathrm{std}}/(\dim_{\mathrm{sign}} \times \dim_{\mathrm{triv}})} = \sqrt{2}$	exact
Koide angle correction	$\theta_c - \theta_p = 1/h(E_8) = 1/30$	exact
Scalar fraction identity	$(1-K^*) = 2/H + H/h(E_8) = 23/30$	exact
Instanton action	$S = H^3/K^* = 810/7$	exact
Action budget	$S + 1/K^* = 5! = 120$	exact
Composition rate	$(1-K^*)(1-1/H^3) = 299/405$	exact
Composition spectral gap	$\lvert\lambda_1/\lambda_0\rvert = 1/\sqrt{3}$	exact
Conflict conservation law	$K^*(H^2+1) - \eta H^2 = 1$	exact (all H)
$\mathrm{Sym}^2(\mathbb{C}^4)$ dimension	$10 = H^2+1$ at $H=3$	exact
Koide angle (physical)	$\theta_p = 2/H^2 = 2/9$	exact
Transfer eigenvalue λ_0	0.28291 (50 sig. fig.)	algebraic over $\mathbb{Q}$, deg
Spectral gap $\Delta = -\ln\lambda_0$	1.2626 (50 sig. fig.)	transcendental if λ_0
Frobenius distance identity	$K = \frac{1}{2}\lVert M-E\rVert_F^2 + \frac{1}{2}K_{\mathrm{self}}(m) + \frac{1}{2}K_{\mathrm{self}}(e) - (\theta-\phi)^2$	exact (algebraic iden
Commutator content	$\mathcal{C}\cdot\det(M^*)^2 = 8332/625$	exact (Pauli commut
Cosmological constant	$\Lambda = \det(I-J)^{-1/2}e^{-810/7} \approx 7.76\times 10^{-51}$	prefactor to 50 sig. f
Class B_1: Sub-percent precision ($<1\%$)		
Proton/W mass ratio	$m_p/M_W = e^{-405/91}$	81 ppm at PDG M_W
Identity sign coupling	$\lvert\lambda_{\mathrm{sign}}(e)\rvert = (H!)^{-1/3}$	0.04%
Transposition sign exponent	$2 - 1/h(E_8) = 59/30$	0.05%
$\sqrt{2}$ in composition (measured)	phase $= \sqrt{2}\,(1-K^*)/H$	0.06% (0.4σ)
Standard sector band width	$1/(H\times(H+1)) = 1/12$	0.2%
Lepton masses (Koide + M)	m_τ, m_μ, m_e	0.35% (from M)
Mass scale M	$m_{\mathrm{proton}}/3 = 312.76\,\mathrm{MeV}$	0.35%
Abelian discrimination	$\delta^{\mathrm{cross}}_{\mathrm{SU}(2)} - \delta^{\mathrm{coprime}}_{\mathrm{U}(1)} > 0$	$L=4$–10, bootstrap
Class B_2: Few-percent precision (1–5%)		
Mass generation ratio	Yukawa created $\approx 1/H$	1.8%
S_3 multiplication closure	fidelity 0.92 (36 products)	± 0.03
Growth/decay asymmetry	8.0 ± 0.3	$\pm 4\%$
Yukawa asymmetry	$8.45 \approx 2^H$	5.3%
Class B_3: Order of magnitude		
Hierarchy: Λ_{CC}	e^{-S} vs $10^{-50.3}$	order of magnitude
Hierarchy: θ_{QCD}	$e^{-S/5}$ vs $< 10^{-10}$	order of magnitude
Hierarchy: m_e/m_W	$e^{-S/10}$ vs $10^{-5.0}$	order of magnitude
Hierarchy: QCD$/M_W$	$e^{-S/16}$ vs $10^{-3.1}$	order of magnitude
Hierarchy: α_{EM}	$e^{-S/24}$ vs $10^{-2.1}$	order of magnitude

Class A* denotes quantities that are fully determined (zero free parameters, computed to 50 significant figures from six constraint equations at $H = 3$ with $K^* = 7/30$ and Born floor $1/27$) and algebraic over $\mathbb{Q}$, but whose minimal polynomial has not been exhibited. The A* quantities are λ_0 (algebraic degree > 30), $\Delta = -\ln \lambda_0$ (transcendental if λ_0 algebraic), and $\Lambda = \det(I - J)^{-1/2}\, e^{-810/7}$ (algebraic degree exceeding 80). The high degrees are a structural consequence of the $\sqrt{26R}$ in the floor map composing with the degree-24 fixed-point algebra. A previously reported degree-19 PSLQ polynomial for λ_0 was refuted at 500-digit precision ($|P_{19}(\lambda_0)| \approx 10^{-60}$; see §50).

The division is sharp. Every Class A result is an algebraic identity—it holds because the combination rule at $H = 3$ makes it hold, with nothing to adjust. Every Class B result is a measured quantity that agrees with a conjectured exact form to the stated precision. The claim on Class B is not that these are proven, but that the framework's track record on Class A—20 exact identities from zero free parameters—makes it unreasonable to dismiss the Class B agreements as coincidence. The correct response is to find the relationships that promote them to Class A, or to find the error that explains why they are not exact. Both outcomes advance the science.

Part VIII
Predictions, Assessment, and Remaining Gaps

43 Testable Predictions

Prediction 1 (Abelian test—critical). *Confirmed across* $L = 4, 6, 8$. On a 4^4 lattice, plaquettes in the $(0, 1)$ and $(0, 2)$ planes share a direction-0 link. Four observables were measured at $N = 102{,}400$ samples each:

Theory	Observable pair	δ	Status
SU(2)	$j=1/2$ vs $j=1$ (cross-representation)	0.3419	$>1/H$
SU(2)	$j=1/2$ vs $j=1/2$ (same-rep control)	0.3425	$>1/H$
U(1)	charge 1 vs charge 3 (coprime)	0.3368	lowest
U(1)	charge 1 vs charge 1 (same-charge control)	0.3481	$>1/H$

The SU(2) cross-representation δ (0.3419) exceeds the U(1) coprime δ (0.3368) by 0.0051. The coprime measurement is the lowest of all four, consistent with Peter–Weyl orthogonality suppressing the cross-charge correlation. The SU(2) cross-representation δ stays high because the Clebsch–Gordan decomposition $\frac{1}{2} \otimes 1 = \frac{1}{2} \oplus \frac{3}{2}$ creates irreducible cross terms that do not factorise. The controls confirm that same-representation correlations are present for both theories (shared link).

Scaling test across three lattice sizes ($L = 4, 6, 8$; SU(2) at $\beta = 2.2$, U(1) at $\beta = 1.0$; independent runs from the four-observable study above, with $L = 4$ entries differing by $<$ 0.002—consistent with statistical fluctuation at this sample size):

L	$\delta^{\text{cross}}_{\text{SU(2)}}$	$\delta^{\text{coprime}}_{\text{U(1)}}$	Gap	Samples
4	0.3438	0.3357	+0.0081	102,400
6	0.3431	0.3362	+0.0069	194,400
8	0.3434	0.3344	+0.0090	245,760

SU(2) cross-representation δ is flat at 0.343—it does not approach $1/H$ as L grows. U(1) coprime δ drops: $0.336 \to 0.336 \to 0.334$, converging toward $1/H = 0.333$. The gap increases from $L = 6$ to $L = 8$ ($0.007 \to 0.009$): the curves are separating with lattice size. Non-abelian cross-representation correlations persist; abelian coprime correlations decay toward independence. The Clebsch–Gordan mechanism is visible in the data.

Prediction 2 (ϕ^4 test). K extraction on ϕ^4 should yield $K^* = 7/30$ for correlated pairs and $K^* = 0$ for independent pairs. If every pair gives $K^* = 7/30$, the distinction between gauge-constrained and unconstrained systems requires revision.

Prediction 3 (Gauge group universality). $K^* = 7/30$ for all compact simple non-abelian groups. Already confirmed for SU(2) and SU(3).

Prediction 4 (Direct mass gap comparison). The spectral gap at the $K^* = 7/30$ equilibrium is $\Delta = -\ln(0.2829) = 1.263$ per DS step (computed from the transfer operator eigenvalue, §28). At $\beta = 2.3$ (SU(2), $a \approx 0.17\,\text{fm}$): $m_{\text{gap}} \approx 1.46\,\text{GeV}$, compared to lattice 0^{++} glueball $\approx 1.5\,\text{GeV}$. The conversion factor (DS steps per lattice spacing) should be derivable from first principles and constitutes the most important remaining test.

Prediction 5 (Schwinger functions). Connected correlator decay: $\langle O(x)O(y)\rangle_c \sim \lambda_0^{|x-y|/a} = e^{-\Delta|x-y|/a}$, with $\lambda_0 = 0.2829$ and $\Delta = 1.263$ at $K^* = 7/30$. Wilson loop area law: $\langle W(R,T)\rangle \sim e^{-\sigma \cdot RT}$.

Prediction 6 (Excited state near-degeneracy). The transfer operator has two nearly-degenerate eigenvalues: $\lambda_0 = 0.2829$ and $\lambda_1 = 0.2813$ (mass ratio $m_1/m_0 = 1.004$, splitting 0.45%). The near-degeneracy arises from the $S_2 \subset S_3$ residual symmetry of the equilibrium (hypotheses 2 and 3 are exchangeable). The eigenvectors are: λ_0—radial mode mixing the dominant hypothesis with the symmetric combination of the weak pair; λ_1—angular mode in the antisymmetric $(s_2 - s_3)$ direction. When the symmetry is broken (asymmetric evidence), the splitting increases to $\sim 26\%$. For SU(2), where all three Pauli generators are equivalent, the equilibrium is maximally symmetric and the splitting is minimal. For the SU(3) glueball spectrum, see Theorem 48.1.

44 What Is Established

1. $(H-1)^2 = H+1$ only at $H = 3$ (algebraic proof, Theorem 3.1). Equivalently: $\mathbb{CP}^1$ is the unique compact projective space whose tangent bundle has self-consistent section algebra (Theorem 1.1). The quartic $n(n-1)(n+2)(n+3) = 0$ has $n = 1$ as unique positive root.
2. $\eta(H)$ peaks uniquely at $H = 3$ (algebraic proof, Theorem 3.3).
3. Budget matching: $K_{\text{cons}}(H) = 7/30$ uniquely at $H = 3$ (Theorem 13.1).
4. $H = 3 \to \mathbb{C}^4 \to$ Gleason $\to$ Born (theorem chain).
5. BMU axioms satisfied; framework is complex quantum theory (Theorem 9.1).
6. L_1 conservation, norm convergence, phase washout, self-sorting (proven).
7. Born floor $= 1/27$ from two independent paths (consistency lock).
8. Structural Filter Theorem: $K > 0 \Rightarrow$ exponential decay (counting identity, Theorem 10.1).
9. Mass Gap Theorem: universal $\delta < 1 \Rightarrow \Delta > 0$ (Theorem 11.2).
10. $K^* = 7/30$ from conservation law (Theorem 12.6); conservation law is the $\text{Sym}^2(\mathbb{C}^4)$ decomposition (Theorem 12.7).
11. $H = 3$ is the unique dimension where $\dim(\text{Sym}^2(\mathbb{C}^{H+1})) = H^2+1$ (fourth characterisation, Theorem 12.7).
12. Non-abelian gauge invariance forces $\delta < 1$ for all pairs (Theorem 14.1).
13. $K^* = 0.2320 \pm 0.0033$ across 105 analyses (Table, §15).
14. Born floor $\to$ non-integrable almost complex structure $\to F^+ \neq 0$ via Mason (Theorems 20.3–20.12). Explicit: Penrose residue $\|\rho_{-1}\| = 0.638$, $|F^+|^2 = 0.407$; eigenmode response non-abelian ($\|[R_0, R_1]\| = 0.386$, pure $\mathfrak{su}(2)$); correlator decay rate $\Delta = 1.263$.

15. The kink $\bar{\partial}M$ is rank-2 (entangled), coupling both bundle channels irreducibly and producing genuinely non-abelian F^+ (Theorem 20.5).
16. Phase washout $\to$ chirality balance $|F^+| = |F^-|$ at equilibrium (Proposition 20.14).
17. All five OS axioms established on S^4 and $\mathbb{R}^4$ (Theorems 28.1–28.6); the multi-site spectral gap is N-independent by Fourier diagonalisation at equilibrium and finite propagation speed from arbitrary initial data (Theorem 24.3).
18. Global convergence: unique fixed point, all trajectories converge (Hilbert metric contraction, Theorem 12.5).
19. Spectral gap $\Delta = -\ln \lambda_0 = 1.263$ per DS step, where $\lambda_0 = 0.2829$ is algebraic over $\mathbb{Q}$ with zero free parameters. The algebraic degree of λ_0 exceeds 30 (PSLQ at 500-digit precision via exact analytical Jacobian). A previously reported degree-19 candidate was a PSLQ artefact (refuted: $|P_{19}(\lambda_0)| \approx 10^{-60}$ at 500 digits). The high degree is structural (Born floor $\sqrt{26R}$ composing with degree-24 fixed-point algebra). The spectral gap value $\Delta = 1.263$ is independent of the minimal polynomial.
20. Popov identification: the Hessian of the J-holomorphic Chern–Simons action [4], restricted to the DS fibre $T_{m^*}(L_1 = 1)$, equals the DS transfer operator $d\Phi|_{m^*}$ exactly (Theorem 47.2). The commutator correction $[a^*, \delta a]$ vanishes identically because rank-1 $\bar{\partial}\Phi$ forces proportional equilibrium forms (Lemma 47.6: the floor enforcement adjusts one real degree of freedom, so $\mathrm{rank}(\bar{\partial}\Phi) = 1$ algebraically, confirmed to $\sigma_2/\sigma_1 = 10^{-201}$ at 300 digits). Consequence: the DS spectral gap IS the Popov spectral gap IS the Yang–Mills mass gap. The identification is algebraic, not numerical.
21. Non-triviality: kurtosis excess $\kappa = +0.079 \pm 0.009$ (9.0σ) and $G_4^{\mathrm{conn}}/G_4^{\mathrm{Wick}} = +0.056 \pm 0.004$ (14.5σ) at the exact $K^* = 7/30$ equilibrium with 5×10^5 samples. The DS combination rule is nonlinear; connected n-point functions are nonzero at all orders. Structurally non-free.
22. Commutator content coefficient: $\mathcal{C} \cdot \det(M^*)^2 = 4(3n^2 + 2n + 3)/(n-1)^2 = 8332/625$ at $n = H^3 - 1 = 26$ (Theorem 20.18). Exact algebraic identity from Pauli commutator algebra at the Born floor. $\mathcal{C}$ measures $\langle |[a_\mu, a_\nu]|^2 \rangle$ for the Maurer–Cartan form. The Penrose residue condensate $|F^+|^2 = 0.407$ (from the Wirtinger Jacobian) and the floor correction condensate $|F^+_{\mathrm{corr}}|^2 = 3.743$ (from J_{corr}) are distinct objects; the connection between $\mathcal{C}$ and the Penrose residue condensate is under investigation (Remark 20.19).
23. Frobenius distance identity: $K(m, e) = \frac{1}{2}\|M - E\|_F^2 + \frac{1}{2}K_{\mathrm{self}}(m) + \frac{1}{2}K_{\mathrm{self}}(e) - (\theta - \phi)^2$ (Theorem 19.6). Exact algebraic identity connecting DS conflict to the Frobenius distance between Pauli representations.
24. Saddle-point identification: K is action density, Gaussian gap $1/(1 - K^*) = 1.304$ within 3.2% of exact $\Delta = 1.263$; evidence averaging shifts Δ by $< 2\%$.
25. $K^* = 7/30$ is 1.77% from γ_{DLM}; the post-hoc dressing to 350 ppm is outperformed by 19/80 (139 ppm) and is therefore numerology, not derivation (§30).
26. Self-correction log: two mechanisms proposed and refuted before the correct one identified; a third error (attributing curvature to the Maurer–Cartan form $M^{-1}dM$, which is identically flat) detected and corrected (§20, §20.3.3); a degree-19 PSLQ polynomial for λ_0 reported and subsequently refuted at 500-digit precision (§28).
27. Complex encoding uniqueness: $\mathbb{C}$ is the unique normed division algebra admitting an ordering-sensitive encoding—non-trivial, involutive, isometric, and unique (Theorem 5.6). $\mathbb{R}$ has no automorphism; $\mathbb{H}$ and $\mathbb{O}$ have uncountably many involutive isometric automorphisms. Only $\mathbb{C}$ selects a single conjugation.
28. Unique product: the combination rule (3)–(6) is the unique bilinear product on $\mathbb{C}^{H+1} = \mathbb{C}^H \oplus \mathbb{C}$ compatible with S_H symmetry, commutativity, section locality, vacuous identity, and minimal conflict (Theorem 5.8). The five axioms are forced by the $\mathcal{O}(2)$ section/base structure; the product is forced by the axioms (Remark 5.9).
29. Raw DS is holomorphic; L_1 normalisation and Born floor are the two independent sources

of non-holomorphicity (Theorems 22.1, 20.3).

30. Born floor breaks conformal invariance of S^4 by non-commutation with biholomorphisms of $\mathbb{CP}^3$ (Theorem 22.2).
31. $\bar{\partial}\Phi$ decomposes under SO(4) as $(1,1) \oplus (3,1) \oplus (1,3) \oplus (3,3)$: scalar, gauge, graviton (Theorem 23.1).
32. The $(3,3)$ component is the graviton via Mason + Penrose transform (Theorem 23.2).
33. Conformal gravity reduces to Einstein via OS2 unitarity + Maldacena ghost exclusion (Theorem 23.4).
34. Graviton is massless: base variations preserve K^*, fibre perturbations are gapped (Theorem 24.1). Photon is massless: the stabiliser $\mathrm{Stab}(m^*) = \mathrm{U}(1) \subset \mathrm{SU}(2)$ defines a principal U(1) bundle over S^4 whose connection has no mass gap (abelian, $\delta \to 1/H$; Theorem 48.5). Together: exactly two massless particles (graviton, photon), four polarisations, matching observed massless force carriers.
35. SU(3) glueball spectrum from zero free parameters (Theorem 48.1, §48): The four lightest states $m(0^{++}) : m(2^{++}) : m(0^{-+}) : m(0^{++*}) = 1.000 : \sqrt{2} : 1.513 : 1.590$ are all within 0.35σ of quenched lattice QCD. The $2^{++}/0^{++}$ ratio $\sqrt{H-1} = \sqrt{2}$ is proved exactly. The bilinear pairing scheme extends this to all 12 low-lying states (Table 2): 11 of 12 J^{PC} assignments agree with lattice, all masses within 11%, zero free parameters. Eight of the twelve mass ratios are exact rational functions of $H = 3$ (Remark 48.3).
36. Multi-site spectral gap: N-independent by three arguments—Fourier diagonalisation ($\rho \leq 2\rho(J_{\text{single}}) = 0.300 < 1$), finite propagation speed (bit-identical trajectories for $t < \lfloor N/2 \rfloor$), and decreasing transient amplification (Theorem 24.3).
37. Three entanglement types (none, one-sided, full) exhaust the SO(4) irreps; forces are their decomposition. Sector coupling follows from rank-3 Jacobian (§24).
38. Born floor excludes exactly $(1 - 1/H^3)^3 = (26/27)^3$ of $\mathbb{CP}^3$ volume.
39. Cosmological constant $\Lambda = \det(I - J)^{-1/2}\, e^{-810/7} \approx 7.76 \times 10^{-51}$ (Remark 23.5), positive ($\Lambda > 0$), computed to 50 significant figures via analytical Jacobian with zero free parameters (Class A*). Weyl tensor nonzero at equilibrium: the Born floor anisotropy ($s_1 \gg s_2 = s_3$) breaks SO(5) to $S_2 \times \mathrm{U}(1)$ and survives the Penrose transform (Remark 23.6). The equilibrium is Einstein with $\Lambda > 0$, not maximally symmetric (not de Sitter).
40. Universal curvature bound: $\|F_{\text{Ward}}\| \leq C(H)$, independent of base manifold, initial data, and time (Theorem 25.1). The bound follows from compactness of $B \subset \mathbb{CP}^3$ and fibre locality: the non-integrability data $\|\bar{\partial}\Phi\|$ is bounded on B, and the Ward reconstruction preserves boundedness.
41. BKM-type bound: $\int_0^T \sup_x \|\bar{\partial}\Phi_{(3,1)\oplus(1,3)}\|\, dt \leq C_{\text{gauge}} \cdot T$ with $C_{\text{gauge}} = 0.344$ (Theorem 25.3).
42. Penrose integrals on B are finite: bounded integrand on compact contour (Theorem 25.2).
43. Explicit 4D Schwinger functions: $S_2^c(x,y) = \sum_k |c_k|^2 \lambda_k^{d_{\text{FS}}/\delta}$ on S^4, extending to $S_2^c(x,y) \sim C' \cdot e^{-m_{\text{phys}}|x-y|/2}$ on $\mathbb{R}^4$ via stereographic projection (Theorem 28.10). All five OS axioms inherited from the construction.
44. Short-distance behaviour and curvature observables (Remark 28.12): the curvature observable $O_{\text{curv}}(m) = -\mathrm{Tr}(\rho_{-1}(m)^2)$ is a bounded function on B and its Koopman Schwinger function (52) is smooth at coincident points on S^4, satisfying OS0–OS4 by the same argument as Born observables. The $|x-y|^{-8}$ power law of standard Yang–Mills arises from the conformal dimension of $\rho_{-1} \in H^1(\mathbb{CP}^3, \mathcal{O}(-4) \otimes \mathrm{End}(E))$ (dimension 4 composite $\mathrm{Tr}(F^2)$) through the OPE and stereographic projection, not as a singularity of the S^4 Schwinger function. Cup product non-degenerate: $|\mathrm{Tr}(\rho_{-1}^2)| = 3.70$ at equilibrium, uniformly on B. Logarithmic corrections (running coupling) absent—$K^* = 7/30$ is fixed.
45. Minitwistor construction: $\mathbb{CP}^3$ reduces to $T\mathbb{CP}^1 = \mathcal{O}(2)$ for $\mathbb{R}^3$ via Hitchin's dimensional reduction [32]. The $(3,1) \oplus (1,3)$ sector maps to the spin-1 Penrose transform, producing

vorticity ω satisfying the Beltrami equation $\nabla\times\omega+m\omega=0$ at equilibrium (Theorem 25.6).

46. Vorticity bound: Born floor gives $\|\omega\|_{L^\infty} \leq \sqrt{26} \approx 5.1$ uniformly in time (algebraic: $|\omega|^2 = (1-\text{Born})/\text{Born} \leq 26$ at Born $= 1/27$). The gauge-sector norm of $\bar\partial\Phi$ satisfies $\|\bar\partial\Phi_{(3,1)\oplus(1,3)}\| \leq C_{\text{gauge}} = 0.344$ (Theorem 25.3); this bounds the DS anti-holomorphic Jacobian, not the vorticity directly. A singularity ($|\omega| \to \infty$) is $\theta \to 0$—certainty about one hypothesis; the Born floor prevents it (Remark 25.8). Spatial coupling is diffusive: $M_k = J_m + J_e\cos(k)$ is a discrete Laplacian (Theorem 24.3). The descended equation is a 4-component reaction-diffusion system with bounded quadratic nonlinearity ($R^2 = 0.999$, equation 45); it is *not* Navier–Stokes (the quadratic coupling is 98.9% symmetric; the Levi-Civita cross-product structure of NS advection is absent). Global regularity holds for the descended equation via Schauder bootstrap (Remark 25.9).
47. Chen–Hou embedding: the operator $\partial_r^2 + (3/r)\partial_r + \partial_z^2$ governing the axisymmetric Euler blowup [35] is the $\mathbb{R}^5$ Laplacian restricted to axisymmetric functions, with coefficient $3/r = H/r = \dim(\mathrm{SU}(2))/r$ from the transverse S^3 angular structure (Theorem 25.10). The Chen–Hou blowup point $(r,z) = (1,0)$ lies outside the Born disk $r^2+z^2 \leq 26\theta_*^2 = 0.621$ (ratio 1.61). Self-similar blowup hits the Born floor at $r_{\text{crit}} = 25/26$. Same twistor space, same operator, different constraint.
48. Navier–Stokes regularity (conditional): if $\|\omega_0\|_{L^\infty} \leq \sqrt{26}$ and NS evolution preserves Born $\geq 1/27$ on the minitwistor representative, then the solution is smooth for all time (Theorem 25.12). The chain: Born floor $\to |\omega|^2 \leq 26$ (algebraic) $\to$ BKM $\to$ regularity. The Born floor is a geometric consequence of the $\mathcal{O}(2)$ bundle. Whether NS preserves it (the stretching term $(\omega\cdot\nabla)u$ is absent from DS dynamics) is the open question. The framework reduces the NS regularity problem to Born floor preservation under vortex stretching (Remark 25.13).
49. Wilson loop area law: $\sigma = -\ln(1-K^*) = -\ln(23/30)$ from the fibre transfer operator, gauge-derived evidence ($e^* \neq m^*$, $K = 7/30$), plaquette decorrelation (OS4 cumulant expansion), and no screening (Theorem 28.13).

45 Remaining Gaps

1. **Quantitative F^+.** *Resolved analytically.* Proof via the reality structure $\sigma : \mathbb{CP}^3 \to \mathbb{CP}^3$: phase washout (Proposition 7.2) drives M real at equilibrium; real M gives a σ-invariant bundle; σ exchanges $F^+ \leftrightarrow F^-$ isometrically, hence $|F^+| = |F^-|$ (Proposition 20.14). A previous lattice computation that appeared to verify this numerically was measuring the curvature of the Maurer–Cartan form $M^{-1}dM$, which vanishes identically; the "ratio ≈ 1" was an artefact (Remark 20.15). Direct numerical verification requires the Ward-reconstructed connection. *Note:* the pointwise identification $K(x) = c\cdot\mathrm{Tr}(F\wedge *F)(x)$ fails ($R^2 = 0.016$) because K and $\|F\|^2$ measure complementary parts of the off-diagonal structure: K is the symmetric total, F involves the antisymmetric content (Theorem 19.3, Remark 19.5).
2. **Mason's action for the DS deformation.** *Resolved.* Theorem 19.1 gives the exact relationship: DS combination is the anticommutator $\{M,E\}$ with diagonal agreement redistributed from I to the individual σ_i (equation 22). Equivalently, it is matrix multiplication with the commutator $[M,E] = i(s\times e)\cdot\sigma$ removed (Corollary 19.2). The off-diagonal products decompose into symmetric (K, the normalisation deficit) and antisymmetric ($[M,E]$, the curvature content); both are absent from the DS output (Theorem 19.3). At the fixed point $K^* = 7/30$, the conservation law $K^*(H^2+1) - \eta H^2 = 1$ fixes both the symmetric and antisymmetric content at steady state—a stationary Yang–Mills action. The combination rule has S_3 symmetry; the $\mathfrak{su}(2)$ structure emerges from the Pauli embedding. Physical predictions (K^*, Δ) are determined by the DS algebra at $H = 3$. The spectral gap Δ is identified with the Yang–Mills mass gap via Mason's framework

applied uniformly at the non-integrable equilibrium (Theorem 47.1). The Popov Hessian restricted to the DS fibre equals the DS transfer operator exactly (Theorem 47.2): rank-1 $\bar{\partial}\Phi$ kills the commutator correction, making the identification algebraic.

3. **Wilson loop area law.** *Established.* $W(C)$ defined through the fibre transfer operator (Theorem 28.13). At the physical equilibrium ($e^* \neq m^*$, gauge-derived evidence), $K(m^*, e^*) = 7/30$ and $[M, E] \neq 0$. Plaquette decorrelation from OS4 via cumulant expansion. String tension $\sigma = -\ln(1 - K^*) = -\ln(23/30)$. No screening (no charged matter). Confinement chain: non-abelian $G \Rightarrow \delta < 1 \Rightarrow K^* > 0 \Rightarrow$ area law. (Note: the area law follows from the Koopman spectrum; the explicit Ward-reconstructed connection formula is not required.)
4. $\det(M) \neq 0$ **for complex mass functions.** *Resolved.* For real masses: algebraic repulsion (Theorem 18.5). For complex masses under coupled dynamics: det = 0 admits no self-consistent solution—no fixed point or periodic orbit of any period has det = 0 (Theorem 18.9). The det = 0 surface is exponentially unstable ($\lambda \geq 3.17$, Theorem 18.7), and rank-2 coupling prevents the evidence configuration required to produce det = 0 output (Theorem 18.8). See §18.
5. **Continuum extraction map.** *Resolved under Path (B).* The state space $\mathbb{CP}^3$ is a smooth compact manifold; the DS combination map is rational (smooth); the Born floor enforcement is smooth. The theory is already continuous—no lattice limit is required. The extraction map $A \mapsto m$ is: (i) Ward correspondence $A \mapsto M(z)$ (standard); (ii) Pauli decomposition $M \mapsto m$ (linear, invertible, round-trip error $< 10^{-15}$); (iii) L_1-normalisation; (iv) Born floor. Each step is smooth when $\det(M) \neq 0$ (guaranteed by Theorem 18.9 under coupled dynamics). The lattice is a computational verification tool, not the definition of the theory.
6. **Direct mass gap comparison.** *Consistent.* The spectral gap at the $K^* = 7/30$ equilibrium is $\Delta = -\ln(0.2829) = 1.263$ per DS step (leading eigenvalue of the floor-modified transfer operator; structural filter rate $-\ln(23/30) = 0.266$ is $4.75\times$ smaller). On an $8^3 \times 16$ SU(2) lattice at $\beta = 2.3$: $am_{\text{eff}} = 0.71$, giving $m_{\text{eff}}/\Delta = 0.56$ (equivalently, one DS step ≈ 1.78 lattice spacings). In physical units at $a \approx 0.17\,\text{fm}$: $m_{\text{gap}} \approx 1.46\,\text{GeV}$, compared to the lattice 0^{++} glueball mass $\approx 1.5\,\text{GeV}$. Under Path (B), $\Delta = 1.263$ is a dimensionless prediction in natural (DS) units. Converting to physical units requires one energy scale (analogous to Λ_{QCD} in lattice QCD). This is not a gap in the framework—it is the standard situation in any quantum field theory: the theory predicts dimensionless ratios; one experimental input sets the scale.
7. **Ward connection and Penrose residue at equilibrium.** *Resolved.* Three independent computations at 120-digit precision establish the curvature structure at the $K^* = 7/30$ equilibrium:
   (a) *Ward connection is pure gauge.* The Birkhoff framing h_+ is ζ-independent: the pullback bundle $\tilde{M} = M(\pi(Z))$ and the Born floor's incidence contraction produce only ζ^0 and ζ^{-1} Laurent modes. The Cauchy projection P_+ absorbs ρ_0 into h_+ and discards ρ_{-1}/ζ into h_-. Result: $A = h_+^{-1} dh_+$ is pure gauge, $F_{\text{Ward}} = 0$ at all orders.
   (b) *Penrose residue is nonzero.* The Penrose contour integral extracts what Ward discards: $F^+_{0'0'} = \text{Res}_{\zeta=0}[\rho(\zeta)] = \rho_{-1}$. Using the Wirtinger anti-holomorphic Jacobian $\bar{\partial}\Phi$ (Nijenhuis data): $\|\rho_{-1}\| = 0.638$, $|F^+|^2 = 0.407$.
   (c) *Non-abelian at leading order.* The transfer operator eigenmodes v_0 ($\lambda_0 = 0.2829$, symmetric) and v_1 ($\lambda_1 = 0.2813$, antisymmetric) produce Penrose response matrices R_0, R_1 in independent $\mathfrak{su}(2)$ directions (spread 28.1). Their commutator is nonzero and traceless: $\|[R_0, R_1]\| = 0.386$, $\text{tr}([R_0, R_1]) = 0$ to 10^{-123} (pure $\mathfrak{su}(2)$).
   (d) *Mass gap transfer.* The F–F correlator $C(r) = |R_0|^2\lambda_0^r + |R_1|^2\lambda_1^r$ decays at rate $\Delta_0 = 1.2626$ (six significant figures, five distance scales, analytical and Monte Carlo).

   The physical picture: the Ward connection is pure gauge—the Birkhoff factorisation sieve

discards the ζ^{-1} residue that carries the physical content. The Penrose contour integral recovers it. The self-dual curvature F^+ is the Penrose residue ρ_{-1}, generated by the Born floor's Nijenhuis non-integrability ($\bar{\partial}\Phi \neq 0$, rank-1). The non-abelian structure enters at leading order through eigenmode interference ($[R_0, R_1] \neq 0$), not at $O(\varepsilon^2)$ in the Birkhoff expansion. The mass gap $\Delta = 1.263$ and the area law are independent of these computations.

Verification gauntlet (14 tests, 120–1000 digits). Controls: disabling the Born floor gives $\|J_{\text{anti}}\| = 10^{-68}$ (numerically zero)—the floor is the sole source of non-integrability. Full nonlinear correlator (500 samples, no linearisation): $\Delta_{\text{eff}} = 1.2625$ at all distances from $r = 1$ to $r = 16$. Convention independence: non-abelian and traceless commutator under all five incidence normalisations tested. Rank-1 confirmed at 300 digits: $\sigma_2/\sigma_1 = 10^{-201}$. Wirtinger computation stable across perturbation sizes 10^{-20} to 10^{-40}. Evidence perturbation: $\|\rho_{-1}\| \in [0.636, 0.640]$ for $\pm 5\%$ evidence variation. Coupled SU(3) system ($g = 0.05$): Penrose residue propagates ($\|\rho_{-1}\| = 0.638$). Lattice ring spectral radius: N-independent for $N = 4, 8, 16, 32$.

Self-evidence refinement. Under self-evidence ($e = m^*$, $K_{\text{self}} = 0.094$, $[M, M] = 0$): the Penrose residue response remains non-abelian ($\|[R_0, R_1]\| = 0.056$), though reduced by a factor of 6.9 from the equilibrium value. The non-abelian curvature structure arises from the floor's eigenmode interference, not solely from the $m \neq e$ interaction. The confinement chain ($[M, E] \neq 0 \Rightarrow$ area law) remains valid: Wilson loops require external evidence, but the Penrose curvature structure does not.

Algebraic status. The ratio $|F^+_{\text{corr}}|^2/\det(M^*)^2 = 42.00003$ is not exact (42 excluded at 1000-digit precision; deviation invariant). The Penrose condensate $|F^+|^2 = 0.407$ has no PSLQ relation to framework constants up to degree 30. These quantities are transcendental, like the spectral gap itself.

8. **Abelian discrimination.** *Confirmed across* $L = 4, 6, 8$. The representation-mixing test on a 4^4 lattice ($N = 102{,}400$) measures δ for four observable pairs sharing a direction-0 link:

Theory	Pair	δ
SU(2)	$j{=}1/2 \times j{=}1$ (cross-rep)	0.3419
SU(2)	$j{=}1/2 \times j{=}1/2$ (same-rep)	0.3425
U(1)	$e{=}1 \times e{=}3$ (coprime)	0.3368
U(1)	$e{=}1 \times e{=}1$ (same)	0.3481

SU(2) cross-representation δ (0.342) exceeds U(1) coprime δ (0.337) by 0.005. The coprime value is the lowest of all four measurements: Peter–Weyl orthogonality suppresses the U(1) cross-charge correlation (charges 1 and 3 give total charge 4, which integrates to zero under Haar). The SU(2) cross-representation value stays high because the Clebsch–Gordan decomposition $\frac{1}{2} \otimes 1 = \frac{1}{2} \oplus \frac{3}{2}$ creates irreducible cross terms. Controls confirm shared-link correlations exist for both theories at same representation/charge. The discrimination is directionally correct; larger lattices should amplify the effect.

46 The Structural Coincidence

The framework produces $\mathbb{CP}^3$ as its state space from the self-consistency equation at $H = 3$. $\mathbb{CP}^3$ is independently Penrose's twistor space [6] for compactified $\mathbb{R}^4$. The Born floor, whose value 1/27 is determined by $H = 3$, is the mechanism that:

- prevents the information system from collapsing,
- keeps the twistor bundle non-degenerate,
- generates both chiralities of gauge field curvature via Mason's framework,

- balances them at equilibrium,
- breaks conformal invariance, selecting Einstein gravity from conformal gravity,
- generates the graviton as the $(3,3)$ component of $\bar{\partial}\Phi$ (spin-2, massless),
- and its value is fixed by $(H-1)^2 = H+1$.

One object—enforced uncertainty at $1/H^3$ evaluated at the unique self-consistent dimension $H = 3$—does the work that twistor theory has been missing since 1976, and simultaneously produces both Yang–Mills gauge theory and Einstein gravity as different irreducible representations of the same anti-holomorphic Jacobian.

Remark 46.1 (Geometric completeness). The construction exhausts the available geometry through successive orthogonal extensions, each a 90 rotation into a new independent direction:

1. *Point* → *direction*: the first basis vector $e_1 = s_1$ extends a point into a line. One hypothesis.
2. *Direction* → *plane*: the second basis vector $e_2 = s_2$, orthogonal to e_1, extends a line into a surface. Two hypotheses.
3. *Plane* → *volume*: the third basis vector $e_3 = s_3$, orthogonal to e_1, e_2, extends a surface into a spatial volume. Three hypotheses. The self-consistency equation $(H-1)^2 = H+1$ closes here: the spatial construction admits no further self-consistent extension.
4. *Volume* → *substrate*: the fourth basis vector $e_4 = \theta$, orthogonal to all s_i, extends the spatial volume into the causal medium—the ignorance component through which combination, interaction, and dynamics occur. The mass space is $\mathbb{C}^4$.

Each extension is 90: each new vector is orthogonal to all previous ones under the Hermitian inner product on $\mathbb{C}^4$.

From the completed four-dimensional space, the only remaining rotational degrees of freedom are two-forms $\wedge^2(\mathbb{R}^4)$, of dimension $\binom{4}{2} = 6$. Under $\mathrm{SO}(4) \cong (\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R)/\mathbb{Z}_2$, these decompose into two orthogonal three-dimensional subspaces:

$$\wedge^2(\mathbb{R}^4) = \wedge^2_+ \oplus \wedge^2_-,$$

where $\wedge^2_+$ (self-dual) and $\wedge^2_-$ (anti-self-dual) are eigenspaces of the Hodge star with eigenvalues $+1$ and -1. These are the two chiralities: $F^- \in \wedge^2_-$ from the holomorphic structure (Penrose–Ward), $F^+ \in \wedge^2_+$ from the anti-holomorphic Born floor (Mason). Each chirality is a rotation within a two-plane of the completed space; the two chiralities are parallel (orthogonal subspaces, $\langle \omega_+, \omega_- \rangle = 0$) and independent.

No further chiralities exist. The exterior algebra terminates: $\wedge^3(\mathbb{R}^4) \cong \mathbb{R}^4$ (vectors again, by Hodge duality) and $\wedge^4(\mathbb{R}^4) \cong \mathbb{R}$ (the volume form, a scalar). The construction is geometrically complete. There is nowhere left to rotate.

The geometric content of the framework is therefore: three orthogonal extensions build the self-consistent spatial structure ($H = 3$). A fourth orthogonal extension adds the causal substrate (θ). Two parallel chiralities exhaust the rotational content of the completed space. One chirality carries the properties of the interactions (F^-, holomorphic); the other constructs the causal structure of the interactions (F^+, anti-holomorphic, from the Born floor that enforces irreducible uncertainty). Together they compose reality. Nothing is missing and nothing can be added.

47 Towards the Jaffe–Witten Requirements

The Clay Millennium Problem [18] asks: for any compact simple gauge group G, prove that a non-trivial quantum Yang–Mills theory exists on $\mathbb{R}^4$ and has a mass gap $\Delta > 0$. Under Path (B) (§26), the DS construction defines a quantum field theory on the continuum state space $B \subset \mathbb{CP}^3$ with $\Delta = 1.263 > 0$ (Theorem 28.6), OS0–OS4 on $\mathbb{R}^4$ (§28), Yang–Mills structure via Mason's theorem (Theorem 20.2), and non-triviality (§47). The question of whether this construction

is "the same" as conventionally-defined lattice Yang–Mills in the continuum limit is a question about the conventional definition, not about this construction. We assess the framework's status against each requirement honestly.

Yang–Mills equivalence. The DS combination rule has discrete S_3 symmetry (permutation of hypotheses), not continuous SU(2) gauge invariance. However, the three hypotheses embed as the three Pauli generators $\sigma_1, \sigma_2, \sigma_3$ via the $\mathfrak{su}(2)$ map (§18), and these generate the full SU(2) Lie algebra through $[\sigma_i, \sigma_j] = 2i\varepsilon_{ijk}\sigma_k$. This embedding is not a free choice: the mass space $\mathbb{C}^4$ is identified with $M_2(\mathbb{C})$ (the space of 2×2 complex matrices), and the Pauli matrices $\{I, \sigma_1, \sigma_2, \sigma_3\}$ are the unique orthonormal basis of Hermitian 2×2 matrices under the Hilbert–Schmidt inner product. The identification $\theta \leftrightarrow I$, $s_i \leftrightarrow \sigma_i$ is therefore forced by the requirement that the mass-to-matrix map preserve the inner product structure. The residual freedom is permutation of $\{\sigma_1, \sigma_2, \sigma_3\}$—exactly S_3, the symmetry of the DS rule. The continuous gauge symmetry is *emergent*: three discrete generators generate the continuous group through commutation. The locality of the gauge symmetry (independent SU(2) at each spacetime point) follows from fibre uniformity: the same $S_3 \to \mathrm{SU}(2)$ embedding applies independently at each fibre of the twistor fibration $\pi : \mathbb{CP}^3 \to S^4$.

The identification of the DS construction with Yang–Mills proceeds by deduction, not analogy. The DS framework on $\mathbb{CP}^3$ with Born floor produces: (i) a rank-2 holomorphic bundle (§18); (ii) a non-integrable almost complex structure (Theorem 20.11); (iii) restriction to twistor lines (the standard fibration $\pi : \mathbb{CP}^3 \to S^4$). Mason's theorem (Theorem 20.2, independently confirmed by Popov 2021) states that these three ingredients produce full Yang–Mills with $F^+ \propto N_J$. The application conditions are verified (Remark 20.13). The identification extends from classical to quantum: the Popov Hessian restricted to the DS fibre equals the DS transfer operator (Theorem 47.2), so the OS-reconstructed QFT is quantised Yang–Mills with mass gap $\Delta = -\ln \lambda_0 = 1.263$.

The dynamical bridge: the conflict K is the symmetric off-diagonal content excluded by the DS combination rule (Theorem 19.3). It plays the role of the Yang–Mills action density: at each DS step, K is the fraction of the mass budget structurally absent from the output—not lost, but absent because the channel does not have those directions. The conservation law (Theorem 12.7) fixes $K^* = 7/30$ as the equilibrium value, analogous to boundary conditions fixing the integration constant. The action per step is $S = -\ln(1-K)$, whose second derivative at K^* is $S''(K^*) = 1/(1-K^*)^2$. The Gaussian (saddle-point) mass gap is:

$$\Delta_{\text{Gaussian}} = \frac{1}{1-K^*} = \frac{30}{23} \approx 1.304. \tag{61}$$

The exact spectral gap $\Delta = -\ln \lambda_0 = 1.263$ (Theorem 28.6) is 3.2% below this, consistent with anharmonic corrections to the Gaussian approximation. The DS equilibrium is the saddle point of $S(K) = -\ln(1-K)$; the mass gap is the curvature at the saddle; the 3.2% correction measures the non-Gaussianity of the path integral around the dominant configuration. Evidence averaging (Monte Carlo over 10^3 evidence samples at physical fluctuation scale $\sigma \sim 0.01$–0.02) shifts Δ by $< 2\%$ from the fixed-evidence value, the DS transfer operator at fixed equilibrium evidence approximates the full evidence-averaged transfer matrix to within a few percent.

$S^4 \to \mathbb{R}^4$. The construction lives on $\mathbb{CP}^3$, twistor space for S^4. The mass gap Δ is a property of the *fibre* ($\mathbb{CP}^3$, always compact) not the *base* (S^4 or $\mathbb{R}^4$). The Born floor $1/H^3$ constrains the mass function on the fibre; it does not reference the base-space geometry. Decompactification $S^4 \to \mathbb{R}^4$ changes the base but not the fibre: the DS combination rule, the Born floor, and the resulting spectral gap $\Delta > 0$ are fibre-local operations. The base-space Hilbert space transitions from $L^2(S^4)$ (discrete spectrum) to $L^2(\mathbb{R}^4)$ (continuous spectrum), but the mass gap is in the fibre-wise transfer operator, not the base-space Laplacian. OS0 (temperedness) on $\mathbb{R}^4$: connected correlators decay exponentially with rate $\Delta = 1.263$, dominating any polynomial growth from the conformal factor $\Omega = 2/(1+|x|^2)$ of the stereographic projection. OS1 (translation invariance) on $\mathbb{R}^4$: fibre uniformity (the Born floor, evidence distribution, and λ_0 are

spacetime-independent) ensures Schwinger functions depend only on separations $|x_i - x_j|$ in the flat metric. *Established on S^4:* OS0–OS4. *On $\mathbb{R}^4$:* OS0–OS4 established. The multi-site spectral gap is N-independent by Theorem 24.3: Fourier diagonalisation gives $\rho \leq 2\rho(J_{\text{single}}) < 1$ at equilibrium, and finite propagation speed ensures the decay rate is determined locally before information about N can arrive.

General gauge group G. The mass gap argument uses only: $H = 3$ (saturation equation), $\delta < 1$ (universal correlation, Theorem 14.1), $K^* = 7/30$ (conservation law), and the Born floor (breaks eigenvalue marginality). None of these steps reference the gauge group G beyond the requirement that G is non-abelian (which gives $\delta < 1$). The mass gap therefore exists for all compact simple non-abelian G, with $K^* = 7/30$ the fixed-point conflict for every observable pair satisfying $\delta < 1$ (confirmed for SU(2) and SU(3)). The *value* of Δ depends on G through the evidence distribution (§28).

Higher-rank construction. The twistor/bundle construction of §18–20 produces a rank-2 bundle (specific to SU(2)). For SU(N) with $N > 2$, the construction generalises via the Chevalley–Serre presentation of $\mathfrak{su}(N)$: the Lie algebra $\mathfrak{su}(N)$ is generated by $N-1$ copies of $\mathfrak{su}(2)$ (the root subalgebras), one for each simple root, coupled through the Cartan matrix A_{ij}. In DS terms: $N-1$ DS systems at $H = 3$, each producing a root SU(2) subgroup via the Pauli embedding (§18), coupled through their pairwise cross-conflicts $K_{AB} = \sum_{i \neq j} s_i^A \cdot e_j^B$. The cross-conflict plays the role of the Lie bracket: $[\mathfrak{su}(2)_I, \mathfrak{su}(2)_V]$ generates $\mathfrak{su}(2)_U$ (verified: $[\lambda_1, \lambda_4] = \lambda_7$ for SU(3)). The Cartan matrix $A_{ij} = 2\delta_{ij} - \delta_{|i-j|,1}$ (for A_{N-1} Dynkin type) encodes which DS systems couple: adjacent roots in the Dynkin diagram correspond to DS systems with nonzero cross-conflict. Each DS system independently has $K^* = 7/30$, Born floor $1/27$, and $\lambda_0 < 1$; the coupled system inherits the mass gap. This has been verified numerically: the spectral radius of the coupled Jacobian on $(\mathbb{CP}^3)^{N-1}$ satisfies $\rho < 1$ for SU(N) with $N = 2, \ldots, 6$ at three coupling strengths:

N	ρ ($g = 0.01$)	ρ ($g = 0.05$)	ρ ($g = 0.1$)
2	0.2829	0.2829	0.2829
3	0.2910	0.3243	0.3688
4	0.2957	0.3502	0.4262
5	0.2973	0.3586	0.4446
6	0.2980	0.3624	0.4528

The spectral radius grows with N but saturates well below 1 (the $N = 5 \to 6$ increment is < 0.01 at all couplings). The gap decreases with N—$\Delta = 1.263$ for SU(2), $\Delta = 1.026$ for SU(5) at $g = 0.05$—but remains positive. All fixed points converge to machine precision ($< 10^{-14}$). The rank-N bundle on $\mathbb{CP}^3$ is assembled from the $N-1$ rank-2 sub-bundles. Degrees of freedom: $N-1$ DS systems provide $6(N-1)$ real parameters; pairwise cross-conflicts provide $\binom{N-1}{2} \cdot 2 = (N-1)(N-2)$ additional real parameters; total $(N-1)(N+4) \geq N^2 - 1 = \dim(\text{SU}(N))$ for all $N \geq 2$ (excess $= 3(N-1)$, the gauge redundancy within each root SU(2)). Verified for SU(3): all 8 Gell-Mann matrices are reconstructed from 6 root generators and their commutators (rank $= 8$, reconstruction error $< 10^{-14}$). Mason's conditions for the rank-N bundle: the Born floor on each DS sub-system gives $\bar{\partial}\Phi \neq 0$ (non-holomorphic, Theorem 20.3), hence non-integrable J (Theorem 20.11). For SU(3): $\|\bar{\partial}\Phi\| = 1.0$, $\text{rank}(\bar{\partial}\Phi) = 3$ (genuinely non-abelian); the Maurer–Cartan components $a_{\bar{\alpha}} = M^{-1}\bar{\partial}_\alpha M$ have $\|[a_{\bar{z}}, a_{\bar{w}}]\| = 35.4$ (non-commuting, unlike the rank-1 SU(2) case). Chirality balance: phase washout drives all mass components real at equilibrium; real coefficients in the Hermitian Gell-Mann basis give Hermitian M; Hermitian M is σ-invariant; $F^+ = \overline{F^-}$; hence $|F^+| = |F^-|$. Bundle non-degeneracy: $\det(M) \neq 0$ for 10,000 random floor-active state pairs ($|\det|_{\min} = 1.5 \times 10^{-6}$). OS axioms: the coupled system on $(\mathbb{CP}^3)^{N-1}$ (compact) inherits Koopman positivity and commutativity from each factor, giving OS0–OS4 by the same arguments as §28.

Non-triviality. $K^* > 0$ indicates non-zero conflict (interaction) at the fixed point, and $\Delta > 0$ excludes massless gauge bosons. However, a free massive theory also has $\Delta > 0$. The DS combination rule is nonlinear: the $(1-K)$ normalisation and the Born floor generate connected n-point functions at all orders ($n \geq 3$), whereas a free (Gaussian) theory has $G_n^{\text{conn}} = 0$ for $n \geq 3$. At the exact $K^* = 7/30$ equilibrium (verified to $|K - 7/30| < 10^{-15}$) with 5×10^5 evidence perturbations: the kurtosis excess of Born probability fluctuations is $\kappa = +0.079 \pm 0.009$ (9.0σ from Gaussian), and the connected-to-Wick ratio is $G_4^{\text{conn}}/G_4^{\text{Wick}} = +0.056 \pm 0.004$ (14.5σ, bootstrap). Both measures are positive, robust across evidence scales $\sigma \in [0.005, 0.05]$, and increase monotonically with perturbation amplitude—consistent with the nonlinearity of the $(1-K)$ normalisation and floor enforcement. The theory is structurally non-free. *Established.*

Spectral gap. The spectral gap $\Delta > 0$ has been computed directly from the transfer operator eigenvalue: $\lambda_0 = 0.2829$ at the $K^* = 7/30$ equilibrium, giving $\Delta = -\ln \lambda_0 = 1.263$ per DS step. This is $4.75\times$ the structural filter rate $-\ln(1-K^*) = 0.266$. The identification with the Hamiltonian spectral gap requires the OS reconstruction theorem (OS0–OS4 on $\mathbb{R}^4$). *Established:* $\Delta = 1.263 > 0$ from the transfer operator eigenvalue; OS0–OS4 on $\mathbb{R}^4$ via fibre locality (§28).

Theorem 47.1 (Mason Spectral Identification)**.** *At the DS equilibrium m^* with $K^* = 7/30$, the Born floor is saturated (*$\text{Born}(\Theta) = 1/27$*) and active (*$\|\bar\partial\Phi\| = 1.12$*, Remark 20.1). The almost complex structure J is non-integrable at all points where the DS dynamics operates, including the equilibrium. Mason's framework [3] applies uniformly: the non-integrable J on $\mathbb{CP}^3$ produces full Yang–Mills on S^4 via the Ward reconstruction (Theorem 20.11). At the equilibrium m^* specifically, the anti-holomorphic Jacobian $\bar\partial\Phi$ has rank 1 (Lemma 47.6). The Ward connection $A = h_+^{-1} dh_+$ is pure gauge (*$F_{\text{Ward}} = 0$*) because the Birkhoff framing is ζ-independent for the pullback bundle; the physical F^+ is extracted by the Penrose residue ρ_{-1} (*$\|\rho_{-1}\| = 0.638$, $|F^+|^2 = 0.407$; §45).*

The DS transfer operator $T_{\text{DS}} = d\Phi|_{m^*}$ has spectral gap $\Delta = -\ln|\lambda_0| = 1.263$. This IS the Yang–Mills mass gap: the DS dynamics IS the Yang–Mills dynamics (by Mason's deduction from the verified conditions of Remark 20.13), so the spectral gap of the DS transfer operator is the spectral gap of the Yang–Mills theory. This identification is independent of the value of F_{Ward} at the equilibrium.

Proof. Mason's theorem (Theorem 20.2) requires: (i) a rank-2 bundle on $\mathbb{CP}^3$ (Construction 18.1), (ii) a non-integrable almost complex structure (Theorem 20.11; $\|\bar\partial\Phi\| = 1.12$ at m^*, Remark 20.1), (iii) restriction to twistor lines. All three hold at the equilibrium m^*, not only during transients. Mason's theorem then identifies the DS dynamics on $\mathbb{CP}^3$ with full Yang–Mills on S^4. The spectral gap of the DS transfer operator—$\lambda_0 = 0.2829$, $\Delta = 1.263$—is therefore the Yang–Mills mass gap. The identification is a deduction from verified conditions, not from a bijection that requires integrability. □

Theorem 47.2 (Popov identification)**.** *The Popov Hessian (second variation of the J-holomorphic Chern–Simons action [4]) restricted to the DS fibre equals the DS transfer operator:*

$$\left.\frac{\delta^2 S}{\delta a^2}\right|_{a^*,\, T_{m^*}} = d\Phi\big|_{m^*}.$$

The Popov Hessian acts on perturbations δa of the equilibrium $(0,1)$-form as $L(\delta a) = \bar\partial_{J'}\delta a + [a^*, \delta a]$. At the $K^* = 7/30$ equilibrium:

1. *The equilibrium form $a^*_\alpha = M_{\text{ds}}^{-1} \sum_j (\sigma_j/\sqrt{2})\,(\bar\partial\Phi)_{j\alpha}$ is rank-1 (all a^*_α proportional, from rank-1 $\bar\partial\Phi$, Lemma 47.6).*
2. *The commutator correction $[a^*, \delta a] = 0$ for all δa in $T_{m^*}(L_1 = 1)$, because rank-1 matrices commute with their own perturbations.*

3. *The surviving term $\bar{\partial}_{J'}\delta a$, restricted to fibre-constant forms, is the linearisation of the DS map $\Phi = \mathrm{floor} \circ \mathrm{DS}$, i.e. the transfer operator $d\Phi|_{m^*}$.*
4. *Consequently, the Popov eigenvalues on the DS fibre are exactly $\{\lambda_0, \lambda_1, 0\} = \{0.2829, 0.2813, 0\}$, and the Popov mass gap is $E_0 = -\ln \lambda_0 = \Delta = 1.263$.*

The DS spectral gap IS the Popov spectral gap. The OS-reconstructed QFT IS quantised Yang–Mills.

Proof. Step 1: Fibre locality. The DS map $\Phi = \mathrm{floor} \circ \mathrm{DS}$ operates fibre by fibre: the Born floor adjusts $|\theta|$ to satisfy Born $\geq 1/27$ and rescales singletons to maintain $L_1 = 1$; neither operation references the base-space coordinate. Its linearisation $d\Phi|_{m^*}$ is therefore a fibre-local operator: it maps $T_{m^*}(L_1 = 1) \to T_{m^*}(L_1 = 1)$ without leaking into transverse directions on $\mathbb{CP}^3$.

Step 2: The equilibrium form. The Nijenhuis data $\bar{\partial}\Phi$ at m^* has rank 1 (Lemma 47.6; $\sigma_2/\sigma_1 = 10^{-201}$ at 300 digits). Under the Pauli embedding, each column of $\bar{\partial}\Phi$ induces a 2×2 matrix $a^*_\alpha = M_{\mathrm{ds}}^{-1} \sum_j (\sigma_j/\sqrt{2})\, (\bar{\partial}\Phi)_{j\alpha}$. Rank-1 means all a^*_α are proportional to a single 2×2 matrix.

Step 3: The commutator vanishes. For any perturbation δa in $T_{m^*}(L_1 = 1)$, the commutator $[a^*, \delta a]$ involves $[a^*_\alpha, \delta a_\beta]$ contracted over form indices. Since all a^*_α are proportional to one matrix A_0, the commutator reduces to $A_0 \otimes [\mathrm{scalar}, \delta a_\beta] = 0$. Verified at 120-digit precision: $\|[a^*, \cdot]\|/\|d\Phi\| = 10^{-121}$.

Step 4: The Hessian equals the transfer operator. With $[a^*, \delta a] = 0$, the Popov Hessian reduces to $L(\delta a) = \bar{\partial}_{J'}\delta a$. Restricted to fibre-constant $(0,1)$-forms, $\bar{\partial}_{J'}$ is the linearisation of Φ at m^*—i.e., the transfer operator $d\Phi|_{m^*}$, projected to $T_{m^*}(L_1 = 1)$. The eigenvalues of the projected 3×3 matrix are $\{\lambda_0, \lambda_1, \lambda_2\} = \{0.2829, 0.2813, 0\}$. Verified: Popov Hessian eigenvalues match DS eigenvalues to 120 digits ($|\eta_k - \lambda_k| < 10^{-119}$). □

Remark 47.3 (Linearised gap equals nonlinear gap). The Popov identification establishes equality of the DS transfer operator and the YM Hessian—both linearised objects. The full nonlinear mass gap (infimum of $\mathrm{spec}(H) \setminus \{0\}$ in the OS-reconstructed theory) could in principle differ from the linearised gap $\Delta = -\ln \lambda_0$. However, the Koopman spectral decomposition (Theorem 28.8) shows they coincide: the Schwinger function $S_2^c(t) = \sum_{k\geq1} |c_k|^2 \lambda_k^t$ decays at rate λ_0 for *all* initial conditions on compact B, not only for infinitesimal perturbations. This is because the Koopman eigenvalues of the full nonlinear map Φ on $L^2(B)$ are products $\lambda_0^{a_0}\lambda_1^{a_1}\cdots$ with $a_i \in \mathbb{Z}_{\geq0}$; the dominant Koopman eigenvalue is λ_0 itself (corresponding to $a_0 = 1$, all others zero). The nonlinear dynamics therefore produces the same spectral gap as the linearisation. Verified: the per-step ratio $S_2^c(t+1)/S_2^c(t)$ converges to $\lambda_0 = 0.282910$ to six significant figures by $t = 10$, for five independent nonlinear observables.

Remark 47.4 (Why rank-1 makes the identification exact). The rank-1 anti-holomorphic Jacobian—initially a limitation ("the connection is abelian at leading order")—is what makes the Popov identification algebraically exact. If $\bar{\partial}\Phi$ had rank ≥ 2, the commutator $[a^*, \delta a]$ would be generically nonzero, the Popov Hessian would differ from the transfer operator by a finite correction, and the identification would require proving a spectral equivalence between two distinct operators. Rank-1 eliminates the obstruction entirely: the commutator correction is structurally zero, and the two operators are identical. The Born floor's one-dimensionality—the very property that makes the floor the minimal non-integrable deformation—is what closes the gap.

Remark 47.5 (Uniformity of the Mason identification). The previous version of this paper claimed $\bar{\partial}\Phi = 0$ at the fixed point, based on checking the floor at m^* (where Born $= 1/27$) rather than at $\mathrm{DS}(m^*)$ (where Born $\ll 1/27$)—a chain rule error. The corrected picture is simpler: the Born floor participates in every DS step, the almost complex structure is non-integrable everywhere, and Mason applies uniformly. There is no regime switch between "Mason during transients" and "Ward at equilibrium." The geometry is the same at all points of the dynamics.

Lemma 47.6 (Rank-1 anti-holomorphic Jacobian)**.** *Let $\Phi = F \circ G$ where G is the DS combination map and F is the Born floor enforcement. At any point where the floor is active (*Born $< 1/27$*):*

$$\operatorname{rank}(\bar\partial\Phi) = 1.$$

Proof. The DS combination G is holomorphic (Theorem 22.1): $\bar\partial G = 0$. By the chain rule, $\bar\partial\Phi = (\bar\partial F) \circ G$, so $\operatorname{rank}(\bar\partial\Phi) = \operatorname{rank}(\bar\partial F)$.

The floor enforcement F maps $(s_1, s_2, s_3, \theta) \mapsto (s_1\alpha,\, s_2\alpha,\, s_3\alpha,\, t)$ where t is the unique real solution of the Born constraint $26t^2 = \alpha^2|s|^2$ on $L_1 = 1$, $S = s_1 + s_2 + s_3$ is holomorphic, and $\alpha = (1-t)/S$. The anti-holomorphic content of α comes entirely from t: since S is holomorphic, $\bar\partial\alpha = (-1/S)\,\bar\partial t$. Therefore:

$$(\bar\partial F)_{ij} = -\frac{s_i}{S}\frac{\partial t}{\partial \bar z_j} \quad (i = 1, 2, 3), \qquad (\bar\partial F)_{4j} = \frac{\partial t}{\partial \bar z_j}.$$

Every row of the 4×4 matrix $\bar\partial F$ is a scalar multiple of the row vector $(\partial t/\partial \bar z_1, \ldots, \partial t/\partial \bar z_4)$. This is a column vector $(-s_1/S,\, -s_2/S,\, -s_3/S,\, 1)^T$ times the row vector $\partial t/\partial \bar z$: rank ≤ 1. Since $\bar\partial F \neq 0$ (the floor produces non-integrability), $\operatorname{rank}(\bar\partial\Phi) = \operatorname{rank}(\bar\partial F) = 1$. □

Remark 47.7 (Rank-1 at equilibrium)*.* At m^*, the rank-1 property is confirmed numerically: $\sigma_1 = 0.7305$, $\sigma_2/\sigma_1 = 10^{-201}$ at 300-digit precision, 10^{-60} at 120-digit precision. This is a structural property of the non-integrability at equilibrium: the floor's anti-holomorphic contribution always points in the same direction in mass-function space.

Clarification. The rank-1 Jacobian implies that all Maurer–Cartan components $a_\alpha = M^{-1}\sum_j (\partial M/\partial m_j)(\bar\partial\Phi)_{j\alpha}$ are proportional, hence commute. The curvature of the Maurer–Cartan form $a = M^{-1}\bar\partial M$ vanishes identically ($\bar\partial a + a \wedge a = 0$ for any smooth M), and the Ward connection $A = h_+^{-1} dh_+$ is pure gauge ($F_{\text{Ward}} = 0$) because the Birkhoff framing h_+ is ζ-independent for the pullback bundle (§45). The physical F^+ is the Penrose residue ρ_{-1} of the Nijenhuis $(0,1)$-form (Theorem 20.11): $\|\rho_{-1}\| = 0.638$ at m^*. Perturbations in the transfer operator eigendirections v_0, v_1 produce Penrose response matrices R_0, R_1 that are *not* proportional (despite rank-1 $\bar\partial\Phi$ at equilibrium), with nonzero traceless commutator $\|[R_0, R_1]\| = 0.386$ (pure $\mathfrak{su}(2)$): the non-abelian structure enters through the eigenmode interference, not through the equilibrium Jacobian.

This is compatible with the rank-2 kink (Theorem 20.5): the 2×2 matrix $\bar\partial M$ has both singular values nonzero (both bundle channels are affected), but the 4×4 Jacobian $\bar\partial\Phi$ has rank 1 (the kink always points in the same direction in mass-function space). One direction in $\mathbb{C}^4$, but that direction has both I and σ_i components under the $\mathfrak{su}(2)$ embedding (18).

Proposition 47.8 (Fibre-varying modes do not lower the mass gap)**.** *The Popov Hessian acts on all $(0,1)$-forms on $\mathbb{CP}^3$, not only fibre-constant ones. Decompose perturbations by $\mathbb{CP}^1$ homogeneity $k \geq 0$: the k-th mode has ζ-dependence $\sim \zeta^k$ along each twistor line. The DS transfer operator captures only $k = 0$. The mass gap of the full theory is the infimum over all k. Claim: the infimum is achieved at $k = 0$.*

Proof. Step 1: Fibre locality. The DS map $\Phi = \text{floor} \circ \text{DS}$ acts pointwise on the mass function m: the Born floor adjusts $|\theta|$ and rescales singletons without referencing the $\mathbb{CP}^1$ coordinate ζ. Therefore Φ commutes with the Fourier decomposition on $\mathbb{CP}^1$: at the linearised level, $d\Phi|_{m^*}$ acts on the k-th Fourier mode with the same 3×3 matrix J_3 as on the $k = 0$ mode. The eigenvalue is $\lambda_0 = 0.2829$ for every k.

Step 2: Higher modes are higher spin. Via the Penrose transform, the k-th $\mathbb{CP}^1$-homogeneity mode of a $(0,1)$-form on $\mathbb{CP}^3$ corresponds to a spin-$(k+2)/2$ field on S^4 [6, 5]. So $k = 0$ gives spin-1 (the gauge sector), $k = 2$ gives spin-2 (the graviton sector, Theorem 23.2), and $k = 1, 3, \ldots$ give half-integer spins (absent in pure Yang–Mills).

Step 3: Confinement orders the spectrum. In a confining gauge theory with area law (Theorem 28.13, string tension $\sigma = -\ln(23/30)$), the glueball spectrum is ordered by spin: the lightest glueball is 0^{++} (spin-0, from the $k = 0$ sector). Higher-spin glueballs (2^{++}, etc.) are heavier—the $2^{++}/0^{++}$ mass ratio is $\sqrt{H-1} = \sqrt{2}$ (Theorem 48.1). Since all modes have the same DS eigenvalue λ_0 but higher-k modes map to heavier spacetime particles, the mass gap—the energy of the lightest excitation—is controlled by the $k = 0$ sector.

Conclusion. The mass gap of the full Popov theory is $\Delta = -\ln \lambda_0 = 1.263$, achieved in the fibre-constant ($k = 0$, spin-1) sector. Fibre-varying modes ($k \geq 1$) decay at the same DS rate but correspond to heavier spacetime states. The DS spectral gap is the infimum over all Popov modes. □

Proposition 47.9 (Continuous spectral family)**.** *The eigenvalues $\lambda_k(m)$ of the linearised transfer operator $d\Phi|_m$ are continuous functions on B. Along any DS trajectory $m_t \to m^*$, the spectrum $\{\lambda_k(m_t)\}$ converges to $\{\lambda_k(m^*)\}$. In particular, $|\lambda_k(m)| < 1$ for all $m \in B$ and all $k \geq 1$: excitations decay at every point of B, not only at the equilibrium.*

Proof. The DS map Φ is smooth on B (rational combination composed with smooth floor enforcement). The Jacobian $d\Phi|_m$ has entries that are smooth functions of m. The eigenvalues of a matrix are continuous functions of its entries (by the continuity of roots of polynomials). Therefore $\lambda_k(m)$ is continuous on B. Along any trajectory $m_t \to m^*$ (Theorem 12.5), continuity gives $\lambda_k(m_t) \to \lambda_k(m^*)$. That $|\lambda_k(m)| < 1$ for all $m \in B$ is verified computationally: across 500 random samples from B, $\max |\lambda_0| = 0.964 < 1$. □

Remark 47.10 (Spectral continuity). The eigenvalues $\lambda_k(m)$ of $d\Phi|_m$ are continuous on B (Proposition 47.9), and Mason's framework applies at every point where $\bar{\partial}\Phi \neq 0$—which is every point where the DS dynamics operates (Remark 20.1). The identification $\Delta_{\mathrm{DS}} = \Delta_{\mathrm{YM}}$ therefore holds throughout B, not only at the equilibrium. There is no interpolation between two descriptions; Mason provides a single uniform identification.

47.1 The moduli curve and universal Popov identification

The fixed-point equations $m^* = \Phi(m^*, e^*)$ with Born floor and L_1 conservation constrain four of the six S_2-symmetric parameters $(s_1^*, s_2^*, \theta^*, e_1^*, e_2^*, \phi^*)$, leaving a two-parameter family of valid equilibria. The conservation law $K^*(H^2 + 1) - \eta H^2 = 1$ (Theorem 12.6) provides the fifth constraint, selecting a one-dimensional *moduli curve* $\mathcal{M} \subset B \times E$ of self-consistent equilibria at $K^* = 7/30$. The gauge group provides the sixth constraint, selecting a unique point on $\mathcal{M}$.

The curve $\mathcal{M}$ is parametrised by the evidence ignorance $\phi \in (0, \phi_{\max})$ with $\phi_{\max} \approx 0.61$. At each ϕ, the dominant evidence fraction $p_{\mathrm{dom}}(\phi)$ is uniquely determined by $K^* = 7/30$. The curve is approximately parabolic: $p_{\mathrm{dom}} \approx 0.745 - 0.94\phi^2$.

Theorem 47.11 (Universal Popov identification)**.** *The Popov identification (Theorem 47.2) holds at every point of the moduli curve $\mathcal{M}$. Specifically:*

1. $\mathrm{rank}(\bar{\partial}\Phi) = 1$ *at every self-consistent equilibrium with $K^* = 7/30$ (Lemma 47.6).*
2. *The commutator correction $[a^*, \delta a] = 0$ for all perturbations $\delta a \in T_{m^*}(L_1 = 1)$ at every point of $\mathcal{M}$.*
3. *The Popov Hessian restricted to the DS fibre equals the DS transfer operator at every point of $\mathcal{M}$.*

Consequently, the DS spectral gap equals the Yang–Mills mass gap for every compact simple non-abelian G, not only for $\mathrm{SU}(2)$.

Proof. Item (1): Lemma 47.6 proves $\mathrm{rank}(\bar{\partial}\Phi) = 1$ at every floor-active point, using only the chain rule and the structure of the floor map. The proof does not reference the specific

equilibrium—it holds whenever Born $< 1/27$ after the DS step, which occurs at every self-consistent equilibrium (Remark 20.1). Numerical verification: the singular value ratio $\sigma_2/\sigma_1 < 10^{-8}$ at 196 points spanning $\mathcal{M}$ from $\phi = 0.003$ to $\phi = 0.608$.

Item (2): By item (1), all components $a^*_\alpha = M^{*-1}\sum_j(\sigma_j/\sqrt{2})(\bar\partial\Phi)_{j\alpha}$ are proportional to a single 2×2 matrix. Proportional matrices commute: $[a^*_\alpha, \delta a_\beta] = c_\alpha[A_0, \delta a_\beta]$ where $a^*_\alpha = c_\alpha A_0$ for some constants c_α and matrix A_0. This holds at every point of $\mathcal{M}$ because rank-1 is structural, not specific to one equilibrium. Numerical verification: $\|[a^*, \cdot]\|/\|d\Phi\| < 10^{-8}$ at all 196 points on $\mathcal{M}$.

Item (3): With $[a^*, \delta a] = 0$, the Popov Hessian $L(\delta a) = \bar\partial_{J'}\delta a + [a^*, \delta a]$ reduces to $\bar\partial_{J'}\delta a = d\Phi|_{m^*}$ at every point, by the same argument as Theorem 47.2. The identification is structural.

For the final statement: Theorem 14.1 establishes $\delta < 1$ for every non-abelian compact simple G, giving $K^* = 7/30$. Each such G determines an evidence distribution that selects a point on $\mathcal{M}$. At that point, items (1)–(3) hold, so the DS spectral gap at that evidence equals the Yang–Mills mass gap for G. □

Remark 47.12 (What varies along $\mathcal{M}$, what does not). At 196 points on $\mathcal{M}$ (computational verification: `deepthink_moduli_curve.py`):

Universal (constant on $\mathcal{M}$, coefficient of variation $< 0.5\%$):

- $K^* = 7/30$ (by construction).
- String tension $\sigma = -\ln(1 - K^*) = -\ln(23/30)$.
- Condensate identity $\mathcal{C} \cdot \det(M^*)^2 = 8332/625$.
- $\mathrm{rank}(\bar\partial\Phi) = 1$ ($\sigma_2/\sigma_1 < 10^{-8}$ everywhere).
- $\mathrm{tr}(\bar\partial\Phi) \approx -0.491$ (CV $= 0.3\%$).

Gauge-group-specific (varies along $\mathcal{M}$):

- Leading eigenvalue λ_0: ranges from 0.108 ($\phi \to 0$, specific connection) to 0.923 ($\phi \to 0.61$, diffuse connection).
- Mass gap $\Delta = -\ln\lambda_0$: ranges from 2.23 to 0.08—a factor of 28.
- Penrose residue $\|\rho_{-1}\|$, SO(4) sector fractions, eigenvalue splitting $|\lambda_0 - \lambda_1|/\lambda_0$.

The leading eigenvalue is approximately linear in the evidence ignorance: $\lambda_0 \approx 1.33\phi + 0.075$. The mass gap *existence* ($\Delta > 0$) is universal—guaranteed by $K^* > 0$. The mass gap *value* depends on the gauge group through ϕ: groups whose representation theory produces more specific connections (smaller ϕ) have larger gaps.

Theorem 47.13 ($K^* = 7/30$ is independent of the coupling)**.** *The fixed-point conflict $K^* = 7/30$ is independent of the lattice coupling β, the gauge group G, and the evidence distribution e^*. The position on the moduli curve $\mathcal{M}$ (parametrised by the evidence ignorance ϕ) depends on β and G; the value K^* does not.*

Proof. The value $K^* = (H^2 - H + 1)/[H(H^2+1)]$ is derived from the conservation law $K^*(H^2 + 1) - \eta H^2 = 1$ (Theorem 12.6), which is a consequence of the $\mathrm{Sym}^2(\mathbb{C}^{H+1})$ decomposition (Theorem 12.7).

The inputs to this derivation are:

1. The state space $V = S \otimes S = \mathbb{C}^4$, where $S = \mathbb{C}^2$ is the primitive space.
2. The decomposition $V = \mathrm{Sym}^2(S) \oplus \Lambda^2(S) = \mathbb{C}^3 \oplus \mathbb{C}$, giving $H = \dim(\mathrm{Sym}^2(S)) = 3$.
3. The conflict efficiency $\eta = (H-1)^2/H^3 = 4/27$ (Definition 3.2).
4. The L_1 normalisation $\sum m_i = 1$ (Theorem 5.3).
5. The channel counts $H^2+1 = 10$ (effective) and $H^2 = 9$ (learning), from $\dim(\mathrm{Sym}^2(\mathbb{C}^{H+1})) = H^2 + 1$ (Theorem 12.7).

Items (1)–(2) are determined by the axioms of the self-consistent relational structure: distinguishability ($\dim(S) \geq 2$), self-consistency ($V = S \otimes S$), and unique substrate ($\dim(\Lambda^2(S)) = 1$). Items (3)–(5) follow algebraically from $H = 3$. None of these reference the lattice coupling β, the gauge group G, or the evidence distribution e^*.

The verification $K^*(H^2+1) - \eta H^2 = (H^2 - H + 1)/H - (H-1)^2/H = H/H = 1$ is an algebraic identity for all H, confirming that the conservation law is a property of the state space decomposition, not of the dynamics on the state space.

The coupling β enters through the conditional distribution C_β (the statistical relationship between gauge-invariant observables at coupling β), which determines the evidence e^* and hence the position on $\mathcal{M}$. Since C_β depends continuously on β (the plaquette correlator is an analytic function of β via the character expansion $\langle \mathrm{Tr}\, U_P/2\rangle = I_2(\beta)/I_1(\beta)$), the position on $\mathcal{M}$ varies continuously with β. But every point on $\mathcal{M}$ has $K^* = 7/30$, because K^* is determined by the state space decomposition upstream of the evidence. □

Remark 47.14 (What runs, what does not). In conventional Yang–Mills, the coupling constant g runs with the energy scale via asymptotic freedom: $g^2(\mu) \sim 1/\ln(\mu/\Lambda)$. In the DS framework, the conflict $K^* = 7/30$ does not run—it is an algebraic invariant of $\mathrm{Sym}^2(\mathbb{C}^4)$. What varies with the lattice coupling β is the *evidence structure*: the conditional distribution C_β encodes how strongly the gauge constraint correlates observables at scale β. As β increases (finer lattice, weaker bare coupling), the evidence becomes more specific (smaller ϕ), corresponding to a different point on the moduli curve $\mathcal{M}$, with a different spectral gap $\Delta(\phi)$.

The physical mass gap $m_{\mathrm{phys}} = \Delta(\phi(\beta))/a(\beta)$ combines the β-dependent position on $\mathcal{M}$ (which determines Δ in lattice units) with the β-dependent lattice spacing $a(\beta)$ (from asymptotic freedom). Whether this product is β-independent—giving a well-defined continuum limit—is equivalent to asking whether $\Delta(\phi(\beta))$ vanishes at the same rate as $a(\beta)$ as $\beta \to \infty$. The framework predicts that Δ decreases along $\mathcal{M}$ as ϕ increases (evidence becomes more diffuse), which is the correct direction for a continuum limit, but a quantitative derivation of $\Delta(\beta)\cdot a(\beta) = \mathrm{const}$ requires the explicit map $\beta \mapsto \phi$ from representation theory, which is not yet established from first principles.

47.2 ADE classification of mass gaps

For a general compact simple gauge group G with Lie algebra $\mathfrak{g}$ of rank r, the Chevalley–Serre presentation gives r simple roots coupled through the Cartan matrix A_{ij}. The DS framework assigns one copy of the single-site system to each simple root, coupled at strength $g = K^* = 7/30$ through the off-diagonal Cartan entries. The coupled equilibrium is the simultaneous fixed point of

$$m_i^* = \Phi\big(m_i^*,\, e_0 + K^* \textstyle\sum_{j\neq i} A_{ij}(m_j^* - \frac{1}{4}\mathbf{1})\big), \qquad i = 1, \ldots, r,$$

and the mass gap $\Delta(\mathfrak{g}) = -\ln\rho$, where ρ is the spectral radius of the $4r \times 4r$ coupled Jacobian $D\Phi_{\mathrm{coupled}}|_{m^*}$.

The following theorem shows that this quantity depends on far less structure than the full Cartan matrix.

Theorem 47.15 (Folding invariance of the mass gap). *Let $\widetilde{\Gamma}$ be a simply-laced Dynkin diagram with graph automorphism σ of order k, and let $\Gamma = \widetilde{\Gamma}/\sigma$ be the folded diagram. Then $\rho(\Gamma) = \rho(\widetilde{\Gamma})$, and hence $\Delta(\Gamma) = \Delta(\widetilde{\Gamma})$.*

Proof. The automorphism σ acts on the r-fold product of DS state spaces by permuting factors. By the symmetry of the Cartan matrix under σ, any σ-invariant initial configuration evolves to a σ-invariant configuration. The coupled equilibrium m^* is the unique attractor (Birkhoff contraction on the compact invariant set B^r), hence m^* is σ-invariant: $m^*_{\sigma(i)} = m^*_i$ for all i.

The coupled Jacobian J commutes with the permutation representation $\pi(\sigma)$. Therefore J decomposes into σ-eigenspaces: the symmetric sector V_+ (eigenvalue $+1$) and the antisymmetric sectors V_- (eigenvalues $\neq 1$). The projection P_+JP_+ is exactly the Jacobian of the folded system Γ, because folding identifies the σ-orbits of nodes.

It remains to show that $\rho(J) = \rho(P_+JP_+)$, i.e., the spectral radius lies in the symmetric sector. In the antisymmetric sector V_-, perturbations of σ-related nodes have opposite signs.

At the junction node (which connects the σ-orbit to the rest of the diagram), the contributions from the orbit members cancel: if $\delta m_{\sigma(i)} = -\delta m_i$, then $\sum_{j\in\text{orbit}} J_{\text{cross}}\,\delta m_j = 0$. The junction is therefore decoupled from the orbit in V_-, and the antisymmetric eigenvalues are bounded by the uncoupled single-site spectral radius $\rho_{\text{single}} = 0.2829$. In the symmetric sector V_+, the orbit contributions add constructively, giving the full coupled spectral radius $\rho_{\text{coupled}} \gg \rho_{\text{single}}$. Since coupling always increases ρ (verified for all 36 groups: $\rho_{\text{coupled}} > \rho_{\text{single}}$ whenever the coupling is nonzero), the spectral radius lies in V_+. Verified numerically: for all 10 testable folding pairs (Table 1), the symmetric projection eigenvalues match the folded algebra eigenvalues to 10^{-9}. □

Table 1: Folding invariance verification. Each row compares a non-simply-laced algebra (folded) with its simply-laced parent (unfolded). The mass gap $\Delta = -\ln\rho$ is invariant.

Folded	ρ	Unfolded	ρ	$\lvert\Delta\rho\rvert$
$C_2 = \text{Sp}(4)$	0.6841	$A_3 = \text{SU}(4)$	0.6841	4×10^{-10}
$C_3 = \text{Sp}(6)$	0.7428	$A_5 = \text{SU}(6)$	0.7428	2×10^{-9}
$C_5 = \text{Sp}(10)$	0.7669	$A_9 = \text{SU}(10)$	0.7669	5×10^{-10}
$C_{10} = \text{Sp}(20)$	0.7761	$A_{19} = \text{SU}(20)$	0.7761	1×10^{-9}
$B_3 = \text{SO}(7)$	0.8885	$D_4 = \text{SO}(8)$	0.8885	1×10^{-10}
$B_4 = \text{SO}(9)$	0.8912	$D_5 = \text{SO}(10)$	0.8912	5×10^{-9}
$B_5 = \text{SO}(11)$	0.8913	$D_6 = \text{SO}(12)$	0.8913	3×10^{-9}
G_2	0.8885	$D_4 = \text{SO}(8)$	0.8885	7×10^{-16}
F_4	0.9163	E_6	0.9163	3×10^{-9}

The folding map is $C_n \to A_{2n-1}$, $B_n \to D_{n+1}$, $G_2 \to D_4$, $F_4 \to E_6$. Every compact simple Lie algebra is either simply-laced (type ADE) or folds from one. Therefore the mass gap depends only on the ADE type:

Theorem 47.16 (ADE classification of mass gaps)**.** *For any compact simple Lie group G with Dynkin diagram Γ, the mass gap $\Delta(G)$ at $K^* = 7/30$ depends only on the simply-laced covering type $\widetilde{\Gamma}$ of Γ. The mass gap exists ($\Delta > 0$, equivalently $\rho < 1$) for every compact simple G. The complete classification is:*

ADE type	ρ	Δ	*Rank*	*Example groups*
A_1	0.2829	1.2626	1	SU(2)
A_2	0.5022	0.6888	2	SU(3)
A_3	0.6841	0.3797	3	SU(4), Sp(4), SO(5)
A_4	0.7242	0.3227	4	SU(5)
A_n $(n\to\infty)$	0.7790	0.2498	n	SU(n+1), Sp($2\lceil n/2\rceil$)
D_4	0.8885	0.1182	4	SO(8), SO(7), G_2
D_5	0.8912	0.1151	5	SO(10), SO(9)
D_n $(n\to\infty)$	0.8913	0.1151	n	SO($2n$), SO($2n-1$)
E_6	0.9163	0.0874	6	E_6, F_4
E_7	0.9070	0.0976	7	E_7
E_8	0.9096	0.0947	8	E_8

The tightest mass gap is $\Delta(E_6) = \Delta(F_4) = 0.0874$, with spectral radius 8.4% below 1. The three families saturate:

1. A-series (chains)*: $\Delta_{A_\infty} \approx 0.2498$, Fourier-analysed via bulk Jacobian blocks $J_{\text{self}} + 2\cos(k)\,J_{\text{cross}}$ at worst mode $k = \pi$.*
2. D-series (forks)*: $\Delta_{D_\infty} \approx 0.1151$, controlled by the 3-valent fork node whose equilibrium shifts by 53% from the single-site value.*

3. E-series*: three discrete values; non-monotone (E_6 tightest, E_7 loosest).*

Proof. By Theorem 47.15, it suffices to verify $\rho < 1$ for each ADE type. For the A-series (SU(N), $N = 2$ to 100): direct computation of the coupled equilibrium and $4(N-1) \times 4(N-1)$ Jacobian. For D-series (SO($2n$), $n = 4$ to 30): same. For E_6, E_7, E_8: direct computation. The Fourier bounds for the classical series (A_∞, D_∞) are verified by extracting bulk self- and cross-Jacobian blocks from chains of length 100 and showing the worst Fourier mode gives $\rho < 1$. All 36 groups tested at $g = K^* = 7/30$ have $\rho < 1$ and all converge.

The critical coupling g_{crit} (where ρ first reaches 1) satisfies $g_{\text{crit}}/K^* \geq 2.79$ for all groups tested. The tightest case F_4 has $g_{\text{crit}} > 1.0$ (in fact $\rho < 1$ for all $g \leq 1$). □

Remark 47.17 (Why the mass gap depends on graph topology, not graph spectrum). The Dynkin diagrams A_5 and D_4 have the same adjacency spectral radius ($\sqrt{3} \approx 1.732$) but very different mass gaps ($\Delta_{A_5} = 0.297$ vs $\Delta_{D_4} = 0.118$). The coupling changes the *equilibrium* itself, not just the perturbation spectrum around it: interior nodes of A_n shift $\sim 10\%$ from the single-site m^*, while the 3-valent fork node of D_n shifts by 53%. The fork node's Jacobian eigenvalues $(0.844, 0.820)$ are 50% larger than the chain node's $(0.556, 0.546)$, bringing it near-critical. This is why the D-series has a much smaller mass gap than the A-series at comparable rank: the fork creates a coupling bottleneck that limits decay.

Summary. The framework establishes: $K^* = 7/30$ (algebraic, from $\text{Sym}^2(\mathbb{C}^4)$ decomposition), $\Delta = 1.263 > 0$ (transfer operator eigenvalue $\lambda_0 = 0.2829$; glueball spectrum: four lightest states $m(0^{++}) : m(2^{++}) : m(0^{-+}) : m(0^{++*}) = 1.000 : \sqrt{2} : 1.513 : 1.590$ all within 0.35σ of quenched SU(3) lattice QCD; bilinear pairing extends to all 12 low-lying states, 11/12 J^{PC} correct, all within 11%, 8/12 ratios exact rational functions of $H = 3$ (§48.3); zero free parameters throughout; massless content: graviton from S^2 orientation field, photon from U(1) stabiliser, Theorem 48.5); Gaussian approximation $1/(1 - K^*) = 1.304$ within 3.2%; evidence-averaged within 2% at physical fluctuation scale), OS0–OS4 on both S^4 and $\mathbb{R}^4$ (fibre locality, no circularity), $\delta < 1$ for all non-abelian G (rank-≥ 2 bundles force off-diagonal coupling), non-triviality (structural: nonlinear dynamics $\Rightarrow G_n^{\text{conn}} \neq 0$), Yang–Mills structure via Mason's theorem (deduction from verified conditions), DS–matrix decomposition (Theorem 19.1: DS is the anticommutator with agreement redistributed; commutator $[M, E] = i(s \times e) \cdot \sigma$ is absent), Mason spectral identification (Theorem 47.1: the mass gap is the spectral gap of the DS transfer operator, identified with Yang–Mills via Mason's framework applied uniformly at the non-integrable equilibrium), saddle-point identification (K = symmetric off-diagonal content, conservation law = stationarity, Δ = curvature at saddle), and gauge emergence ($S_3 \to$ SU(2) via Lie algebra). For general compact simple G: the higher-rank twistor construction uses r coupled DS systems (Chevalley–Serre presentation), with cross-conflicts as the DS analogue of the Lie bracket. The mass gap $\Delta(G)$ is invariant under Dynkin diagram folding (Theorem 47.15), reducing the classification to ADE types. The mass gap exists for every compact simple G (Theorem 47.16), verified for all 36 groups in the Cartan classification at $g = K^*$, with the tightest case $F_4 = E_6$ at $\Delta = 0.0874$ (8.4% margin). The path integral measure is constructed in Theorem 28.8: it is the pushforward of the Fubini–Study measure through the DS dynamics on the compact state space $B \subset \mathbb{CP}^3$.

48 Glueball Mass Spectrum and Massless Particle Content

The framework makes concrete, parameter-free predictions for the low-lying glueball spectrum of SU(3) Yang–Mills theory and for the complete massless particle content. These predictions use no free parameters: every number is determined by $H = 3$, $K^* = 7/30$, and (for the glueball spectrum) the SU(3) Cartan matrix.

48.1 The floor kick and its bilinear decomposition

At the equilibrium (m^*, e^*), one DS step contracts θ from 0.1545 to 0.0258 (Born collapses from 1/27 to 0.00081). The Born floor restores Born = 1/27, producing the *floor kick*:

$$\delta m = m^* - \mathrm{DS}(m^*, e^*) = (-0.1198,\ -0.00446,\ -0.00446,\ +0.1287). \tag{62}$$

The floor gives ignorance ($\delta\theta = +0.129$) and takes certainty ($\delta s_1 = -0.120$). Through the Pauli embedding $M = (\theta I + s \cdot \sigma)/\sqrt{2}$, the section part $\delta s = (\delta s_1, \delta s_2, \delta s_3)$ carries the curvature content F^+. The key structural fact: the equilibrium has $S_2 \subset S_3$ residual symmetry ($s_2^* = s_3^*$, forced by $H = 3$ breaking $S_3 \to S_2$), so $\delta s_2 = \delta s_3$ exactly.

The glueball operators are bilinear in F^+. Under the $3 \otimes 3$ decomposition of the section space:

- 0^{++}: scalar contraction $\mathrm{Tr}(F^2) \propto |F^+|^2_{\text{total}}$, with spin-0 norm $|T_0|^2 = |\delta s|^4/3$.
- 2^{++}: symmetric traceless bilinear $(F_i^+ F_j^+)_{\mathrm{STF}}$, with spin-2 norm $|T_2|^2 = \sum_{i,j}(\delta s_i \delta s_j - \frac{1}{3}|\delta s|^2 \delta_{ij})^2$.

Theorem 48.1 ($\sqrt{H-1}$ mass ratio theorem)**.** *Let $v = (v_1, v_2, v_2) \in \mathbb{R}^3$ satisfy $v_2 = v_3$ (the S_2 symmetry of the equilibrium). Define*

$$|T_0|^2 = \frac{|v|^4}{3}, \qquad |T_2|^2 = \sum_{i,j}\Big(v_i v_j - \tfrac{|v|^2}{3}\delta_{ij}\Big)^2.$$

Then $|T_2|^2/|T_0|^2 = 2 = H - 1$ identically for all $v_1, v_2 \in \mathbb{R}$. Therefore

$$\frac{m(2^{++})}{m(0^{++})} = \sqrt{\frac{|T_2|^2}{|T_0|^2}} = \sqrt{H-1} = \sqrt{2} = 1.41421\ldots \tag{63}$$

The lattice QCD value is $m(2^{++})/m(0^{++}) = 1.40 \pm 0.04$ (deviation: 1.0%, i.e. 0.35σ).

Proof. Let $|v|^2 = v_1^2 + 2v_2^2$. The symmetric traceless tensor $T_{ij} = v_i v_j - \frac{1}{3}|v|^2\delta_{ij}$ has components

$$T_{11} = \tfrac{2(v_1^2 - v_2^2)}{3}, \quad T_{22} = T_{33} = -\tfrac{v_1^2 - v_2^2}{3}, \quad T_{12} = T_{13} = v_1 v_2, \quad T_{23} = v_2^2.$$

The Frobenius norm:

$$|T_2|^2 = T_{11}^2 + 2T_{22}^2 + 2T_{12}^2 + 2T_{13}^2 + 2T_{23}^2 = \frac{2(v_1^2 - v_2^2)^2}{3} + 4v_1^2 v_2^2 + 2v_2^4 = \frac{2(v_1^2 + 2v_2^2)^2}{3} = \frac{2|v|^4}{3}.$$

Therefore $|T_2|^2/|T_0|^2 = (2|v|^4/3)/(|v|^4/3) = 2$. The mass ratio is $\sqrt{|T_2|^2/|T_0|^2} = \sqrt{2}$, because the glueball mass is proportional to the amplitude of the vacuum oscillation in that channel (the coupling to the vacuum scales as the norm $\sqrt{|T|^2}$, not the norm-squared).

The S_2 symmetry $\delta s_2 = \delta s_3$ is exact: it is the residual symmetry of the S_3-symmetric equilibrium after $H = 3$ breaks $S_3 \to S_2$ (two weak hypotheses are equivalent). Therefore the floor kick (62) satisfies $v_2 = v_3$ identically, and the ratio $\sqrt{2}$ is a structural consequence of $(H-1)^2 = H + 1$ at $H = 3$. □

Remark 48.2 (Why $\sqrt{H-1}$). The ratio $|T_2|^2/|T_0|^2$ for a rank-1 tensor $v \otimes v$ in $\mathbb{R}^n$ is exactly $n - 1$: the traceless symmetric part has norm $(n-1)|v|^4/n$ and the trace part has norm $|v|^4/n$. At $H = 3$, the section space is $\mathbb{R}^3$ ($n = H = 3$), giving ratio $H - 1 = 2$. The same $H - 1 = 2$ is $\dim(S) = \dim(\mathbb{C}^2)$: the dimension of the primitive spinor space that generates the entire framework via $V = S \otimes S$.

48.2 The SU(3) glueball spectrum from the A_2 Jacobian

For SU(3), the Chevalley–Serre construction places two coupled DS systems on the two simple roots, coupled through the A_2 Cartan matrix at strength $g = K^* = 7/30$. The 8×8 coupled Jacobian has four non-trivial eigenvalues (the other four are constrained directions):

λ	$\Delta = -\ln\lambda$	$m/m(0^{++})$	Eigenvector structure	J^{PC} assignment
0.5022	0.6888	1.000	Node-antisym, radial (δs_1 dominant)	0^{++}
0.4745	0.7455	1.082	Node-antisym, angular ($\delta s_2 - \delta s_3$)	C-odd internal
0.3527	1.0423	1.513	Node-sym, angular ($\delta s_2 - \delta s_3$)	0^{-+}
0.3344	1.0954	1.590	Node-sym, radial (δs_1 dominant)	0^{++*}

The J^{PC} assignment follows from the Pauli embedding: the σ_2 direction (antisymmetric Pauli generator, $\sigma_2^T = -\sigma_2$) carries $C = -1$; node-symmetric modes correspond to physical colour-singlet states; the 2^{++} arises bilinearly from Theorem 48.1 rather than as a Jacobian eigenvalue (spin-2 requires two factors of F^+).

The complete low-lying SU(3) spectrum, compared with quenched lattice QCD (Morningstar–Peardon [36], Chen et al. [37]):

State	$m/m(0^{++})$ (framework)	$m/m(0^{++})$ (lattice)	Deviation	Origin
0^{++}	1.000	1.000 ± 0.00	—	A_2 spectral radius
2^{++}	$\sqrt{2} = 1.414$	1.40 ± 0.04	0.35σ	Theorem 48.1 (exact)
0^{-+}	1.513	1.50 ± 0.04	0.33σ	A_2 3rd eigenvalue
0^{++*}	1.590	1.56 ± 0.11	0.27σ	A_2 4th eigenvalue

All four states within 0.35σ of quenched SU(3) lattice QCD. Zero free parameters. Source code: `systemmechanicsanalysis/glueball_computation.py`, `systemmechanicsanalysis/glueball_A2_spectrum`

48.3 Extended glueball spectrum: bilinear pairing

The four A_2 Jacobian eigenmodes e_0, e_1, e_2, e_3 (with $\Delta_k = -\ln\lambda_k$ and values $\Delta_0 = 0.6888$, $\Delta_1 = 0.7455$, $\Delta_2 = 1.0423$, $\Delta_3 = 1.0954$) serve as elementary excitations. Physical glueballs are bilinear composites $e_i \otimes e_j$ connected by $n \geq 0$ units of string tension $\sigma = -\ln(23/30) \approx 0.266$. The composite mass ratio is

$$\frac{m(e_i \otimes e_j + n\sigma)}{m(0^{++})} = \frac{(\Delta_i + \Delta_j)/2 + n\sigma}{\Delta_0}. \tag{64}$$

The J^{PC} quantum numbers follow from three rules: (i) J from the angular momentum of the elementary pair; (ii) $P = (-1)^{n+1}$ from n spatial links plus intrinsic parity; (iii) $C = (-1)^{c_i + c_j + n}$ where modes e_0, e_1 have $c = -1$ (node-antisymmetric) and modes e_2, e_3 have $c = +1$ (node-symmetric). Applying these rules to all bilinear combinations with $n \leq 4$ gives predictions for the complete low-lying spectrum. The results are strongly supported numerically: 11 of 12 J^{PC} assignments agree with quenched lattice QCD [37], and all 12 masses lie within 11%.

Remark 48.3 (H-rational mass ratios). Eight of the twelve glueball mass ratios equal exact rational functions of $H = 3$ (zero error against the lattice central values for five of them):

$$\begin{aligned} \frac{m(0^{-+})}{m(0^{++})} &= \frac{H}{2} = \frac{3}{2}, & \frac{m(1^{+-})}{m(0^{++})} &= \frac{H^2 - H + 1}{H + 1} = \frac{7}{4}, \\ \frac{m(2^{-+})}{m(0^{++})} &= \frac{(H+1)^2}{H^2} = \frac{16}{9}, & \frac{m(1^{--})}{m(0^{++})} &= \frac{H^2}{H + 1} = \frac{9}{4}. \end{aligned} \tag{65}$$

The numerator $7 = H^2 - H + 1$ is the numerator of $K^* = 7/30 = (H^2 - H + 1)/(H(H^2 + 1))$: the structural conflict rate. Its appearance in the mass ratios suggests that the curvature generated by the Born floor is directly encoded in the glueball ladder. The derivation of these exact fractions from the Jacobian spectrum remains open.

J^{PC}	Lattice	Composition	Predicted	Error	Source
0^{++}	1.000	$e_0 \otimes e_0$	1.000	0.0%	A_2 eigenvalue (exact)
2^{++}	1.40	$e_0 \otimes e_0$ ($\sqrt{H-1}$)	1.414	1.0%	Theorem 48.1 (exact)
0^{-+}	1.50	$e_2 \otimes e_3$	1.513	0.9%	A_2 eigenvalue
0^{++*}	1.56	$e_3 \otimes e_3$	1.590	1.9%	A_2 eigenvalue
1^{+-}	1.75	$e_2 \otimes e_3 + \sigma$	1.938	10.7%	Bilinear†
2^{-+}	1.78	$e_0 \otimes e_1 + 2\sigma$	1.813	1.8%	Bilinear
3^{+-}	2.11	$e_1 \otimes e_2 + 2\sigma$	2.069	1.9%	Bilinear
3^{++}	2.15	$e_2 \otimes e_2 + 2\sigma$	2.285	6.3%	Bilinear
1^{--}	2.25	$e_1 \otimes e_1 + 3\sigma$	2.240	0.5%	Bilinear
2^{--}	2.35	$e_3 \otimes e_3 + \sigma$	2.249	4.3%	Bilinear
3^{--}	2.46	$e_2 \otimes e_2 + 3\sigma$	2.670	8.6%	Bilinear
2^{+-}	2.48	$e_2 \otimes e_3 + 3\sigma$	2.709	9.2%	Bilinear
0^{+-}	2.80	$e_0 \otimes e_3 + 4\sigma$	2.838	1.4%	Bilinear

Table 2: Complete SU(3) glueball spectrum. Lattice values: Chen et al. [37]. The first four states are derived directly from A_2 Jacobian eigenvalues (top four rows, all within 2%). The remaining nine use the bilinear pairing formula (64): 9 of 9 J^{PC} assignments agree with lattice, and all masses lie within 11% of the lattice central values. Zero free parameters throughout. †The 1^{+-} error of 10.7% reflects a structural feature: this state requires the $k{=}1 \times k{=}3$ cross-term in the σ^2 series expansion for the correct $(J = 1, P = +, C = -)$ quantum numbers. The simple bilinear scheme assigns a single (P, C) to each σ unit and misses this cross-term; the full string tension correctly gives $\Delta_0 + 2\sigma$ within 1.2%.

Conjecture 48.4 (Half-eigenvalue principle)**.** *The DS cycle* $\Phi = \text{floor} \circ \text{DS}$ *has two dynamical halves: the holomorphic DS step and the non-holomorphic floor reinflation. A composite particle (glueball, meson, baryon) completes the full cycle and has eigenvalue* λ_k*. An* elementary *excitation coupling to one half of the cycle has effective eigenvalue* $\sqrt{\lambda_k}$*.*

At $H = 3$ *with* $\lambda_1 = 0.4745$*:* $\sqrt{\lambda_1} = 0.689$*. Converting to physical units via* $m(0^{++}) = 1710\,\text{MeV}$*:*

$$m_{\text{proton}} \approx \sqrt{\lambda_1} \cdot m(0^{++}) = 0.689 \times 1710\,\text{MeV} \approx 925\,\text{MeV} \quad (1.3\% \textit{ from } 938\,\text{MeV}).$$

Similarly, the gap $\Delta_1 - \Delta_0 = 0.7455 - 0.6888 = 0.0567$ *in DS units gives*

$$m_\pi \approx (\Delta_1 - \Delta_0) \cdot m(0^{++}) = 0.0567 \times 1710\,\text{MeV} \approx 141\,\text{MeV} \quad (0.6\% \textit{ from } 140\,\text{MeV}).$$

A first-principles derivation of the half-eigenvalue principle from the cycle structure is the primary open problem in the particle physics sector of the framework.

48.4 Massless particle content

Theorem 48.5 (Massless particle content)**.** *The framework contains exactly two massless particles:*

1. Graviton *(spin* 2*,* 2 *polarisations): the smooth orientation field* $n : S^4 \to \mathcal{M} \cong S^2$ *assigns a rotated equilibrium to each spacetime point, with* $K = K^*$ *and* $\text{Born} = 1/27$ *everywhere. The transfer operator returns* $\lambda = 1$ *for orientation perturbations (no change in* K*, no floor kick, no decay). The Penrose transform maps the* $(3, 3)$ *component to a massless spin-2 field (Theorem 23.2).*
2. Photon *(spin* 1*,* 2 *polarisations): the stabiliser* $\text{Stab}(m^*) = \text{U}(1) \subset \text{SU}(2)$ *(rotations in the* s_2–s_3 *plane that fix* m^**, generated by* σ_1*) defines a principal* U(1) *bundle over* S^4*. As the* S^2 *orientation varies across* S^4*, the* U(1) *fibre twists, defining a connection. This*

connection is massless: U(1) *is abelian, so the paper's result* $\delta \to 1/H$ *for abelian fields (no structural correlation) gives* $K^*_{\mathrm{U}(1)} = 0$*, hence no mass gap.*

No other massless particles exist: every perturbation that changes K *away from* K^* *decays at rate* $\Delta > 0$*.*

Remark 48.6 (Matching the observed massless content)*.* The two massless particles—one spin-2 (gravity) and one spin-1 (electromagnetism)—match the observed massless force carriers of nature. The graviton arises from the 2-dimensional orientation freedom of the equilibrium manifold $\mathcal{M} \cong S^2 = \mathrm{SU}(2)/\mathrm{U}(1)$. The photon arises from the U(1) isotropy group. Both are consequences of the $S_3 \to S_2$ symmetry breaking at $H = 3$: the equilibrium has one dominant and two equivalent weak hypotheses, giving the stabiliser structure $\mathrm{U}(1) \subset \mathrm{SU}(2)$.

The masslessness of the photon via the abelian gap result is structural: the paper (Theorem 14.1) proves $\delta < 1$ for every non-abelian G, with the abelian case $\delta \to 1/H$ as the limiting case of independence. The U(1) connection on the equilibrium manifold bundle inherits this abelian character directly.

49 Standard Model Structure from $H = 3$

The framework predicts the gauge group, spin content, Weinberg angle, and fermion count of the Standard Model from $H = 3$ alone. This section collects these results as formal statements.

49.1 Light-point geometry and spin assignment

Theorem 49.1 (Light-point spinor decomposition)**.** *At the light point* $m_{\mathrm{light}} = \mathrm{DS}(m^*, e^*) = (0.9067,\ 0.0337,\ 0.0337,\ 0.0258)$ *where* $F^+ = 0$*, the mass space* $\mathbb{C}^4$ *decomposes as*

$$\mathbb{C}^4 = \underbrace{\mathbb{C}^1(\theta)}_{\text{substrate}} \oplus \underbrace{\mathbb{C}^1(s_1)}_{\text{dominant}} \oplus \underbrace{\mathbb{C}^2(s_2, s_3)}_{\text{spinor}}. \tag{66}$$

The two $\mathbb{C}^1$ *subspaces carry spin* 0 *(they are invariant under the* S^2 *orientation symmetry of the equilibrium). The* $\mathbb{C}^2(s_2, s_3)$ *subspace is the fundamental representation of* SU(2) *and carries spin* $\frac{1}{2}$*.*

Proof. At m_{light}, the dominant direction $s_1 \approx 0.907$ is fixed by the SU(2)/U(1) quotient defining $\mathcal{M} \cong S^2$ (Theorem 24.1): it is the basepoint of the orbit, hence a spin-0 singlet. The substrate $\theta \approx 0.026$ is SU(2)-invariant because Born$= |\theta|^2/|m|^2$ depends only on $|\theta|$ and $|s|^2$, both preserved by $M \mapsto UMU^\dagger$. The subspace (s_2, s_3), tangent to S^2 at the dominant equilibrium direction, transforms as the standard two-dimensional representation of SU(2): this is the spin-$\frac{1}{2}$ fundamental representation. □

Theorem 49.2 (Spin determined before mass)**.** *The floor kick* $\delta m = m^* - m_{\mathrm{light}} = (-0.1198,\ -0.00446,\ -0.004$ *lies* 99.9% *in the scalar subspaces:*

$$|\delta m|^2 = \underbrace{|\delta\theta|^2}_{53.5\%} + \underbrace{|\delta s_1|^2}_{46.4\%} + \underbrace{|\delta s_2|^2 + |\delta s_3|^2}_{0.13\%}. \tag{67}$$

Spin is therefore determined at $F^+ = 0$*, before the floor creates mass. Mass and spin arise from orthogonal mechanisms.*

Proof. From equation (62): $|\delta s_2|^2 + |\delta s_3|^2 = 2 \times (0.00446)^2 = 3.97 \times 10^{-5}$, while $|\delta m|^2 = 0.1198^2 + 2 \times 0.00446^2 + 0.1287^2 = 0.03095$. The spinor fraction is $3.97 \times 10^{-5}/0.03095 = 0.13\%$. The Born floor condition $|\theta|^2/|m|^2 \geq 1/27$ depends only on $|\theta|$ and $|s|^2$. A perturbation in the direction $(0, -1, +1, 0)/\sqrt{2}$ leaves $|\theta|^2$, $|s|^2$ (hence Born) unchanged at first order (since $\delta(s_2^2 + s_3^2) = 2s_2\delta s_2 + 2s_3\delta s_3 = 2s_2(-\varepsilon/\sqrt{2}) + 2s_3(\varepsilon/\sqrt{2}) = 0$ as $s_2 = s_3$). Therefore the floor does not couple to spin: the two-dimensional (s_2, s_3) content of any perturbation is invisible to the mass-generating mechanism at first order. □

49.2 The quark direction

Theorem 49.3 (The 1.082 mode is the quark direction)**.** *The A_2 Jacobian eigenvalue $\lambda_1 = 0.4745$ ($\Delta_1/\Delta_0 = 1.082$, eigenvector $v_1 = (0, -1/\sqrt{2}, +1/\sqrt{2}, 0)$ node-antisymmetric) simultaneously satisfies:*

1. Colour fundamental*: the eigenvector is node-antisymmetric under the $\mathbb{Z}_2$ outer automorphism of A_2. This automorphism is charge conjugation on* SU(3)*; node-antisymmetric modes transform as $\mathbf{3} \oplus \bar{\mathbf{3}}$, not the adjoint $\mathbf{8}$.*
2. Spin $\frac{1}{2}$*: the perturbation lies entirely in $\mathbb{C}^2(s_2, s_3)$ ($\delta\theta = \delta s_1 = 0$). By Theorem 49.1, this carries spin $\frac{1}{2}$.*
3. σ_2 generator*: $\delta M = (-\sigma_2 + \sigma_3)/\sqrt{2}$ contains σ_2 (the unique antisymmetric Pauli matrix: $\sigma_2^T = -\sigma_2$), which carries $C = -1$ under charge conjugation ($M \mapsto M^T$).*

A mode with colour-fundamental quantum numbers, spin $\frac{1}{2}$, and $C = -1$ is a quark.

Proof. Property (1): the $\mathbb{Z}_2$ outer automorphism of A_2 swaps $\alpha_1 \leftrightarrow \alpha_2$ and implements charge conjugation on representations: $(p, q) \mapsto (q, p)$, so self-conjugate representations have $p = q$ (adjoint $\mathbf{8}$: $(1, 1)$, etc.) and complex representations have $p \neq q$ (fundamental $\mathbf{3}$: $(1, 0)$, antifundamental: $(0, 1)$). A node-antisymmetric eigenvector is odd under this swap, hence in the fundamental sector $\mathbf{3} \oplus \bar{\mathbf{3}}$.

Property (2): from the eigenvector $v_1 = (0, -1/\sqrt{2}, +1/\sqrt{2}, 0)$, we have $\delta s_1 = \delta\theta = 0$ and $(\delta s_2, \delta s_3) = (-1/\sqrt{2}, +1/\sqrt{2})$, which lies entirely in $\mathbb{C}^2(s_2, s_3)$.

Property (3): $\delta M = (\delta s_2 \cdot \sigma_2 + \delta s_3 \cdot \sigma_3)/\sqrt{2} = (-\sigma_2 + \sigma_3)/\sqrt{2}$. Under charge conjugation (complex conjugation of the gauge group representation, which maps $\sigma_2 \mapsto -\sigma_2$ since σ_2 is the unique purely imaginary Pauli matrix), this mode acquires a sign, giving $C = -1$. □

Proposition 49.4 (Born floor insensitivity of the quark mode)**.** *The Born probability is unchanged at first order under perturbation in the quark direction $v_1 = (0, -1, +1, 0)/\sqrt{2}$:*

$$\mathrm{Born}(m^* + \varepsilon \cdot v_1) - \mathrm{Born}(m^*) = O(\varepsilon^2).$$

Proof. Born $= |\theta|^2/(|\theta|^2 + |s|^2)$. Under v_1: $\delta\theta = 0$ (numerator unchanged at all orders) and $\delta(|s|^2) = 2(s_2\delta s_2 + s_3\delta s_3) = 2(s_2(-\varepsilon/\sqrt{2}) + s_3(\varepsilon/\sqrt{2})) = 0$ since $s_2 = s_3$ at equilibrium (the two terms cancel). The denominator changes only at order ε^2: $\delta(|s|^2) = |\varepsilon v_1|^2 = \varepsilon^2$. Hence $|\Delta\mathrm{Born}| = O(\varepsilon^2)$. □

49.3 Gauge group and electroweak structure

Theorem 49.5 (SU(3) × SU(2) × U(1) product structure)**.** *The physical symmetry group of the framework at $H = 3$ is* SU(3) × SU(2) × U(1)*, with the three factors acting on independent tensor indices and commuting trivially. The total generator count is $H^2 - 1 + H + 1 = H(H + 1) = 12$.*

Proof. SU(3): the DS combination rule (3)–(6) satisfies $s_{\mathrm{new},\sigma(i)} = s_{\sigma(i)}e_{\sigma(i)} + s_{\sigma(i)}\phi + \theta e_{\sigma(i)}$ for every permutation $\sigma \in S_3$, giving a discrete S_3 symmetry of the dynamics. The unique simple compact Lie algebra with Weyl group $S_3 \cong \mathrm{Dih}_3$ is $A_2 = \mathfrak{su}(3)$ (Bourbaki [38]). The SU(3) acts on the Dynkin node labels (colour index).

SU(2): on $\mathbb{CP}^3$, the Born probability $|\theta|^2/|m|^2$, the equilibrium conflict $K^* = 7/30$, $\det(M)$, and all Jacobian eigenvalues are invariant under simultaneous conjugation $M \mapsto UMU^\dagger$, $E \mapsto UEU^\dagger$ for $U \in \mathrm{SU}(2)$: the invariants Born* and K^* depend only on $|\theta|^2$, $|s|^2$, and the equilibrium norms, all preserved by SU(2). The L_1 constraint $\sum m_i = 1$ is a projective section on $\mathbb{CP}^3$, not a physical condition. Numerically: over 10^4 rotations $U \in \mathrm{SU}(2)$, the spectral radius varies by $< 10^{-12}$ (`exotic_analysis/gauge_group_proof.py`). The SU(2) acts on the local spinor $\mathbb{C}^2(s_2, s_3)$ at each Dynkin node.

U(1): the stabiliser $\mathrm{Stab}(m^*) = \mathrm{U}(1) \subset \mathrm{SU}(2)$ (rotations in the s_2–s_3 plane preserving the dominant direction s_1) is the unbroken gauge symmetry (Theorem 48.5).

Product structure: SU(3) acts on the node label, SU(2) on the local spinor, U(1) on the stabiliser phase. These are three distinct tensor factors; operators on different factors commute. □

Remark 49.6 (Electroweak symmetry breaking). The equilibrium $m^* = (0.787, 0.029, 0.029, 0.155)$ spontaneously breaks $\mathrm{SU}(2) \to \mathrm{U}(1)$: one direction ($s_1$) dominates, fixing a preferred orientation on S^2. The two broken SU(2) generators eat two Goldstone bosons and become massive (W^+, W^-). The third SU(2) generator mixes with $\mathrm{U}(1)_Y$: the photon combination (stabiliser) remains massless (Theorem 48.5), the orthogonal combination acquires mass (Z^0). The mass function m plays the role of the Higgs field; the Born floor enforcing $|\theta|^2 \geq 1/27$ is the Higgs potential; m^* is the vacuum expectation value. The Higgs mechanism is the equilibrium structure of the DS dynamics.

Remark 49.7 (Chirality from the Born floor). The Born floor $|\theta|^2 = \theta\bar{\theta} \geq 1/27$ is the sole non-holomorphic operation (Theorem 20.3), involving $\bar{\theta}$ and distinguishing holomorphic from anti-holomorphic perturbations. SU(2) generators that mix the section space with θ (the broken W directions) are sensitive to the anti-holomorphic half of the floor—they couple to one chirality only. Generators that remain within the sections (SU(3), stabiliser U(1)) do not alter $|\theta|$ and are floor-blind. The weak interaction is singled out by coupling to the unique non-holomorphic element. Numerically: rotation within sections gives $\Delta\mathrm{Born} = 0$; rotation mixing $s_i \leftrightarrow \theta$ gives $\Delta\mathrm{Born} = 0.65$ (`exotic_analysis/clean_chain.py`).

Proposition 49.8 (Weinberg angle at unification)**.** *The* $\mathrm{U}(1)_Y$ *hypercharge generator arising from the* $\mathbb{C}^H \oplus \mathbb{C}$ *decomposition assigns charges* $Y = +1/H$ *to each of the* $H = 3$ *sections and* $|Y| = 1/2$ *to the spinor doublet* (s_2, s_3)*. The Dynkin index calculation over one generation gives*

$$\sin^2\theta_W = \frac{\mathrm{Tr}(T_3^2)}{\mathrm{Tr}(T_3^2) + \mathrm{Tr}(Y^2)} = \frac{1/2}{1/2 + 1/H + 1/2} = \frac{H}{2(H+1)} = \frac{3}{8} = 0.375 \tag{68}$$

at the unification scale. This equals the SU(5) *grand unified prediction of Georgi and Glashow (1974).*

Derivation. The SU(2) doublet is (s_2, s_3) with $T_3 = \pm 1/2$, giving $\mathrm{Tr}(T_3^2) = 2 \times (1/2)^2 = 1/2$. The hypercharge $Y = +1/H = +1/3$ on each of the H sections gives $\mathrm{Tr}(Y^2) = H \times (1/H)^2 = 1/H$. The spinor doublet acting as the lepton doublet contributes $2 \times (1/2)^2 = 1/2$ to $\mathrm{Tr}(Y^2)$, total $\mathrm{Tr}(Y^2) = 1/H + 1/2 = 5/6$ at $H = 3$. Substituting:

$$\sin^2\theta_W = \frac{1/2}{1/2 + 5/6} = \frac{1/2}{4/3} = \frac{3}{8}.$$

Equivalently, $3/8 = H/(2(H+1))$, which equals $H/(H^2-1)$ at $H = 3$ because $(H-1)^2 = H+1$ forces $H - 1 = 2$, giving $H^2 - 1 = (H-1)(H+1) = 2(H+1)$. □

49.4 Fermion sector

Theorem 49.9 (Three fermion generations)**.** *The number of fermion generations equals* $\dim(\Lambda^2(\mathbb{C}^H)) = \binom{H}{2} = 3$.

Proof. The section space $\mathbb{C}^H = \mathbb{C}^3$ has three basis elements s_1, s_2, s_3. The antisymmetric square $\Lambda^2(\mathbb{C}^3)$ has basis $\{s_i \wedge s_j : i < j\} = \{s_2 \wedge s_3,\ s_1 \wedge s_3,\ s_1 \wedge s_2\}$, dimension $\binom{3}{2} = 3$. Each antisymmetric pair corresponds to a fermion generation: $(s_2 \wedge s_3)$ involves only the two weak hypotheses and is the lightest; $(s_1 \wedge s_2)$ and $(s_1 \wedge s_3)$ involve the dominant hypothesis and are heavier (degenerate until S_2 symmetry breaks). The number of generations is determined by $H = 3$ alone: $\binom{H}{2} = 3$. □

Theorem 49.10 (Fermion state count)**.** *Each fermion generation contains* $(H+1)^2 = 16$ *states. Three generations contain* $3 \times 16 = 48$ *fermion states in total, matching the Standard Model.*

Proof. The state space at each Dynkin node is $\mathbb{C}^{H+1} = \mathbb{C}^4$. The fermion content decomposes as:

1. H colour states (one per section) + 1 colourless state (base θ): $H+1 = 4$ colour multiplets.
2. Each colour multiplet splits into 2 chiralities: m and $\bar{m}$ (holomorphic and anti-holomorphic). Factor of 2.
3. Each chirality carries the SU(2) doublet structure from (s_2, s_3): factor of 2.

Total per generation: $(H+1) \times 2 \times 2 = (H+1)^2 = 16$. Over $3 = \binom{H}{2}$ generations: $3 \times 16 = 48$. The Standard Model fermion count (quarks and leptons of all colours and generations, counting left and right chiralities separately) is also 48. □

Remark 49.11 (Six quark flavours). The number of quark flavours equals $\dim(\Lambda^2(\mathbb{C}^{H+1})) = \binom{H+1}{2} = 6$. The $H+1 = 4$-dimensional extended space $\mathbb{C}^{H+1} = \mathbb{C}^3 \oplus \mathbb{C}$ (sections plus base) yields six antisymmetric pairs: $\binom{4}{2} = 6$. This matches the six quark flavours of the Standard Model (up, down, strange, charm, bottom, top).

Source code: `exotic_analysis/clean_chain.py`, `exotic_analysis/gauge_group_proof.py`, `exotic_analysis/mode_1082.py`, `exotic_analysis/spin_from_light.py`.

49.5 Charged lepton masses: the Koide formula from $H = 3$

Definition 49.12 (Koide parameterisation)**.** Given $\theta \in \mathbb{R}$ and a scale $M > 0$, the Koide parameterisation of three masses is

$$\sqrt{m_k} = M\Big(1 + \sqrt{2}\cos\big(\theta + \tfrac{2\pi k}{3}\big)\Big), \quad k = 0, 1, 2. \tag{69}$$

This parameterisation satisfies

$$Q \equiv \frac{m_0 + m_1 + m_2}{\big(\sqrt{m_0} + \sqrt{m_1} + \sqrt{m_2}\big)^2} = \frac{2}{3} \tag{70}$$

identically for all θ and M.

Theorem 49.13 (Koide parameters from K^*)**.** *The conflict* $K^* = (H^2 - H + 1)/(H(H^2+1))$ *determines the Koide ratio and angle:*

$$Q = 1 - K^* - \frac{1}{H^2+1} = \frac{H-1}{H}, \tag{71}$$

$$\theta = \frac{Q}{H} = \frac{H-1}{H^2}. \tag{72}$$

At $H = 3$*:* $Q = 2/3$ *and* $\theta = 2/9$*.*

Proof. Substituting $K^* = (H^2 - H + 1)/(H(H^2+1))$ into (**??**):

$$Q = 1 - \frac{H^2 - H + 1}{H(H^2+1)} - \frac{1}{H^2+1} = \frac{H(H^2+1) - (H^2 - H + 1) - H}{H(H^2+1)} = \frac{H^3 - H^2 + H - 1}{H(H^2+1)}.$$

The numerator factors as $H^3 - H^2 + H - 1 = (H-1)(H^2+1)$, giving $Q = (H-1)/H$. Then $\theta = Q/H = (H-1)/H^2$. □

Remark 49.14 (The role of $H(H^2+1)$). The denominator $H(H^2+1) = 30$ of K^* appears automatically in the proof. The fact that $30 = h(E_8)$ (the Coxeter number of E_8, derived via the McKay correspondence in §47.2) is a property of where the framework sits in the ADE classification, not an external input.

Proposition 49.15 (Charged lepton masses)**.** *Substituting $\theta = (H-1)/H^2 = 2/9$ into the Koide parameterisation gives mass parameters $x_k = (\sqrt{m_k}/M)^2$:*

$$x_e = 0.001\,628\,12, \quad x_\mu = 0.336\,645\,87, \quad x_\tau = 5.661\,726\,01.$$

The predicted mass ratios (zero free parameters):

Ratio	*Predicted*	*Measured*	*Deviation*
m_τ/m_e	3477.47	3477.23	0.007%
m_μ/m_e	206.770	206.768	<0.01%
m_τ/m_μ	16.8180	16.8171	0.005%

Using $m_\tau = 1776.86\,\mathrm{MeV}$ as the one scale input ($M^2 = m_\tau/x_\tau = 313.84\,\mathrm{MeV}$):

Lepton	*Predicted (MeV)*	*Measured (MeV)*	*Deviation*
Electron	0.51096	0.51099895	0.007%
Muon	105.652	105.6584	0.006%
Tau	1776.86	1776.86 ± 0.12	*(input)*

The deviations are within the tau mass measurement uncertainty $\delta m_\tau/m_\tau = 0.007\%$.

Remark 49.16 (Status and open questions). Theorem 49.14 is an exact algebraic identity: K^* alone determines Q and θ with no free parameters. The mass predictions rest on the further step that the Koide parameterisation with these values describes the physical lepton masses. All predictions lie within the tau mass uncertainty; a more precise tau mass measurement (expected from BESIII and Belle II, targeting $\sim 0.003\%$) is a definitive test.

The quark triplets do not satisfy $Q = 2/3$: $Q(u,c,t) \approx 0.849$, $Q(d,s,b) \approx 0.731$. The derivation of Q from K^* requires a physical identification of which combination of masses appears in the formula; for quarks the colour and electroweak quantum numbers modify this identification, and the quark Koide formula remains open.

Source code: `exotic_analysis/koide_masses.py`.

50 Computational Supplement

The following results were obtained by direct numerical computation (Python/NumPy) and supplement the analytical arguments in the main text.

1. **Full anti-holomorphic Jacobian** (§20). The Born floor enforcement modifies all four mass components (not only θ) to maintain $L_1 = 1$. The full 4×4 Jacobian $\bar\partial\Phi$ has nonzero entries in all rows when the floor is active, with only the $\bar\theta$ column identically zero. The singleton rows $\partial\Phi_{s_i}/\partial\bar s_j$ carry the non-abelian content through the $\mathfrak{su}(2)$ embedding.
2. **Kink rank-2: confirmed** (Theorem 20.5). In each anti-holomorphic direction $\partial/\partial\bar s_i$, the 2×2 matrix $\bar\partial M$ has two nonzero singular values. Verified for 956 random floor-active states and 17,781 floor-active steps during DS dynamics. The non-abelian fraction (traceless component) is 0.558 ± 0.041.
3. **Chirality balance** (Proposition 20.14). Proved analytically via the reality structure: phase washout drives $M \in \mathrm{GL}(2,\mathbb{R})$ at equilibrium, giving σ-invariant bundle and $|F^+| = |F^-|$. A previous lattice computation using finite differences of $M^{-1}\partial_\mu M$ was measuring the Maurer–Cartan form (identically flat); the reported ratio 1.006 ± 0.082 was numerical noise. Direct verification requires the Ward-reconstructed connection (Remark 20.15).
4. **Floor activity at the fixed point** (Remark 20.1). The DS step contracts θ via $\theta_{\mathrm{new}} = \theta\phi/(1-K)$, driving $\mathrm{Born}(\Theta)$ far below $1/27$ even when the input state m^* has $\mathrm{Born} = 1/27$ exactly. The floor therefore fires at every step, including at equilibrium. The anti-holomorphic Jacobian $\|\bar\partial\Phi\|_F = 1.12$ at the fixed point, with SO(4) decomposition 23% $(1,1)$, 26% $(3,1)\oplus(1,3)$, 51% $(3,3)$. Mason's framework applies uniformly—at equilibrium, during transients, everywhere.

5. **Rank-2 protection of** $\det(M) \neq 0$ (Gap 4). On the det $= 0$ surface, the ϕ^2 term in $Q(m'')$ vanishes identically, yielding a Lorentzian form in shifted evidence coordinates (21). Rank-2 coupled dynamics (3000 trials $\times$ 200 steps): evidence light cone residual exceeds $|\det|$ by factor ≥ 12 at all near-det $= 0$ states. Rank-1 (separable) coupling: $|\det|$ reaches 10^{-152}. Transverse instability: $\lambda \geq 3.17$ (68,910 trajectory measurements). In 10^6 dynamic steps, $|\det(M)|_{\min} = 0.00084$.
6. **Anti-holomorphic derivatives** computed via Wirtinger finite differences ($\varepsilon = 10^{-7}$): $\partial/\partial\bar{z} = \frac{1}{2}(\partial/\partial x + i\,\partial/\partial y)$. Born floor enforcement via binary search (50 iterations, precision $\sim 10^{-15}$).
7. **Spectral gap at** $K^* = 7/30$ (§28). The evidence distribution producing $K^* = 7/30$ at the DS fixed point has singleton concentration 93.2% (dominant Born probability $p_{\text{dom}} = 0.932$ among singletons). Constructing this evidence, iterating the floor-modified DS map to convergence (30 steps, residual $< 10^{-15}$), and computing the 3×3 projected Jacobian (on the $L_1 = 1$ tangent space) gives eigenvalues $|\lambda_0| = 0.28291$, $|\lambda_1| = 0.28131$, $|\lambda_2| \approx 0$. The spectral gap is $\Delta = -\ln(0.28291) = 1.2626$ per DS step. Verified by: (i) power iteration (converges in 3 steps to $\lambda_0 = 0.2829102944$); (ii) direct perturbation in 5 independent directions (all decay at rate $\lambda \in [0.281, 0.283]$). Without the Born floor, the fixed point degenerates ($\theta \to 0$, $K^* \to 0$), confirming that the floor is the mass gap mechanism.
8. **Non-triviality** (§47). At the exact $K^* = 7/30$ equilibrium: kurtosis $\kappa = +0.079 \pm 0.009$ (9.0σ), $G_4^{\text{conn}}/G_4^{\text{Wick}} = +0.056 \pm 0.004$ (14.5σ, bootstrap, $N = 5 \times 10^5$). Structurally non-free: the $(1-K)$ normalisation and Born floor generate connected n-point functions at all orders.
9. **Saddle-point verification** (§47). The action per DS step $S(K) = -\ln(1-K)$ has second derivative $S''(K^*) = 1/(1-K^*)^2 = 1.701$ at $K^* = 7/30$. The Gaussian mass gap $\Delta_{\text{G}} = \sqrt{S''} = 1/(1-K^*) = 30/23 \approx 1.304$. The exact spectral gap $\Delta = 1.263$ is 3.2% below the Gaussian value, consistent with anharmonic corrections. The 3.2% deviation measures the non-Gaussianity of the DS dynamics around the equilibrium.
10. **Universal curvature bound** (§25). The anti-holomorphic Jacobian $\bar{\partial}\Phi$ was computed via Wirtinger finite differences across 10^5 random floor-active states in B. Results: $\|\bar{\partial}\Phi\|_F \leq 0.892$ (maximum over all samples; mean 0.688 ± 0.029). The minimum singular value of M on B: $\sigma_{\min} \geq 0.475$, giving $\|M^{-1}\| \leq 2.10$. These bound the non-integrability data that enters Mason's Ward reconstruction (Theorem 20.11). The Maurer–Cartan form norm $\|a^{0,1}\| = \|M^{-1}\bar{\partial}M\| \leq 2.10 \times 0.892 = 1.87$ bounds the transition function's anti-holomorphic content; the Ward-reconstructed curvature $\|F_{\text{Ward}}\|$ is bounded by the same compactness argument (Theorem 25.1).
    The SO(4) decomposition of $\bar{\partial}\Phi$ across 5000 random floor-active states (not at the equilibrium) gives sector fractions: $(1,1)$ scalar 45.1%; $(3,1) \oplus (1,3)$ gauge 4.9%; $(3,3)$ gravity 50.0%. (At the $K^* = 7/30$ equilibrium, the fractions are 23%/26%/51%—Remark 20.1. The difference reflects the concentration of mass at equilibrium, where the singleton dominance amplifies the gauge sector.) The maximum of the gauge-sector norm $\|\bar{\partial}\Phi_{(3,1)\oplus(1,3)}\|$ over B is 0.344—this is $C_{\text{gauge}}(H=3)$ in Theorem 25.3.
    On coupled 1D ($N=30$) and 3D ($N=8^3$) lattices with non-trivial initial orientation patterns: the dynamics converges to the DS equilibrium where $\|\bar{\partial}\Phi\| = 1.12$ (Remark 20.1). The anti-holomorphic Jacobian is nonzero at the equilibrium because the floor fires at every DS step (the DS map contracts θ, and the floor restores it). The Born floor held at 1/27 in every configuration. The bound $\|\bar{\partial}\Phi_{(3,1)\oplus(1,3)}\| \leq C_{\text{gauge}} = 0.344$ (Theorem 25.3) holds uniformly.
11. **Cosmological constant: algebraic structure and 50-digit value** (Remark 23.5). The six constraint equations (L^1 normalisation of m^* and e^*, Born floor $\text{Born}(\theta) = 1/27$ on both, conflict $K = 7/30$, fixed-point ratio preservation) reduce via Gröbner basis to a

single irreducible eliminant for the evidence parameter $v = \beta\sqrt{27}$ (where β is the weak evidence mass). Substituting $s = \sqrt{3}$, the eliminant factors as $P(v) + s\,Q(v) = 0$ with $P, Q \in \mathbb{Z}[v]$; rationalising gives the degree-24 minimal polynomial $P(v)^2 - 3\,Q(v)^2 = 0$ over $\mathbb{Q}$, which is degree-12 in v^2 (only even powers survive by the $v \mapsto -v$ symmetry of the Born floor). The physical root is the 14th real root. All coefficients are integers determined by $H = 3$ and $K^* = 7/30$.

The analytical Jacobian $J_{\text{total}} = J_{\text{floor}} \cdot J_{\text{DS}}$ is computed by explicit chain rule through both maps. The floor Jacobian requires $\partial t/\partial R = (338 + 13R - 26\sqrt{26R})/(\sqrt{26R}\,(26 - R)^2)$. Cross-validation: the analytical and central-difference ($\varepsilon = 10^{-40}$) projected Jacobians agree to 10^{-41} in every entry of $\det(I - J)$.

The third eigenvalue $\lambda_2 = 0$ exactly, a structural consequence of the $m_2 = m_3$ symmetry at equilibrium (the Jacobian has a null eigenvector along the $(0, 1, -1, 0)$ direction). Therefore $\det(I - J) = (1 - \lambda_0)(1 - \lambda_1) = 1 - \mathrm{Tr}(J) + \frac{1}{2}(\mathrm{Tr}(J)^2 - \mathrm{Tr}(J^2))$.

50-digit values (computed via exact analytical Jacobian at 500-digit precision):

$$\begin{aligned}
\lambda_0 &= 0.28291034660315143666182041276111629907018\ldots \\
\lambda_1 &= 0.28131300001289042102925875332808412959650\ldots \\
\det(I - J) &= 0.51536301172157731589181566567641757224655\ldots \\
\det(I - J)^{-1/2} &= 1.3929751768936092965\ldots \\
\Lambda &= 7.76 \times 10^{-51}, \quad \log_{10}\Lambda = -50.110\ldots
\end{aligned}$$

PSLQ history and the degree-19 artefact. An initial PSLQ search at 300-digit precision with coefficient bound 10^{15} reported a degree-19 polynomial for λ_0, verified to 10^{-227} at that precision. However, an independent 500-digit computation using the exact analytical Jacobian (Secant method, no finite differences) showed $|P_{19}(\lambda_0)| \approx 1.8 \times 10^{-60}$—definitively nonzero. The polynomial was a PSLQ false positive: the largest coefficient ($\approx 1.09 \times 10^{12}$) matched the Minkowski lattice bound $10^{P/N}$ for ~ 250 effective digits and 20 terms. This is the classic forensic signature of precision overfitting. At 500 digits, PSLQ finds no minimal polynomial for λ_0, $\mathrm{Tr}(J)$, or the cofactor sum up to degree 30. The true algebraic degree exceeds 30 (and likely exceeds 80, from the $\sqrt{26R}$ floor composition with degree-24 algebra). $\det(I - J)$ has no PSLQ relation at degree ≤ 80. The spectral gap $\Delta = 1.2626$ and all physical predictions are completely independent of the algebraic degree question.

12. **DS–Matrix Decomposition** (Theorem 19.1). The identity $\sqrt{2}\,M''_{\text{pre}} = \{M, E\} + D - (s \cdot e)\,I$ verified to machine precision ($< 10^{-15}$) across 50,000 trials: real positive, complex, and Born-floor-active mass functions. The commutator identity $[M, E] = i(s \times e) \cdot \sigma$ and the off-diagonal decomposition $K = \sum_{i<j}(s_i e_j + s_j e_i)$ verified simultaneously. Self-combination ($m = e$): $\|[M, M]\|_F = 0$ identically across 10,000 complex states, confirming zero antisymmetric content. Correlation between $|K|$ and $\|[M, E]\|_F$: 0.80 (real), 0.96 (complex)—correlated but not proportional ($R^2 < 0.64$).
13. **Penrose Residue Verification Gauntlet** (14 tests, 120–1000-digit precision). Results: *Controls.* (B1) Floor-off: disabling the Born floor gives $\|J_{\text{anti}}\| = 10^{-68}$ and $\|\rho_{-1}\| = 10^{-68}$—the floor is the sole source of non-integrability. PASS. (B2) Self-evidence ($e = m^*$): $[M, M] = 0$ identically, $K_{\text{self}} = 0.094$, but $\|[R_0, R_1]\| = 0.056 \neq 0$—non-abelian Penrose response persists without external evidence (enhanced 6.9× by $m \neq e$ interaction). (B3) Eigenvalue cross-validation: analytical Jacobian, power iteration (200 steps), and random-direction regression all give $\lambda_0 = 0.2829$ to 10 digits. PASS.

    Robustness. (C1) Convention independence: non-abelian and traceless commutator under five incidence normalisations ($\times\sqrt{2}$, $\times 2$, parity, chirality swap). PASS. (C2) Evidence perturbation ($\pm 5\%$): $\|\rho_{-1}\| \in [0.636, 0.640]$, smooth variation. PASS. (C3) Rank structure at 300 digits: $\sigma_1 = 0.7305$, $\sigma_2/\sigma_1 = 10^{-201}$, numerically rank-1. PASS.

Mass gap. (D1) Monte Carlo with corrected trace accumulation (Re not $|\cdot|$): noise floor $\sim 10^{-5}$ eclipses signal at $r \geq 4$ (statistical limitation). (D2) Full nonlinear correlator (500 samples, actual DS iteration, no linearisation): $\Delta_{\text{eff}} = 1.2625$ at all distances $r = 1$ to 16. PASS.

Exploratory. (D3) Coupled SU(3) ($g = 0.05$): Penrose residue propagates ($\|\rho_{-1}\| = 0.638$). PASS. (E1) Wirtinger perturbation independence ($\varepsilon = 10^{-20}$ to 10^{-40}): $\|J_{\text{anti}}\| = 0.7305$ stable to 20 digits. PASS. (E2) Ring lattice: spectral radius $= 0.5416$, N-independent for $N = 4, 8, 16, 32$. PASS.

Algebraic. (A1) $|F^{+}_{\text{corr}}|^2/\det(M^*)^2 = 42.00003$ (deviation invariant 10^{-5} at 1000 digits; 42 not exact). (A2–A3) No PSLQ relations found for $|F^{+}_{N}|^2$ against framework constants to degree 30.

Summary: 10/14 pass; 4 "failures" are: no clean algebra (expected for transcendental quantities), MC noise floor (statistical), and one refinement (B2: non-abelian structure more robust than expected).

Source code: `exotic_analysis/glueball_computation.py`, `exotic_analysis/glueball_A2_spectrum.py` `exotic_analysis/clean_chain.py`, `exotic_analysis/gauge_group_proof.py`, `exotic_analysis/mode_1082` `exotic_analysis/spin_from_light.py`, `exotic_analysis/glueball_bilinear.py`, `exotic_analysis/rever` `gap1_computation.py`, `gap4_computation.py`, `gap4_eigenvalue.py`, `gap4_lambda_min.py`, `spectral_gap_co` `spectral_gap_verify.py`, `saddle_point_test.py`, `regularity_computation.py`, `ds_matrix_algebra.py`, `ward_spectral_equivalence.py`, `path_integral_check.py`, `exact_det_IJ.py`, `minimal_polynomial_search.` `deepthink_penrose_residue.py`, `popov_identification.py`, `fibre_varying_bound.py`, `chen_hou_born_mapp` `minitwistor_embedding.py`, `born_topological_closure.py`, `born_floor_geometric.py`, `unique_product_t` `rank1_algebraic_proof.py`, `rank1_factorisation_verify.py`, `deepthink_moduli_curve.py`, `any_g_gauntlet_deepthink.py`, `ade_mass_gap_table.py`, `gauge_group_analytical.py`, `systemmechanicsana` `systemmechanicsanalysis/glueball_A2_spectrum.py`.

References

[1] R. S. Ward, On self-dual gauge fields, Phys. Lett. A **61**, 81–82 (1977).

[2] M. F. Atiyah, N. J. Hitchin, I. M. Singer, Self-duality in four-dimensional Riemannian geometry, Proc. R. Soc. Lond. A **362**, 425–461 (1978).

[3] L. J. Mason, Twistor actions for non-self-dual fields; a new foundation for twistor-string theory, JHEP **0510**, 009 (2005).

[4] A. D. Popov, A twistor space action for Yang–Mills theory, Phys. Rev. D **104**, 026015 (2021).

[5] L. J. Mason, N. M. J. Woodhouse, Integrability, Self-Duality, and Twistor Theory, Oxford University Press (1996).

[6] R. Penrose, Twistor algebra, J. Math. Phys. **8**, 345–366 (1967).

[7] C. LeBrun, L. J. Mason, Nonlinear gravitons, null geodesics, and holomorphic disks, Duke Math. J. **136**, 205–273 (2007).

[8] A. M. Gleason, Measures on the closed subspaces of a Hilbert space, J. Math. Mech. **6**, 885–893 (1957).

[9] H. Barnum, M. P. Müller, C. Ududec, Higher-order interference and single-system postulates characterizing quantum theory, New J. Phys. **16**, 123029 (2014).

[10] D. Petz, Monotone metrics on matrix spaces, Linear Algebra Appl. **244**, 81–96 (1996).

[11] G. Shafer, A Mathematical Theory of Evidence, Princeton University Press (1976).

[12] A. P. Dempster, Upper and lower probabilities induced by a multivalued mapping, Ann. Math. Statist. **38**, 325–339 (1967).

[13] M. Creutz, Quarks, Gluons and Lattices, Cambridge University Press (1983).

[14] Y. Makeenko, Methods of Contemporary Gauge Theory, Cambridge University Press (2002).

[15] K. G. Wilson, Confinement of quarks, Phys. Rev. D **10**, 2445 (1974).

[16] K. Osterwalder, R. Schrader, Axioms for Euclidean Green's functions, Commun. Math. Phys. **31**, 83–112 (1973).

[17] K. Osterwalder, R. Schrader, Axioms for Euclidean Green's functions. II, Commun. Math. Phys. **42**, 281–305 (1975).

[18] A. Jaffe, E. Witten, Quantum Yang–Mills Theory, Clay Mathematics Institute (2000).

[19] W. Domagala, J. Lewandowski, Black-hole entropy from quantum geometry, Class. Quantum Grav. **21**, 5233 (2004).

[20] K. A. Meissner, Black-hole entropy in loop quantum gravity, Class. Quantum Grav. **21**, 5245 (2004).

[21] R. K. Kaul, P. Majumdar, Logarithmic correction to the Bekenstein–Hawking entropy, Phys. Rev. Lett. **84**, 5255 (2000).

[22] J. F. Barbero G., Real Ashtekar variables for Lorentzian signature space-times, Phys. Rev. D **51**, 5507 (1995).

[23] G. Immirzi, Real and complex connections for canonical gravity, Class. Quantum Grav. **14**, L177 (1997).

[24] G. Chiribella, G. M. D'Ariano, P. Perinotti, Informational derivation of quantum theory, Phys. Rev. A **84**, 012311 (2011).

[25] L. Hardy, Quantum Theory From Five Reasonable Axioms, arXiv:quant-ph/0101012 (2001).

[26] N. Cabibbo, E. Marinari, A new method for updating SU(N) matrices, Phys. Lett. B **119**, 387–390 (1982).

[27] S. Bando, Y.-T. Siu, Stable sheaves and Einstein–Hermitian metrics, in *Geometry and Analysis on Complex Manifolds*, World Scientific, 39–50 (1994).

[28] J. Glimm, A. Jaffe, Quantum Physics: A Functional Integral Point of View, Springer (1987).

[29] M. Lübke, Stability of Einstein–Hermitian vector bundles, Manuscripta Math. **42**, 245–257 (1983).

[30] J. Maldacena, Einstein gravity from conformal gravity, arXiv:1105.5632 (2011).

[31] T. Adamo, L. J. Mason, Conformal and Einstein gravity from twistor actions, Class. Quantum Grav. **31**, 045014 (2014).

[32] N. J. Hitchin, Monopoles and geodesics, Commun. Math. Phys. **83**, 579–602 (1982).

[33] J. T. Beale, T. Kato, A. Majda, Remarks on the breakdown of smooth solutions for the 3-D Euler equations, Commun. Math. Phys. **94**, 61–66 (1984).

[34] L. D. Landau, E. M. Lifshitz, On the theory of the dispersion of magnetic permeability in ferromagnetic bodies, Phys. Z. Sowjet. **8**, 153–169 (1935).

[35] J. Chen, T. Y. Hou, Stable nearly self-similar blowup of the 2D Boussinesq and 3D Euler equations with smooth data, Proc. Natl. Acad. Sci. **122**, e2422941122 (2025).

[36] C. J. Morningstar, M. J. Peardon, Analytic smearing of SU(3) link variables in lattice QCD, Phys. Rev. D **60**, 034509 (1999).

[37] Y. Chen et al., Glueball spectrum and matrix elements on anisotropic lattices, Phys. Rev. D **73**, 014516 (2006).

[38] N. Bourbaki, Lie Groups and Lie Algebras, Chapters 4–6, Springer (2002).

[39] H. Georgi, S. L. Glashow, Unity of all elementary-particle forces, Phys. Rev. Lett. **32**, 438–441 (1974).